

Health & Research Journal

Vol 6, No 4 (2020)

Volume 6 Issue 4 October - December 2020



Volume 6 Issue 4 October - December 2020

EDITORIAL

THE EFFECT OF EMTRICITABINE AND TENOFOVIR USE IN THE PRE-EXPOSURE PROFYLAXIS (PREP) FOR HIV TRANSMISSION

BRIEF REPORT

DEPRESSION IN DIABETIC FOOT ULCERS

SPECIAL ARTICLES

DEVELOPING SUPPORT NETWORKS FOR SUCCESSFUL IMPLEMENTATION OF INNOVATIVE EDUCATIONAL PROGRAMMES: THE CASE OF "EVIDENCE-BASED PRACTICE IN NURSING" CONTINUING EDUCATION PROGRAMME IN GREECE

RESEARCH ARTICLES

EXPLORATION THE QUALITY OF LIFE AND THE RELATED FACTORS IN WOMEN WITH RECENT DIAGNOSIS OF GYNECOLOGICAL CANCER, BEFORE THE SURGICAL TREATMENT, IN GREECE

ADAPTATION OF "NICHOLSON MCBRIDE RESILIENCE QUESTIONNAIRE" (NMRQ) IN GREEK. A RELIABILITY AND VALIDITY STUDY IN AN EPIDEMIOLOGICAL GREEK SAMPLE

Published in cooperation with the Postgraduate Program "Intensive Care Units", the Hellenic Society of Nursing Research and Education and the Helerga

Depression in diabetic foot ulcers

Stavros Tsirigotis

doi: [10.12681/healthresj.25626](https://doi.org/10.12681/healthresj.25626)

To cite this article:

Tsirigotis, S. (2020). Depression in diabetic foot ulcers. *Health & Research Journal*, 6(4), 105–106. <https://doi.org/10.12681/healthresj.25626>

BRIEF REPORT

DEPRESSION IN DIABETIC FOOT ULCERS

Stavros Tsirigotis

Msc in Applied Clinical Nursing, University of West Attica, Department of Nursing, Athens, Greece

Cite as: Tsirigotis, S. (2020). Depression in diabetic foot ulcers. *Health and Research Journal*, 6(4), 105-106.
<https://ejournals.epublishing.ekt.gr/index.php/HealthRes/>

Corresponding Author: Tsirigotis Stavros, e-mail: stavtsiri8@hotmail.gr

Diabetic foot ulcer (DFU) is a common complication of diabetes mellitus. According to estimates, 15% of diabetic patients may experience a DFU through their life.¹ The majority (60–80%) of DFUs will heal, while 10–15% of them will remain active, and 5–24% of them will finally lead to limb amputation within a period of 6–18 months after the first evaluation. The risk to develop a DFU increases with age and the duration of diabetes mellitus.²

A DFU entails a heavy physical and emotional burden for each individual since it involves impaired mobility and dependency on others, feelings of fear about the clinical outcome, changes in roles, powerlessness, need to follow complex medical procedures, failure to engage in social activities or in social interactions, change in family roles and poor quality of life.^{1,3,4} Therefore, depression may be prevalent among this vulnerable population.

Interestingly, DFU patients are more likely to experience depression than diabetic patients without foot complications.^{3,5,6} Furthermore, depression among DFUs patients is associated with hospital admissions³ and with a two-fold increase of mortality over 5 years after their first foot ulcer.⁴

Depression and DFUs seem to share a bi-directional relationship. More, in detail, on the onset of diabetes mellitus treatment, depression is associated with an increased risk of developing DFU.⁷ High levels of depression are associated with recurrence of DFU,³ with delay in DFU healing⁸ and with increased risk of amputation.³ Additionally more, psychological factors seem to have direct effects on endocrine and immune function, thus affecting healing.⁹ Moreover, depression consists an impede for regular monitoring and follow-up, thus posing a high risk for treatment disruption or failure to adhere to therapy.

Tsirigotis S.

Several socio-demographic and clinical factors may trigger depression among DFU patients. A cross-sectional study among 195 DFUs outpatients in Greece showed high levels of depression in 20% of participants¹ while Udovichenko et al.,¹⁰ illustrated depression in 39% of 285 DFU outpatients, with women to be more likely to be depressed and anxious than men (48% and 46% versus 27% and 25%, respectively). Ahmad et al.,² showed depression in 39.6% of DFUs patients which was more prevalent in those aged <50 years, in female, in those suffering more than three co morbid diseases and those who were current smokers.

Another relevant study in Greece showed moderate levels of depression among 180 DFUs patients while the depressed participants were more likely to be older, single/divorced, of primary education, pensioners, to have some other disease and to be current smokers.¹¹ It is noteworthy that patients' attitudes, perceptions and beliefs may trigger depression. Delays to seek for health care due to beliefs may have detrimental effects on DFU.¹²

Nurses need to early recognize depression, which is a major risk factor for diabetes related complications. More in detail, symptoms of major depression include lack of interest in activities, increased or decreased appetite, insomnia, lack of energy, feelings of guilt and suicidal thoughts or behaviors. These symptoms frequently overlap with anxiety while concentration problems and fatigue are symptoms of both anxiety and depression. Irritability can also manifest in forms of anxiety or depression (instead of low mood).^{3,13,14}

Nurses need to expand their knowledge and understanding of depression with ultimate goal to facilitate better diabetes self-management. Also, need to include early screening of depression in their caring programs.^{3,13,14}

On the other end of the spectrum, patients have to combat with depression and adjust to the demands of DFU in their daily lives. Developing psychological interventions will allow patients to manage their illness and modify their health behavior as well as to follow appropriate emotional and behavioral changes in order to gain a sense of control over their lives. Clinically important is to understand which type of psychological intervention could be most successful when developing intervention tailored to DFU needs.¹⁴

REFERENCES

1. Polikandrioti M, Vasilopoulos G, Koutelekos I, Panoutsopoulos G, Gerogianni G, Babatsikou F, et al. Quality of Life in Diabetic Foot Ulcer: Associated Factors and the Impact of Anxiety/Depression and Adherence to Self-Care. *Int J Low Extrem Wounds*. 2020;19(2):165-179. doi: 10.1177/1534734619900415
2. Alexiadou K, Doupis J. Management of diabetic foot ulcers. *Diabetes Ther*. 2012;3(1):4. doi:10.1007/s13300-012-0004-9
3. Ahmad A, Abujbara M, Jaddou H, Younes NA, Ajlouni K. Anxiety and Depression Among Adult Patients With Diabetic Foot: Prevalence and Associated Factors. *J Clin Med Res*. 2018;10(5):411-418. doi:10.14740/jocmr3352w
4. Ismail K, Winkley K, Stahl D, Chalder T, Edmonds M. A cohort study of people with diabetes and their first foot ulcer: the role of depression on mortality. *Diabetes Care*. 2007;30(6):1473-1479. doi: 10.2337/dc06-231
5. Prinz N, Ebner S, Grunerbel A, Henkeludecke U, Hermanns N, Hummel M, et al. Female sex, young age, northern German residence, hypoglycemia and disabling diabetes complications are associated with depressed mood in the WHO-5 questionnaire - A multicenter DPV study among 17.563 adult patients with type 2 diabetes. *J Affect Disord*. 2017;208:384-391. doi: 10.1016/j.jad.2016.08.077.
6. Chapman Z, Shuttleworth CM, Huber JW. High levels of anxiety and depression in diabetic patients with Charcot foot. *J Foot Ankle Res*. 2014;7:22. doi: 10.1186/1757-1146-7-22.
7. Iversen MM, Tell GS, Espehaug B, Midthjell K, Graue M, Rokne B, et al. Is depression a risk factor for diabetic foot ulcers? 11-years follow-up of the Nord-Trøndelag Health Study (HUNT). *J Diabetes Complications*. 2015;29(1):20-25.
8. Monami M, Longo R, Desideri CM, Masotti G, Marchionni N, Mannucci E. The diabetic person beyond a foot ulcer: healing, recurrence, and depressive symptoms. *J Am Podiatr Med Assoc*. 2008;98(2):130-136.
9. Situm M, Kolić M, Spoljar S. Quality of life and psychological aspects in patients with chronic leg ulcer. *Acta Med Croatica*. 2016;70(1):61-63.
10. Udovichenko OV, Maximova NV, Amosova MV, Yunilaynen OA, Berseneva EA, Starostina EG. Prevalence and prognostic value of depression and anxiety in patients with diabetic foot ulcers and possibilities of their treatment. *Curr Diabetes Rev*. 2017;13(1):97-106.
11. Polikandrioti M, Vasilopoulos G, Koutelekos I, Panoutsopoulos G, Gerogianni G, Alikari V, et al. Depression in diabetic foot ulcer: Associated factors and the impact of perceived social support and anxiety on depression. *Int Wound J*. 2020 Mar 26. doi: 10.1111/iwj.13348
12. Hjelm K, Beebwa E. The influence of beliefs about health and illness on foot care in ugandan persons with diabetic foot ulcers. *Open Nurs J*. 2013;7:123-132.
13. Douki S, Marvaki C, Toulia G, Stavropoulou A. Attitudes and perceptions of health professionals towards mental patients attending emergency room. *Health & Research Journal*. 2019;2(4):261-276. doi:https://doi.org/10.12681/healthresj.19846
14. McGloin H, Devane D, McIntosh CD, Winkley K, Gethin G. Psychological interventions for treating and preventing recurrence of foot ulcers in people with diabetes. *Cochrane Database Syst Rev*. 2017;2017(10):CD012835 doi:10.1002/14651858.CD012835