

Health & Research Journal

Vol 11, No 3 (2025)

Volume 11 Issue 3 July - September 2025



Volume 11 Issue 3 July – September 2025

EDITORIAL

«CO-EXISTENCE» IN A HEALTHCARE ENVIRONMENT

RESEARCH ARTICLES

PARENTAL EXPERIENCE OF CHILDREN WITH SICKLE CELL DISEASE: A QUALITATIVE STUDY

KNOWLEDGE, VIEWS AND ATTITUDES OF HEALTHCARE PROFESSIONALS TOWARDS THE VARIOUS FORMS OF DOMESTIC VIOLENCE

DETERMINANTS OF SCHOOL PERFORMANCE IN A SAMPLE OF ADOLESCENTS IN GREECE

PERIPHERAL MICROCIRCULATION ADAPTATIONS IN RESPONSE TO THE ADDITION OF INSPIRATORY MUSCLE TRAINING IN HEART FAILURE CARDIAC REHABILITATION REGIMEN

FAMILIAL HYPERCHOLESTEROLAEMIA IN GREEK FEMALES. AN EPIDEMIOLOGICAL STUDY

REVIEWS

KNOWLEDGES, BELIEFS AND PRACTICES ON RADIATION PROTECTION OF NON-RADIOLOGISTS PHYSICIANS WHO USE IONIZING RADIATION AND PARTICIPATE IN RADIOSCOPICALLY GUIDED PROCEDURES

APPLICATION OF THE HIGH-FLOW NASAL CANNULA IN PATIENTS WITH ACUTE RESPIRATORY DISTRESS SYNDROME

THE VALUE OF MUSCULOSKELETAL ULTRASOUND IMAGING IN PHYSIOTHERAPY CLINICAL ASSESSMENT AND PRACTICE

SPECIAL ARTICLES

VALIDATION OF THE GREEK VERSION OF EUTHANASIA ATTITUDE SCALE IN THE GENERAL POPULATION: A QUANTITATIVE STUDY

CARE AND SUPPORT OF PATIENTS WITH END-STAGE RESPIRATORY DISEASE ON HOME MECHANICAL VENTILATION – ETHICAL AND LEGAL ISSUES



Knowledges, beliefs and practices on radiation protection of non-radiologists physicians who use ionizing radiation and participate in radioscopically guided procedures

Markos Kapsampelis, George Intas, Eleftherios Lavdas, Pantelis Stergiannis, Athanasios Bakas

doi: [10.12681/healthresj.40148](https://doi.org/10.12681/healthresj.40148)

To cite this article:

Kapsampelis, M., Intas, G., Lavdas, E., Stergiannis, P., & Bakas, A. (2025). Knowledges, beliefs and practices on radiation protection of non-radiologists physicians who use ionizing radiation and participate in radioscopically guided procedures. *Health & Research Journal*, 11(3), 259–266. <https://doi.org/10.12681/healthresj.40148>

REVIEW ARTICLE

KNOWLEDGES, BELIEFS AND PRACTICES ON RADIATION PROTECTION OF NON-RADIOLOGISTS PHYSICIANS WHO USE IONIZING RADIATION AND PARTICIPATE IN RADIOSCOPICALLY GUIDED PROCEDURES

Markos Kapsampelis¹, George Intas², Eleftherios Lavdas³, Pantelis Stergiannis⁴, Athanasios Bakas⁵

1. Radiographer Diagnostic Imagine, MSc, PhD(c), Department of Biomedical Sciences, University of West Attica
2. RN, MSc, MHSc, PhD, General hospital of Nikaia Agios Panteleimon
3. Professor of Radiology and Radiotherapy sector, Department of Biomedical Sciences, University of West Attica
4. Assistant Professor, Department of Nursing, National and Kapodistrian University of Athens
5. Associate Professor of Radiology and Radiotherapy sector, Department of Biomedical Sciences, University of West Attica

Abstract

Background: Technical innovations have spread the use of ionizing radiation in daily clinical practice. About one third to half medical decisions are guided by radiological examinations. This review aims to evaluate current literature relating non-radiologists physicians' knowledge and practices of radiation safety.

Method and Material: A literature search from the MEDLINE and CINAHL databases was performed where original articles were retrieved, published in the English language from the past 10 years.

Results: While almost all physicians use lead aprons not all of them use thyroid protectors and very few of them use lead gowns and goggles. Also the majority of physicians do not use a dosimeter badge. The major reason for not using radiation protection equipment is their unavailability. Physicians have low levels of knowledge about radiological safety. Longer length of service, having attended a radiation protection and safety course, increased frequency of ordering imaging per day and specialty are factors that affect positively physicians' level of knowledge. Although occupational radiation exposure has been associated with leukemia and nonmalignant thyroid nodular disease, increased risk of cancer, cataract, headache and fatigue, eye symptoms with most common the red eyes, very few physicians have read articles on radiation safety or have received training.

Conclusions: The inadequate level of radiation knowledge can lead to increased radiation exposure and risk for complications to healthcare professionals and patients. Healthcare organizations must ensure the adequate procurement of personal protective equipment so it is available to physicians who use ionizing radiation.

Keywords: Knowledges, beliefs, practices, radiation protection, non-radiologists physicians, ionizing radiation.

Corresponding Author: Markos Kapsampelis, Email: mkapsampelis@uniwa.gr

Cite as: Katsampelis, M., Intas, G., Lavdas, E., Stergiannis, P., Bakas, A. Knowledges, beliefs and practices on radiation protection of non-radiologists physicians who use ionizing radiation and participate in radioscopically guided procedures. (2025). Health and Research Journal, 11(3),259-266. <https://ejournals.epublishing.ekt.gr/index.php/HealthRes/>

INTRODUCTION

The first usage of ionizing radiation in healthcare setting was more than 100 years ago. The ionizing radiation is used by physicians for different purposes. Technical innovations have spread the use of ionizing radiation in daily clinical practice. About one third to half medical decisions are guided by radiological examinations.¹ For example, neurosurgeons use radiation to visualize blood vessels and implants intraoperatively.²

Physicians routinely are exposed to ionizing radiation during their daily practice, which has been associated with a variety of complications. Despite this, the utilization of ionizing radiation in healthcare settings increases continuously. In this line, healthcare professionals should be compliant with protection guidelines when ordering imaging or when they use ionizing radiation. There are three principles for the protection of healthcare professionals and patients from ionizing radiation: dose limits, optimization and justification.³ It is well known that personal protective equipment reduces occupational radiation exposure.⁴

Ionizing radiation exposure leads to impairment of cellular function or cell death, which can contribute to the onset of cancer. The effects of ionizing radiation are dependent on the dose of radiation applied, the length of time spent in the radiation field, and the amount of protection employed.⁵ Because the correct use of ionizing radiation is very important for the safety of both healthcare professionals and patients, the efforts of reducing their exposure to ionizing radiation has led to the development of the As Little As Reasonably Achievable (ALARA) principle. The International Radiation Commission strongly recommend healthcare professionals not exceed 20 mSv of exposure per year. Due to the fact that many physicians who are not specialized in radiology use ionizing radiation in their daily clinical practice, they have to know the radiation dosage of these exams and all the equipment and techniques they have to use to protect themselves, their colleagues and the patient.⁶ Physicians do not know the exact dose of radiation that patient are expose to. In particular, patients receive 16 times greater dose of radiation than physicians believe.⁷ Inadequate knowledge for radiation may limit the physicians` ability to protect themselves and pa-

tients. So, according to the International Commission on Radiological Protection, healthcare professionals must have all the necessary knowledge and be aware about the danger of radiation exposure.^{3,8}

This review aims to evaluate current literature relating non-radiologists physicians' knowledge and practices of radiation safety.

METHODOLOGY

The search of literature was performed using MEDLINE and CINAHL. Keyword search with MeSH, and citation analysis were conducted. Inclusion criteria were applied for population (non-radiologists physicians who use ionizing radiation and participate in radioscopically guided procedures), exposure (radiation safety), and outcome (practices and knowledge of participants). The following key words were used: "Orthopedic Surgeon" OR "Orthopedic" OR "Cardiologist" OR "Urologist" OR "Neurosurgeon" OR "Physician" AND "knowledge" OR "awareness" OR "practices" AND "radiation safety" OR "radiation risks" OR "radiation exposure" OR "radiation". The final articles selected were original ones from peer reviewed journal, which were published the last 10 years (2015-2024) and written in the English language. Exclusion criteria included qualitative research, which were older than 10 years old, and the study population being radiologists. The process of literature review is shown in the following flowchart.

RESULTS

Totally, the review included 22 articles. Out of all of them, 15 were for practices of occupational radiation exposure, 12 for knowledges of occupational radiation exposure and 11 for implications of occupational radiation exposure.

Practices of occupational radiation exposure

The exposure to ionizing radiation differs among physicians. According to a US study, most neurosurgeons performing surgeries weekly using radiation, do not use a dosimeter but do wear a lead apron and a thyroid shield.²

Urologists, especially endourologists, is a specialty of physicians with high frequency of radiation exposure with more than 15 times per week. Usually, they use lead aprons and thyroid shields

but fewer of them use lead eyeglasses and gloves.⁹ In Turkey, all urologists use lead aprons but half of them use thyroid protectors. However, very few of them use a combination of lead gowns, goggles, and thyroid protectors together. At the same time the majority of them do not use a dosimeter, since they believe they are not at risk due to the short period of exposure.¹⁰ According to the ESUT/EULIS survey, the majority of urologists use lead apron and thyroid shield during their practice, about half of them use lead shield and very few use glasses and gloves.¹¹

Another specialty of physicians that use ionizing radiation very frequently is orthopedic. They use ionizing radiation either for diagnostic practices or for operative management or for follow-up of patients. All orthopedics in the UK and Scotland use lead aprons/gowns and about half of them use thyroid protection when using x-rays.¹² A study showed that the protective equipment an orthopedic surgeon use is lead apron and thyroid shield for every case but only few of them wear lead glasses.⁴ All orthopedics and general surgeons in the USA wear thyroid shields when using ionizing radiation. Most of them, also, report that they want to wear lead but they can't due to lack of availability.¹³ According to a study, about half of orthopedics in Brazil use lead apron and very few of them use thyroid protector. More than 10% of orthopedics do not use personal protective equipment, with the main reason being that personal protective equipment hinders the surgical procedure. On the other hand very few physicians do not use personal protective equipment because the hospital does not provide it. Although, physicians were concerned about the complications of ionizing radiation, almost all of them did not use a dosimeter.¹⁴

Almost all spinal surgeons in Brazil wear lead aprons when working with ionizing radiation, more than half of them use thyroid protection and very few of them wear lead glasses and lead gloves. The majority of physicians though, reported that they do not use a dosimeter badge. Regarding the practice they use to protect themselves and patients from exposure to ionizing radiation it mainly includes standing one or two steps away from the fluoroscope, removing their hand from the field during fluoroscopy, using the fluoroscope in pulse mode and standing behind the radiation source. On the contrary, neurosurgeons use the

thyroid shield, navigation systems, and stay behind the X-ray source more often than orthopedic surgeons.¹⁵

In Tunisia all cardiologists wear lead apron and thyroid shield but only few of them use lead glasses and cap. At the same time a result worth noting is that most of cardiologists never wear a dosimeter due to unavailability. Also, the majority of cardiologists never visit or consult other doctors for health issues related to the exposure of ionizing radiation.¹⁶ In India, cardiologists who performed coronary interventions in the catheterization laboratory try to reduce the area of exposure of the patient and once every 6 months calibrate the equipment for optimal frequency. However, they appear to be neglectful of selecting the frame rate for fluoroscopy, terminating timely the Cine recording, and moving away from the x-ray unit. Cardiologists tend to position the image detector closer to the patient's chest but only half of them use a dosimeter and most of them place it on the chest under the lead apron. Regarding personal protective equipment, all cardiologists wear lead apron, most of them wear thyroid shields and few of them wear lead goggles and lead caps.¹⁷

Almost all cardiologists in Africa use a lead apron when they are in the catheterization laboratory, and most of them use thyroid protection. Half of cardiologists never use radiation protection eyeglasses and none of them uses radiation protection gloves, while the major reason for not using radiation protection tools being their unavailability.¹⁸

Most emergency physicians discuss with their patients about the potential dangers of radiation before conducting the exam, and one third of them observe the benefits and drawbacks of the diagnostic radiological exam on a pregnant patient requesting for a lead vest to be worn by the patient.¹⁹

Most females do not wear female-specific lead due to lack of availability and discomfort.⁴ In recent years there has been an increased concern about fetus radiation dose and fetus damage for women who are pregnant and due to that fear, female invasive cardiologists stop working when they get pregnant. When females use lead aprons of 0.5 mm Pb, the levels of radiation that their abdominal is exposed to are very low. Thus, lead aprons are an appropriate radiation safety practice for female physicians who use ionizing radiation.²⁰

Consultant physicians often fail to inquire whether patients have undergone previous radiographic imaging. The purpose of the question regarding the previous X-rays is to report the clinical necessity of the exams. The majority of them lacked awareness of the concept of ALARA because they failed to take part in courses that promote radiation safety. Only one out of three physicians evaluates the patients' effective dose and its measurement unit. Most physicians supported that the lack of education regarding the biological effects of radiation during medical school had caused the increase in demand for radiography and CT-scopes.²¹

Knowledges of occupational radiation exposure

Physicians in Saudi Arabia have low levels of knowledge about radiological safety. The highest percentage of correct answers of physicians is about modalities that have more radiation followed by most common complication of radiation exposure and the lowest percentage of correct answers is about the safety of the patient's relative to enter the CT room with the patient during the imaging process.²² In Canada, physicians are not educated on radiation safety. As a result, their knowledge about radiation is very low.⁹

Almost none of the neurosurgeons in the USA could identify the safety limit for occupational radiation and about half of neurosurgeons identify the relationship between distance and radiation dose reduction.²

Orthopedics in the UK and Scotland have low levels of radiation training and low levels of knowledge.¹² In USA, orthopedics, general surgeons and emergency physicians underestimated the effective radiation doses of all imaging and especially pelvic X-ray, hip X-ray, and pelvic CT scan.¹³ Also, orthopedics from Brazil have low levels of knowledge of ionizing radiation. That's the reason they are afraid of it, only 1 in 3 physicians feel safe using ionizing radiation, move away from the equipment during the imaging and keep their hand out of the primary beam.¹⁴

Cardiologists found to have medium levels of knowledge since they gave higher percentages of correct answers regarding practices' influence on operator exposure over time and the risk of radio-induced cataracts after X-ray exposure for more than 10 years and lower percentages of correct answers regarding the

minimum distance from which X-rays become safe and the suitability of cataract a physician to work in the catheterization room.¹⁶

Additionally, urologists did not know the appropriate fluoroscopy doses and claimed that there is no relevant literature. Their overall knowledge of fluoroscopy was moderate and half of them had not read literature about the fluoroscopy protection and the potential damage it can cause to them and to patients.¹⁰

A study conducted in Iraq found that 71.9% of non-radiologists physicians answered correctly to the definition of radioactive radiation, 40.6% of participants were aware of the ALARA principle, 21.9% knew the unit of Radioactivity, and 15.6% knew the radiation dose of a single chest X-Ray.²³ In Iran, 87.44% of 41 urologists, neurologists, surgeons and orthopedics used dosimeter in the right place during procedures with ionizing radiation. The physicians had moderate level of knowledge about operator time (52%) and distance (50.6%). More than half of physicians didn't wear personal protective equipment for eyes (66.1%), thyroid (77.4%), and aprons (64.5%). The majority of physicians were aware of the sensitivity of fetuses or pregnant mothers (88.3%) and only 16% of them answered the questions about organ radiosensitivity correctly.²⁴ Similarly, another study from Iran found that physicians who work with ionizing radiation have a moderate level of knowledge.³

Emergency physicians underestimate the overall dose of ionizing radiation and they were found to have moderate level of knowledge of ionizing radiation. Only one third of physicians answered that a one-time abdominal CT increases the lifetime risk of developing cancer in children. One third of physicians responded correctly that the exam with the lowest risk to the fetus is the chest X-ray, as opposed to the lumbar and abdominal x-rays and the pelvic and abdominal ones.¹⁹

Consultant physicians have moderate level of knowledge about ionizing radiation. Just one to three physicians believed that MRI radiation exposes patients to ionizing radiation. The same percentage of physicians knew how to evaluate the radiation dose of patients and how to calculate the risk of cancer in patients following medical physics and radiobiology classes. A lot of the participants advocated for the necessity of retraining programs for radiation protection, the presence of a radiation safety officer

in hospitals and educational centers were considered essential.²¹ Training seems to improve the level of knowledge physicians have about radiation protection. Surgeons and anesthesiologists who attended a training program about radiation protection answered correctly significantly more questions about the ALARA principle, the distance from the source of radiation, the dosimeter, the annual dose limit and the dose limit for a pregnant woman.¹⁹

Longer length of service,^{3,19,22} having attended a radiation protection and safety course,²² increased frequency on ordering imaging per day²² and specialists¹⁹ are factors that affect positively the level of knowledge of physicians.

Implications of occupational radiation exposure

Fluoroscopy is commonly used for diagnostic and therapeutic purposes. The occupational radiation exposure has been associated with leukemia and nonmalignant thyroid nodular disease,² increased risk of cancer,^{9,11,21} cataract,¹⁶ headache, fatigue and eye symptoms with the most common one being the red eyes.¹⁰ Cardiologists seem to be aware of the cancer risk of radiation use. Surgeons and anesthesiologists feel at risk of a cataract.²⁵ Very few physicians, less than 1 to 10, have read articles on radiation safety²⁶ or have received recent training.^{11,14,19,27} The only training they have received took place more than 4 years ago.¹⁴ For these reasons, education on radiation safety is crucial.

CONCLUSIONS

Physicians utilize ionizing radiation for both diagnostic and treatment purposes. The level of knowledge and practice of the physicians who use ionizing radiation and are not radiologists, in terms of radiation protection is insufficient. The inadequate level of radiation knowledge can lead to increased radiation exposure and risk for complications to healthcare professionals and patients. ALARA is the foundation of radiation safety along with optimal use of protective equipment. There is a need for increased awareness and training on radiation safety for all doctors' specialties. Educational programs targeted to physicians will increase the levels of compliance of to radiation safety practices. Also, there is unavailability of most radiation protective

equipment and this increases the physician's exposure to ionizing radiation. Healthcare organizations must procure a sufficient amount of personal protective equipment so it is available to physicians who use ionizing radiation.

REFERENCES

1. Selmi M, Natarajan M. Radiation awareness amongst junior doctors. *J Adv Radiol Med Image*. 2016;1(2):205.
2. McCloskey K, Gajjar AA, Salem MM, Susca L, Sioutas GS, Srinivasan VM, Jankowitz BT, Burkhardt JK. Intraoperative Ionizing Radiation Exposure Awareness and Associated Morbidity in Neurosurgery: A Nationwide Survey. *World Neurosurgery*. 2024 May 27.
3. Shabani F, Hasanzadeh H, Emadi A, Mirmohammadkhani M, Bitarafan-Rajabi A, Abedelahi A, Bokharaeian M, Masoumi H, Seifi D, Khani T, Sanchooli M. Radiation protection knowledge, attitude, and practice (KAP) in interventional radiology. *Oman medical journal*. 2018 Mar;33(2):141.
4. Mengers S, Emara A, Maheshwer B, Lopresti J, Mistovich RJ. Knowledge and Utilization of Sex-Specific Lead Aprons Among Pediatric Orthopaedic Surgeons. *Journal of Pediatric Orthopaedics*. 2024 Oct 1;44(9):e858-62.
5. Burgio E, Piscitelli P, Migliore L. Ionizing radiation and human health: Reviewing models of exposure and mechanisms of cellular damage. An epigenetic perspective. *International journal of environmental research and public health*. 2018 Sep;15(9):1971.
6. Clement C, Ruehm W, Harrison J, Applegate K, Cool D, Larson CM, Cousins C, Lochard J, Bouffler S, Cho K, Kai M. Keeping the ICRP recommendations fit for purpose. *Journal of Radiological Protection*. 2021 Dec 6;41(4):1390.
7. Campanella F, Rossi L, Giroletti E, Micheletti P, Buzzi F, Villani S. Are physicians aware enough of patient radiation protection? Results from a survey among physicians of Pavia District-Italy. *BMC health services research*. 2017 Dec;17:1-6.
8. Shaw PV, Crouail P, Paynter R, Coeck M. Education and training in radiation protection: improving ALARA culture. *Journal of Radiological Protection*. 2015 Feb 19;35(1):223.

9. Smith M, Thatcher MD, Davidovic F, Chan G. Radiation Safety Education and Practices in Urology: A Review. *Journal of Endourology*. 2024 Jan 1;38(1):88-100.
10. Altintas E, Bahceci T, Batur AF, Kaynar M, Kilic O, Akand M, Goktas S, Gul M. A survey analysis of knowledge levels of urologists about radiation safety and fluoroscopy use. *International Journal of Clinical Practice*. 2021 Apr;75(4):e13862.
11. Tzelves L, Somani B, Knoll T, Kamphuis G, Sarica K, Seitz C, Liatsikos E, Skolarikos A. Level of knowledge on radiation exposure and compliance to wearing protective equipment: where do endourologists stand? An ESUT/EULIS survey. *World Journal of Urology*. 2020 Mar;38:761-8.
12. Snowden G, Jabbal M, Akhtar A. Radiation safety awareness and practices amongst orthopaedic surgeons in Scotland. *Scottish Medical Journal*. 2022 Aug;67(3):103-8.
13. Hijji FY, Schneider AD, Thomas MD, Lyons JG, Bohl DD, Jerele JL, Prayson MJ. Knowledge of radiation exposure associated with common trauma imaging modalities among orthopaedic surgeons, emergency medicine physicians, and general surgeons in the United States. *World Journal of Emergency Medicine*. 2023;14(4):294.
14. Barbosa HB, Silva AM. Perception of Orthopedics Surgeons about the Risks of Radiation Exposure in Fluoroscopy. *Revista Brasileira de Ortopedia*. 2022 Sep 2;57(04):546-51.
15. Falavigna A, Ramos MB, Iutaka AS, Menezes CM, Emmerich J, Taboada N, Riew KD. Knowledge and attitude regarding radiation exposure among spine surgeons in Latin America. *World neurosurgery*. 2018 Apr 1;112:e823-9.
16. Hammami R, Jmaa AB, Bahloul A, Charfeddine S, Ellouze T, Mallek S, Mrad IB, Abid L, Kammoun S, Jdidi J. Évaluation des pratiques et connaissances en radioprotection des cardiologues en Tunisie. *Pan African Medical Journal*. 2021 Mar 23;38(1).
17. Uthirapathy I, Dorairaj P, Ravi S, Somasundaram S. Knowledge and practice of radiation safety in the Catheterization laboratory among Interventional Cardiologists—An online survey. *Indian Heart Journal*. 2022 Sep 1;74(5):420-3.
18. Tefera E, Qureshi SA, Gezmu AM, Mazhani L. Radiation protection knowledge and practices in interventional cardiologists practicing in Africa: a cross sectional survey. *Journal of Radiological Protection*. 2020 Feb 21;40(1):311.
19. Barnawi RA, Alrefai WM, Qari F, Aljefri AA, Hagi SK, Khafaji M. Doctors' knowledge of the doses and risks of radiological investigations performed in the emergency department. *Saudi Medical Journal*. 2018;39(11):1130.
20. Fetterly KA, Schueler BA, Mihailovic JM, Fiedler JH, Sturchio GM, Cabalka A, Best PJ, Gulati R, Guerrero M. Radiation exposure and protection for (assumed) pregnant interventional cardiologists and electrophysiologists. *Journal of the Society for Cardiovascular Angiography & Interventions*. 2024 Sep 1;3(9):102239.
21. Azmoonfar R, Faghirnavaz H, Younesi H, Morovati E, Ghorbani ZH, Tohidnia M. Physicians' knowledge about radiation dose in radiological investigation in Iran. *Journal of biomedical physics & engineering*. 2016 Dec;6(4):285.
22. Wally SF, Sabir SA, Alharbi SM, Albalawi IA, Alharbi KM, Binsulaiman N, Albalawi NM, Alshareef AM. Radiation Safety Awareness Among Non-radiology Staff at Tabuk Hospitals, Saudi Arabia. *Cureus*. 2024 Oct 1;16(10):e70603.
23. Rowantree SA, Currie C. Orthopaedic surgeons' knowledge and practice of radiation safety when using fluoroscopy during procedures: A narrative review. *Radiography*. 2024 Jan 1;30(1):274-81.
24. Shafiee M, Rashidfar R, Abdolmohammadi J, Borzoueisileh S, Salehi Z, Dashtian K. A study to assess the knowledge and practice of medical professionals on radiation protection in interventional radiology. *Indian Journal of Radiology and Imaging*. 2020 Jan;30(01):64-9.
25. Giaccardi M, Anselmino M, Del Greco M, Mascia G, Paoletti Perini A, Mascia P, De Ferrari GM, Picano E. Radiation awareness in an Italian multispecialist sample assessed with a web-based survey. *Acta Cardiologica*. 2021 May 4;76(3):307-11.
26. Tunçer N, Kuyucu E, Sayar Ş, Polat G, Erdil İ, Tuncay İ. Orthopedic surgeons' knowledge regarding risk of radiation exposure: a survey analysis. *Sicot-j*. 2017;3.

-
27. Pires RE, Reis IG, de Faria ÂR, Giordano V, Labronici PJ, Bel-
angero WD. The hidden risk of ionizing radiation in the op-
erating room: a survey among 258 orthopaedic surgeons in
Brazil. *Patient Safety in Surgery*. 2020 Dec;14:1-7

ANNEX

FIGURE 1.

