



Health & Research Journal

Vol 11, No 2 (2025)

Volume 11 Issue 2 April - June 2025



To cite this article:

Paspaliari, S. (2025). The antihypertensive effect of plant foods. *Health & Research Journal*, *11*(2), 95–99. https://doi.org/10.12681/healthresj.41017

Editorial Article

THE ANTIHYPERTENSIVE EFFECT OF PLANT FOODS

Hypertension, the invisible health risk, according to WHO is a widespread health condition affecting over a billion people across the globe, with considerable burden on the healthcare system. Worryingly, 46% of adults with hypertension are unaware of it, while only 1 in 5 adults (21%) have it under control.¹ Traditionally, the main approach to managing hypertension has been a pharmacological intervention. However, in recent years, there has been a paradigm shift towards incorporating lifestyle modifications, with particular emphasis on dietary strategies, as part of a comprehensive approach to managing this condition.²

According to the National Health Institute, a public health strategy, which decreases the BP level in the general population by even slight amounts, has the potential to substantially reduce morbidity and mortality or at least delay the onset of hypertension. For example, it has been estimated that a 5 mmHg reduction of Systolic Blood Pressure (SBP) in the population would result in a 14 percent overall reduction in mortality due to stroke, a 9 percent reduction in mortality due to CHD, and a 7 percent decrease in all-cause mortality.³ Therefore, there is a healthcare urgency in underlining the importance of applying healthy dietary patterns to patients, as part of the holistic management of this condition, which leads to a decrease of this magnitude.

The DASH dietary pattern, acronym of Dietary Approaches to Stop Hypertension, is a diet promoted by The United States Association's National Institute of Health's National Heart, Lung, and Blood Institute (NHBLI) to combat this silent killer. This food pattern emphasizes daily consumption of whole grains, fruits and vegetables, low fat dairy products, nuts, beans and seeds, lean meat, poultry and fish, meanwhile restricting intake of saturated and total fat. In the original DASH Diet Study, a significant reduction in systolic and diastolic blood pressure (DBP) by 11.4 and 5.5 mmHg, respectively, in patients following the DASH diet vs the control diet was found although sodium intake was held constant, more evident in the hypertensive subjects.⁴

In Europe, the recently revised recommendations of hypertension by the ESH 2023 guidelines, advise the adoption of plant-based healthy diets limiting the consumption of animal-based foods and emphasize the increased consumption of potassium, preferably via dietary modification, in alignment with the dietary recommendations for BP regulation from the AHA guidelines.^{2,5}

The Mediterranean diet is based on the traditional dietary pattern of ancient Greece and southern Italy, which is rich in fruits, vegetables, legumes, cereals, dairy products (cheese, yogurt), fish, olive oil, and with a small amount of wine and lean meats. Although the main differences with the DASH diet are restricted in emphasizing the seasonality of fresh foods, the large number of foods containing polyphenols (especially olives and extra virgin olive oil) and the use of vegetable oils rather than animal oils, the antihypertensive impact of this diet seems less pronounced, although this might not be the case. Plenty of data from reviews and meta-analyses show that MD acts protectively reducing SBP and DBP by 1,39 to 2.35 mmHg and 1.53 to 1.58 mmHg, respectively, effect able to impact significantly on public health, while the odds of devel-

oping hypertension are 13% lower with higher rather than lower adherence to MD.^{6,7} In the Hellenic National Nutrition and Health Survey (HNNHS), overweight and obese individuals with stricter adherence to the MD with a score > 23 had a 36% lower probability of developing hypertension, and when combined with body weight status, this probability increased to 60%.⁸ Systematic reviews and meta-analyses of RCTs and observational studies show that vegetarian dietary patterns, including vegan diets and diets which incorporate specific animal products (eggs and dairy), show significant reducing action in SBP of 5-7 mmHg.^{9,10}

The impact of fruits and vegetables on BP has been imprinted in several studies, indicating that increased consumption relates with a significant drop in BP. In a 6-month interventional trial, participants who were educated to increase their fruit and vegetable consumption had a 4 mm Hg average drop in their SBP when compared with the control group.¹¹ In the prospective Chicago Western Electric Study, it was shown that men who consumed 14-42 cups of vegetables and fruits per month (0.5-1.5 cup per day), in comparison with lower consumers, showed significantly lower increase in SBP per year for the next seven years (-2.8 mmHg and -2.2 mm Hg totally), with a concomitantly significant reduction in the increase of the DBP.12

Nut consumption is known to reduce blood cholesterol and is associated with reduced cardiovascular disease, but effects on BP are inconsistent. Findings from the Walnuts and Healthy Aging Study, where the intervention group consumed 15% of the energy needs as walnuts for 2 years, indicated that walnut consumption reduces mean office SBP in elderly –4.6 mm Hg (95% CI, –7.4 to –1.8 mm Hg) in the walnut group vs –0.6 mm Hg (95% CI, –3.4 to 2.2 mm Hg) in the control group, especially in those with mild hypertension, reducing also the need for antihypertensive medication.¹³

Dietary fiber, in which food with complex carbohydrates are rich in, constitutes an ingredient which seems to have a considerable antihypertensive effect, lowering both SBP and DBP at a magnitude of -1.13 to -1.56 mmHg and -0.39 to -1.65, respectively, more profoundly in hypertensive subjects. So, the increased consumption of dietary fiber, contributes to the prevention and treatment of hypertension, which seems so important for cardiovascular health.^{14,15,16}

Nitric oxide (NO) is an omnipresent molecule, proposing its important role in health maintenance. It is a potent vasodilator and relaxant of the vascular endothelium, and a decrease in the availability and/or bioactivity of NO is associated with the development of hypertension, endothelial dysfunction, atherosclerosis, and CAD. NO, is mainly endogenously produced by the amino acid L-arginine, with the presence of oxygen and the NO synthase enzymes, however, the diet can provide an alternative source, contributing significantly to the availability of NO in the human body via the nitrate-nitric-NO pathway. Dietary patterns rich in nitrates like the Mediterranean, the DASH diet and other plant-based diets, contribute to the higher production of NO in our organisms in comparison with the Western dietary pattern, exerting their favorable effects in the cardiovascular system. The main dietary sources of inorganic nitrates, precursors of NO, are the green leafy vegetables, among which the beetroot holds the lead. Meta-analyses of inorganic nitrate in CVD risk factors, have shown that its consumption- mainly from beetroot juice- can decrease SBP and DBP significantly in short-term^{17,18} and lower the risk of hypertension incident after 3 years of consuming a diet rich in nitrate-containing vegetables in the Tehran and Lipid Glucose Study.¹⁹ The aforementioned plant-based diets have significant potential to enhance NO bioavailability because they contain variety of foods rich in I-arginine and nitrates—the two key substrates for endogenous and exogenous NO production, respectively. In addition, these dietary patterns are rich in vitamin C, polyphenols, and marine long-chain ω 3 fatty acids, which may enhance NO production and reduce NO degradation in the body.²⁰

WHO recommends increasing dietary potassium intake to reduce BP and the risk of CVD, stroke, and coronary heart disease in adults (strong recommendation). It recommends a potassium intake of at least 3,510 mg/day for adults (under recommendation conditions).²¹ A forest plot meta-analysis comparing BP levels in the K supplement group and the control group identified a mean difference of -3.9 (95% Cl, -5.2 to -2.6) and -2.4 (95% Cl, -3.8 to -1.1) mm Hg for SBP and DBP, respectively. A dose-response meta-analysis of RCTs on potassium intake and BP, showed that there exists a nonlinear relationship between potassium intake and SBP and DBP. Greater reduction in BP was seen in patients with hypertension and individuals with higher levels of sodium intake receiving K supplementation. Increased BP with high potassium excretion was noted in participants with hypertension taking antihypertensive medications and in individuals with low to intermediate Na intake, indicating that these groups should avoid excess intake of potassium supplements.²² Consequently, the sodium to potassium ratio may be more important than the specific amounts of sodium and potassium an individual consumes. The data support the view that an increase in the potassium to sodium ratio would reduce BP in the general population and would significantly reduce CV mortality. Compared to diets based on whole foods, diets based on processed foods are high in sodium and low in potassium.²³ Siani A et al showed that increasing dietary potassium intake from natural foods is a feasible and effective measure to reduce antihypertensive therapy, since 38% and 81% of the intervention group patients with hypertension after 1 year of following a high potassium diet stopped the treatment and reduced the treatment to less than 50% vs 9% and 29% in the control group of the customary diet, respectively.²⁴

According to the position document of the ESH, there are specific dietary components, called nutraceuticals, the use of which in low-risk individuals with normal-to-high BP levels, could decrease BP. Beetroot juice has the most convincing evidence of antihypertensive effect. Antioxidant rich beverages (teas, coffee) could be considered. Among nutrients, magnesium, potassium and vitamin C supplements could also improve BP.²⁵

It is, therefore, significant that healthcare professionals encourage their patients with pre-hypertension or hypertension to comply with the principles of a healthy diet which promotes the systematic consumption of whole plant-derived foods which through a variety of mechanisms act protectively and therapeutically against this health hazard.

References

- 1. World Health Organization: hypertension (2023) https://www.who.int/news-room/factsheets/detail/hypertension)
- 2023 ESH Guidelines for the management of arterial hypertension The Task Force for the management of arterial hypertension of the European Society of Hypertension: Endorsed by the International Society of Hypertension (ISH) and the European Renal Association (ERA) J Hypertens. 2023 Dec 1;41(12):1874-2071

- 3. The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. National Institutes of Health National Heart, Lung, and Blood Institute
- 4. A clinical trial of the effects of dietary patterns on blood pressure. DASH Collaborative Research Group, N Engl J Med 1997 Apr 17;336(16):1117-24
- 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. Hypertension. 2018;71:e13–e115
- 6. The effect of Mediterranean diet on metabolic syndrome and its components: a meta-analysis of 50 studies and 534.906 individuals. J Am Coll Cardiol . 2011 Mar 15;57(11):1299-313 doi: 10.1016/j.jacc.2010.09.073
- 7. Effects of a Mediterranean diet on blood pressure: a systematic review and meta-analysis of randomized controlled trials and observational studies. J Hypertens. 2021 Apr 1;39(4):729-739. doi: 10.1097/HJH.00000000002667.
- 8. Presence of Hypertension Is Reduced by Mediterranean Diet Adherence in All Individuals with a More Pronounced Effect in the Obese: The Hellenic National Nutrition and Health Survey (HNNHS). Nutrients 2020, 12, 853
- 9. Vegetarian diets and blood pressure: a meta-analysis. JAMA Intern Med 2014 Apr;174(4):577-87
- 10. The effect of plant-based dietary patterns on blood pressure: a systematic review and meta-analysis of controlled intervention trials. J Hypertens. 2021 Jan;39(1):23-37. doi: 10.1097/HJH.00000000002604.
- 11. Effects of fruit and vegetable consumption on plasma antioxidant concentrations and blood pressure: a randomized controlled trial. Lancet 2002 Jun 8;359(9322):1969-74. doi: 10.1016/s0140-6736(02)98858-6.
- 12. Relation of vegetable, fruit, and meat intake to 7-year blood pressure change in middle-aged men: the Chicago Western Electric Study. 2004 Mar 15;159(6):572-80.
- 13. Effect of a Walnut Diet on Office and 24-Hour Ambulatory Blood Pressure in Elderly Individuals. Hypertension. 2019 May;73(5):1049-1057.
- 14. Dietary Fiber and Blood Pressure. Arch Intern Med. 2005;165:150-156
- 15. Effect of dietary fiber intake on blood pressure: a meta-analysis of randomized, controlled clinical trials. J Hypertens. 2005 Mar;23(3):475-81.
- 16. The effect of viscous soluble fiber on blood pressure: A systematic review and meta-analysis of randomized controlled trials. Nutr Metab Cardiovasc Dis. 2018 Jan;28(1):3-13.
- 17. Inorganic nitrate and beetroot juice supplementation reduces blood pressure in adults: a systematic review and metaanalysis. J Nutr. 2013 Jun;143(6):818-26.
- 18. The role of inorganic nitrate and nitrite in cardiovascular disease risk factors: a systematic review and meta-analysis of human evidence. Nutrition Reviews Vol. 76(5):348-371
- 19. Consumption of nitrate-containing vegetables is inversely associated with hypertension in adults: a prospective investigation from the Tehran Lipid and Glucose Study. J Nephrol. 2016 Jun;29(3):377-384.

- 20. Nitric Oxide Boosting Effects of the Mediterranean Diet: A Potential Mechanism of Action. Journals of Gerontology: BIOLOGICAL SCIENCES, 2018, Vol. 73, No. 7
- 21. Guideline: Potassium intake for adults and children. World Health Organization, 2012 (Reprinted, 2014).
- 22. Potassium Intake and Blood Pressure: A Dose-Response Meta-Analysis of Randomized Controlled Trials. J Am Heart Assoc 2020 9:e015719.
- 23. Should we eat more potassium to better control blood pressure in hypertension? Nephrol Dial Transplant (2019) 34: 184–193 doi: 10.1093/ndt/gfx340
- 24. Increasing the dietary potassium intake reduces the need for antihypertensive medication. Ann Intern Med 1991 Nov 15;115(10):753-9. doi: 10.7326/0003-4819-115-10-753.
- 25. Nutraceuticals and blood pressure control: a European Society of Hypertension position document. Journal of Hypertension 2020, 38:799–812

Sofia Paspaliari, Dietitian-Nutritionist, Msc Clinical Nutrition, Chief of Nutrition & Dietetics Department, General Hospital of Athens "Ippokrateio", Greece