


Social Cohesion and Development

Vol 9, No 1 (2014)



Primary Health Services Utilization in Greece: Studying the Past for Planning the Future

Dimitris Zavras, Mary Geitona, John Kyriopoulos

doi: [10.12681/scad.8925](https://doi.org/10.12681/scad.8925)

Copyright © 2016, Dimitris Zavras, Mary Geitona, John Kyriopoulos



This work is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike 4.0](https://creativecommons.org/licenses/by-nc-sa/4.0/).

To cite this article:

Zavras, D., Geitona, M., & Kyriopoulos, J. (2016). Primary Health Services Utilization in Greece: Studying the Past for Planning the Future. *Social Cohesion and Development*, 9(1), 49–61. <https://doi.org/10.12681/scad.8925>

Primary Health Services Utilization in Greece: Studying the Past for Planning the Future

Dimitris Zavras, *National School of Public Health*
Mary Geitona, *University of Peloponnese*
John Kyriopoulos, *National School of Public Health*

Χρήση Πρωτοβάθμιων Υπηρεσιών Υγείας στην Ελλάδα: Μελέτη του Παρελθόντος για το Σχεδιασμό του Μέλλοντος

Δημήτρης Ζάβρας, *Εθνική Σχολή Δημόσιας Υγείας*
Μαρία Γείτονα, *Πανεπιστήμιο Πελοποννήσου*
Γιάννης Κυριόπουλος, *Εθνική Σχολή Δημόσιας Υγείας*

ABSTRACT

The objective of this paper is to study healthcare utilization in Greece. The data were derived from a national survey conducted in 2006, and analysed through Logistic and Linear Regression. The likelihood of primary healthcare utilization is determined by the gender, the existence of a chronic disease, the self-rated health and the age, while the logarithm of the visits to primary health services is determined by the gender, the existence of a chronic disease, the income, the geographical region and the perceived threat from the health condition. Thus, primary healthcare utilization is determined by beliefs, health need as well as socioeconomic factors.

KEY WORDS: Primary health services, healthcare utilization, change over time, Greece

ΠΕΡΙΛΗΨΗ

Σκοπός του συγκεκριμένου άρθρου είναι η μελέτη της χρήσης των πρωτοβάθμιων υπηρεσιών υγείας στην Ελλάδα. Για το σκοπό αυτό αναλύθηκαν, μέσω των μεθόδων Logistic και Linear Regression, δεδομένα από πανελλαδική έρευνα η οποία υλοποιήθηκε το έτος 2006. Η πιθανότητα χρήσης υπηρεσιών υγείας προσδιορίζεται από το φύλο, την ύπαρξη χρόνιας νόσου, το αυτό-αξιολογούμενο επίπεδο υγείας και την ηλικία, ενώ ο λογάριθμος του αριθμού των επισκέψεων προσδιορίζεται από το φύλο, την ύπαρξη χρόνιας νόσου, τη γεωγραφική περιφέρεια, το εισόδημα και την αντίληψη σχετικά με την απειλή από τα ιατρικά προβλήματα. Η χρήση πρωτοβάθμιων υπηρεσιών υγείας συσχετίζεται επομένως με τις αντιλήψεις των ατόμων αλλά και με παράγοντες οι οποίοι εκφράζουν τις ανάγκες υγείας ή το κοινωνικοοικονομικό επίπεδο.

ΛΕΞΕΙΣ-ΚΛΕΙΔΙΑ: Πρωτοβάθμιες υπηρεσίες υγείας, χρησιμοποίηση υπηρεσιών υγείας, μεταβολές στο χρόνο, Ελλάδα

1. Introduction

The utilization of health services is determined mainly by the health needs and by several geographical, demographic, socioeconomic and structural factors of the healthcare systems (Phelps and Newhouse, 1974; Newhouse and Marquis, 1978; Wagstaff, 1986; Kasper, 1986; Feldstein, 1988; McGuire et al., 1988; Marmot and Wilkinson, 1999). More specifically, perceived morbidity, self-rated health (SRH), degree of urbanization, geographical region, age, gender, marital status, education, income, occupation and insurance coverage, are considered as the most important determinants of health services use (Pappa and Niakas, 2006; Economou, 2006, Geitona et al., 2007, Alexopoulos and Geitona, 2009).

Since medical care is an input of the health production process (Grossman, 1972), the understanding of the mechanism through which the factors mentioned above affect health services utilization, is crucial in understanding differences presented in population health. However, the relative importance of health services utilization's predictors, as well as health status's predictors, may change over time (Lix et al., 2002; Bernstein et al., 2003), making imperative the need for examination of findings on changes over time relative to social and economic policies (Braveman, 2003). Since the negative impact of current economic crisis on SRH and health care needs' satisfaction in Greece (Kentikelenis et al., 2011; Zavras et al., 2012), is the outcome of the economic policies adopted during the last three years, particular attention should be given on past health services use (Parkhurst, 2008). In addition, since past healthcare utilization predicts, in many cases, future utilization (Doran, 2011), studying the past may contribute to a clearer understanding of the present and this may affect the future (Porter, 1995). Thus, research which examines past experience empowers policy analysis that should be focused on the future (Klarman, 1980).

In this context, the aim of this article is to identify factors affecting primary health services utilization in Greece during 2006, attempting to provide decision-makers with insights into how to prioritize healthcare resources and manage utilization by studying the past.

2. Material and Methods

This paper is based on the data of a cross-sectional survey conducted in Greece between May 2006 and July 2006. Sample selection was based on the 2001 Census of the Hellenic Statistical Authority, following random stratified sampling by age, gender, degree of urbanization and geographical region of place of residence (NUTS II). Due to lack of information related to the distribution of health services utilization with the stratification variables, the formula of simple random sampling was used:

$$n > \frac{p(1-p)z_{a/2}^2}{d^2}$$

where $a=0.05$, $d=1.55\%$, and $p_{\text{users-non users}}=50\%$. The sample size was calculated at $n=4003$ adult individuals (18+) and distributed in 364 strata (strata_{gender}=2, strata_{region}=13, strata_{degree of urbanization}=2, strata_{age}=7). For the data collection the method of personal face to face interviews

with a structured questionnaire was applied. The questionnaires were piloted before the main data collection phase (June 2006). The interviews were carried out by specially trained professional interviewers and standard quality assurance procedures were used for reconfirming 15% of the interviews.

The survey instrument was based on the methodology of the World Health Organization (WHO) (Üstün et al., 2001), but it was adapted to the characteristics of the Greek health care system (e.g. type of health services provided, health insurance fund). In order to identify ambiguities in the questionnaire and to control the degree of difficulty in completing it, 150 interviews were conducted in the area of Athens by experienced trained researchers. The audit showed that there were no questions creating problems to the respondents and therefore the 150 completed questionnaires of the pilot survey were included in the study.

Per capita visits which are the key variable of the present study correspond to a recall period of one (1) year according to Jones (2006) and O'Donnell & Van Doorslaer (2007). Given that the minimum of this variable is 0, the analysis of the primary health services utilization was performed in two stages: a) on the first, the annual number of visits to doctors was dichotomized (0: 0 visits to doctors per year, ie non-utilization of health services during the year and 1: 1-maximum number of visits to doctors per year, ie utilization of health services during the year) and b) in the second stage, the variable representing the number of annual visits and had a minimum value of 1 (Buntin and Zaslavsky, 2004) was used. In the first stage given that the dependent variable is binary the Logistic Regression (LR) method was applied. In the second stage the variable under consideration is discrete quantitative so the selection of the appropriate statistical method was based on the methodology of Manning and Mullahy (2001), who suggest that the analysis should begin with a Generalized Linear Model (GLM) and if the coefficient of kurtosis of the ln scale residuals is greater than 3, the Ordinary Least Squares (OLS) method should be applied with the logarithm of the variable under consideration as the dependent variable. With this methodology we used a GLM with Gamma Family and Log Link. However, since the ln scale residuals of this model had coefficient of kurtosis $5,038 > 3$, the transformation of the variable under consideration as ln (utilization of primary health services) and the application of the OLS method was necessary. Both the LR Model and the OLS Model have been evaluated based on diagnostics tests as to their goodness of fit to the data and the violation of regression assumptions e.g. normality and homoscedasticity of residuals.

The independent variables used in the analysis are: a) gender (1: female, 2: male), b) age (1: 18-30, 2: 31-45, 3: 46-55, 4: 56 -65, 5: 66 +), c) self-rated health (1: very bad and bad, 2: moderate, 3: very good and good), d) the existence of chronic disease (0: no, 1: yes) e) education level (1: no education or primary education, 2: secondary education, 3: tertiary education), f) income (1: no income, 2: 1-500€, 3: 501-1000€, 4: 1001-1500€, 5: 1501-2000€, 6: 2001€ +), g) employment (1: employed, 2: unemployed, 3: retired, 4: housewife 5: school student, university student, soldier, 6: other), h) public health insurance (0: no, 1: yes), i) private health insurance (0: no, 1: yes), j) degree of urbanization (1: rural, 2: urban), k) geographical region (1: rest of Greece, 2: Athens, 3: South Attica, 4: West Attica, 5: North Attica, 6: Piraeus), l) perceived threat from the health condition (1: moderate or low at all times, 2: serious sometimes, 3: serious most of the times).

3. Results

Regarding the sample characteristics, the participants mean age was equal to 45.71 years ($n = 3983$, missing values $= 20$, standard deviation $= 17.52$), the percentages of men and women 49.29% ($n = 1973$) and 50.71% ($n = 2030$), respectively, and the percentages of urban and rural areas 63.53% ($n = 2543$) and 36.47% ($n = 1460$), respectively. Table 1 shows the percentage of geographical distribution of the sample.

According to the LR Model, the likelihood of utilization of the primary health services is determined by: a) age, b) self-rated health, c) gender and d) the existence of chronic disease. More specifically, as the OR of age is equal to $0.88 < 1$, the elderly, compared with younger people, have lower likelihood to use primary health services. Additionally, as the OR of the gender equals $0.66 < 1$, men compared to women, have lower likelihood to use primary health services. The same holds for people with higher levels of self-rated health, compared to people with lower levels of self-rated health, since the OR for this variable is equal to $0.62 < 1$. However, persons who suffer from a chronic disease, compared to healthy individuals, have a higher likelihood to use primary health services, since the OR is equal to $6.52 > 1$ (Table 2).

The results of the Link Test are satisfactory. The hat is statistically significant, while the hat² is not. Therefore this model does not suffer from specification error (Table 3).

The Pearson and Hosmer & Lemeshow goodness of fit criteria correspond to p equal $0.56 > 0.05$ and $0.61 > 0.05$ respectively, indicating that the model has a good fit to the data. In addition, the assumptions of normality of the deviance residuals and of homoscedasticity of standardized deviance residuals are not violated since $p_{\text{Jarque-Bera}} = 0.27 > 0.05$ and $p_{\text{Brown \& Forsythe}} = 0.37 > 0.05$.

Finally, as the area under the ROC curve is equal to 0.72, which is relatively high (Graph 1), this model has high ability of prediction.

With regard to the study of the frequency of primary health services utilization, according to the OLS model the logarithm of the number of visits to primary health services is determined by: a) the geographical region, b) the gender, c) the existence of a chronic disease, d) the perceived threat from the medical condition and e) the income. More specifically, the South East Attica and North Attica, presents a higher number of visits to primary health services compared to the city of Athens. However people who suffer from a chronic disease have higher rates of health services utilization than healthy people. The same holds for individuals who perceive the threat from a health condition as high. On the contrary, men compared to women and high income people compared to low income people have a lower utilization of health services (Table 4).

According to the results of the Link Test, this model does not suffer from a specification error as the hat is statistically significant while the hat² is not (Table 5).

Moreover, as the p of the Modified Hosmer & Lemeshow criterion equals $0.069 > 0.05$, this model has a good fit to the data.

Additionally, the p of the Ramsey Reset Test is equal to $0.34 > 0.05$, indicating that there are no omitted variables.

Finally, the standardized residuals follow the normal distribution ($p_{\text{Jarque-Bera}} = 0.068 > 0.05$) with equal variances ($p_{\text{Brown \& Forsythe}} = 0.063 > 0.05$), therefore the assumptions of the regression are not violated.

4. Discussion

According to the results of the analysis, the likelihood of primary health services utilization is determined by the gender, the age, the existence of a chronic disease and the self-rated health. In addition, the logarithm of the frequency of visits to primary health services is determined by the gender, the existence of a chronic disease, the income, the geographical region, and the perceived threat from the health condition.

The results of our analysis are consistent with international literature (Fylkesnes, 1993; Newbold et al., 1995; Murphy and Hepworth, 1996; Travassos et al., 2000; Van Der Heyden et al., 2003). More specifically, since women live longer than men, their health status is poorer (Bertakis et al., 2000). As a consequence, due to the increased morbidity that they experience, women have higher rates of utilization of health services than men (Redondo-Sendino et al., 2006). In our study, the differences of the primary health services utilization between men and women, may be, regardless of any differences in morbidity, a consequence of the difference in health behaviours that both genders adopt (women usually adopt healthier behaviours) (Zemp and Ackermann-Liebrich, 1988) or an effect of psychosocial factors (Green and Pope, 1999). Also, gender specific utilization differences can be considered as a consequence of women's tendency to use health services (Koopmans and Lamers, 2007). Verbrugge (1986) and Gijsbers van Wijk et al. (1991) suggested that women are more sensitive to symptoms and argued that women recall symptoms more readily than men and that they may be more willing to report their health problems. In addition, it is considered that women's greater experience of illness arises, at least in part, as a function of their social roles and position (Woods and Hulka, 1979). Psychosocial factors such as psychological distress, life stress, social support etc. predict health services utilization (Dorn et al., 2010). More specifically, persons undergoing stress tend to use more health services and the relationship between psychological distress and primary health care utilization is positive (Gortmaker and Eckenrode, 1982). In addition, the interaction of high life stress exposure and low social support is consistently linked to increased rates of health services utilization (Counte and Glandon, 1991).

Due to biological deterioration, age is associated with the utilization of health services, and it seems that this association is positive (Nie et al., 2008). However, as in our paper, some studies claim that the association between the age and the utilization of health services is negative (Economou et al., 2008). The negative association between the age and the likelihood of utilization of primary health services, regardless of the fact that health status may appear to be improved even in advanced age (Mitnitski et al., 2007), can be attributed to differences in access to health services across the various age groups (Whitehead, 1992), since psychological and physical barriers affect access to care among the elderly which is also influenced by poverty (Fitzpatrick et al., 2004).

The utilization of health services is also determined to a large extent by the self-reported morbidity (Foets et al., 1985). More specifically, self-reported morbidity is associated with higher probability of health services utilization. Thus, the health needs reflected through the self-rated health are negatively associated with the utilization of health services, as persons with lower self-rated health have a higher probability of using health services (Fernandez de la Hoz and Leon, 1996). The effect of the self-rated health in the specific study confirms the findings of similar studies which claim that a better health status relates to a lower number of doctor visits (Lima and Kopec, 2005). The effect of the existence of a chronic disease in the utilization of primary health services shows that the individuals' choices on health care seeking are a response to the

disease symptoms experienced. These symptoms include physical and emotional distress and impairments on various functions of the body and are not a mechanistic process (Lynch and Gardner, 2009). Since the association between the existence of health conditions and health care seeking behaviour is positive, and the association between the health status and the utilization of health care is negative (Shibusawa and Mui, 2008; Simmons et al., 2008), these findings are consistent with the international literature.

The income is also an important factor affecting the utilization of health services (Louvison et al., 2008), as it seems that people of low income have lower probability of utilization of health services (Geitona et al., 2007; Lemstra et al., 2009). The association between the income and the utilization of primary health services, which according to the analysis of this study is negative may reflect the fact that people with higher income adopt healthier behaviours (Leganger and Kraft, 2003). There is an association between common health practices like smoking, drinking, regularity of meals, physical activity and physical health status (Joung et al., 1995). Except for the positive association between income and high alcohol intake, measures of "unhealthy" behaviours are inversely associated with the socio-economic indices, suggesting that individuals in lower socio-economic groups are at an increased risk for health problems (Pomerleau et al., 1997). Consequently, health-related behaviors act as risk factors of poor health and lead to increased utilization of health services (Ruigomez et al., 1995). In addition, differences in health behaviours observed among different socioeconomic groups, are largely due to the fact that people of these groups perceive differently the random nature of health and the impact that the adoption of healthy lifestyles has in shaping the health status (Lantz et al., 1998; Kieffer et al., 2006; McGuire et al., 2006). It is not surprising, therefore, that the income as a determinant of health behaviour acts as a determinant of health services utilization (Strain, 1991). In addition, the negative association between the income and the utilization of health services is explained by the fact that people of higher socioeconomic status have better health (Adler and Ostrove, 1999; Molarius et al., 2007; Alexopoulos and Geitona, 2009; Zavras et al., 2011) and visit more frequently specialists (Dunlop et al. 2000). Additionally, the increased utilization of primary health services by people with lower income, based on a psychosocial interpretation, reflects the fact that people with lower income are more exposed to adverse psychosocial events compared to higher income people (Davey-Smith et al., 1998; House, 2001; Pikhart et al., 2003).

The differences in the utilization of primary health services across geographical regions may be partially explained by the impact of the physical environment on health, and specifically by the existence and the types of hazards (biological, physical, chemical, etc.), their source (natural, industrial, agricultural, etc.), 3) their origin (air, water, food, etc.) and their space of exposure (home, community, school, work) (Young, 2005). Additionally, the "sociobiological translation", is a phenomenon that explains the impact of society and the socio-economic environment on people's health as it considers that the way in which the social characteristics are perceived leads to biological signals which in turn are converted into disease (Tarlov, 1996). However, it can also be attributed to differences in access which is determined by the distance from health services or by spatial factors (Brooks, 1973; Meade and Earickson, 2005). These differences are explained by the distance from health services and the lack of access to the means of transport (Arcury et al., 2005a; Arcury et al., 2005b).

Finally, the results of our study are also consistent with the fact that the ways in which people perceive the threat from the disease plays a dominant role in decisions related to health care seeking (Rogers and Elliott, 1997; Rosenstock, 2005).

The findings of this study may contribute to effective planning of health services in Greece in times of economic crisis since they provide evidence from the past. The importance of this point lies in the fact that much of what we live within the present is a direct result of decisions made in the past (Tosh, 2000; Merriman, 2000; Ion and Beer, 2003). Understanding the past is a useful way of opening up the possibilities that may exist in the present and the future, especially when the economy slows down as in our days.

Bibliographical References

- Adler N. E. and Ostrove J. M., (1999), "Socioeconomic Status and Health: What we Know and What we Don't", *Annals of the New York Academy of Sciences* 896:3-15.
- Alexopoulos E. and Geitona M., (2009), "Self-Rated Health: Inequalities and Potential Determinants" *International Journal of Environmental Research and Public Health* 6(9):2456-69.
- Arcury T. A., Gesler W. M., Preisser J. S., Sherman J., Spencer J. and Perin J., (2005 a), "The Effects of Geography and Spatial Behavior on Health Care Utilization Among the Residents of Rural Region", *Health Services Research* 40(1):135-155.
- Arcury T. A., Preiser J. S., Gesler W. M. and Powers J. M., (2005 b), "Access to Transportation and Health Utilization in a Rural Region", *The Journal of Rural Health* 21(1):31-38.
- Bernstein A.B., Hing E., Moss A.J., Allen K.F., Siller A.B. and Tiggle R.B., (2003), *Health Care in America: Trends in utilization*. Maryland: National Center for Health Statistics.
- Bertakis K. D., Azari R., Helms L. J., Callahan E. J. and Robbins J. A., (2000), "Gender Differences in the Utilization of Health Care Services", *The Journal of Family Practice* 49(2):147-152.
- Braveman P.A., (2003), "Monitoring Equity in Health and Healthcare: A Conceptual Framework", *Journal of Health, Population & Nutrition* 21(3):181-192.
- Brooks C.H., (1973), "Associations Among Distance, Patient Satisfaction and Utilization of Two Types of Inner-city Clinics", *Medical Care* 11:373-383.
- Buntin M. B. and Zaslavsky A. M., (2004), "Too Much Ado About Two-Part Models and Transformation? Comparing Methods of Modeling Medicare Expenditures", *Journal of Health Economics* 23:525-542.
- Counte M. A., Glandon G. L., (1991), A Panel Study of Life Stress, Social Support, and the Health Services Utilization of Older Persons, *Medical Care* 29(4): 348-261.
- Davey Smith G., Hart C., Hole D., MacKinnon P., Gillis C., Watt G., Blane D. and Hawthorne V., (1998), "Education and Occupational Social Class: Which is the More Important Indicator of Mortality Risk?", *Journal of Epidemiology and Community Health* 52:153-160.
- Doran D. M., (2011), *Nursing Outcomes: The State of the Science*, London: Jones & Bartlett.
- Dorn S. D, Hernandez L., Minaya M. T., Morris C. B., Hu Y., Lewis S., Leserman J., Bangdiwala S. I., Green P. H. R., Drossman D. A., (2010), "Psychosocial Factors Are More Important Than Disease Activity in Determining Gastrointestinal Symptoms and Health Status in Adults at a Celiac Disease Referral Center", *Digestive Diseases and Science* 32(4): 354-357.
- Dunlop S., Coyte P.C., McIsaac V., (2000), "Socioeconomic Status and the Utilization of Physician Services: Results from the Canadian National Population Health Survey", *Social Science and Medicine* 51(1): 123-133.

- Economou A., Nikolaou A. and Theodossiou I., (2008), "Socioeconomic Status and Health-Care Utilization: A Study of the Effects of Low Income, Unemployment and Hours of Work on the Demand for Health Care in the European Union", *Health Services Management Research* 21(1):40-59.
- Economou, C. (2006) "Inequalities in Health, Poverty and Social Exclusion: Theoretical Approaches and Strategic Choices", in: Economou C. and Feronas A., (eds), *Poverty and Social Exclusion in Modern Societies*, p.p 313-347, Athens: Dionikos (in Greek).
- Feldstein P. J., (1988), *Health Care Economics*, New York: John Wiley.
- Fernandez de la Hoz K. and Leon D. A., (1996), "Self-Perceived Health Status and Inequalities in Use of Health Services in Spain", *International Journal of Epidemiology* 25: 593-603.
- Fitzpatrick A. L., Powe N. R., Cooper L. S, Ives D. G., Robbins J A., (2004), "Barriers to Health Care Access Among the Elderly and Who Perceives Them", *American Journal of Public Health* 94(10): 1788-1794.
- Foets M., Berghmans F. and Janssens L., (1985), "The Primary Health Care Project in Belgium: A Survey on the Utilization of Health Services", *Social Science & Medicine* 20(3):181-190.
- Fylkesnes K., (1993), "Determinants of Health Care Utilization Visits & Referrals", *Scandinavian Journal of Social Medicine* 21:40-50.
- Geitona M., Zavras D. and Kyriopoulos J., (2007), "Determinants of Healthcare Utilization in Greece: Implications for Decision-Making", *European Journal of General Practice* 13(3): 144-150.
- Gijsbers van Wijk, C.M.T., van Vliet, K.P., Kolk, A.M., Everaerd, W.T.A.M., (1991), "Symptom Sensitivity and Sex Differences: A Review of Health Surveys in the United States and the Netherlands", *Women and Health* 17 (1), 91-124.
- Gortmaker S. L., Eckenrode J., (1982), "Stress and the Utilization of Health Services: A Time Series and Cross-Sectional Analysis", *Journal of Health and Social Behavior* 23(1): 25-38.
- Green C. A. and Pope C. R., (1999), "Gender, Psychosocial Factors and the Use of Medical Services: A Longitudinal Analysis", *Social Science and Medicine* 48(10):1363-1372.
- Grossman M., (1972), *The Demand for Health: A Theoretical & Empirical Investigation*, Cambridge MA: National Bureau of Economic Research.
- House J. S., (2001), "Understanding Social Factors and Inequalities in Health: 20th Century Progress and 21th Century Prospects", *Journal of Health and Social Behavior* 43: 125-142.
- Ion R. M. and Beer M. D., (2003), "Valuing the Past: The Importance of an Understanding of the History of Psychiatry for Healthcare Professionals, Service Users and Carers", *International Journal of Mental Health Nursing* 12:237-242.
- Jones A. M., (2006), *The Elgar Companion to Health Economics*, Cheltenham Glos: Edward Elgar Publishing.
- Joung I. M. A., Stronks K., Van de Mheen H., Mackenbach J. P., (1995), "Health Behaviours Explain Part of the Difference in Self Reported Health Associated with Partner/Marital Status in the Netherlands, *Journal of Epidemiology and Community Health* 49:4820488.
- Kasper J. D., (1986), "Health Status & Utilization: Differences by Medicaid Coverage & Income", *Health Care Financing Review* 7:1-17.

- Kentikelenis A. Kranikolos M., Papanicolas I., Basu S., McKee M. and Stuckler D., (2011), "Health Effects of Financial Crisis: Omens of a Greek Tragedy", *The Lancet* 378(9801):1457-1458.
- Kieffer E. C., Sinco B. R., Rafferty A., Spencer M. S., Palmisano G., Watt E. E. and Heisler M., (2006), "Chronic Disease-Related Behaviors and Health Among African Americans and Hispanics in the Reach Detroit 2010 Communities, Michigan and the United States", *Health Promotion Practice* 7(3):256S-264S.
- Klarman H. E., (1980), "Observations on Health Services Research & Health Policy Analysis", *The Milbank Memorial Fund Quarterly* 58(2):201-216.
- Koopmans G. T. and Lamers L. M., (2007), "Gender and Health Care Utilization: the Role of Mental Distress and Help-Seeking Propensity", *Social Science and Medicine* 64(6):1216-1230.
- Lantz P. M., House J. S., Lepkowski J. M., Williams D. R., Mero P. R. and Chen J., (1998), "Socioeconomic Factors, Health Behaviors and Mortality: Results from a Nationally Representative Prospective Study of US Adults", *Journal of the American Medical Association* 279(21):1703-1708.
- Leganger A. and Kraft P., (2003), "Control Constructs: Do they Mediate the Relation Between Educational Attainment and Health Behavior?", *Journal of Health Psychology* 8(3):361-372.
- Lemstra M., Mackenbach J., Neudorf C. and Nannapaneni U., (2009), "High Health Care Utilization and Costs Associated with Lower Socio-Economic Status: Results from a Linked Dataset", *Canadian Journal of Public Health* 100(3):180-183.
- Lima V. D and Kopec J. A., (2005), "Quantifying the Effect of Health Status on Health Care Utilization Using a Preference-Based Health Measure", *Social Science and Medicine* 60(3):515-524.
- Lix L.M., Newburn-Cook C.V., Roos N.P. and Derksen S., (2002), "Trends in Health and Health-care Utilization in Manitoba", *Healthcare Management Forum* Winter Supplement:35-38.
- Louvison M. C., Lebrão M. L., Duarte Y. A., Santos J. L., Malik A. M. And Almeida E. S., (2008). "Inequalities in Access to Health Care Services and Utilization for the Elderly in Sao Paulo, Brazil", *Revista de Saude Publica* 42(4):733-740.
- Lynch W. D. and Gardner H. H., (2009), *Aligning Incentives, Information and Choice: How to Optimize Health and Human Capital Performance*, Health as a Human Capital Foundation, Wyoming: Cheyenne.
- Manning, W.G. and Mullahy, J., (2001), "Estimating Log Models: To Transform or Not to Transform?", *Journal of Health Economics* 20:461-494.
- Marmot M. and Wilkinson R., (1999), *Social Determinants of Health*, Oxford: Oxford University Press.
- McGuire A., Henderson J. and Mooney G., (1988), *The Economics of Health Care*, London: Routledge.
- McGuire L. C., Ahluwalia I. B. and Strine T W., (2006), "Chronic Disease-Related Behaviors in U.S. Older Women: Behavioral Risk Factor Surveillance System", *Journal of Women's Health* 15(1):3-7.

- Meade M. S. and Earickson R. J., (2005), *Medical Geography*, New York: Guilford Press.
- Merriman D., (2000), *Economic Conditions and State Tax Policy: Experience over the Last Decade and Implications for the Future*, Washington, D.C.: The Urban Institute.
- Mitnitski A., Song X. and Rockwood X., (2007), "Improvement and Decline in Health Status From Late Middle Age: Modeling Age-Related Changes in Deficit Accumulation", *Experimental Gerontology* 42(11):1109-1115.
- Molarius A., Berqlund K., Eriksson C., Lambe M., Nordstrom E., Eriksson H. G. and Feldman I., (2007), "Socioeconomic Conditions, Lifestyle Factors and Self-Rated Health Among Men and Women in Sweden", *European Journal of Public Health* 17(2):125-133.
- Murphy J. F. and Hepworth J. T., (1996), "Age & Gender Differences in Health Services Utilization", *Research in Nursing & Health* 19:323-329.
- Newhouse J. P. and Marquis S. E., (1978), "The Norms Hypothesis & the Demand for Medical Care", *Journal of Human Resources* 13:159-182.
- Nie J. X., Wang L., Tracy C. S., Moineddin R. and Upshur R. E., (2008), "Health Care Services Utilization Among the Elderly: Findings from the Study to Understand the Chronic Condition Experience of the Elderly and the Disabled", *Journal of Evaluation in Clinical Practice* 14(6):1044-1049.
- O'Donnell O. and Van Doorslaer E., (2007), *Analyzing Health Equity Using Household Survey Data: A Guide to Techniques and Their Implementation*, Washington DC.: World Bank Institute.
- Pappa E. and Niakas D., (2006), "Assessment of Health Care Needs and Utilization in a Mixed Public-Private System: The Case of the Athens Area", *BMC Health Services Research* 6:146 doi:10.1186/1472-6963-6-146.
- Parkhurst J., (2008), "Understanding Determinants of Health Service Use from a Systems Perspective", *Journal of Health Services Research & Policy* 13(2): 10.1258/jhsrp.2008.008019.
- Phelps C. E. and Newhouse J. P., (1974), *Coinurance and the Demand for Medical Services*, Santa Monica: RAND Corporation.
- Pikhart, H., Bobak, M., Rose, R. and Marmot, M., (2003), "Household Ownership and Self-Rated Health: Material and Psychosocial Explanations", *BMC Public Health* 3:38-44.
- Pomerleau J., Pederson L. L., Ostbye T., Speechley M., Speechley K. N., (1997), "Health Behaviours and Socio-Economic Status in Ontario", Canada, *European Journal of Epidemiology* 13(6): 613-622.
- Porter D., (1995), "The Mission of Social History of Medicine: a Historical View", *Social History of Medicine* 8(3):345-359.
- Redondo-Sentino A., Guallar-Castillon P., Banegas J. R. and Rodriguez-Artalejo E., (2006), "Gender Differences in the Utilization of Health-care Services Among the Older Adult Population in Spain", *BMC Public Health* 16;6:155.
- Rogers A. and Elliott H., (1997), *Primary Care: Understanding Health Need and Demand*, New York: Radcliffe Medical Press.
- Rosenstock I. W., (2005), "Why People Use Health Services", *The Milbank Quarterly* 83(4):1-32.

- Ruigomez A., Alonso J., Anto J. M. (1995), "Relationship of Health Behaviours to Five-year Mortality in an Elderly Cohort", *Age and Ageing* 24(2): 113-11.
- Shibusawa T. and Mui A. C., (2010), "Health Status and Health Services Utilization Among Older Asian Indian Immigrants", *Journal of Immigrant and Minority Health* 12(4):527-533.
- Simmons L. A., Anderson E. A. and Braun B., (2008), "Health Needs and Health Care Utilization Among Rural, Low-Income Women", *Women & Health* 47(4):53-69.
- Strain L. A., (1991), "Use of Health Services in Later Life: The Influence of Health Beliefs", *Journal of Gerontology* 46(3):S143-S150.
- Tarlov, A. R., (1996), "Social Determinants of Health: The Sociobiological Translation", in: Blane D., Brunner E. and Wilkinson R., (eds), *Health and Social Organisation*, p.p 71-93, London: Routledge.
- Tosh, J., (2000), *The Pursuit of History*, London: Pearson Education.
- Travassos C., Viacava F., Pinheiro R. and Brito A., (2000). "Utilization of Health Care Services in Brazil: Gender, Family Characteristics & Social Status", *Rev Panam Salud Publica* 11:365-373.
- Van Der Heyden J. H., Demarest S., Tafforeau J. and Van Oyen H., (2003), "Socioeconomic Differences in the Utilization of Health Services in Belgium", *Health Policy* 65:153-165.
- Verbrugge, L.M., (1986), "Role Burdens and Physical Health of Women and Men", *Women and Health* 11(1), 47-77.
- Üstün T. B, Chatterji S., Villanueva M., Bendib L., Çelik, C., Sadana R., Valentine N., Ortiz J., Tandon A., Salomon, J., Cao Y., Wan Jun X., Özaltin E., Mathers C. and Murray C. J. L., (2001), "WHO Multi-country Survey Study on Health and Responsiveness 2000-2001", *GPE Discussion Paper 37*, WHO (available at <http://www.who.int>).
- Wagstaff A., (1986), "The Demand for Health: Some New Empirical Evidence", *Journal of Health Economics* 5:195-233.
- Whitehead M., (1992), "The Concepts and Principles of Equality and Health", *International Journal of Health Services* 22(3):429-445.
- Woods, N.F., Hulka, B.S., (1979), Symptom Reports and Illness Behavior Among Employed Women and Homemakers, *Journal of Community Health* 5 (1), 36-45.
- Young T., K., (2005), *Population Health: Concepts and Methods*, Oxford: Oxford University Press
- Zavras D., Tsiantou V., Pavi E., Mylona K. and Kyriopoulos J., (2012), "Impact of Economic Crisis and Other Demographic and Socio-Economic Factors on Self-Rated Health in Greece", *The European Journal of Public Health* 1-5.
- Zavras, D., Geitona, M. and Kyriopoulos, J. (2011) "Determinants of Self-Rated Health in Greece", in: Kyriopoulos J. and Dimitrova D., (eds), *Public Health and Health Care in Greece and Bulgaria: The Challenge of the Cross-border Collaboration in Times of Financial Crisis*, p.p. 423-428, Athens: Papazissis.
- Zemp E. And Ackermann-Lieblich U., (1988), "Sex Differences in Health and Health Behavior", *Sozial und Praventivmedizin* 33(3):186-19.

Table 1: Geographical distribution

Geographical region	Frequency	Percentage (%)	Cumulative Percentage (%)
Rest of Greece	2561	63.98	63.98
Athens	301	7.52	71.50
Southeast Attica	463	11.57	83.06
West Attica	338	8.44	91.51
North Attica	140	3.50	95.00
Piraeus	200	5.00	100.00
Total	4003	100.00	

Table 2: LR Model on the Likelihood of Primary Health Services Utilization

Utilization of primary health Services	OR	Std. Err.	z	P>z	95% Confidence Interval
Age	0.8892932	.0304208	-3.43	0.001	0.8316243 -0.950961
Self-Rated Health	0.6230192	.0632894	-4.66	0.000	0.5105429-0.760275
Gender	0.6615056	.0523913	-5.22	0.000	0.566393-0.772589
Existence of Chronic Disease	6.520939	.8659601	14.12	0.000	5.026588-8.459545

Table 3: Link Test

Utilization of primary health services	Coefficient	Std. Err.	z	P>z	95% Confidence Interval
hat	1.050813	.2105878	4.99	0.000	0.6380683-1.463557
hat ²	-.0170314	.0682786	-0.25	0.803	-0.150855 -0.1167922
Constant	-.0210482	.1030794	-0.20	0.838	-0.2230801-0.1809836


Table 4: OLS Model on the Logarithm of the Number of Visits to Primary Health Services

Logarithm of the Number of Visits to Primary Health Services	Coefficient	Std. Err.	t	P>t	95% Confidence Interval
Rest of Greece	.0376427	.0697444	0.54	0.589	-.099136- .1744213
Southeast Attica	.1859269	.0828103	2.25	0.025	.0235241- .3483298
West Attica	.0664586	.0916526	0.73	0.468	-.1132854- .2462025
North Attica	.2590377	.1200437	2.16	0.031	.0236148- .4944606
Piraeus	.0771966	.1012301	0.76	0.446	-.1213301- .2757234

Gender	-.158914	.0379153	-4.19	0.000	-.2332713- .0845568
Existence of chronic disease	.8824502	.0416865	21.17	0.000	.800697- .9642034
Perceived threat of health condition	.1635113	.0246316	6.64	0.000	.1152053- .2118174
Income	-.0331781	.0158608	-2.09	0.037	-.0642834- -.0020728
Constant	.6400628	.1091232	5.87	0.000	.4260566- .854069

Table 5: Link Test

Logarithm of the number of visits to primary health services	Coefficient	Std. Err.	t	P>t	95% Confidence Interval
hat	.8100787	.2861558	2.83	0.005	.2488874 - 1.37127
hat ²	.0752244	.1124919	0.67	0.504	-.1453878 - .2958366
Constant	.098163	.1543703	0.64	0.525	-.2045787- .4009046



Biographical Notes

Dr. Dimitris Zavras is a Senior Researcher in the Department of Health Economics and Faculty Member in the Graduate Program in Healthcare Management at the National School of Public Health, Athens, Greece. He is in the Faculty of the Hellenic Open University (Graduate Program in Healthcare Management) and Neapolis University Pafos (Graduate Program in Public Administration). Dr. D. Zavras' research interests focus around the demand and utilization of healthcare services, patient satisfaction, and health inequalities. His latest publication is entitled "Investigating Factors of Self-Care Orientation and Self-Medication Use in a Greek Rural Area" (Rural & Remote Health, 2014). Address: Department of Health Economics, National School of Public Health, 196 Alexandras Avenue, 11521 Athens, Greece. E-mail: dzavras@gmail.com

Mary Geitona is Associate Professor in economic analysis of social policy and health economics at the University of Peloponnese. She is the founder of the Hellenic Society for Pharmacoeconomics (HELSPOR) and was the President of HELSPOR and editor-in-Chief of the Greek journal "Pharmacoeconomia" since 1990. Additionally, she is the founder and the president of the Greek Chapter of the International Society of Pharmacoeconomics & Outcomes Research (ISPOR). Her main interest is health technology assessment (HTA) and she is strongly involved in its implementation in the decision making in Greece. Address: Department of Social and Educational Policy, Damaskinou & Kolokotroni str., 20100 Corinth, Greece. E-mail: geitona@uop.gr

John Kyriopoulos is Professor Emeritus of Health Economics of the Department of Health Economics of the National School of Public Health. He has also served in long terms as the Head of the Department of Health Economics, Dean of the National School of Public Health, and Director of the Postgraduate Program in Health Services Administration. As an expert advisor he has developed extensive scientific activity in the Balkan Peninsula and the Black Sea region. He has received many awards from international organizations and universities. His research interests focus on Health Economics, Health Policy, and Social Policy and Insurance. His work has been widely presented in national and international conferences, and is published extensively in scientific and scholarly publications in the field of health economics and health policy. Address: Department of Health Economics, National School of Public Health, 196 Alexandras Avenue, 11521 Athens, Greece.