


ENTOMOLOGIA HELLENICA

Vol 19, No 2 (2010)



The Bougainvillea mealybug *Phenacoccus peruvianus*, a rapid invader from South America to Europe

A. Beltrà, A. Soto, J.-F. Germain, D. Matile-Ferrero, G. Mazzeo, G. Pellizzari, A. Russo, J.C. Franco, D.J. Williams

doi: [10.12681/eh.11581](https://doi.org/10.12681/eh.11581)

Copyright © 2017, A. Beltrà, A. Soto, J.-F. Germain, D. Matile-Ferrero, G. Mazzeo, G. Pellizzari, A. Russo, J.C. Franco, D.J. Williams



This work is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike 4.0](https://creativecommons.org/licenses/by-nc-sa/4.0/).

To cite this article:

Beltrà, A., Soto, A., Germain, J.-F., Matile-Ferrero, D., Mazzeo, G., Pellizzari, G., Russo, A., Franco, J., & Williams, D. (2010). The Bougainvillea mealybug *Phenacoccus peruvianus*, a rapid invader from South America to Europe. *ENTOMOLOGIA HELLENICA*, 19(2), 137–143. <https://doi.org/10.12681/eh.11581>

The Bougainvillea mealybug *Phenacoccus peruvianus*, a rapid invader from South America to Europe

A. BELTRÀ¹, A. SOTO^{1*}, J.-F. GERMAIN², D. MATILE-FERRERO³,
G. MAZZEO⁴, G. PELLIZZARI⁵, A. RUSSO⁴, J.C. FRANCO⁶ AND
D.J. WILLIAMS⁷

¹Institut Agroforestal Mediterrani, CPI Acceso 8E, Universitat Politècnica de València, 46022 Valencia, Spain

²Laboratoire National de la Protection des Végétaux, station de Montpellier, CBGP Campus International de Baillarguet, CS 30016, 34988 Montferrier-sur-Lez, France

³Muséum National d'Histoire Naturelle, Département de Systématique & Évolution, UMR 7205 MNHN-CNRS: Origine, Structure et Évolution de la Biodiversité case postale N° 50 75231 PARIS Cedex 05, France

⁴Dipartimento di Scienze e Tecnologie Fitosanitarie, University of Catania, Via S. Sofia 100, 95123 Catania, Italy

⁵Department of Environmental Agronomy and Crop Production - Entomology, University of Padua, Viale dell'Università 16, 35020 Legnaro, Italy

⁶Departamento de Protecção de Plantas e de Fitoecologia, Instituto Superior de Agronomia, Universidade Técnica de Lisboa, 1349-017 Lisboa, Portugal

⁷Department of Entomology, The Natural History Museum, Cromwell Road London SW7 5BD, UK

ABSTRACT

Scale insects are frequent invaders of new territories. The Bougainvillea mealybug, *Phenacoccus peruvianus* Granara de Willink, 2007, was recorded in Europe for the first time in 1999 in Spain (Almeria) and later in 2002 in Italy (Sicily). Initially, this unknown species was identified provisionally as *Phenacoccus* sp. Records of this species became frequent later when it was found in other localities in Spain (including the Balearic Islands), Great Britain, France (including Corsica), Monaco, and Portugal. The host plants of this mealybug were in most cases species of *Bougainvillea*.

KEYWORDS: Alien species, Pseudococcidae, *Phenacoccus peruvianus*, *Bougainvillea glabra*, invasive species, ornamental plants.

Introduction

In the last few decades, mealybugs (Hemiptera: Pseudococcidae) have been involved in several pest outbreaks in the tropics, causing severe damage to agricultural crops and ornamental plants: *Phenacoccus manihoti*

Matile-Ferrero on cassava throughout tropical Africa and *Rastrococcus invadens* Williams on fruit trees in West Africa (Herren and Neuenschwander 1991, Han et al. 2007), *Maconellicoccus hirsutus* (Green) in the Caribbean region (Matile-Ferrero et al. 2000), *Paracoccus marginatus* Williams and

*Corresponding author, e-mail: asoto@eaf.upv.es

Granara de Willink in the Caribbean and Pacific region (Matile-Ferrero et al. 2000, Muniappan et al. 2008), and *Phenacoccus solenopsis* Tinsley damaging cotton in India and Pakistan (Hodgson et al. 2008). Biological control is an efficient method for the management of these pests, and many of these outbreaks are now under control by natural enemies imported from the countries of origin of each species.

The increase of worldwide trade of plants has facilitated the introduction and spread of many pests in Europe and in the Mediterranean Basin. Scale insects represent the second largest group of alien insects in Europe after aphids (DAISIE 2008). This is especially true of mealybugs which are typical invasive species (Miller et al. 2005) due to their little size and cryptic behaviour. Since the 1990s, several mealybug species have invaded the Mediterranean area, such as *Phenacoccus solani* Ferris (Mazzeo et al. 1999), *Phenacoccus parvus* Morrison (Bendov et al. 2005) and *Pseudococcus comstocki* (Kuwana) (Pellizzari 2005). In the present communication we report for the first time the presence of *Phenacoccus peruvianus* Granara de Willink causing problems on ornamental plants in different countries in Europe.

Materials and Methods

Mealybug specimens were collected from diverse host plants of Portugal, Spain, France, Italy and UK. Samples were slide-mounted (Fig. 1) according to the procedure detailed by Williams and Granara de Willink (1992) with a few modifications made by the different authors depending on their own preferences. Specimens were identified using the keys of Granara de Willink and Szumick (2007). Communication between European specialists was entailed during the identification process.



FIG. 1. Microscopic permanent slide of adult female *Phenacoccus peruvianus*.

Results and Discussion

Records of the *Bougainvillea* mealybug in Europe

The *Bougainvillea* mealybug was first reported in 1999 in Almeria, Spain. Since then, this insect has been recorded in several areas of the Mediterranean coast of Spain, including the Balearic Islands, the south coasts of Portugal and France, including Corsica, and Sicily in Italy. It was also discovered in England (Wisley) under glass (Table 1). The fast dispersal of this insect indicates that the ornamental trade might be involved.

Mealybug identification

The systematic position of the *Phenacoccus* species found in the survey was studied by several specialists and finally clarified in 2008, after Granara de Willink

TABLE 1. First records of *P. peruvianus* in the different European countries and regions.

| Year | Country | Region | Host-plants | Host-plant family |
|-------------|----------------|----------------------------|---|---|
| 1999 | Spain | Andalusia | <i>Bougainvillea glabra</i> | Nyctaginaceae |
| 2002 | Italy | Sicily | <i>Bougainvillea glabra</i> | Nyctaginaceae |
| 2004 | Spain | Valencian Community | <i>Bougainvillea glabra</i> | Nyctaginaceae |
| 2005 | Great Britain | South East England | <i>Bougainvillea</i> sp. | Nyctaginaceae |
| | France | Corsica | <i>Bougainvillea</i> sp. | Nyctaginaceae |
| 2006 | Portugal | Algarve | <i>Bougainvillea</i> sp. | Nyctaginaceae |
| | Portugal | Lisbon | <i>Bougainvillea</i> sp. | Nyctaginaceae |
| | Spain | Catalonia | <i>Bougainvillea glabra</i> | Nyctaginaceae |
| 2008 | France | Provence-Alpes-Côte d'Azur | <i>Bougainvillea glabra</i> | Nyctaginaceae |
| | Spain | Valencian Community | <i>Araujia sericifera</i> , <i>Aucuba japonica</i> , <i>Myoporum laetum</i> | Asclepiadaceae Aucubaceae Myoporaceae |
| | Monaco | Monaco | <i>Bougainvillea</i> sp. | Nyctaginaceae |
| | France | Languedoc-Roussillon | <i>Buddleja</i> sp. | Scrophulariaceae |
| 2009 | Spain | Majorca | <i>Justicia suberecta</i> , <i>Solanum vespertilio</i> | Acanthaceae Solanaceae |

and Szumik (2007) had published a comprehensive paper on the Phenacoccinae of Central and South America, with descriptions of several new species. According to their identification key, description and illustration, the species recorded in Europe on *Bougainvillea* sp. is *Phenacoccus peruvianus* Granara de Willink, 2007, described from Peru and two provinces of Argentina (Santiago del Estero and Tucumán). This identification was confirmed by M.C. Granara de Willink (pers. commun.). Preliminary molecular studies demonstrate that the populations from Spain and France belong to the same species (Beltrà, unpublished work). Further studies are in progress to compare Neotropical populations with the European ones.

Phenacoccus peruvianus can be added to the list of 29 species (Ben-Dov et al. 2009) of the genus *Phenacoccus* reported from the western European Mediterranean area (Table 2). Although most of them are native to this area, species such as *P. madeirensis* Green

TABLE 2. List of *Phenacoccus* species identified in the Western European Mediterranean Basin, with location and year of first record (according to Ben-Dov et al. 2009).

| Species | Country | Date |
|---|----------------|-------------|
| <i>P. aceris</i> (Signoret) | France | 1875 |
| <i>P. asphodeli</i> Goux | France | 1942 |
| <i>P. avenae</i> Borchsenius | Italy | 1990 |
| <i>P. balagnus</i> Balachowsky | Corsica | 1933 |
| <i>P. defectus</i> Ferris | France | 2006 |
| <i>P. eschscholtziae</i> McKenzie | Italy | 1987 |
| <i>P. evelinae</i> (Tereznikova) | Italy | 1989 |
| <i>P. ferulae</i> Borchsenius | Italy | 1984 |
| <i>P. formicarum</i> Leonardi | Italy | 1908 |
| <i>P. gossypii</i> Townsend & Cockerell | Spain | 1946 |
| <i>P. graminicola</i> Leonardi | Italy | 1908 |
| <i>P. hordei</i> (Reuter) | France | 1933 |
| <i>P. hystrix</i> (Baerensprung) | Spain | 1985 |
| <i>P. incertus</i> (Kiritshenko) | Sardinia | 1992 |
| <i>P. interruptus</i> Green | France | 1935 |
| <i>P. longoi</i> Russo | Italy | 1994 |
| <i>P. madeirensis</i> Green | Italy | 1990 |
| <i>P. meridionalis</i> Gómez-Menor Ortega | Spain | 1965 |
| <i>P. neohordei</i> Marotta | Italy | 1992 |
| <i>P. parietariae</i> (Lichtenstein) | France | 1881 |
| <i>P. parietaricola</i> Goux | France | 1938 |
| <i>P. phenacoccoides</i> (Kiritchenko) | France | 1941 |
| <i>P. piceae</i> (Low) | France | 1931 |
| <i>P. pumilus</i> Kiritchenko | France | 1948 |
| <i>P. quadridauata</i> (Signoret) | France | 1875 |
| <i>P. silvanae</i> Longo & Russo | Italy | 1989 |
| <i>P. solani</i> Ferris | Sicily | 1999 |
| <i>P. yerushalmi</i> Ben-Dov | Sicily | 1994 |

TABLE 3. List of *Phenacoccus peruvianus* host plant species in Europe.

| Family | Plant species |
|------------------|----------------------------|
| Acanthaceae | <i>Justicia suberecta</i> |
| Amaranthaceae | <i>Alternanthera sp.</i> |
| Asclepiadaceae | <i>Araujia sericifera</i> |
| Asteraceae | <i>Baccharis sp.</i> |
| | <i>Eupatorium sp.</i> |
| Aucubaceae | <i>Aucuba japonica</i> |
| Myoporaceae | <i>Myoporum lateum</i> |
| Nyctaginaceae | <i>Bougainvillea spp.</i> |
| Scrophulariaceae | <i>Buddleja sp.</i> |
| Solanaceae | <i>Cestrum sp.</i> |
| | <i>Solanum vespertillo</i> |

and *P. solani* are exotic and have a Neotropical origin (Granara de Willink and Szumik 2007).

Host plants

Phenacoccus peruvianus feeds mainly on plant species of the genus *Bougainvillea*, which are native to tropical and subtropical regions of South America. These plants are widely grown in gardens in the Mediterranean region and are traded frequently as potted plants. However, the mealybug has been reported on many other plant species in the region and therefore must be regarded as a polyphagous species (Table 3).

Acknowledgements

We are grateful to M. C. Granara de Willink for discussing and confirming the identification.

References

Ben-Dov, Y., Y. Gottlieb and T. Sando. 2005. First record of *Phenacoccus parvus*

Morrison (Hemiptera: Coccoidea: Pseudococcidae) from the Palaearctic region. *Phytoparasitica* 33: 325-326.

Ben-Dov, Y., D.R. Miller and G.A.P. Gibson. 2009. ScaleNet, Scales in a Region/Country/Subunit Query. <http://www.sel.barc.usda.gov/scalecgi/region.exe?region=®name=&country=&ctryname=&subunit=&action=Pick+Region&querytype=&family=&genus=&intro=A> (last accessed on 3 January 2010).

[DAISIE] Delivering Alien Invasive Species Inventories for Europe. 2008. European Invasive Alien Species Gateway. <http://www.europe-alien.org/> (last accessed on 3 January 2010).

Granara de Willink, M.C. and C. Szumik. 2007. Phenacoccinae de Centro y Sudamérica (Hemiptera: Coccoidea: Pseudococcidae): Sistemática y Filogenia. *Rev. Soc. Entomol. Argen.* 66: 29-129.

Han, S.H., A.B. Ndiaye and J.L. Hemptinne. 2007. Plantes-hôtes et prédateurs de la cochenille farineuse du Manguier *Rastrococcus invadens* Williams, 1986, nouvellement introduite à Dakar, Sénégal (Hemiptera: Pseudococcidae). *Bull. Soc. Entomol. France* 112: 121-125.

Herren, H.R. and P. Neuenschwander. 1991. Biological control of cassava pests in Africa. *Annu. Rev. Entomol.* 36: 257-283.

Hodgson, C., G. Abbas, M.J. Ari, S. Saeed and H. Karar. 2008. *Phenacoccus sole-nopsis* Tinsley (Sternorrhyncha: Coccoidea: Pseudococcidae), an invasive mealybug damaging cotton in Pakistan and India, with a discussion on seasonal morphological variation. *Zootaxa* 1913: 1-35.

Matile-Ferrero, D., J. Étienne and G. Tiego. 2000. Introduction de deux ravageurs d'importance pour la Guyane française: *Maconellicoccus hirsutus* et *Paracoccus marginatus* (Hem., Coccoidea, Pseudococcidae). *Bull. Soc. Entomol. France* 105: 485-486.

Mazzeo, G., A. Russo and P. Suma. 1999. *Phenacoccus solani* Ferris (Homoptera:

- Coccoidea) on ornamental plants in Italy. Boll. Zool. Agrar. Bachic. 31: 31-35.
- Miller, D.R., G.L. Miller, G.S. Hodges and J.A. Davidson. 2005. Introduced scale insects (Hemiptera: Coccoidea) of the United States and their impact on US agriculture. Proc. Entomol. Soc.Wash. 107: 123-158.
- Muniappan, R., B.M. Shepard, G.W. Watson, G.R. Carner, D. Sartiami, A. Rauf and M.D. Hammig. 2008. First report of the Papaya Mealybug, *Paracoccus marginatus* (Hemiptera: Pseudococcidae), in Indo-nesia and India. J. Agric. Urban Entomol 25: 37-40.
- Pellizzari, G. 2005 Cocciniglie nuove o poco note potenzialmente dannose per l'Italia: Fiorinia pinicola Maskell, *Pseudococcus comstocki* (Kuwana), *Peliococcus turanicus* (Kiritshenko). Inf. Fitopatol. 55: 20-25.
- Williams, D.J. and M.C. Granara de Willink. 1992. Mealybugs of Central and South America. C.A.B. International, Wallingford, 1st ed. 644 pp.

Εισβολή του κοκκοειδούς *Phenacoccus peruvianus* από την Νότια Αμερική στην Ευρώπη

A. BELTRÀ¹, A. SOTO¹, J.-F. GERMAIN², D. MATILE-FERRERO³,
G. MAZZEO⁴, G. PELLIZZARI⁵, A. RUSSO⁴, J.C. FRANCO⁶ KAI
D.J. WILLIAMS⁷

¹Institut Agroforestal Mediterrani, CPI Acceso 8E, Universitat Politècnica de València, 46022 Valencia, Spain

²Laboratoire National de la Protection des Végétaux, station de Montpellier, CBGP Campus International de Baillarguet, CS 30016, 34988 Montferrier-sur-Lez, France

³Muséum National d'Histoire Naturelle, Département de Systématique & Évolution, UMR 7205 MNHN-CNRS: Origine, Structure et Évolution de la Biodiversité case postale N° 50 75231 PARIS Cedex 05, France

⁴Dipartimento di Scienze e Tecnologie Fitosanitarie. University of Catania, Via S. Sofia 100, 95123 Catania, Italy

⁵Department of Environmental Agronomy and Crop Production - Entomology, University of Padua, Viale dell'Università 16, 35020 Legnaro, Italy

⁶Departamento de Protecção de Plantas e de Fitocologia, Instituto Superior de Agronomia, Universidade Técnica de Lisboa, 1349-017 Lisboa, Portugal

⁷Department of Entomology, The Natural History Museum, Cromwell Road London SW7 5BD, UK

ΠΕΡΙΛΗΨΗ

Τα κοκκοειδή έντομα εμφανίζονται συχνά ως εισβολείς σε νέες περιοχές. Το είδος *Phenacoccus peruvianus* Granara de Willink, 2007, γνωστό και ως ψευδόκοκκος της βουκαμβίλιας παρατηρήθηκε για πρώτη φορά στην Ευρώπη το 1999 στην Ισπανία (Αλμερία) και αργότερα το 2002 στην Ιταλία (Σικελία). Αρχικά το είδος είχε αναγνωριστεί ως *Phenacoccus* sp. Καταγραφές του είδους έγιναν συχνότερες και σε άλλες περιοχές της Ισπανίας (συμπεριλαμβανομένων και των Βελεαρίδων νήσων), στη Μ. Βρετανία, τη Γαλλία (συμπεριλαμβανομένης και της Κορσικής), το Μονακό και την Πορτογαλία. Τα φυτά ξενιστές ήταν στις περισσότερες περιπτώσεις του γένους *Bougainvillea*.