ENTOMOLOGIA HELLENICA

Vol 7 (1989)

First records of the bayberry whitefly, *Parabemisia myricae* (Kuwana) in Greece

G. Michalopoulos

doi: [10.12681/eh.13968](https://doi.org/10.12681/eh.13968)

Copyright © 2017, G. Michalopoulos

This work is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike 4.0](https://creativecommons.org/licenses/by-nc-sa/4.0/).

To cite this article:

Michalopoulos, G. (1989). First records of the bayberry whitefly, *Parabemisia myricae* (Kuwana) in Greece. *ENTOMOLOGIA HELLENICA*, 7, 43–45. <https://doi.org/10.12681/eh.13968>

First Records of the Bayberry Whitefly, *Parabemisia myricae* (Kuwana) in Greece¹

G. MICHALOPOULOS

ICI HELLAS S.A., Syngrou 231,
171 21 N. Smirni, Athens, Greece

Citrus groves of Corfu had been attacked during the last decade by citrus whitefly *Dialeurodes citri* (Ashmead) (Pappas 1981). It presumably expanded later, and it was found to cause a serious infestation on lemon trees of Achaia region during summer of 1986, while it disappeared later existing today only in very small pockets on ornamental citrus species in Attica area.

Late in August 1988, the leaves of citrus trees, and to a lesser extent of grapevines, in the region of Corinthia, NE Peloponnesos, in South Greece were covered by larvae, pupae and adults of a whitefly that was different from *D. citri*. In autumn of the same year, it was noticed that the infestation of this new whitefly had expanded to other parts of Peloponnese such as Argos, Patras and Ilia region.

Samples of citrus and grapevine leaves with nymphs of the insect were sent for identification to British Museum of Natural History. Dr. J. H. Martin identified the species as *Parabemisia myricae* (Kuwana) (Homoptera: Aleyrodidae), making the first record of the species in Greece. The species is a native of Japan and possibly of other eastern asian countries such as Taiwan and West Malaysia (Rose et al. 1981). Today, the insect has been recorded in USA (California), Cyprus, Israel and Turkey (Martin 1987), Egypt (Martin personal communication) and it is a pest of woody plants, especially citrus trees and grapevines.

The damage caused to the plants by this insect, as a feeder, can be serious in cases of large populations accompanied with problems associated with the development of sooty mould on the excreted honeydew, as shown in

FIG. 1. Infestation and damage of a leaf on autumn flash of vegetation by bayberry whitefly nymphs that subsequently survived the winter.

Fig. 1. In the autumn of 1988, infestation was high (more than 20 nymphs/leaf) in Achaia and Corinthia regions of N. Peloponnese, causing great concern to citrus growers.

This species of whitefly was not included in the lists of known plant pests of Greece, Isaakides (1935, 1936, 1939), Pelekasis (1962), Stathopoulos et al. (1967), Anonymous (Volos, 1963), Buchelos and Soueref (1962), Buchelos et al. (1963, 1965), and Mourikis and Vassilaina-Alexopoulou (1975). It is probable that it invaded Greece from Israel.

P. myricae is a newly introduced pest in citrus growing countries and there is little information available on its biology and control. Most of the published data comes from Israel and California on the biology and biological control of the insect (Rose et al. 1981, De Bach and Rose 1982, Rose and De Bach 1982, Swirski et al. 1986, 1988). Additional data have been published on flight behaviour (Meyerdick and Moreno 1984), oviposition behaviour, and survival of young nymphs on leaves of different age citrus leaves (Walker

¹ Received for publication May 18, 1989.

FIG. 2. Oviposition of bayberry whitefly on young citrus leaf.

1985, 1987 and 1988, Walker and Aitken 1985).

According to Rose et al. (1981), the adults of *P. myricae* are smaller than *D. citri* adults and they have a "dusty blue gray or lavender appearance".

The females oviposit on very young leaves, often in circles (Fig. 2). Preference for oviposition on very young leaves (Walker 1987) seems to be related to properties of the leaf cuticle (Walker 1988). Leaf age is a critical factor for survival of nymphs (Walker and Aitken 1985). This gives a good explanation for the observed high rates of mortality in spring for nymphs growing on rapidly maturing leaves (Papasotiriou et al. 1989).

It has been observed that on orange trees less eggs are laid on the lower surface of the leaves, while on lemon trees they tend to be laid equally on both surfaces.

Initially eggs are off-white, while after about one day they gradually turn to brownish and finally black, before they hatch. First instar crawlers settle on the surface of the young leaves with a preference to the lower surface.

First flights of adults were noticed in this area in late February of 1989. By end of March early April the emergence of adults of the first 1989 generation had been completed.

It seems that it produces a quite large number of generations during the year, as it requires only 21 days at temperatures fluctuating between 21.0° C and 17.3 °C and 65-100 percent relative humidity, to complete its cycle in glasshouse (Rose et al. 1981). Being a pest of *Citrus* and grapevines (*Vitis vinifera*) it has also been found on hosts like *Ficus*, *Persea*, *Prunus*, *Psidium* and *Thea* (Martin 1987). As a matter of fact it has also been found in small colonies on *Prunus cerasi* in Northern Peloponnese in autumn 1988. Work on the control of *Parabemisia myricae* with the insect growth regulator P-p-rofezin has started in Greece in 1988 and is expected to be concluded in 1990.

Acknowledgment

I would like to express my thanks to Dr. Martin of British Museum (Nat. History) for the identification of the species on the samples from Greece. I am also grateful to Dr. L. C. Argyriou for her invaluable help in the preparation of the manuscript.

References

- Anonymous, 1963. Determination des ennemis animaux des plantes cultivées et conseils pour leur lute. Les acts de la periode 1954-1963. Station Phytopathologique de Volos.
- Buchelos, T. C. and S. T. Soueref. 1962. A list of the principal pests of crops recorded during the year 1961. Patras Plant Prot. Res. Sta., Ann. Rep. 3-9.
- Buchelos, T. C., S. T. Soueref and A. Tsoka. 1963. List of the principal pests of crops recorded during the year 1962. Patras Plant Prot. Res. Sta., Ann. Rep. 1963: 135 (in Greek).
- Buchelos, T. C., S. T. Soueref and A. Tsokas-Thanasouloupoulos. 1965. List of the principal crop pests recorded during the years 1963-1965. Patras Plant Prot. Res. Sta., Ann. Rep. 3-6.
- DeBach, P. and M. Rose. 1982. Biological control research on bayberry whitefly. Sunkist Pest Control Circ. No 528.
- Isaakides, C. A. 1935. List I des insectes et autres animaux nuisibles aux plantes cultivées et des insectes auxiliaires de la Grèce. Annls Inst. Phytopath. Benaki 1(2): 1-12.
- Isaakides, C. A. 1936. Liste II des insectes et autres animaux nuisibles aux plantes cultivées et des insectes auxiliaires de la Grèce. Annls Inst. Phytopath. Benaki 2(1): 5-7.
- Isaakides, C. A. 1939. List III des insectes et autres animaux nuisibles aux plantes cultivées et des insectes auxiliaires de la Grèce. Annls Inst. Phytopath. Benaki 3(1): 5-8.
- Martin, J. H. 1987. An identification guide to common

- whitefly pest species of the world (Homoptera: Aleyrodidae). Tropical Pest Management. 33: 298-322.
- Meyerdirk, D. E. and D. S. Moreno. 1984. Flight behavior and color-trap preference of *Parabemisia myricae* (Kuwana) (Homoptera: Aleyrodidae) in a citrus orchard. Environmental Entomology 13: 167-170.
- Mourikis, P. A. and P. Vassilaina-Alexopoulou. 1975. Report on the most important pests observed on cultivated plants in Greece from 1936 to 1966. Annls. Inst. Phytopath. Benaki (N.S.) 11: 141-150.
- Papasotiriou, K., G. Michalopoulos and E. Vlasserou. 1989. *Parabemisia myricae*, a new pest of citrus and vines in Greece, and first indications for its control. Proceedings of the 2nd Congress of the Greek Entomological Society (in press).
- Pappas, S. 1981. The presence of *Dialeurodes citri* (Ashmead) (Homoptera, Homoptera, Aleyrodidae) in Corfu. Ministry of Agriculture of Greece, Bulletin of Agricultural Research 5(2): 179-181.
- Pelekasis, C. E. D. 1962. List of the most important insects and other animals injurious to the cultivated plants in Greece, during the last thirty years. Annls Inst. Phytopath. Benaki (N.S.) 5: 5-104.
- Rose, M., P. DeBach and J. Wooley. 1981. Potential new citrus pest: Japanese bayberry whitefly. Calif. Agric. 35 (3-4): 22-24.
- Rose, M. and P. DeBach. 1982. A native parasite of the bayberry whitefly. Citograph 67: 272-276.
- Stathopoulos, D. G., I. A. Mentzelos and S. D. Savvidis. 1967. Survey of insects and other pests on crops of Macedonia and Thrace. II. Annu. Rept. Plant Prot. Agric. Res. Sta., Thessaloniki, Greece, 3 (1965): 102-106 (in Greek).
- Swirski, E., Y. Izhar, M. Wysoki and D. Blumberg. 1986. Overwintering of the Japanese bayberry whitefly, *Parabemisia myricae*, in Israel. Phytoparasitica 14: 281-286.
- Swirski, E., D. Blumberg, M. Wysoki and Y. Izhar. 1988. Phenology and biological control of the Japanese bayberry whitefly, *Parabemisia myricae*, on citrus in Israel. Proceedings of the Sixth International Citrus Congress: 1163-1167.
- Walker, G. P. 1985. Stylet penetration by the bayberry whitefly, as effected by leaf age in lemon, *Citrus limon*. Entomologia Experimentalis et Applicata 39: 115-121.
- Walker, G. P. and D. C. G. Aitken. 1985. Oviposition and survival of bayberry whitefly, *Parabemisia myricae* (Homoptera: Aleyrodidae) on lemon as a function of leaf age. Environmental Entomology 14: 254-257.
- Walker, G. P. 1987. Probing and oviposition behavior of the bayberry whitefly (Homoptera: Aleyrodidae) on young and mature lemon leaves. Ann. Entomol. Soc. Am. 80: 524-529.
- Walker, G. P. 1988. The role of leaf cuticle in leaf age preference by bayberry whitefly (Homoptera: Aleyrodidae) on lemon. Ann. Entomol. Soc. Am. 81 (2): 365-369.

KEY WORDS: Aleyrodidae, *Parabemisia myricae*, Bayberry whitefly, Citrus, Vines

Πρώτη Αναφορά στην Ελλάδα του Αλευρώδη των Εσπεριδοειδών *Parabemisia myricae* (Kuwana)

Γ. ΜΙΧΑΛΟΠΟΥΛΟΣ

ICI HELLAS S.A., Συγγρού 231,
171 21 Ν. Σμύρνη, Αθήνα

ΠΕΡΙΛΗΨΗ

Το καλοκαίρι του 1988 παρατηρήθηκε έξαρση της προσβολής των εσπεριδοειδών, και της σουλτανίνας σε μικρότερο βαθμό, από αλευρώδη. Η κύρια εστία εντοπίστηκε στην βορειοδυτική Πελοπόννησο. Στα δείγματα που λήφθηκαν από τις περιοχές Ξυλοκάστρου και Πύργου, διαπιστώθηκε η παρουσία του είδους *Parabemisia myricae* (Kuwana), για πρώτη φορά σε Ευρωπαϊκή χώρα. Ο αλευρώδης αυτός που ενδημεί σε γειτονικές χώρες της Βόρειας Αφρικής και της Μέσης Ανατολής προσβάλλει ξυλώδη φυτά προκαλώντας κυρίως έμμεσες ζημιές από τις οποίες η σημαντικότερη είναι η ποιοτική υποβάθμιση των καρπών από την ανάπτυξη μυκήτων πάνω στα εκκρίματά του.

Η βιολογία του έχει μελετηθεί σχετικά λίγο. Είναι γνωστό ότι τα θηλυκά γεννούν αποκλειστικά σε πολύ τρυφερή βλάστηση. Βρέθηκε ότι η γρήγορη ωρίμανση των φύλλων προκαλεί θνησιμότητα των ατελών μορφών. Αναφέρεται ότι ο βιολογικός του κύκλος στη διάρκεια του καλοκαιριού διαρκεί περί τις τρεις εβδομάδες.