


ENTOMOLOGIA HELLENICA

Vol 12 (1998)



First Record of the South American Leafminer *Liriomyza huidobrensis* (Blanchard) (Diptera: Agromyzidae) in Greece

N. E. Roditakis

doi: [10.12681/eh.14023](https://doi.org/10.12681/eh.14023)

Copyright © 2017, N. E. Roditakis



This work is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike 4.0](https://creativecommons.org/licenses/by-nc-sa/4.0/).

To cite this article:

Roditakis N. E. (1998). First Record of the South American Leafminer *Liriomyza huidobrensis* (Blanchard) (Diptera: Agromyzidae) in Greece. *ENTOMOLOGIA HELLENICA*, 12, 81–83. <https://doi.org/10.12681/eh.14023>

First Record of the South American Leafminer *Liriomyza huidobrensis* (Blanchard) (Diptera: Agromyzidae) in Greece

N. E. RODITAKIS

Plant Protection Institute of Heraklion,
71 110 Heraklion, Crete, Greece

In mid autumn 1992 the South American leafminer *Liriomyza huidobrensis* (Blanchard) was firstly recorded on greenhouse cucumbers, melons, beans and outdoor beans, broad beans and lettuce in the following locations of Crete: 1) Mires, Tymbaki, Antiskari at Messara valley of Southern Crete 2) Filiissia in the midland and 3) Platanias and Kokini Chani in Northern Crete. The following year *L. huidobrensis* had spread all over Crete (Ierapetra, Stomion, Koutsoura, Chania etc.) while melon and potato leaves which were sent to our laboratory from mainland Greece, Pirgos (Peloponnissos) and Chalkida (Evia island), were found heavily damaged by the same leafminer. The different types of mines (it usually mines the leaf alongside the veins), the colour of pupae (blackish) and the sudden attack of some plants (lettuce, broad beans, onions) which have never been damaged by the known local leafminers as yet, indicated that it was a newly introduced species. The new leafminer alerted the growers due to the high crop losses on outdoor and greenhouse vegetables (lettuce, broad beans, beans etc) despite the frequent application of registered insecticides. In the following year an extensive survey started to investigate its distribution and host plant in the greenhouses and outdoor vegetables and ornamentals in Crete.

L. huidobrensis, a quarantine insect (Anony-

mous 1980, 1992), is a polyphagous leafminer distributed in most biogeographical regions (Spencer 1973, 1990, Martinez 1993). It is native in South American countries from which it was spread soon in North America, Asia and Europe. In Europe it was first noticed in Holland, England and France in 1989 causing considerable damage to vegetables and ornamentals (Bartlett 1991, de Goffau 1991, Trouve et al., 1993). It mines the spongy mesophyll reducing the photosynthesis more than the other leafminers (Parella et al. 1985). This type of mines are not easily visible, unless the leaf is observed through transmitted sunlight or artificial light. This method was applied to ensure accurate detection of larvae and their parasitoids under a stereoscope. In our observations we found mines in leaves and cucumber fruits as well. Studies of its life cycle on greenhouse beans and melons revealed that most pupae (98%) remain on the bean leaves outside a hole in the autumn, while a few fall on the ground. The number of pupae collected from ten leaves per 24h were: 5.7 ± 1.6 on lettuce, 17.07 ± 4.1 on beans, 24.2 ± 7.3 on melon, 6.0 ± 2 on broad bean leaves. During the survey the following host plants were recorded:

- Solanaceae: *Lycopersicum esculentum*,
Solanum nigrum, *Solanum melongena*, *Solanum tuberosum*, *Capsicum annuum*
- Chenopodiaceae: *Spinacia oleracea*
- Cruciferae: *Brassica oleracea*
- Scrofulariaceae: *Antirrhinum sp.*
- Cucurbitaceae: *Cucumis sativus*,
Cucumis melo, *Cucumis pepo*
- Compositae: *Lactuca sativa*, *Cerbera sp.*,
Cynara scolymus,
Chichorium sp.
- Papilionaceae: *Vicia faba*, *Phaseolus vulgaris*,
Trifolium pratense
- Malvaceae: *Malva sp.*
- Liliaceae: *Allium cepa*
- Leguminosae: *Melilotus indica*
- Caryophyllaceae: *Dianthus caryophyllus*

Chemical control could be effective by applying certain insecticides (abamectin, triazophos, imidacloprid, heptenophos etc.) while biological control seems to be rather effective by the known parasitoids, which are already used against the

tomato leafminer *Liriomyza bryoniae* Kalt (Cheek et al. 1993, Linden 1993). Linden (1993) found that biological control of *L. huidobrensis* was effective on lettuce by means of repeated releases of *Dacnusa sibirica* Telenga and *Diglyphus isaea* (Walker) but so far native parasitoids proved to be able to control the pest. Mass trapping with coloured sticky traps seems to be also a potential method in IPM programmes (Bartlett 1991). The mortality of pupae collected from heavily treated greenhouse plants with insecticides varied from 65 to 74% while that of untreated plants was between 18-25%. No pupal parasitoids were recorded but *D. isaea* and *D. sibirica* were both recorded as larval parasitoids. *D. isaea* was abundant all over the year while *D. sibirica* only in spring-summer period. These two parasitoids were able, in certain cases, to control sufficiently the leaf miner in untreated greenhouse cucumber and bean plants. The neem seed extract (*Azadirachta indica*) applied on outdoor tomatoes in Antiskari (Southern Crete) proved harmless both on hymenoptera and on the mirid predator *Macrolophus caliginosus* Wagner. Despite its weak larvicide action on *L. huidobrensis* it was very effective in conjunction with parasitoids consisting of a promising candidate in IPM programmes (Roditakis and Roditakis 1994).

Aknowledgments

Many thanks to: a) Dr. D. Bordat (CIRAD-CA) who assisted my contact with Dr. M. Martinez, b) Dr. Martinez for the identification of species (INRA-Station de Zoologie, Laboratoire de Biosystematique de l'Insecte, Versailles), and c) Dr. E. Tzortzakakis for useful suggestions and corrections of the manuscript.

References

- Anonymous, 1980. The import and export order 1980. Statutory instrument 1980 No 420 HMSO London.
- Anonymous, 1992. Commission Directive 92/103/EEC Official Journal of the European Community No 1 363/1-65.
- Bartlett, P.W. 1991. The forceout and starve strategy. The Grower 115 : 9-10.
- Cheek, S., O. K. Macdonald and P. W. Bartlett, 1993. Statutory action against *Liriomyza huidobrensis* (Blanchard) in the United Kingdom. «Liriomyza» Conference on leaf mining flies in cultivated plants. Montpellier, France (24-26 March 1993).
- De Goffau, L. J. W. 1991. *Liriomyza huidobrensis* (Diptera: Agromyzidae) a new economically important leaf miner in the Netherlands. Proceedings Exp. & Appl. Entom. N.E.V. Amsterdam 2 : 41-45.
- Linden, A. van der. 1993. Biological control of leafminers in glasshouse lettuce. «Liriomyza» conference on leaf

mining flies in cultivated plants. Montpellier, France (24-26 March 1993).

- Martinez, M. 1993. Liste des 'especies de *Liriomyza* d'importance agronomique, leur synonymes et leur regions biogeographiques. «Liriomyza» Conference on leaf-mining flies in cultivated plants. Montpellier, France (24-26 March 1993): 79-86.
- Parrella, M.P., V.P. Jones, R.R. Youngman & L.M. Lebeck. 1985. Effect of leafmining and leaf stippling of *Liriomyza* spp. on photosynthetic rates of chrysanthemum. Ann. Entom. Soc. America 78 : 90-93.
- Roditakis, N.E. and E.N. Roditakis. 1994. Effect of neem seed extract (Neemark) on South American leafminer *Liriomyza huidobrensis* (Blanchard) (Diptera: Agromyzidae). Proc. of the Meeting «Integrated control in protected crops. Mediterranean Climate», IOBC/WPRS Bul. 17 Lisboa, Portugal (21-23 September 1994).
- Spencer, K.A. 1973. Agromyzidae (Diptera) of Economic Importance Dr. W. Junk B.V. The Hague. Series Entomologica 418 pp.
- Spencer, K.A. 1990. Host specialization in the world Agromyzidae (Diptera) Kluwer Academic Publish. (Dordrecht/London). Series Entomologica 45: 1-444.
- Trouve, C., M. Phalip and M. Martinez. 1993. Situation en France de *Liriomyza huidobrensis* (Blanchard). «Liriomyza» Conference on leafmining flies in cultivated plants Montpellier, France (24-26 March 1993): 49-57.

KEY WORDS: Agromyzidae, *Liriomyza huidobrensis*, South American leafminer, vegetables, ornamentals, greenhouse plants, hosts, damage, Greece, Crete

**Πρώτη Διαπίστωση του
Υπονομευτή της Ν. Αμερικής
Liriomyza huidobrensis
(Blanchard) (Diptera:
Agromyzidae) στην Ελλάδα**

N. E. ΡΟΔΙΤΑΚΗΣ

*Ινστιτούτο Προστασίας Φυτών Ηρακλείου,
71 110 Ηράκλειο Κρήτης*

ΠΕΡΙΛΗΨΗ

Ο υπονομευτής φύλλων της Ν. Αμερικής *Liriomyza huidobrensis* (Blanchard) επισημάνθηκε για πρώτη φορά το 1992 σε θερμοκηπιακές και υπαίθριες καλλιέργειες κηπευτικών και ανθέων στη Κρήτη και σε δείγμα φύλλων πεπο-

νιάς από τον Πύργο (Πελοπόννησος) και φύλλων πατάτας από τη Χαλκίδα (Εύβοια). Το έντομο αυτό προκάλεσε σοβαρές ζημιές στις φασολιές, κουκιά, μαρούλια, πεπόνια και αγγουριές ενώ προσέβαλε επίσης σε μικρότερο βαθμό πατατόφυτα, κρεμμύδια, αγγινάρες και σπανάκι. Σήμερα αποτελεί το επικρατέστερο είδος υπονομευτή και εξαπλώθηκε σε όλη την Κρήτη. Αντιμετωπίζεται δύσκολα με τα συνήθως χρησιμοποιούμενα εντομοκτόνα για τους άλλους υπονομευτές. Η αντιμετώπισή του μπορεί να γίνει μόνο από ένα πολύ περιορισμένο αριθμό εντομοκτόνων (abamectin, triazophos, imidacloprid, heptenophos κλπ.). Τα υμενόπτερα παράσιτα *Diglyphus isaea* (Walter) και *Dacnusa sibirica* Telenga υπάρχουν ως ιθαγενή στην πανίδα της Κρήτης και ελέγχουν αποτελεσματικά τον νέο αυτό υπονομευτή.