


ENTOMOLOGIA HELLENICA

Vol 31, No 2 (2022)

Entomologia Hellenica 31(2)



First occurrence of the black field earwig, *Nala lividipes* (Dermaptera: Labiduridae) in Greece

Evangelos Koutsoukos, Jakovos Demetriou, Konstantinos Kalaentzis, Christos Kazilas

doi: [10.12681/eh.30836](https://doi.org/10.12681/eh.30836)

Copyright © 2022, Evangelos Koutsoukos, Jakovos Demetriou, Konstantinos Kalaentzis, Christos Kazilas



This work is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike 4.0](https://creativecommons.org/licenses/by-nc-sa/4.0/).

To cite this article:

Koutsoukos, E., Demetriou, J., Kalaentzis, K., & Kazilas, C. (2022). First occurrence of the black field earwig, *Nala lividipes* (Dermaptera: Labiduridae) in Greece. *ENTOMOLOGIA HELLENICA*, 31(2). <https://doi.org/10.12681/eh.30836>



First occurrence of the black field earwig, *Nala lividipes* (Dermaptera: Labiduridae) in Greece

E. KOUTSOUKOS^{1,2,*}, J. DEMETRIOU³, C. KAZILAS⁴ AND K. KALAENTZIS^{4,5}

¹Section of Ecology and Systematics, Department of Biology, National and Kapodistrian University of Athens, 15784 Athens, Greece. E-mail: vag18000@gmail.com

²Museum of Zoology, National and Kapodistrian University of Athens, 15784 Athens, Greece

³Laboratory of Vector Ecology and Applied Entomology, Joint Services Health Unit, British Forces Post Office 57, Royal Air Force Akrotiri, Cyprus.

⁴Naturalis Biodiversity Center, PO Box 9517, 2300 RA, Leiden, The Netherlands.

⁵Hydrobiological Station of Rhodes, Hellenic Centre for Marine Research, Cos Street, 85100 Rhodes, Greece.

ABSTRACT

The cosmopolitan black field earwig, *Nala lividipes* (Dufour, 1820) (Dermaptera: Labiduridae) is observed for the first time in Greece. A photographic record was obtained by the first author in 2016, from Salamis Island followed by a photographic citizen-science observation from Rhodes in 2021. Despite frequent inspections and efforts to raise public awareness, no additional samples or observations have been recovered. The poor study of Dermaptera in Greece and the absence of early warning systems for alien insects, may have delayed the detection of this alien species in the country. However, given the agricultural significance of *N. lividipes*, a short description is given to assist further data collection and monitoring. Finally, the potential establishment of the earwig in Greece, along with the impacts on the native flora and fauna are discussed.

KEY WORDS: Agricultural pest, alien species, cosmopolitan, early warning systems.

Introduction

Globalization and the constant rise of international trade have led to an unprecedented rise of species translocations and a great distortion of their native ranges (Hulme 2009). Many of these alien species have been associated with socioeconomic, human health and environmental impacts (Simberloff et al. 2013; Schindler et al. 2015; Vilà and Hulme 2016), something that highlights the need to monitor their occurrence and dynamic shifts in distribution ranges.

The reported cases of introductions and translocation of earwigs (Dermaptera) are rather uncommon, supported by only seven alien earwig species being recorded in Europe (Zafeiriou et al. 2021). The small number of reported alien Dermaptera could be related to both their low species richness as well as their cryptic nature. Until now, only two alien earwigs have been recorded from Greece, namely *Euborellia femoralis* (Dohrn, 1863) and *Forficula lucasi* Dohrn, 1865 raising the Greek earwig fauna to 18 species (Haas 2015, 2018; Demetriou et al.

*Corresponding author: vag18000@gmail.com

2021; Fontana et al. 2021; Zafeiriou et al. 2021; Kalaentzis et al. 2022).

The black field earwig, *Nala lividipes* (Dufour, 1820) is a cosmopolitan species, presumably of Afrotropical origin, which has gradually spread across all biogeographic realms. The species is widely distributed along the Mediterranean basin with records from Algeria, France, Italy (including Sardinia and Sicily), Malta, Morocco, Portugal, Spain (including the Balearic and Canary Islands), Tunisia, and Turkey (Albouy and Caussanel 1990; Mifsud and Taglianti 2008; Rasplus and Roques 2010; Anlaş and Kočárek 2012; Pages 2012) (Fig. 1).

The alien insect-fauna of Greece has been recently updated, holding a total of 469 species (Demetriou et al. 2021) in

anticipation of further additions. Citizen scientists have been supplementing our knowledge around the presence and distribution of alien species, providing photographic observations and specimens of new alien to the country insects such as the feather-legged fly *Trichopoda pictipennis* Bigot, 1876 and lantana plum moth *Lantanophaga pusillidactylus* (Walker, 1864) (Demetriou et al. 2020; Kazilas et al. 2020; Dios et al. 2021). Alientoma, an online database for the alien insects of Greece, promoting public participation in scientific research (Kalaentzis et al. 2021b, c), was utilised to track the presence of *N. lividipes* in the country. In this publication, the black field earwig *N. lividipes* is reported for Greece for the first time.



FIG. 1: Known distribution of *Nala lividipes* (Dufour, 1820) in the Mediterranean basin. Countries where the species has been previously reported are shaded red, while the new records from Greece are depicted with a dot. The map was created with QGIS, v.3.14.16. Inset: The adult, female individual observed on Salamis island, Greece.

Materials and Methods

A female individual was observed and photographed in Greece: Saronic Gulf, Salamis island, Paloukia [37.962°N, 23.514°E], vii/2016, alt. 25 m, observed by E. Koutsoukos, in a house garden (Fig. 2A). The observed female was identified as *N. lividipes* following the identification key of Anlaş and Kočárek (2012). The species identification was also confirmed by taxonomic expert (Dr. Petr Kočárek, pers. comm., 2021). A second observation of a

female *N. lividipes* was recorded in Greece: Dodecanese, Rhodes island, [36.3848°N, 28.1312°E], 23/vii/2021, alt. 50 m, observed by Mr. Benoît Segerer in a hotel garden, uploaded to iNaturalist platform (iNaturalist 2021) (Fig. 2B).

Online educational material regarding the taxonomy, distribution, morphology, ecology and impacts of *N. lividipes* was uploaded to the Alientoma webpage and social media platform (Kalaentzis et al. 2021a), in order to investigate the presence of further photographic observations.




FIG. 2: Adult, female individual of *Nala lividipes* (Dufour, 1820) observed and photographed by Evangelos Koutsoukos on Salamis (A) and Mr. Benoît Segerer on Rhodes island (B).

Results and Discussion

These are the first reported occurrences of the black field earwig in Greece, which comprises the third alien species of the order Dermaptera in the country. Taking into consideration its broad distribution along the Mediterranean basin, its report in

Greece comes as no surprise, although, the species is considered rather rare in Europe (Rasplus and Roques 2010). Despite of further inspections on the observation sites as well as the educational material uploaded to Alientoma, no additional material has been recovered. Consequentially, any speculations regarding the presence of established populations in the country seem

to be rather uncertain. However, the long distance between the two observation sites may indicate that the species has long been present in the country, probably remaining unnoticed due to the insufficient study and public interest in Dermaptera on a national level.

A short diagnosis of the species is provided to assist further data collection by citizen-scientists and researchers: Head and thorax dark brown, antennae with >20 segments and wings fully developed. Tegmina and wing scales brown, opaque, rugose; tegmina with a ridge along the costal margin. Abdomen darker, black with sparse pale pubescence. Overall body length with forceps <15 mm (8.5-11 mm) (Fig. 2). Female forceps slender and almost straight; male forceps curved holding small teeth basally and a larger tooth towards their apex (Mifsud and Taglianti 2008; Anlaş and Kočárek 2012).

The black field earwig is a polyphagous insect feeding mostly on decaying organic matter or plant material (i.e. seeds, stems, roots), while it has also been reported as carnivorous (Albouy and Caussanel 1990; Simpson 1993). On one hand, it is often described as a crop pest, affecting beetroot, cotton, maize, sorghum, soybean, sunflower, and winter cereal cultivations, particularly when high population density is

reached (Hargreaves 1970; Radford and Allsopp 1987; Simpson 1993). On the other hand, *N. lividipes* has also been utilised in classical biological control programs, preying on crop-damaging pests (i.e. aphids and moths) (Albouy and Caussanel 1990; Binns et al. 2020). For example, in Australia *N. lividipes* is acknowledged as the main predator of *Pieris rapae* (Linnaeus, 1758) (Kapuge et al. 1987), which is a serious, accidentally introduced pest of crucifers (CABI 2021). To our best knowledge, no damages have been reported by the black field earwig in Europe. However, given the socioeconomic effects associated with the species in question, it is important to monitor its distribution and feeding habits in the invaded territories, as well as to assess its diverse effects on agriculture and native fauna.

In conclusion, this article renders *N. lividipes* the third alien earwig reported from Greece, raising the number of its Dermaptera fauna to 19 species. This manuscript is intended to alert the scientific community and the public of the potential threats of a new alien insect in Greece, using citizen science towards the implementation of early warning systems for alien and invasive alien species.

Acknowledgements

We would like to wholeheartedly thank Mr. Benoît Segerer for his provision of photographic material and data of *N. lividipes* from Rhodes island. We would also like to thank Dr. Petr Kočárek (University of Ostrava, Czechia) for verifying the identification of the observed individual and his revision of the manuscript. We are also grateful to Dr. Leonid N. Anisyutkin (Zoological Institute of Russian Academy of Sciences) and Dr. Dimitrios N. Avtzis (Forest Research Institute, Hellenic Agricultural Organization Demeter) for their valuable corrections to the manuscript as well as to the two anonymous reviewers for their comments and corrections during the review process.

References

- Albouy, V. and C. Caussanel. 1990. *Dermatères ou Perce-oreilles*. Faune de France 75. Fédération Française des Sociétés de Sciences Naturelles, Paris, 245 pp.
- Anlaş, S. and P. Kočárek. 2012. Current status of Dermaptera (Insecta) fauna of Turkey and Cyprus. *Türkiye Entomoloji Dergisi* 36(1): 43-58.
- Binns, M., A. Hoffmann, M. van Helden, T. Heddle, L. Kirkland and P. Umina. 2020. Earwigs - an appetite for destruction or are they beneficial? <https://grdc.com.au/resources-and-publications/grdc-update-papers/tab-content/grdc-update-papers/2020/02/earwigs-an-appetite-for-destruction-or-are-they-beneficial>; accessed 15 August 2021
- CABI 2021. Invasive Species Compendium. *Pieris rapae* (cabbage white butterfly). <https://www.cabi.org/isc/datasheet/41167>; accessed 15 August 2021
- Demetriou, J., C. Kazilas, E. Koutsoukos and K. Kalaentzis. 2020. A new tropical invader in Greece: The lantana plume moth *Lantanophaga pusillidactylus* (Lepidoptera: Pterophoridae). *Entomologia Hellenica* 29: 1-7. <https://doi.org/10.12681/eh.21907>
- Demetriou, J., K. Kalaentzis, C. Kazilas, E. Koutsoukos, D. N. Avtzis and C. Georgiadis. 2021. Revisiting the non-native insect fauna of Greece: Current trends and an updated checklist. *NeoBiota* 65: 93-108. <https://doi.org/10.3897/neobiota.65.64686>
- Dios R. V. P., J. Ziegler and T. Zeegers. 2021. The American genus *Trichopoda* (Diptera: Tachinidae) in Europe – decades of a misidentified invasive species. *Contributions to Entomology* 71(2): 221-225. <http://dx.doi.org/10.21248/contrib.ento.mol.71.2.221-225>
- Fontana, P., F. Marangoni, P. Kočárek, P. Tirello, G. Giovagnoli and L. Colacurcio. 2021. Updated knowledge on Italian Dermaptera with the report of a new alien species: *Forficula smyrnensis* Audinet-Serville, 1838. *Memorie della Società Entomologica Italiana* 97: 261–270. <https://doi.org/10.4081/memoriesei.2020.261>
- Haas, F. 2015. Earwig Research Centre – ERC – The site on earwig biology. <http://www.earwigs-online.de/GR/gr.html>; accessed 15 August 2021
- Haas, F. 2018. Biodiversity of Dermaptera. In: Footitt, R.G. and P.H. Adler (Eds). *Insect Biodiversity: Science and Society*. Volume II. John Wiley & Sons Ltd., Hoboken. pp. 315-334.
- Hargreaves, J.R. 1970. The black field earwig. *Queensland Agricultural Journal* 96: 391-392.
- Hulme, P.E. 2009. Trade, transport and trouble: Managing invasive species pathways in an era of globalization. *Journal of Applied Ecology* 46: 10-18. <https://doi.org/10.1111/j.13652664.2008.01600.x>
- iNaturalist 2021. iNaturalist - A Community for Naturalists. <https://www.inaturalist.org>; accessed 31 August 2021
- Kalaentzis, K., C. Kazilas, J. Demetriou, E. Koutsoukos, C. Georgiadis and D.N. Avtzis. 2021a. *Alientoma - Nala lividipes*. <https://alientoma.myspecies.info/el>; accessed 15 August 2021
- Kalaentzis, K., C. Kazilas, J. Demetriou, E. Koutsoukos, C. Georgiadis and D.N. Avtzis. 2021b. Engaging citizen-scientists in mapping alien species: Introducing Alientoma, a dynamic database for alien insects in Greece. The 1st International Electronic Conference on Entomology (IECE). DOI: 10.3390/IECE-10512 (poster presentation).

- Kalaentzis, K., C. Kazilas, J. Demetriou, E. Koutsoukos, C. Georgiadis and D.N. Avtzis. 2021c. AlienToma, a Dynamic Database for Alien Insects in Greece and Its Use by Citizen Scientists in Mapping Alien Species. *Insects* 12: 1101. <https://doi.org/10.3390/insects12121101>
- Kalaentzis, K., C. Kazilas, G. Agapakis and P. Kočárek. 2022. Hidden in plain sight: First record of the alien earwig *Euborellia femoralis* (Dohrn, 1863) in Europe. *BioInvasions Records* 10(4): 1022-1031. <https://doi.org/10.3391/bir.2021.10.4.27>
- Kapuge, S.H., W. Danthanarayana and N. Hoogenraad. 1987. Immunological investigation of prey-predator relationships for *Pieris rapae* (L.) (Lepidoptera: Pieridae). *Bulletin of Entomological Research* 77 (2): 247-254. <https://doi.org/10.1017/S000748530001172X>
- Kazilas, C., J. Demetriou and K. Kalaentzis. 2020. Filling the gaps in the distribution of an alien species: The case of the feather-legged fly *Trichopoda pennipes* (Diptera: Tachinidae) in the Western Palearctic. *Entomologia Hellenica* 29(1): 8-16. <https://doi.org/10.12681/eh.21774>
- Mifsud, D. and V. Taglianti. 2008. *Nala lividipes* (Dufour, 1828), a new earwig for the Maltese Islands (Dermaptera: Labiduridae). *Bulletin of the Entomological Society of Malta* 1: 11-13.
- Pages, J. 2012. Présence confirmée de *Nala lividipes* (Dufour, 1828) en France (Dermaptera: Labiduridae). *Revue de l'Association roussillonnaise d'Entomologie* 21: 41–42.
- Radford, B.J. and P.G. Allsopp. 1987. Use of insecticides and a press wheel to control soil insects affecting sorghum and sunflower establishment in South Queensland. *Journal of the Australian Entomological Society* 26: 161-167. <https://doi.org/10.1111/j.1440-6055.1987.tb00279.x>
- Rasplus, J.Y. and A. Roques. 2010. Dictyoptera (Blattodea, Isoptera), Orthoptera, Phasmatodea and Dermaptera. *BioRisk* 4 (2): 807-831. <https://doi.org/10.3897/biorisk.4.68>
- Schindler, S., B. Staska, M. Adam, W. Rabitsch and F. Essl. 2015. Alien species and public health impacts in Europe: A literature review. *NeoBiota* 27: 1-23. <https://doi.org/10.3897/neobiota.27.5007>
- Simberloff, D., J.L. Martin, P. Genovesi, V. Maris, D.A. Wardle, J. Aronson, F. Courchamp, B. Galil, E. García-Berthou, M. Pascal, P. Pyšek, R. Sousa, E. Tabacchi and M. Vilà. 2013. Impacts of biological invasions: What's what and the way forward? *Trends in Ecology and Evolution* 28: 58-66. <https://doi.org/10.1016/j.tree.2012.07.013>
- Simpson, G.B. 1993. Effects of temperature on the development, longevity and fecundity of *Nala lividipes* (Dufour) (Dermaptera: Labiduridae). *Journal of the Australian Entomological Society* 32: 265-272. <https://doi.org/10.1111/j.1440-6055.1993.tb00585.x>
- Vilà, M. and P.E. Hulme. 2016. Impact of biological invasions on ecosystem services. Springer, Cham. 354 pp. <https://doi.org/10.1007/978-3-319-45121-3>
- Zafeiriou, S., P. Kočárek and K. Kalaentzis. 2021. First record of the desert earwig *Forficula lucasi* Dohrn, 1865 (Dermaptera: Forficulidae) in Greece: A hitchhiker among the refugees or a seldom encounter? *Journal of Insect Biodiversity* 21: 15-17. <https://doi.org/10.12976/jib/2021.21.1.2>

Πρώτες παρατηρήσεις του Δερμαπτέρου *Nala lividipes* στην Ελλάδα

Ε. ΚΟΥΤΣΟΥΚΟΣ^{1,2,*}, Ι. ΔΗΜΗΤΡΙΟΥ³, Χ. ΚΑΖΙΑΛΑΣ⁴ ΚΑΙ Κ. ΚΑΛΑΕΝΤΖΗΣ^{4,5}

¹Τομέας Οικολογίας και Συστηματικής, Τμήμα Βιολογίας, Εθνικό και Καποδιστριακό Πανεπιστήμιο Αθηνών, 15784 Αθήνα, Ελλάδα. E-mail: vag18000@gmail.com

²Μουσείο Ζωολογίας, Εθνικό και Καποδιστριακό Πανεπιστήμιο Αθηνών, 15784 Αθήνα, Ελλάδα.

³Laboratory of Vector Ecology and Applied Entomology, Joint Services Health Unit, British Forces Post Office 57, Royal Air Force Akrotiri, Cyprus.

⁴Naturalis Biodiversity Center, PO Box 9517, 2300 RA, Leiden, The Netherlands.

⁵Υδροβιολογικός σταθμός Ρόδου, Ελληνικό Κέντρο Θαλάσσιων Ερευνών, οδός ΚΩ, 85100 Ρόδος, Ελλάδα.

ΠΕΡΙΛΗΨΗ

Το κοσμοπολίτικο δερμάπτερο, *Nala lividipes* (Dufour, 1820) (Dermaptera: Labiduridae) παρατηρείται για πρώτη φορά στην Ελλάδα. Το είδος φωτογραφήθηκε από τον πρώτο συγγραφέα το 2016, από το νησί της Σαλαμίνας ακολουθούμενο από μια φωτογραφική παρατήρηση ενός πολίτη-επιστήμονα από τη Ρόδο, το 2021. Συχνές επιθεωρήσεις στον τόπο παρατήρησης και προσπάθειες ευαισθητοποίησης του κοινού, δεν έχουν μέχρι στιγμής παράσχει επιπλέον υλικό. Η ελλιπής μελέτη των δερμάπττερων στην Ελλάδα και η απουσία συστημάτων έγκαιρης προειδοποίησης για τα ξενικά έντομα, πιθανώς να καθυστέρησαν την παρατήρηση του ξενικού αυτού είδους στην χώρα. Παρόλα αυτά, δεδομένης της γεωργικής σημασίας του είδους, δίνεται μια σύντομη περιγραφή του προς υποβοήθηση της περαιτέρω συλλογής δεδομένων αλλά και της παρακολούθησης του. Τέλος, η πιθανή εδραίωση του δερμάπττερου στην Ελλάδα, καθώς και οι επιπτώσεις στην αυτόχθονη χλωρίδα και πανίδα, είναι υπό διερεύνηση.