



ENTOMOLOGIA HELLENICA

Vol 33, No 2 (2024)

Entomologia hellenica 33(2)



To cite this article:

ElSayed, W., Kondoh, M., Hori, M., Nagasawa, A., Nakamura, K., & Shahenda, A. E. (2024). Feeding Categorisation of some Caeliferan and Ensiferan species (Insecta: Orthoptera) collected from selected grasslands in Sendai City, Japan. *ENTOMOLOGIA HELLENICA*, *33*(2), 90–109. Retrieved from https://ejournals.epublishing.ekt.gr/index.php/entsoc/article/view/37715

Received 08 May 2024

Accepted 10 November 2024

Available online 06 December 2024

Feeding Categorisation of some Caeliferan and Ensiferan species (Insecta: Orthoptera) collected from selected grasslands in Sendai City, Japan

Abu ElEla Shahenda, A. A.^{1, 3}, Michio Kondoh², Masatoshi Hori³, Atsuhiko Nagasawa³, Koji Nakamura⁴, Wael M. ElSayed¹

¹Department of Entomology, Faculty of Science, Cairo University, Egypt 12613 ²Graduate School of Life Sciences, Tohoku University, 6-3 Aoba, Aramaki, Aoba-ku, Sendai 980-8578, Japan

³Graduate School of Agricultural Science, Tohoku University, Sendai, Miyagi 980-8572, Japan ⁴Ishikawa Prefectural Natural History Museum, Kanazawa, Ishikawa 920-1147, Japan

ABSTRACT

Although orthopteran species have often been regarded as polyphagous herbivores, most of them show variable degrees of diet selectivity and particular food preferences. Still some species possess peculiar feeding categories which need further investigation. An assemblage of 43 orthopteran species, 7 families, 12 subfamilies and 13 tribes was surveyed from five grasslands in Sendai City, Japan to investigate their feeding preferences and were classified into seven main feeding categories based on examination of morpho-mandibular characteristics and postmortem gut content analyses. The forbivorous category was the most dominant, whereas the herbivorous the least common.

KEY WORDS: Orthoptera, mandibular characteristics, gut contents analysis, feeding categories.

Introduction

Orthopterans have emerged as a crucial group of invertebrates for environmental monitoring and assessment. The Orthoptera order, comprising a diverse array of species, holds a prominent position amongst insect orders (Zhang 2011; Bidau, 2014). With approximately 28,000 identified species worldwide and ranked as the sixth largest order in Class Insecta, trailing only the Hemiptera order as declared by Cigliano et al. (2022). Species belonging to Orthoptera are present in every terrestrial environment and exhibit remarkable diversity (ElEla et al., 2010; Yadav and Kumar 2017).

Taxonomists classified Orthoptera into two distinct suborders: Ensifera, encompassing crickets, katydids, and their relatives, and Caelifera, which include grasshoppers and their allies (Song, 2018). Orthopteran species play considerably vital roles in terrestrial food webs, serving as a valuable protein source for various animals, including amphibians, birds, small reptilian species, as well as some mammalian species. Consequently, their scarcity could disrupt the delicate trophic structure within an ecosystem (Soliman et al., 2017). They play a central role in food webs, as they are mostly primary herbivores and constitute an abundant food resource for other groups of carnivorous organisms (Parr and Chown, 2003). The composition of orthopteran assemblage is considerably highly responsive to environmental changes, play a crucial role in the ecosystems function and can serve as a valuable environmental

indicator (O'Neill et al., 2003). Indeed, these species potentially serve as useful bioindicators for land disturbance (Saha et al. 2011) providing valuable insights into qualities of ecosystems and the effectiveness of ecological networks (Zhang et al., 2019).

Orthopteran species have attracted significant attention due to the extensive damage they inflict by their herbivory action on crops and various forms of green vegetation (Dakhel et al., 2020). Aspects of feeding strategies, including selection of food, preference of food, diverse feeding patterns and diet specialization have been studied by many researchers for diverse orthopteran species (e.g., Isely, 1944; Williams, 1954; Mulkern, 1967; Otte and Joern, 1976; Bernays and Chapman, 1978; Uvarov, 1977; Ohabuike, 1979, Joern; 1983, 1985; Bernavs and Bright, 1993, ElSayed, 2005; ElShazly and ElSayed, 2006; ElEla et al., 2010, 2012; Kuřavová and Kočárek, 2016; ElSaved et al., 2020).

Adaptation to food intake has led to behavioral and morphological specialization of mouthparts (Snodgrass, 1935). Frv et al. (1978) found that caeliferan species. especially Acridomorpha, are often phytophagous, and the adaptations of their mandibular structures are associated with different types of food and different species of ecological groups of plants, grasses, forbs, flowers, and seeds (Isely, 1944; Patterson, 1983, 1984; Kang and Chen, 1994; Bernays, 1998; Gangwere et al. 1998; ElEla, 2011; ElEla et al. 2010, 2012; Di Russo et al. 2014; Kuřavová and Kočárek, 2016; ElSayed et al. 2020).

Considering the paucity of information on the assemblage of orthopteran species inhabiting the five selected grasslands in Sendai city, especially on the variation of mandibles in accordance with food preference, our team was stimulated to examine the morpho- mandibular variations of orthopteran species and evaluate the gut contents analysis as evidence supporting these morpho-adaptations. Moreover, there is a lack of information regarding the diet of tetrigid species which are relatively among the least-studied groups of Orthoptera (Flook and Rowell, 1997; Hochkirch et al. 2000 & 2006) especially those recorded in some parts of Asia (Song et al. 2015) and more specifically from Sendai City, Japan (ElEla et al. 2010).

Materials and Methods

Climate and topography. The study was carried out in Sendai City, Japan (lat. $38^{\circ}16'05''$ N, long. $140^{\circ}52'11''$ E). The area of the city is ca. 788.09 km² and stretches from the Pacific Ocean to the Ōu Mountains, which are the east and west borders of Miyagi Prefecture. As a result, the geography of the city is quite diverse. Eastern Sendai is a plain, the center of the city is hilly, and western areas are mountainous.

According to the Japan Meteorological Agency (<u>http://www.data.jma.go.jp</u>), meteorological records showed that Sendai City has a humid subtropical climate, which features warm and wet summers, and cool and dry winters. The summer season starts in June and ends at the end of September with significant seasonal variations in temperature and rainfall. Summers are warm, with an August average temperature of 24.1 °C and the majority of the annual precipitation is delivered during summer.

The city is rarely hit by typhoons, and experiences only 6 days with more than 10 centimeters of precipitation on average.

Monsoon season usually begins in late April to early October, which is later than in most cities in Japan. During this season, cold winds from the Okhotsk air mass, which are called "Yamase", blow in and depress daytime highs.

Extremes of temperature degrees range from -11.7 to 37.2 °C and Sendai City experiences fewer days of extreme temperatures (highs outside 0 - 30 °C at 19.6 days/year compared to Tokyo's average of 49 days/year.

Winters are cool and relatively dry, with the January temperature averaging at

1.5 °C. Concerning snow, snowfalls are much lower than at the cities on the Sea of Japan coast, such as Niigata and Tottori (Climate data for Sendai normals, extremes [1926–present], Japan Meteorological Agency, <u>http://www.data.jma.go.jp</u>).



FIG. 1.: Map of the study site: (*i*) Map of Japan showing Miyagi Prefecture and Sendai City, Japan, (*ii*) Map of Sanjo-machi with the grassland plots (*iii*) aerial photograph showing the five grassland plots within Sanjo-machi (marked by white capital letters). (Source: Tohoku University https://www.tohoku.ac.jp/english/profile/campus/01/access/)

Study site. The study site comprised of five main plots of grasslands (Fig. 1). Grassland A and B were longitudinal, L-shaped transects, covered by short and long grasses, with a presence of several shrubs. Grassland C was adjacent to a parking lot. It was a longitudinal quadrate plot covered by short grasses, with long grasses on the edges. Grassland D was a T-shaped plot, with a canopy almost dominated by relatively short grasses. Grassland E was almost trapezoid with very few trees scattered all over (Fig. 1). The canopy was composed of mixed grasses and diverse flowering plant species. A small bond was located at the northern edge and was surrounded by long grasses.

In general, on analyzing the selected sampling sites for their floral composition, it was found comparatively rich with several, wildly grown, monocot and dicot species. The most dominant grass species was Bermuda grass, *Cynodon dactylon* (L.).

The five selected grasslands were almost intact, not subjected to any kind of land-use effects, reclaiming or being influenced by severe anthropogenic stresses. The grasslands were only mowed twice per year, the first mow being undertaken after the end of Spring (during the last week of April) and the second at the beginning of Autumn (during the first week of November) and before initiation of snow falls. Each mow lasted 2-3 days for all five grasslands. It should be noted that a negative effect on Orthoptera presence due to mowing (Humbert et al. 2010; Cizek et al. 2012; Chisté et al. 2016) was not possible since sampling was performed during a period not associated with the postmowing period, as orthopterans start to appear at the end of June and completely disappear by the beginning of October.

Sampling protocol. Orthopteran assemblage was collected from the study sites, from the beginning of July until the end of September, over two consecutive years, 2022 and 2023, using standard sweep

net. Concerning the time effort per unit, the sampling schedule was adjusted to be performed between 1000 and 1400 h. To minimize the bias in sampling, only one of was responsible researchers for the sampling the individuals of different orthopteran species. The collected individuals were promptly killed in the field and stored in proper sampling containers. Details of the sampling technique are given by ElEla (2011).

The collected specimens were brought to the laboratory for systematic identification using the key of Ichikawa et Individuals al. (2006).of different orthopteran species were counted, sorted and kept in individually marked clean glass vials for further analyses. Additional identified specimens were stored individually in secure vials which then freezed (- 25 °C).

Feeding category. Feeding categories of different orthopteran species were elucidated from mandibular morphological characteristics and compared with previous when information data was reports. available. Laboratory dissections were followed immediately after bringing the collected specimens to the lab. Mandibles of adult orthopterans were exposed by lifting the labrum and hauling the maxillae. The mandibles were lightly brushed with 80% ethyl alcohol and distilled water to remove adhering debris (ElSayed, 2005; Smith and Capinera, 2005). Mandibles were then carefully detached from the head capsule by loosening the abductor muscles connecting the mandibles to the head capsule by regular and gentle back and forth hauling of each mandible using forceps, stereomicroscope (Olympus under а SZX12, Hamburg, Germany). Mandibles examined then at 20-40x were magnification, under direct illumination and photographs were taken from both ventral and dorsal views of each mandible with a digital camera (Sony HDR-CX590V, Japan). Morpho- mandibular characteristics were carefully examined and recorded in attempts to predict the feeding category of each orthopteran species.

Due to the rather wide gap in the relative abundance amongst orthopteran species, five adult male individuals from each species were subjected to gut contents analysis (total n = 215). As in many instances, females of certain orthopteran species feed more often than males (Hochkirch, 1999), it is necessary to clarify that only males were processed in the analyses since food strategies often differ between sexes, where it has been shown that females could consume plant species richer in nutrients (Chapman, 1990) and fertilized females tend to consume more protein, so even in purely herbivorous species, arthropod remains in the gut could appear during this period for relatively higher demand in protein requirements for production of eggs (ElSayed, 2005). For species with more than five adult males, the five individuals were randomly selected among collected males.

Adult male individuals of the collected orthopteran species were dissected by dissecting scissors and the alimentary tract was exposed, then longitudinally opened the contents of the and gut were permanently mounted on а glass microscope slides (Mulkern and Anderson, 1959: Brusven and Mulkern. 1960: Ohabuike, 1979; Kang and Chen, 1994; LeGall et al., 1998, 2003; ElSayed, 2005; ElSayed et al., 2020). Drops of distilled water wereadded whenever required to minimize dryness of the collected gut contents. These slides were examined under a light binocular microscope (Olympus® CX41RF, Hamburg, Germany) at 40× magnification. Microscopic examinations of fragments including different plant species (monocots and dicots), arthropod body parts and/or other ingested matters including debris particles were performed

(Kang and Chen, 1994; ElSayed, 2005; ElEla et al. 2010; ElSayed et al. 2020).

Qualitative records of the gut contents were made following ElEla et al. (2010) and ElSayed et al. (2020) to categorize each orthopteran species into a proper feeding category. The proposed seven feeding categories were:

- 1. Herbivorous feeders (H): The frequency of fragments of dicots species \cong frequency of fragments of monocots species with absence of fragments from arthropod parts.
- **2.** Herbivorous-mixed feeders (Hm): the same as herbivorous category with arthropod parts recorded in the gut.
- **3.** Graminivorous feeders (G): Frequency of fragments of monocots species exceeding 75% of the gut contents and no arthropod matter detected.
- **4.** Forbivorous feeders (F): Frequency of fragments of dicots species exceeding 75% of the gut contents with absence of fragments from arthropod parts.
- **5.** Forbivorous-mixed feeders (Fm): resembles the forbivorous category with arthropod parts recorded in the gut.
- **6.** Mixed-feeders (M): plant matters (including tubers roots, and subterranean and arthropod parts) and/or earthworm (Oligochaeta) parts encountered almost in equal proportions.
- 7. Detrito-bryophagous feeders (Db): detritus, mosses, algae and lichens were mainly encountered in the gut beside other plant matters.

Results

Taxonomic composition

The entire study area, for all five sites pooled, a total of 43 species (28 and 35 in 2022 and 2023, respectively), representing 25 genera, 7 families, 12 subfamilies and 13 tribes (Table 1) were recorded over the two consecutive years. The total number of collected individuals was 988 and 1324 during 2022 and 2023, respectively. The most dominant orthopteran species was Atractomorpha lata (Mochulsky, 1866). Overall, five orthopteran species were dominating the assemblage. These species (Mochulsky, Α. lata 1866). were Conocephalus maculatus (Le Guillou, 1841), Loxoblemmus equestris (Saussure, 1877), Mecopoda niponensis (Haan, 1843), Acrida cinerea (Thunberg, 1815) (Table 1). These species constituted 59.72% of the total assemblage during 2022 and 52.34% of the total assemblage during 2023 (Table 1). The sex ratio varied considerably among the different species and even between years (Table 1). Moreover, an absence of males or females of relatively rare species. e.g. Locusta migratoria (Linnaeus, 1758) and Loxoblemmus tsushimensis (Ichikawa, 2001) was recorded, as well as the absence of both sexes of other species e.g. Euconocephalus varius (Walker, 1869) during a specific year, 2022, was also recorded (Table 1).

Feeding category

Based on the observations of the variability of mandibular characteristics and analysis of contents of the alimentary canal, seven different feeding categories were identified (Table 1). Out of 43 recorded species sampled over the study period, 12 species (ca. 27.91%) were mixed-feeders (M) (Table 1).

Gryllidae and Trigonididae showed mandibles with comparatively sharp incisors and relatively long knife-shape terebral ridge. These mandibular characteristics could delineate a predacious feeding behavior. Based on postmortem gut content analyses, guts of dissected species showed fragmentary parts of plant materials including roots, occasionally debris matters and subterranean arthropod body parts. Consequently, the feeding category of these eight species could be positioned in the mixed-feeding category (M).

On the second rank of our recorded feeding categories, nine detritobryophagous feeders (Db), which comprised approximately 20.93% of the categories, were recorded in all tetrigid species (Table 1). Dissected guts of these species showed peculiar contents in which their guts were mainly loaded with detritus, algal parts, mosses and fungal hyphae.

Graminivorous (G) and forbivorousmixed feeders (Fm) were positioned in the third rank (ca. 16.28% for each category). The main graminivorous feeders frequently showed mandibular abrasions (Fig. 2). Both incisor and molar areas were subjected to obvious mechanical wear as a result of feeding habit, as is the case for *Acrida cinerea* (Thunberg, 1815) (Fig. 2).

On the other hand, species which are members in Conocephalini and Mecopodini were forbivorours-mixed feeders (Fm) where the incisor area was prominent and robust comprising a comparatively larger portion of the mandible and was associated with sharp terebral ridge (Fig. 2). In comparison, the molar area was small with two short and parallel molar slats.

Three species were forbivorous (F) in which the mandibular characteristic features were associated with the processing of forb resources (Table 1). The incisor area is equipped with robust and sharp teeth. The molar area consists of a small molar ridge forming a trituration area with molar slats for grinding forbs thoroughly (Fig. 2). The gut content analyses of these species showed that the contents were loaded with irregular fragments of forbs.

The three Phaneropterini species, *Hexacentrus japonicus* (Karny, 1907), *Phaneroptera nigroantennata* (Brunner von Wattenwyl, 1878) and *P. falcate* (Poda, 1761) were herbivourous-mixed feeders (Hm), with prominent and sharp incisors, without prominent wearing, and a comparatively smaller molar area (Fig. 2). **TABLE 1.** Orthopteran species collected from the study sites in Sendai, Japan, with their families, subfamilies, tribes, number of individuals and their feeding category.

Family	Subfamily	Tribe	Orthopteran species	No. of individuals ($3:2$)		Feeding category*
				2022	2023	
Acrididae	Acridinae	Acridini	Acrida cinerea (Thunberg, 1815)	54 (23: 31)	47 (19:28)	G
			A. conica (Fabricius, 1781)	33 (18:15)	39 (24:15)	G
	Melanopilinae	Podismini	Parapodisma mikado (Bolivar, 1890)	0	11 (8:3)	F
	Oedipodinae	Aiolopini	<i>Aiolopus thalassinus tamulus</i> (Fabricius, 1798)	23 (10:13)	10 (6:4)	Н
			Oedaleus infernalis (Saussure, 1884)	10 (3:7)	19 (12:7)	G
		Locustini	Locusta migratoria (Linnaeus, 1758)	0	7 (5:2)	Ğ
			Gastrimargus marmoratus (Thunberg, 1815)	0	9 (5:4)	G
	Oxyinae	Oxyini	Oxya japonica (Thunberg, 1815)	12 (4:8)	22 (15:7)	G
			O. yezoensis (Shiraki, 1910)	0	8 (5:3)	G
Pyrgomorphidae	Pyrgomorphinae	Atractomorphini	Atractomorpha lata (Mochulsky, 1866)	308 (131:177)	458 (168:290)	F
			A. sinensis (Bolivar, 1905)	0	14 (8:6)	F
Tetrigidae			Euparatettix tricarinatus (Bolívar, 1887)	13 (8:5)	0	Db
			Formosatettix niigataensis (Storozhenko & Ichikawa, 1993)	0	15 (7:8)	Db
			Tetrix akagiensis (Uchida & Ichikawa, 1991)	13 (9:4)	0	Db
			T. japonica (Bolivar, 1887)	21 (15:6)	29 (13:16)	Db
	Tetriginae	Tetrigini	T. kantoensis (Uchida & Ichikawa, 1991)	29 (18:11)	28 (14:14)	Db
			T. larvatus (Bei-Bienko & Mishchenko,1951)	16 (6:10)	37 (16:21)	Db
			T. macilenta (Ichikawa, 1993)	18 (6:12)	22 (14:8)	Db
			T. minor (Ichikawa, 1993)	7 (2:5)	0	Db
			T. silvicultrix ichikawa (Ichikawa, 1993)	20 (6:14)	21 (12:9)	Db
Eneopteridae	Oecanthinae	Oecanthini	<i>Oecanthus similator ichikawa</i> (Ichikawa, 2001)	12 (6:6)	0	F
			Loxoblemmus aomoriensis (Shiraki, 1930)	23 (9:14)	48 (22:26)	М

www.entsoc.gr

© 2024 Hellenic Entomological Society

			-			
			L. equestris (Saussure, 1877)	68 (39:29)	69 (45:24)	М
Gryllidae			L. tsushimensis (Ichikawa, 2001)	0	12 (7:5)	М
	Gryllinae	Gryllini	L. sylvestris (Matsuura, 1988)	6 (5:1)	0	Μ
			Teleogryllus emma (Ohmachi & Matsuura,	0	12 (5:7)	М
			1951)			
			T. infernalis (Saussure, 1877)	8 (2:6)	20 (6:14)	Μ
			Trigonidium pallipes (Stål, 1861)	7 (7:0)	0	Μ
			Velarifictorus aspersus (Walker, 1869)	0	7 (5:2)	Μ
			V. mikado (Saussure, 1877)	43 (17:26)	90 (36: 54)	Μ
			V. ornatus (Shiraki, 1911)	14 (8:6)	0	Μ
			Conocephalus maculatus (Le Guillou, 1841)	75 (43:32)	94 (59:35)	Fm
	Conocephalinae	Conocephalini	C. melaenus (Haan, 1843)	21 (14:7)	14 (5:9)	Fm
Tettigoniidae			C. japonicus (Redtenbacher, 1891)	0	9 (6:3)	Fm
			Euconocephalus varius (Walker, 1869)	0	11 (8:3)	Fm
			Ruspolia dubia (Redtenbacher, 1891)	0	12 (10:2)	Fm
			R. lineosa (Walker, 1869)	38 (22:16)	59 (27:32)	Fm
	Mecopodinae	Mecopodini	Mecopoda niponensis (Haan, 1843)	85 (39:46)	25 (7:18)	Fm
			Hexacentrus japonicus (Karny, 1907)	0	11 (7:4)	Hm
	Phaneropterinae	Phaneropterini	Phaneroptera falcate (Poda, 1761)	0	20 (15:5)	Hm
			<i>P. nigroantennata</i> (Brunner von Wattenwyl, 1878)	0	8 (6:2)	Hm
Trigonididae	Nemobiinae	Pteronemobiini	Dianemobius furumagiensis (Ohmachi & Furukawa, 1929)	0	7 (5:2)	М
			Pteronemobius fascipes (Walker, 1869)	11 (6:5)	0	М
Total number of individuals (♂:♀)				988 (476:512)	1324 (632:692)	

*Feeding category: Db: Detrito-bryophagous feeder, F: Forbivorous feeder, Fm: Forbivorous-mixed feeder, G: Graminivorous feeder, H: Herbivorous feeder, Hm: Herbivorous-mixed feeder M: Mixed feeder.

Table (1) showed that only one acridid species, *Aiolopus thalassinus tamulus* (Fabricius, 1798), possessed a purely herbivorous feeding category (H) which comprised approximately the least proportions of the total categories (ca. 2.33%). The incisor and molar areas were almost of uniform proportions. The incisors were parallel and sharp with no clear abrasions either in molar or incisor areas of the mandible (Fig. 2).

It was clear that the major differences in mandibular structure were related to differences in feeding behavior rather than the taxonomic position of the species as family Acridinae and Tettigoniidae possessed more than feeding category (Table 1).

Discussion

In the majority of phytophagous insect orders, Chapman (1982) noted that oligophagy or monophagy was observed in more than 50% of the species whereas the Orthoptera, stand apart from the other orders, more than 60% of the species have been classified as polyphagous and 25% as graminivorous (Picaud et al. 2003).

Orthopteran species are considered as one of the more diverse taxa with thousands of extant described species possessing prominent diversity in forms and habitats (Grimaldi and Engel, 2005; Bidau, 2014; Eades et al. 2014; Song et al. 2015) and representing crucial links in food chains (Badenhausser, 2012). Orthopteran species are widespread and important herbivores in natural grassland ecosystems and agroecosystems as they consume a considerable portions of grasses and forbs (Köhler et al. 1987, Blumer and Diemer 1996; ElSayed, 2005; Bharamal and Koli 2014) and regulate plant community structure (Zhang et al. 2011). Indeed, Order Orthoptera plays an influential and functional role in the recycling and equilibrium of natural ecosystems by enhancing the conversion of biomass through their easily transformable excrement (Samways, 1994; Hao et al.

2015; Ngoute et al. 2021). All the previous aspects triggered the urge to study the variations in feeding categories of different orthopteran species through mandibular morpho- characteristic variations and gut contents analysis rather than relying on field or laboratory observations or food choice tests.

The aspects of food strategies including selection of food, feeding patterns, food preferences, specialization in dietary requirements have been studied by many authors in a diverse orthopteran species (e.g., Williams, 1954; Gangwere, 1961; Mulkern, 1967; Otte and Joern, 1976; Bernays and Chapman, 1978; Uvarov, 1977; Joern, 1983, 1985; Bernays and Bright, 1993). Still the food strategy based on the structure of the mouthparts, habitat preferences, or feeding behavior was observed in only a few species (ElSayed et al. 2020).

Morphological variations in mouthparts, including mandibles, were subjected to further thorough research and itemized investigations have been directed by numerous authors in various regions, significant among them were Snodgrass (1928), Gangwere (1965, 1966), Gangwere et al. (1976, 1998) and Patterson (1984). Smith and Capinera (2005) in North America, Liebermann (1968), Gangwere and Ronderos (1975) in South America, Williams (1954); Kaufmann (1965),Gangwere and Morales (1973) in Europe, Gangwere and Spiller (1995); Gangwere et al. (1998) in the Mediterranean islands, Gapud (1968), Feroz and Chaudhry (1975), Kang et al. (1999), ElEla et al. (2010, 2012), ElSayed et al. (2020) in Asia, Chapman (1964), Le Gall et al. (1998, 2003) and ElSayed (2005) in Africa.

The strong relationship with diet makes morphological characteristics of mouthparts an important trait for insect evolutionary biologists (Snodgrass, 1928; Brues, 1939) and systematists (Mulkern, 1967). One of the first who coped with this topic was Isely (1944) who studied the de-



FIG. 2.: Images of morpho-mandibular variability representing different feeding categories. a. *Tetrix japonica* (Bolivar, 1887), a representative of Forbivorous species (F); b. *Hexacentrus japonicus* (Karny, 1907), a representative of Herbivourous-mixed species (Hm); c. *Conocephalus melaenus* (Haan, 1843), a representative of Forbivorous-mixed species (Fm); d. *Velarifictorus mikado* (Saussure, 1877), a representative of Mixed-feeder species (M); e. *Aiolopus thalassinus tamulus* (Fabricius, 1798), a representative of Herbivorous species (H); f. *Acrida cinerea* (Thunberg, 1815), a representative of Graminivorous species (G).

tailed morphological and structural features of mouthparts and correlated these characteristics with various feeding habits. Initial results and information offered by Isely (1944) on categorizing mandibles according to mandibular groups have since been shown to be prevalent, especially, in grasshoppers and other entomological taxa.

Isely (1944), on his research on 89 species, proclaimed two extreme forms of mandibles. i.e. graminivorous and forbivorous, with a series of intermediate forms associated with a mixed diet. Generally, Isely (1944) defined three groups of mandibles according to the overall structure and distinctive diet: (i) graminivorous (grass feeding type) with molar area adapted for grinding and incisors typically merged into a scythe like edge, (ii) forbivorous(forb or broad leaf plant feeding type) having a molar area composed of a depression encircled by elevated teeth and a strong interlocking incisor, (iii) herbivorous (mixed- feeding type) that have features of both of the previously mentioned groups.

On the other hand, Joern (1979) proposed that orthopteran species possess a broad spectrum in diversity of diet and consequently feeding behavior ranged from strict specialist to extreme generalist.

accordance, several researchers In (Gangwere, 1965; Chapman, 1990; Smith and Capinera, 2005) agreed with Isely (1944) and they had merely classified three main categories based on the plant categories. These categories were graminivorous. forbivorous and herbivorous (mixed-feeding). However, due to variations in mandibular morphocharacteristics, other studies provided broader classification of feeding categories (Kuřavová and Kočárek, 2016; ElEla et al. 2010; ElSayed et al. 2020).

Comparisons were performed by Patterson (1983) concerning the morphological structure of the mandibles with their niche variations. Hypothesis of niche variations predicts greater overall mandibular characteristic variability in orthopteran species having relatively the broadest trophic niches. In this study, mandibular characteristics of orthopteran species vary in different groups and these variations are therefore believed to meet the different feeding habits and in correspondence specifying а definite feeding category. These were in accordance with findings of several studies (Gangwere, 1965; ElEla, 2011; ElSayed et al. 2020).

Apparent observed similarities among members belonging to the same subfamily and striking differences between subfamilies assume a strong phylogenetic component to feeding adaptations as suggested by Otte and Joern (1977) and Joern (1979).

Ensiferan species, on the other hand, feeding categories ranged from forbivorous category which was observed in a single eneopterid species to strictly mixed feeding category. Ensiferans tended to feed on taxonomically diverse plant species (Palmer et al. 2024) beside frequent proportions of arthropod parts which could suggest predaceous or cannibalistic habits where some ensiferan species specialized on rather unusual food sources (ElEla, 2010; ElSayed et al. 2020).

The consumption of a specific diet is commonly associated with the shape and structure of the mandibles. Graziella et al. (2015) showed significant differences in the shapes of the mandibles among the variant forms of grasshopper species which may be due to the role of trophic diversification in the morphological differentiation of insect. The apparent diverse variations among mandibular characteristic features of different orthopteran species were remarkable to consider. In our research, more types of mandibles were suggested in association with analyses of gut contents in an attempt to position each collected species in the proper feeding category.

Although orthopteran species have often been regarded as polyphagous herbivores (ElSayed et al. 2020), most of these species show degree of dietary specialization (Mulkern, 1967; Uvarov, 1977; ElSaved et al. 2020). Orthopteran species possessed a dietary specialization was reflected in patterns of mandibular variations, and yet this diversification in morpho-mandibular characteristics could be utilized to catalogue orthopteran species in proper feeding category.

In this study, caeliferan species were strictly not engaged in cannibalistic behavior as confirmed by the absence of arthropod parts in their guts. However, other studies have recorded cannibalistic behavior in some caeliferan species only under crowded conditions (Bomar and Lockwood, 1994; ElEla, 2010).

It was interesting to spot the light on the peculiar detrito-bryophagy which was restricted to the tetrigid species with mosses, algal hyphae and debris matters were frequently encountered in their guts (Kuřavová and Kočárek, 2016).

Studies performed on tetrigids are considerably rare and restricted to a few species (Paranjape and Bhalerao, 1985; Blackith and Blackith, 1987; Reynolds et al. 1988; Hochkirch et al. 2000, Gröning et al. 2007; Kuřavová and Kočárek 2015; Kuřavová and Kočárek, 2016; Musiolek and Kočárek 2016), and even fewer and fragmentary in Japan (ElEla et al. 2010; ElEla, 2011; Tan et al. 2017; ElSayed et al. 2020).

The feeding habit of Tetrigidae has been described, ranging from observational to quantitative data, in only a few species (Tan et al. 2017). However, the detritobryophagy has been observed in almost all species (Verdcourt, 1947; Paranjape and Bhalerao, 1985; Hochkirch et al., 2000; Bidau, 2014; Kuřavová and Kočárek, 2015).

In a related species, Tetrix undulata (Sowerby, 1806). Verdcourt (1947)performed the faecal analysis and found a variety of mosses species in 80% of the fecal pellets. Lock (1996) analyzed crop content of three specimens of T. subulata (Linnaeus, 1758) and observed the presence of algae, detritus and sand grains with no traces of mosses or higher plants. Dietary habit of T. undulata (Sowerby, 1806) was studied by Hodgson (1963) and he observed feeding of the species on grasses (monocots), mosses (variety of species), lichens and algae, and on humus. Besides that, he observed also feeding on the dead bush cricket Pholidoptera griseoaptera (De Geer, 1773) in insectarium. Further studies were conducted by Hochkirch et al. (2000) who found that males of T. undulata (Sowerby, 1806) were exclusively feeding on mosses and algae, while females also included grasses and forbs in their diet. On other related tetrigid species, Reynolds et al. (1988) studied the diet of two tropical tetrigids in Sulawesi by analyzing their crop contents and they observed the presence of mosses or vascular plants in 82% of specimens of Scelimena celebica (Bolivar, 1887) and in 100% of specimens of Diotarus pupus (Bolivar, 1887). Indeed, it is far from precise to extend the detritobryophagy among all members of Tetrigidae since more documentation of dietary preference for each species is required in different spots and among different sexes.

Feeding category could be considered as an additional important ecological trait used to analyze the structure of orthopteran fauna. The collected orthopteran species were relatively dominated by forbivorous (F) and mixed (M) feeding species. However, mixed-feeding habit also comprised a relatively major percentage in the feeding categories of the assemblage. Although Min and Min (2008), in their taxonomic studies, mentioned that *Aiolopus thalassinus* is an important graminivorous pest on different plants and grasses; our results from mandibular structure and gut contents analysis revealed a forbivorous feeding category for this species.

Comparisons in our research have shown that orthopteran species of the family Acrididae exhibited the broadest trophic niches with three feeding categories patterns which support diverse of mandibular characteristics and dietary preferences. Herbivorous and forbivorous species with mixed feeding habits (Hm and Fm) have more complex mandibular characteristics and relatively more heterogeneous food intake. These associations, in regard of morphological characteristics and dietary preference in each group, support Patterson's niche variation hypothesis Patterson (1983).

Possibly, species with mixed feeding categories (M, Fm and Hm) circumvent the defense systems by feeding only small portions of many plant species and other organic parts thus keeping the toxic levels of secondary metabolites (allelochemicals) below some critical level (Freeland and Janzen, 1974). However, since this is a per se mechanism and is species-specific, more investigations are required.

Although little information is known about the chemical properties of most grasses in the study sites, grasses accumulate relatively high levels of silica in their tissue which could be an alternative tactic, with potential benefits, including antiherbivore defense (Vicari and Bazely 1993; ElSayed et al. 2020; Quigley et al. 2020).

Conclusion

The gathered data on the comparatively small assemblage give only a hint concerning the whole story of evolution of

Graminivorous species differ in their dietary choices when feeding on graminoids (monocots) in natural grassland vegetation when compared to forbivorous or herbivorous species and this considered as species-specific (Joern, 1985). Bemays et al. (1989) have also reported that tannins are often considered a major class of allelochemical defense in dicotyledonous plant species (dicots) and they have also considered deterrents been as to graminivorous orthopteran species (Bemays and Chapman, 1977) which could partly explain the selection of grasses by graminivorous species.

Mandibular abrasion in both incisor and molar areas in graminivorous species (G) could be attributed to the fact that these species feed regularly on silica-rich grasses. These silica particles could act as a sand paper that wear the mandibles of these graminivorous species (ElSayed et al. 2020). It was reported that silica causes wear to insect mandibles which could potentially impact on the performance of herbivory (Baker et al., 1959; Vicari and Bazely, 1993; Massey and Hartley, 2009; ElSayed et al. 2020).

Our analysis of the gut contents of orthopteran species with mixed-feeding category (M) or even species of mixed feeding habit beside main category (i. e. Hm and Fm) could suggest that arthropod parts and other detritus of plant matters are not incidentally consumed but they are main part of their dietary preference. These could intentionally species feed on arthropod parts which were predominant in the gut of the entire dissected individuals. However, more detailed investigations and analyses on this aspect are required.

functional food strategies among orthopeteran species. A general scheme for explicating the diet of a given orthopteran species, or other taxa group, could be started with detailed inspections of their mandibular morphological characteristics. Although most species with forb feeding mandibles could feed on mixture of grasses and forbs, one of the confirmation avenue is associating mandibular characteristics with observations and data gathered from analyzing the gut contents.

It is hoped that more researchers will find that the study of morpho-mandibular characteristics and gut contents analysis is rewarding will contribute to the advancement of our knowledge concerning feeding strategies in Orthoptera. This research tries to open the door for more studies to analyze more mandibles from more taxa (Patterson, 1983).

Acknowledgments

The authors wish to acknowledge the staff at the Laboratory of Ecological Integration and Laboratory of Insect Science and Bioregulation, Tohoku University for their sincere assistance and prompt provided facilities. Also, sincere and warm gratitude is extended to Prof. Dr. Urabe Jotaro (Graduate School of Life Science, Tohoku University, Japan) for his keen and endless assistance.

Conflict of interest

The authors declare no conflict of interest associated with this article and Prof. Dr. Ames Cheryl Lynn (International Integrative Research and Instruction Unit, Tohoku University, Japan) for her persistent cooperation and continuous encouragement.

References

Badenhausser. L 2012. Estimation d'Abondance des Criquets (Orthoptera: Acrididae) dans les Ecosystèmes Société Prairiaux. Annales de la_ Entomologique de France 48: 397-406. https://doi.org/10.1080/00379271.2012.1 0697787

- Baker, G., L. H. P. Jones and I. D. Wardrop. 1959. Cause of wear in sheeps teeth. Nature. 184: 1583–1584. https://doi.org/10.1038/1841583b0
- Bernays, E. A. 1998. Evolution of feeding behavior in insect herbivores. BioScience. 48: 35-44.

https://doi.org/10.2307/13 13226

- Bernays, E. A. and K. L. Bright. 1993. Dietary mixing in grasshoppers: a review. Comparative Biochemistry and Physiology. A 104: 125–131. https://doi.org/10.1016/0300-9629(93)90019-Z
- Bernays, E. A., G. Cooper Driver and M. Bilgener. 1989. Herbivores and plant tannins. Advances in Ecological Research. I9:263-302.
- Bernays, E. A. and R. F. Chapman. 1977. Deterrent chemicals as a basis of oligophagy in *Locusta migratoria* (L) Ecological Entomology. 2:1-18. https://doi.org/10.1111/j.1365-2311.1977.tb00861.x
- Bernays, E. A. and R. F. Chapman. 1978. Plant chemistry and acridoid feeding behavior. In: Harborne HB, editor. Biochemical Aspects of Plant and Animal Coevolution: Annual Proceedings of the Phytochemical Society of Europe. No.15. New York: Academic Press; pp. 99-141.
- Bharamal, D. L., and Y. J. Koli. 2014. Grasshoppers as multidisciplinary model organism in changing environment. International Journal of Current Microbiology and Applied Science. 3: 372–380.
- Bidau, C. J. 2014. Patterns in Orthoptera biodiversity. I. Adaptations in ecological and evolutionary contexts. Journal of Insect Biodiversity. 2 (20): 1–39. https://doi.org/10.12976/jib/2014.2.20
- Blackith, R. E. and R. M. Blackith. 1987. Tridactylids and pygmy grasshoppers (Orthoptera) from Sulawesi, Indonesia. Overgedrukt uit het Tijdschrift voor Entomologie. 130: 1–10.

- Blumer, P. and M. Diemer, 1996. The occurrence and consequences of herbivory in an grasshopper alpine grassland, Swiss Central Alps, Arctic and Research. Alpine 28: 435-440. https://doi.org/10.1080/00040851.1996.12 003196.
- Bomar, C. R. and J. A. Lockwood. 1994. Olfactory basis of cannibalism in grasshoppers (Orthoptera: Acididae): I. laboratory assessment of attractants. Journal of Chemical Ecology. 20(9):2249-60.

http://doi.org/10.1007/ BF02033200

- Brues, C. T. 1939. Food, drink, and evolution. Science. 90: 145-149.
- Brusven, M. A. and G.B. Mulkern. 1960. The Use of Epidermal Characteristics for the Identification of Plants Recovered in Fragmentary Condition from the Crops of Grasshoppers. Vol. 3. North Dakota, USA: North Dakota Agricultural Experimental Station Research Reports; p. 11
- Chapman, R. F. 1964. The structure and wear of the mandibles in some African grasshoppers. Proceedings of the Zoological Society of London. 142:107-121.
- Chapman R. F. 1982. Chemoreception: the significance of receptor numbers. Advances in Insect Physiology. 16: 247– 356. https://doi.org/10.1016/S0065-2806(08)60155-1
- Chapman, R. F. 1990. Food selection. In: Chapman, R. F. and Joern, A. (eds), Biology of grasshoppers. Wiley, pp. 39-72.
- Chisté, M. N., K. Mody, M. M. Gossner, N. K. Simons, G. Köhler, W. W. Weisser and N. Blüthgen. 2016. Losers, winners, and opportunists: How grassland land-use intensity affects orthopteran communities. Ecosphere. 7(11): e01545.

https://doi.org/10.1002/ecs2.1545

Cigliano, M. M., H. Braun, D.C. Eades and D. Otte. 2022. Orthoptera Species File. Version 5.0/5.0.

- Cizek, O., J. Zamecnik, R. Tropek, P. Kočárek and M. Konvicka. 2012. Diversification of mowing regime increases arthropods diversity in species-poor cultural hay meadows. Journal of Insect Conservation. 16: 215–226. http://doi.org/10.1007/s10841-011-9407-6.
- Dakhel, W. H., S. T Jaronski and S. Schell. 2020. Control of pest grasshoppers in North America. Insects. 11, 566. https://doi.org/10.3390/insects11090566
- Di Russo, C., M. Rampini, L. Latella and M. Cobolli. 2014. Measurements of the diet in two species of *Troglophilus* Krauss, 1879 cave crickets from Italian subterranean habitats (Orthoptera, Rhaphidophoridae). Subterranean Biology. 13: 45-54.

https://doi.org/10.3897/subtbiol.13.6719

- Eades, D. C., Otte, D., Cigliano, M. M. and H. Braun. 2014. Orthoptera Species File. Version 5.0/5.0. [8/1/2014]. http://Orthoptera.SpeciesFile.org.
- ElEla, A. S. 2011. Anthropogenic disturbance and structure of Orthopteran assemblages in Satoyama. Ph.D. Thesis. 146 pp. Institute of Natural Science and Technology, Kanazawa University.
- ElEla, S. A., W. ElSayed and K. Nakamura. 2010. Mandibular structure, gut contents analysis and feeding group of orthopteran species collected from different habitats of Satoyama area within Kanazawa City, Japan. Journal of Threatened Taxa. 2(5): 849- 857. https://doi.org/10.11609/JoTT.o2346.849-

57 ElEla, A. S., W. ElSayed and K. Nakamura. 2012. Incidence of orthopteran species (Insecta: Orthoptera) among different sampling sites within Satoyama area, Japan. Journal of Threatened Taxa. 4: 2409-2480.

<u>http://doi.org/10.11609/JoTT.O2775.2476</u> <u>- 80</u>.

ElSayed W. M. 2005. Grasshoppers (Orthoptera: Acrididae) communities in Abu-Raûwash district (Giza, Egypt) [MSc. Thesis]. Cairo University; pp. 169.

ElSayed, M. Wael, Shahenda Abu ElEla and Koji Nakamura. 2020. Mandibular Structure, Gut Contents Analysis and Feeding Group of Orthopteran Species Collected from Different Habitats of Satoyama Area within Kanazawa City, Japan. Pests, Weeds and Diseases in Agricultural Crop and Animal Husbandry Production. IntechOpen.

http://doi.org/10.5772/intechopen.88705.

- ElShazly, M. and W. M. ElSayed. 2006. The structure and host plant selection of an acridid community on the edge between a desert- and an agro-ecosystem in Egypt. Environmental Informatics Archives. 4: 1-25.
- Feroz, M. and M. A. Chaudhry. 1975. Studies on mandibles of some grasshoppers of Lahore. Biologia. (Lahore). 21: 211-225.
- Flook, P. K. and C. H. F. Rowell. 1997. The Phylogeny of the Caelifera (Insecta, Orthoptera) as Deduced from mtrRNA Gene Sequences. Molecular Phylogenetics and Evolution. 8 (1): 89-103. https://doi.org/10.1006/mpev.1997.0412
- Freeland, W. J. and D. H. Janzen. 1974. Strategies in herbivory by mammals: the role of plant secondary compounds. The American Naturalist. 108:269–289. https://doi.org/10.1086/282907
- Fry, B., A. Joern and P. L. Parker. 1978. Grasshopper food web analysis: use of carbon isotope ratios to examine feeding relationships among terrestrial herbivores. Ecology. 59: 498-506. https://doi.org/10.2307/1936580
- Gangwere, S. K. 1961. A monograph on food selection in Orthoptera. Transactions of the American Entomological Society. 87: 67-230.
- Gangwere, S. K. 1965. Food selection in the Oediponidae grasshopper *Arphia sulphurea*. American Midland Naturalist. 74: 67-75.

Gangwere, S. K. 1966. Relationships between mandibles, feeding behavior and damage inflicted on plants by the feeding of certain acridids (Orthoptera). Michigan Entomology. 1: 13-16.

https://doi.org/10.22543/0090-0222.1006

- Gangwere, S. K. and A. Morales. 1973. Food selection and feeding behavior in Iberian Orthopteroidea. Anales del Instituto Nacionalde Investigaciones Agrarias. Serie proteccion vegetal. 3: 251-337.
- Gangwere, S. K. and R. A. A. Ronderos. 1975. Synopsis of food selection in argentine Acridoidea. Acrida. 4: 173-194.
- Gangwere, S. K., F. C. Evans and M. L. Nelson. 1976. The food habits and biology of Acrididae in an old-field community in Southeastern Michigan. Great Lakes Entomology. 9: 83-123. https://doi.org/10.22543/0090-0222.1275
- Gangwere, S. K. and D. O. Spiller. 1995. Food selection and feeding behavior in selected Orthoptera sen. lat. of the Balearic Islands, Spain. Journal of Orthopteran Research. 4:147-160.
- Gangwere, S. K., J. C. Mc Kinney, M. A. Ernemann and R. G. Bland. 1998. Food selection and feeding behavior in selected Acridoidea (Insecta: Orthoptera) of the Canary Islands, Spain. Journal of Orthopteran Research. 7: 1-21.
- Gapud, V. P. 1968. The external morphology of the head and mouthparts of some Philippine Orthoptera. Philippine Entomology. 1: 11-32.
- Graziella, M. Barcebal, Kimverly Hazel I. Coronel, Mark Anthony J., Torres Cesar G. Demayo. 2015. Variability in the shape of the mandibles of grasshopper (Orthoptera: Acrididae) from selected places in Mindanao, Philippines. Advances in Environmental Biology. 9 (19): 123-126.
- Grimaldi, D. and M. S. Engel. 2005. Evolution of the Insects. Cambridge University Press, New York.

- Gröning, J., S. Krause and A. Hochkirch. 2007. Habitat preference of an endangered insect species, Cepero's groundhopper (Tetrix ceperoi). Ecological Research. 22: 767–773. https://doi.org/10.1007/s11284-006-0315-2
- Hao, S., S. Wang, A. Cease and Le Kang. 2015. Landscape Level Patterns of Grasshopper's Communities in Inner Mongolia: Interactive Effects of Livestock and Precipitation Gradient. Journal of Landscape Ecology. 30: 1657–1668. https://doi.org/10.1007/s10980-015-0247-8
- Hochkirch, A. 1999. Leaf or stem? sexspecific differences in the locations of *Rhainopomma usambaricum* (Ramme, 1929) (Orthoptera: Acridoidea: Lentulidae). Mitteilungen der Deutschen Gesellschaft für Allgemeine und Angewandte Entomologie. 12: 459-462.
- Hochkirch, A., J. Groöning, T. Loos, C. Reichelt. Metzing and M. 2000. Specialized diet and feeding habits as key factors for the habitat requirements of the grasshopper species Tetrix subulata (Orthoptera: Tetrigidae). Entomologia Generalis. 25:39-51. http://doi.org/10.1127/entom.gen/25/2000

/39

- Hochkirch, A., J. Deppermann and J. Gröning. 2006. Visual communication behaviour as a mechanism behind reproductive interference in three pygmy grasshoppers (genus *Tetrix*, Tetrigidae, Orthoptera). Journal of Insect Behavior. 19: 559-571. http://doi.org/10.1007/s10905-006-9043-2
- Humbert, J.-Y., J. Ghazoul, N. Richner and T. Walter. 2010. Hay harvesting causes high orthopteran mortality. Agriculture, Ecosystems and Environment. 139: 522–527. https://doi.org/10.1016/j.agee.2010.09.01 2
- Ichikawa A, Y. Kano, Y. Kawai, M. Kawai, O. Tominago and T. Murai. 2006. Orthoptera of the Japanese Archipelago in Color. 2nd ed. pp. 687. (In Japanese)

- Isley, F.B. 1944. Correlation between mandibular morphology and food specificity in grasshoppers. Annals of Entomological Society of America. 37: 47-67.
- Joern, A. 1979. Feeding patterns in grasshoppers (Orthoptera: Acrididae): factors influencing diet specialization. Oecologia (Berl.). 38:325-347.
- Joern, A. 1983. Host plant utilization by grasshoppers (Orthoptera: Acrididae) from a sandhills prairie. Journal of Range Management. 36: 793-797.
- Joern, A. 1985. Grasshopper dietary (Orthoptera: Acrididae) from a Nebraska Sand Hills Prairie. Transactions of the Nebraska Academy of Sciences. 8: 21-32.
- Kang, L. and Y. L. Chen1994. Trophic niches of grasshoppers within steppe ecosystem in Inner Mongolia. Acta Entomologica Sinica. 37: 178–189 (in Chinese with English summary).
- Kang, L., Y. Gan and S. L. Li. 1999. The structural adaptation of mandibles and food specificity in grasshoppers on Inner Mongolian grasslands. Journal of Orthopteran Research. 8: 257-269.
- Kaufmann, T. 1965. Biological studies on some Bavarian Acridoidea (Orthoptera), with special reference to their feeding habits. Annals of the Entomological Society of America. 58: 791-801.
- Köhler, G., H.-P. Brodhun and G. Schäller.1987. Ecological energetics of Central European grasshoppers (Orthoptera: Acrididae). Oecologia. 74: 112–121.
- Kuřavová, Katerina and Peter Kočárek. 2016. Mandibular morphology and dietary preferences in two pygmy mole crickets of the genus *Xya* (Orthoptera: Tridactylidae). Turkish Journal of Zoology. 40 (5): 720-728. http://doi.org/10.3906/zoo-1510-19
- Kuřavová, K., Š. Grucmanová, Z. Filipcová, V. Plášek, P. Drozd and P. Kočárek. 2017.
 Is feeding on mosses by groundhoppers in the genus *Tetrix* (Insecta: Orthoptera) opportunistic or selective? Arthropod-

Plant Interactions. 11: 35–43. https://doi.org/10.1007/s11829-016-9461-9

- Le Gall, P., E. Mingouolo and G. Bani. 1998. Diets of *Zonocerus variegates* (L.) (Orthoptera: Acrididae) in cassava fields in Congo. Journal of Applied Entomology. 122: 9-13. https://doi.org/10.1111/j.1439-0418.1998.tb01453.x
- Le Gall, P., Z. Djihou, G. Tchenga and C. J. Lomer. 2003. Diet of *Zonocerus* variegatus (Linné, 1758) (Orthoptera: Acrididae) in cassava field in Bénin. Journal of Applied Entomology. 127 (7): 435-440. http://doi.org/ 10.1046/j.1439-0418.2003.00762.x
- Liebermann, J. 1968. The mandibles of grasshoppers of the subfamily Chilacridinae. Revista de Investigaciones Agropecuarias. (5): 53-62. (In Spanish).
- Massey, F. P., A. R. Ennos and S. E. Hartley. 2006. Silica in grasses as a defence against insect herbivores: Contrasting effects on folivores and a phloem feeder. Journal of Animal Ecology. 75(2), 595–603. https://doi.org/10.1111/j.1365-2656.2006.01082.x
- Massey, F. P. and S. E. Hartley. 2009. Physical defences wear you down: progressive andirreversible impacts of silica on insect herbivores. Journal of Animal Ecology. 78 (1): 281-91.

https://doi.org/10.1111/j.1365-2656.2008.01472.x

- Min, Z. Z. and S. H. Min. 2008. A taxonomic study of the genus *Aiolopus* Fieber (Orthoptera, Oedipodinae) from China, with description of new species, Acta Zootaxanomica Sinica. 33(4): 660-663. http://doi.org/10.3969/j.issn.1000-0739.2008.04.005
- Mulkern, G. B. 1967. Food selection by grasshoppers. Annual Review of Entomology. 12: 59-78. https://doi.org/10.1146/annurev.en.12.010 167.000423

- Mulkern, G. B. and J. F. A. Anderson. 1959. A technique for studying to the food habits and preferences of grasshoppers. Journal of Economic Entomology. 52: 342.
- Musiolek, D. and P. Kočárek. 2016. Weatherdependent microhabitat use by *Tetrix tenuicornis* (Orthoptera: Tetrigidae). The Science of Nature. 103: 68. https://doi.org/10.1007/s00114-016-1393-9.
- Ngoute, C. O., D. Hunter and M. Lecoq. 2021. Perception and knowledge of grasshoppers among indigenous communities in tropical forest areas of southern Cameroon: Ecosystem conservation, food security, and health. Journal of Orthoptera Research. 30(2): 117- 130. https://doi.org/10.3897/jor.30.64266
- Ohabuike, J. E. 1979. Grass availability and food preference of the African migratory locust, *Locusta migratoria migratoriodes* (R. and F.). Zeitschrift für Angewandte Entomologie. 88: 354-363. https://doi.org/10.1111/j.1439-0418.1979.tb02513.x
- O'Neill, K. M., B. E. Olson, M. G. Rolston, R. Wallander, D. P. Larson and C. E. Seibert. 2003. Effects of livestock grazing on rangeland grasshopper (Orthoptera: Acrididae) abundance. Agriculture, Ecosystems & Environment. 97(1-3): 51-64. https://doi.org/10.1016/S0167-

8809(03)00136-1

- Otte, D. and A. Joern. 1976. On feeding patterns in desert grasshoppers and the evolution of specialized diets. Proceedings of the Academy of Natural Sciences of Philadelphia. 128: 89-126.
- Palmer, C. M., N. L. Wershoven, S. J. Martinson, H. M. ter Hofstede, W. J. Kress and L. B. Symes. 2024. Patterns of Herbivory in Neotropical Forest Katydids as Revealed by DNA Barcoding of Digestive Tract Contents. Diversity. 14, 152. https://doi.org/10.3390/d14020152

Paranjape S. Y. and A. M. Bhalerao. 1985. Bioecological observations on a pigmy locust, *Potua sabulosa* Hancock (Tetrigidae: Orthoptera). Psyche. 92(2-3): 331–336.

https://doi.org/10.1155/1985/30570

- Parr, C. L. and S. L. Chown. 2003. Burning issues for conservation: a critique of faunal fire research in Southern Africa. Austral Ecology. 28: 384–395. https://doi.org/10.1046/j.1442-9993.2003.01296.x
- Patterson, B. D. 1983. Grasshopper mandibles and the niche variation hypothesis. Evolution. 37: 375-388. https://doi.org/10.1111/j.1558-5646.1983.tb05546.x
- Patterson, B. D. 1984. Correlation between mandibular morphology and specific diet of some desert grassland Acrididae (Orthoptera). American Midland Naturalist. 111: 296-303.
- Picaud, F., E. Bonnet, V. Gloaguen and D. Petit. 2003. Decision Making for Food Choice by Grasshoppers (Orthoptera: Acrididae): Comparison Between a Specialist Species on a Shrubby Legume and Three Graminivorous Species, Environmental Entomology. 32(3): 680– 688. https://doi.org/10.1603/0046-225X-32.3.680
- Quigley, K. M., D. M. Griffith, G. L. and T. M. Anderson. 2020. Soil nutrients and precipitation are major drivers of global patterns of grass leaf silicification. Ecology. 101(6):e03006. http://doi.org/10.1002/ecy.3006
- Reynolds, J. D., R. E. Blackith and R. M. Blackith. 1988. Dietary observations on some tetrigids (Orthoptera: Caelifera) from Sulawesi (Indonesia). Journal of Tropical Ecology. 4: 403–406. https://doi.org/10.1017/S02664674000030 72.
- Saha, H. K., A. Sarkar and P. Haldar. 2011. Effect of Anthropogenic disturbance on the diversity and composition of the

acridid fauna of sites in the dry deciduous forest of West Bengal, India. Journal of Biodiversity and Ecological Sciences. 1 (4): 313–320.

- Samways, M. J. 1994. Insect conservation biology. Chapman & Hall, London, UK.
- Smith, T. R. and J. L. Capinera. 2005. Mandibular morphology of some Floridian grasshoppers (Orthoptera: Acrididae). Florida. Entomologist. 88: 204–207. https://doi.org/10.1653/0015-4040(2005)088[0204:MMOSFG]2.0.CO; 2
- Snodgrass, R. E. 1928. Morphology and evolution of the insect head and its appendages. Smithsonian Miscellaneous Collection. 81: 1-158.
- Snodgrass, R. E. 1935 Principles of insect morphology. McGraw-Hill Publishing Co., New York.
- Soliman, M. M., A. A. Haggag and M. M. El-Shazly. 2017. Assessment of grasshopper diversity along a pollution gradient in the Al-Tebbin region, South Cairo. Journal of Entomology and Zoology Studies. 5(1): 298-306.
- Song, H. 2018. Biodiversity of Orthoptera. pp. 245–279. In: R.G. Foottit and P. H. Adler (eds), Insect Biodiversity. Science and Society. Volume 2. Hoboken: Wiley.
- Song, H., C. Amédégnato, M. M. Cigliano, L. Desutter-Grandcolas, S. W. Heads, Y. Huang, D. Otte and M. F. Whiting. 2015. 300 million years of diversification: elucidating the patterns of orthopteran evolution based on comprehensive taxon and gene sampling. Cladistics. 31:621-651. https://doi.org/10.1111/cla.12116
- Tan, M. K., H. Yeo and W. S. Hwang. 2017. Ground dwelling pygmy grasshoppers (Orthoptera: Tetrigidae) in Southeast Asian tropical freshwater swamp forest prefer wet microhabitats. Journal of Orthoptera Research. 26(1): 73-80. https://doi.org/10.3897/jor.26.14551

- Uvarov, B. 1977. Grasshoppers and Locusts. A Handbook of General Acridology. Vol.2. Cambridge, United Kingdom, London: Centre for Overseas Pest Research; pp. 613.
- Verdcourt, B. 1947. A note on the food of *Acridium* Geoff. (Orthopt.) The Entomologist's Monthly Magazine. 83:190.
- Vicari, M. and D. R. Bazely. 1993. Do grasses fight back – the case for antiherbivore defenses. Trends in Ecology & Evolution.
 8: 137–141. https://doi.org/10.1016/0169-5347(93)90026-L
- Williams, L. H. 1954. The feeding habits and food preferences of Acridinae and the factors which determine them. Transaction

of the Royal Entomological Society of London. 105: 423-454.

- Yadav, R. S. and D. Kumar. 2017. Biodiversity of pyrgomorphid grasshoppers from eastern uttar Pradesh. Journal of Experimental Zoology of India. 20 (1): 1467-1470.
- Zhang, Z. Q. 2011. Animal biodiversity: An outline of higher-level classification and survey of taxonomic richness. Zootaxa. 3148: 1–237. https://doi.org/10.11646/zootaxa.3148.1.1
- Zhang, L., M. Lecoq, A. Latchininsky and D. Hunter. 2019. Locust and grasshopper management. Annual Review of Entomology. 64: 15–34. https://doi.org/10.1146/annurev-ento-011118-112500

Διατροφική Κατηγοριοποίηση ορισμένων ειδών Caelifera και Ensifera (Insecta: Orthoptera) από επιλεγμένους αγρούς στην πόλη Sendai της Ιαπωνίας

Abu ElEla Shahenda, A. A.^{1, 3}, Michio Kondoh², Masatoshi Hori³, Atsuhiko Nagasawa³, Koji Nakamura⁴, Wael M. ElSayed¹

¹Department of Entomology, Faculty of Science, Cairo University, Egypt 12613 ²Graduate School of Life Sciences, Tohoku University, 6-3 Aoba, Aramaki, Aoba-ku, Sendai 980-8578, Japan

³Graduate School of Agricultural Science, Tohoku University, Sendai, Miyagi 980-8572, Japan. Email: masatoshi.hori.a3@tohoku.ac.jp

⁴Ishikawa Prefectural Natural History Museum, Kanazawa, Ishikawa 920-1147, Japan

ΠΕΡΙΛΗΨΗ

Αν και τα είδη των ορθοπτέρων έχουν συχνά θεωρηθεί ως πολυφάγα φυτοφάγα, τα περισσότερα από αυτά παρουσιάζουν ποικίλους βαθμούς επιλεκτικότητας και ιδιαίτερες προτιμήσεις ενδιαιτήματος. Επιπλέον, ορισμένα είδη παρουσιάζουν ιδιόμορφες προτιμήσεις διατροφής που χρήζουν περαιτέρω διερεύνησης. Δείγματα από ένα σύνολο 43 ειδών ορθοπτέρων, από 7 οικογένειες, 12 υποοικογένειες και 13 φυλές, από πέντε αγρούς στην πόλη Σεντάι της Ιαπωνίας, συλλέχθηκαν για να διερευνηθούν οι διατροφικές τους προτιμήσεις, και ταξινομήθηκαν σε επτά κύριες διατροφικές κατηγορίες με βάση την εξέταση των μορφολογικών χαρακτηριστικών της γνάθου και αναλύσεις του μεταθανάτιου περιεχομένου του εντέρου.