



# **ENTOMOLOGIA HELLENICA**

Vol 33, No 2 (2024)

#### Entomologia hellenica 33(2)



To cite this article:

LEBBAL, S., BENHIZIA, T., PÉREZ-SÁNCHEZ, R., & Remedios Morales Corts, M. (2024). Insecticidal activity of garden waste compost tea against Aphis craccivora (Koch) aphids : In vitro and in silico study. *ENTOMOLOGIA HELLENICA*, *33*(2), 78–89. https://doi.org/10.12681/eh.37918

# Insecticidal activity of garden waste compost tea against *Aphis* craccivora (Koch) aphids: In vitro and in silico study

## SALIM LEBBAL<sup>\*1</sup>, TOUFIK BENHIZIA<sup>1</sup>, RODRIGO PÉREZ-SÁNCHEZ<sup>2</sup> & MARÍA REMEDIOS MORALES-CORTS<sup>2</sup>

<sup>1</sup>Abbas Laghrour University, Department of Agricultural Sciences, Laboratory of Management, Conservation and Valorization of Agricultural and Natural Resources (LMCVANR), Khenchela/Algeria

<sup>2</sup>University of Salamanca, Faculty of Agricultural and Environmental Sciences, Salamanca/ Spain

## ABSTRACT

Aphids are important pests of many crops. However, the use of chemical insecticides has provoked ecological and health problems, thus the valuation of natural products becomes an interesting alternative. Compost teas are organic products that are viewed as substitutes for common pesticides. The present study aims to screen the potential insecticidal effect of garden waste compost tea, through *in vitro* and *in silico* approaches. Five concentrations of a compost tea (20, 40, 60, 80 and 100%) were tested against *Aphis craccivora*, in comparison to negative and positive controls. The repellency and toxicity tests were conducted under laboratory conditions. Moreover, an evaluation of the inhibitor capacity of some compost compounds against acetylcholinesterase was carried out using molecular docking. Results revealed that compost had a very weak insecticidal effect against *A. craccivora* (where the corrected mortality did not exceed 24%) compared to the tested chemical pesticide. Furthermore, the repellency test showed that the compost had some repellency effect in comparison to the tested chemical pesticide which had an attractant effect. Concerning the results of the molecular docking, pirimicarb (active molecule of pesticides) recorded a better S score than the three compost compounds.

KEY WORDS: Insect pests, repellency, toxicity, bioinsecticides, molecular docking.

#### Introduction

Biotic stress in plants caused by insect pests is one of the most significant problems, leading to yield losses (Tlak Gajger & Dar, 2021). These insect pests belong to different groups, including Aphididae. Among its important species, *Aphis craccivora* attacks about 50 crops in 19 different plant families (Blackman & Eastop, 2007). It causes damage by direct feeding as well as by transmission of plant viruses causing diseases such as broad bean yellow mosaic and bean leafroll (Weigand & Bishara, 1991).

Synthetic pesticides are being widely used for the management of pests to avoid losses both in field and post-harvest storage (Kumar et al., 2022). The estimated 30% losses from pests that would occur in the absence of pesticides, would spell economic and human disaster for many developing countries around the world (Saxena et al., 2014). However, global concern has been raised in recent times against the utility of synthetic insecticides in households and fields (Barua et al., 2016). Pesticides accumulate toxic residues on food grains used for human consumption and this may lead to health issues, in addition to the very high worldwide mortality rate due to

\*Corresponding authors: <a href="mailto:salim-leb@hotmail.com">salim-leb@hotmail.com</a>

pesticide poisoning (Singh & Kaur, 2018). Besides, pesticide residues may constitute a significant source of contamination of air, water and soil, and a large quantity of pesticides is released into the atmosphere during application, thereby inducing adverse climatic changes (Saxena et al., 2014). That's why for the future, it is necessary to develop a more environmentally friendly agriculture that will decrease inputs in chemicals and generate fewer harmful such pesticide residues outputs as (Wójcicka, 2010).

Consequently, identification and exploring for nature-originating pest control agents could become a possible substitute for the synthetic chemical pesticides (Kumar et al., 2022). Natural insecticides are chemical compounds or substances obtained from living organisms (Barua et al., 2016). Natural compounds known for their bioactivity against insects are considered safe, economical, biodegradable and easy to use (Singh and Kaur 2018). Among these, compost and compost teas have been shown, in previous studies, to possess control properties against a wide range of different plant pests and diseases (Bandara et al., 2010; Edwards et al., 2010; Alao et al., 2011; Shalaby et al., 2012; Pane et al., 2012; Martín, 2014; López-Martín et al., 2018; Morales-Corts et al., 2018; Suwandi et al., 2020: González-Hernández et al., 2021, 2022, among others). Concretely, compost teas are organic solutions obtained by the fermentation of compost in a liquid phase for a few days, with or without aeration (Al-Dahmani et al., 2003). The physical and biochemical quality of compost teas depends on the characteristics of the starting compost, as well as on other parameters that affect its production i.e., compost-to-water and aeration (Ingham, 1999: ratio. Scheuerell & Mahaffee, 2002; Martin et al., 2012).

On the other hand, the application of computational modeling guided the experimental procedure by elucidating the mechanisms of ligand-enzyme binding and greatly reducing the research time and costs (Badawy et al., 2022). Molecular docking is an in silico method which predicts the placement of small molecules or ligands within the active site of their target protein (Surabhi & Singh, 2018). The nature of the interaction between ligand and receptor balance depends in the on а chemical/physical forces between them and the forces between each of these molecules and the solvent or environment (Krumrine et al., 2003).

Thus, the current study's objective is to evaluate *in vitro* the effectiveness of compost tea solutions against *Aphis craccivora*, as well as to screen the insecticidal activity of its components, using molecular docking approach.

## **Materials and Methods**

#### **Compost tea preparation**

The compost tea (CT) derived from composting gardening wastes (mainly a mixture of green and pruning debris, i.e., mainly leaves and branches of cypress (Cupressus sp.), willow (Salix sp.) and poplar trees (Populus alba L.), reaching a C/N ratio of 30), which were obtained from public gardens of the Province of Salamanca (Spain). This process was performed in aerated piles of  $15 \times 2 \times 2$  m for 180 days in a garden center located in Salamanca (Spain) (40°57'23" N; 5°41'8" W, 775 m a.s.l.). Piles were turned twice per week for eight weeks and once a week during the rest of the bioxidative process. Moreover, the moisture of the piles was controlled once a week, and when it dropped below 55%, water irrigation was applied up to maintain the moisture average around 60% during the whole process. The mature compost was obtained under ambient conditions in March. The main characteristics of the garden waste compost were: pH 7.8, C/N 11.4, 41.4% total organic matter, 2.1% total N, 10.90% humic acids and 0.33 dS/m Electrical conductivity (Registration No. F0004957/2031, Viveros El Arca enterprise, Salamanca, Spain).

Compost was mixed with tap water in a ratio of 1:5 (v/v) in polyethylene nondegradable 1000 L containers at room temperature (20 °C) for a brewing period of five days. Water had been previously aerated for 8 h to reduce chlorines concentration. The mixture was aerated for five hours every day by applying circular stirring and making fine bubbles of air using a pump (750 W-300 rpm). Then, it was filtered with a doublelavered cheesecloth, and the aerated compost tea (CT) was stored in a dark container (50 L capacity at room temperature) until use.

# Chemical and microbiological properties of the compost tea

The chemical properties of the CT were directly analyzed; pH. electrical conductivity (EC) and C:N ratio were determined as described by Morales-Corts et al. (2021). Furthermore, assimilable nutrient contents (NO<sup>3-</sup>, PO<sub>4</sub><sup>3-</sup>, K<sub>2</sub>O, SO<sub>4</sub><sup>2-</sup>, Ca<sup>2+</sup> and Mg<sup>2+</sup>) were analyzed with the nutrient analysis photometer HANNA HI 83225. Finally, humic acids were determined following the alkali-acid method described by Pant et al. (2012). Salicilic acid and indoleacetic acid (IAA) were quantified by mass spectrometry (HPLC).

Microbial analysis of the CT was estimated using the serial dilution spread plate method. To determine the microbial population, different selective culture media CT dilutions were and used for microorganism isolation: nutrient agar and  $10^{-3}$  dilution for total aerobic bacteria, Ashby medium, and 10<sup>-2</sup> dilution for Nfixing bacteria, ISP<sup>-2</sup> medium and 10<sup>-1</sup> dilution for actinobacteria and modified potato dextrose agar medium but no dilution for total fungi and Trichoderma ssp. quantification (Wickerham, 1951; Sanchis Solera, 1996; Stella & Suhaimi, 2010; Vargas-Gil et al., 2006). Then, plates were inoculated by depositing on the agar surface

0.1 mL of the CT dilution, which was spread on the media surface with sterile glass beads. Moreover, non-inoculated plates were included as a negative control. Petri dishes were incubated in the dark at 28 °C for 3 to 15 days, depending on the medium. After this time, colony-forming units (CFU) were estimate the counted to cultivable microorganism's population. This experiment was conducted using five replications.

#### In vitro assays

Five CT concentrations (20, 40, 60, 80 and 100%) were tested, in addition to the negative control (distilled water) and a positive control represented by a chemical aphicide 'Lazer' (produced by Ortiva, represented in Algeria by Phytoplus and registered under the homologation number 07 45 036) containing two active molecules (Lambda-cvhalothrine and Pirimicarb. 5%+10%, which are sodium channel acetylcholinesterase modulators and inhibitor, respectively (IRAC, 2022)).

Faba bean leaves, bearing *Aphis craccivora* aphids (of the same colony), were collected from a commercial field, that had not undergone any chemical treatment. The leaves along with wingless insects were left for few hours in the plastic box. Aphids were then collected by means of a small brush and used for the bioassay.

For screening the insecticidal activity under laboratory conditions (T = 25 °C; Relative Humidity= 36%), cut pieces of leaves were dipped in each concentration of the compost for 2-5 seconds and then the solution was drained out and pieces were kept for drying. Three replications of each concentration along with positive and negative controls were taken for bio-assay studies. Treated leaves were kept in the Petri dish with 10 aphids in each replicate. Mortality was observed after 24 h of treatment. Percent mortality was corrected through Abbots formula (1925):

Corrected mortality rate =  $[(Tmp - Cmp)/(100 - Cmp)] \times 100$ 

where Tmp: mortality rate of the treatment and Cmp: mortality rate on the control.

Furthermore, ANOVA analysis followed by Student-Newman-Keuls test were performed by the means of SPSS software, to compare mean aphid mortalities.

Concerning the test of repellency in vitro. three replications for each concentration. in addition the to conventional pesticide, were considered. For each replicate, two leaf discs free of aphids were cut. After that, one leaf disc was dipped for 2-5 seconds in the corresponding solution. Each Petri dish was divided into two equal parts. Leaf discs dipped in treatment solution were placed in one side and untreated ones on the other side. 10 wingless aphids were placed at the center of the dish and left for 24 hours and then the number of aphids on each side was recorded. The repellency percentage was calculated according to the following equation:

 $PR = [(C - T) / (C + T)] \times 100 \text{ (Pavela et al.,} 2009)$ 

where: C = the number of aphids on control disc, and T = the number of aphids on treated disc.

#### In silico study

On the other hand, the evaluation of the insecticidal activity by molecular docking was carried out using the Molecular Operating Environment (MOE) software, version 2014.0901, designed by Chemical Computing Group Inc. (Canada). We studied *in silico* the ability of four ligands, including three compounds of the examined compost (salicylic acid, humic acid and indole-3-acetic acid), in addition to an active molecule of chemical pesticide (Pirimicarb), to inhibit a target enzyme of insects called acetylcholinesterase (AChE), having the PDB code 1QON.

#### Results

#### **Compost tea properties**

The studied characteristics (pH, EC, C-to-N ratio, assimilable macro and micronutrients, and humic acids) are shown in Table 1. Highlighted are the high NO3– and K2O concentrations (2240.4 and 2851.2 ppm, respectively) and the low C-to-N ratio of the extract, as well as the relevant humic acids amount.

Concerning the microbiological analysis of the compost tea, its microbiological composition was: total aerobic bacteria  $2.7 \times 107$  cfu mL-1; N-fixing bacteria  $2.7 \times 107$ cfu mL-1; actinobacteria  $7.4 \times 104$  cfu mL-1; Trichoderma sp. and fungi were valued between 2.7 and  $8.7 \times 102$  cfu mL-1.

#### In vitro assays

Results revealed that for the test of toxicity in vitro, there was a significant difference between the studied treatments. It seems that compost had very weak insecticidal effect against A. craccivora comparatively with the tested chemical pesticide. The corrected mortality did not exceed 24% for the different concentrations of compost tea (Table 2).

Results of the test of repellency in vitro showed that compost had some repellency effect against A. craccivora in comparison with the tested chemical pesticide which seems to have attractant effect (Table 3). All the tested concentrations of the compost were ranged in the class I of scale described by McDonald et al. (1970).

#### In silico study

Concerning the results of the molecular docking, the best S score was recorded for Pirimicarb (active molecule of pesticides) followed by indole-3-acetic acid (Table 4). This result confirms that obtained for in vitro test, i.e. the tested pesticide was more toxic than compost solutions.

81

<sup>\*</sup>Corresponding authors: <a href="mailto:salim-leb@hotmail.com">salim-leb@hotmail.com</a>

| pН       | EC      | C/  | NO <sub>3</sub> <sup>-</sup> (ppm) | $P_2O_5$  | K <sub>2</sub> O (ppm) | $SO_4^2$ - | Ca <sup>2+</sup> | Mg <sup>2+</sup> | Humic  | Salicilic   | IAA    |
|----------|---------|-----|------------------------------------|-----------|------------------------|------------|------------------|------------------|--------|-------------|--------|
|          | (dS/m)  | N   |                                    | (ppm)     |                        | (ppm)      | (ppm)            | (ppm)            | acids  | acid (mg/L) | (ng/L) |
|          |         |     |                                    |           |                        |            |                  |                  | (mg/L) |             |        |
| 7.16±0.1 | 1.2±0.0 | 7.1 | 2240.4±186.                        | 149.7±18. | 2851.2±234.            | 43±16      | 280±2            | 20±13            | 198±3  | 5.85±1.23   | 80±10  |
| 4        | 9       |     | 6                                  | 3         | 5                      |            | 0                |                  | 7      |             |        |

**TABLE 1:** Chemical characteristics of the studied compost tea. *Results are indicated as mean* ± standard deviation

TABLE 2: Corrected mortality percentages of aphids on different solutions

| Corrected mortality (M $\pm$ SE) |  |  |
|----------------------------------|--|--|
| $0.00 \pm 0.00$ a                |  |  |
| $0.00\pm0.00~\mathrm{a}$         |  |  |
| $23.70 \pm 7.73$ a               |  |  |
| $4.63 \pm 3.34$ a                |  |  |
| $9.72 \pm 9.72$ a                |  |  |
| $100.00 \pm 0.00 \; b$           |  |  |
| 0,000 *                          |  |  |
|                                  |  |  |

\* Signification at P < 0.05

| Treatments         | Percentages of Repellency $(M \pm SE)$ |
|--------------------|--|
| 20 %               | $8.15 \pm 39.34$                       |
| 40 %               | $17.04 \pm 11.92$                      |
| 60 %               | $6.67\pm48.07$                         |
| 80 %               | $18.52 \pm 29.60$                      |
| 100 %              | $10.37 \pm 5.78$                       |
| Chemical pesticide | $-14.07 \pm 2.96$                      |

**TABLE 3:** Percentages of repellency of aphids by different solutions

#### **TABLE 4:** S score of the studied ligands with the target protein 1QON

|              |                      |                                   | S score                   |
|--------------|----------------------|-----------------------------------|---------------------------|
| PubChem code | Ligands              | Properties                        | (Kcal mol <sup>-1</sup> ) |
| 31645        | Pirimicarb           | Inhibitor of acetylcholinesterase | - 6.42                    |
| 802          | Indole-3-acetic acid | Compound of compost tea           | - 5.23                    |
| 90472028     | Humic acid           | Compound of compost tea           | - 4.30                    |
| 338          | Salicylic acid       | Compound of compost tea           | - 4.64                    |

83

#### Discussion

The extract's low C:N ratio, high NO<sup>3-</sup> and K<sub>2</sub>O concentrations, and high humic acid content were important characteristics of the studied compost tea. There are numerous factors that affect the quality of vermicompost tea, which in-turn affect its ability for disease and pest suppressiveness, among them are included compost grade. compost maturity, aeration, temperature, microbial inoculants, and compost to water ratio (Yatoo et al., 2021). For instance. Nur et al. (2023) indicated differences in the C:N ratio and humic acid concentration between aerated and non-aerated compost tea made from wild lotus. Orosz et al. (2021) reviewed many mechanisms of action of different compost extract components allowing control of plant diseases. There is a lot of published literature concerning studies regarding the effects of humic acids and different biostimulants on the growth, productivity, and protection of vegetable species is available, whereas literature concerning the study of the effects of compost teas on vegetable crops is more limited (Pilla et al., 2023).

On the other hand, the present investigation revealed that the compost had weak insecticidal effect against A. *craccivora*, in comparison to the chemical pesticide containing pirimicarb, as well as limited repellency activity. Similarly, Abdu-Allah (2012) indicated that the toxicity index of pirimicarb shows higher aphicidal activity than three petroleum ether plant extracts. Additionally, Cantelo (1985) found that compost treated with experimental insecticides affecting acetylcholinesterase (such as diazinon and dimethoate) were effective against the dipteran pest Megaselia halterata.

In contrast, Bandara et al. (2010) found that brinjal aphids, mealy bugs and thrips were significantly reduced in treated plants with different compost solutions, compared to the controls. Besides, other results have shown that organic compost liauid formulations acted as antifeedants for Podagrica sp., Zonocerus variegatus and Bemisia tabaci. while maize stover compost had a very good insect control (Alao et al., 2011). The compost extract containing amino acids showed a remarkable potential to develop into an effective biostimulant for protection from virus disease and its insect vector. Aphis gossypii (Suwandi et al., 2020).

Moreover, insect pests of the plant species Telfairia occidentalis that received foliar spray of compost extracts (from cassava peel and tithonia plant) were minimal compared with non-fertilized plants and those that received soil incorporated NPK fertilizer (Akanbi et al., 2007). Furthermore, Shalaby et al. (2012) mentioned that compost tea used as soil drench applied after sowing, followed by foliar application of compost tea proved to be the best treatment against insect infestations by Pegomia mixta Vill, Scrobipalpa ocellatella Boyd and Cassida vittata Vill. Moreover, Edwards et al. (2010) indicated that all examined vermicompost extracts suppressed pest establishment on the plants, and their rates of reproduction for all three species of pests, significantly. The same researchers concluded that the most possible cause for the unacceptability of the plants to pests the uptake of soluble phenolic was materials from the vermicompost extracts into the plant tissues and these compounds are known to make plants unattractive to pests.

Likewise, previous studies confirmed the influence of compost as soil fertilizer on aphid populations (Ponti et al., 2007; Stafford et al., 2012).

On the other hand, the results of the molecular docking were consistent with the in vitro toxicity assay. The pirimicarb molecule proved to be the more effective in

inhibition of acetylcholinesterase, the recording the best S score (-6.42 kcal mol<sup>-1</sup>), compared to the compost compounds. Other in silico studies showed not very different S score results. For instance, Lebbal et al. (2021) noticed that  $\alpha$ -Thuiene ligand has the best score with 1QON target enzyme (-4.77), among seven compounds contained in Thymus algeriensis and T. numidicus, whereas, Badawy et al. (2022) indicated the docking energies in the AChE enzyme of -9.52, -9.38, and -8.52 kcal mol-1 for chlorpyrifos-methyl, diazinon. and malathion, respectively. Furthermore, Malak et al. (2022) found that among 38 studied bioactive phytochemicals, some compounds

#### References

- Abbott, W.S. (1925). A method of computing the effectiveness of an insecticide. Journal of Economic Entomology, 18, 265-267.
- Abdu-Allah, G. (2012). Aphicidal activity of imidacloprid and primicarb compared with certain plant extracts on *Brevicoryne brassicae* L. and *Aphis craccivora* Koch. Assiut Journal of Agricultural Sciences, 43(1), 108-118.
- Alao, F.O., Adebayo, T.A., Olaniran, O.A., & Akanbi, W.B. (2011). Preliminary evaluation of the insecticidal potential of organic compost extracts against insect pests of Okra (*Abelmoschus esculentus* (L.) Moench). Asian Journal of Plant Science and Research, 1(3), 123-130.
- Al-Dahmani, J.H., Abbasi, P.A., Miller, S.A., & Hoitink, H.A.J. (2003). Suppression of bacterial spot of tomato with foliar sprays of compost extracts under greenhouse and field conditions. Plant Diseases, 87, 913– 919.

recorded docking scores of 9.11 to 7.14 against cattle tick *Rhipicephalus microplus* acetylcholinesterase protein.

#### Conclusion

In summary, the present study confirmed through in vitro and in silico screenings that the compost had low aphicidal effect. In addition, it has better repellent activity compared with the tested chemical aphicide. Nevertheless, its use as fertilizer should be combined with other control methods to reduce efficiently the negative impact of aphid populations.

- Akanbi, W. B., Adebayo, T. A., Togun, O. A., Adeyeye, A. S., & Olaniran, O. A. (2007). The use of compost extract as foliar spray nutrient source and botanical insecticide in *Telfairia occidentalis*. World Journal of Agricultural Sciences, 3(5), 642-652.
- Badawy, M. E., Abd-Elnabi, A. D., & Saad, A. F. S. (2022). Insecticidal activity of nanoemulsions of organophosphorus insecticides against cotton leafworm (*Spodoptera littoralis*) and molecular docking studies. International Journal of Tropical Insect Science, 42(1), 293-313.
- Bandara, P., Wijayagunasekara, H. N. P., & Sirisena, U. G. A. I. (2010).
  Evaluation of the efficacy of compost extracts incorporated with insecticidal plant materials. Research Symposium, 30.
- Barua, S., Gopalakrishnan, R., & Veer, V. (2016). Extraction, purification and characterisation of insecticidal compounds from plants. In V. Veer & R. Gopalakrishnan (Eds.), Herbal Insecticides. Repellents and Biomedicines: Effectiveness and Commercialization (pp. 101-115). Springer, New Delhi.

- Blackman R.L., & Eastop V.F. (2007). Taxonomic issues. In: van Emden H.F. &
- Harrington R. (eds.), Aphids as Crop Pests, CAB International, UK, pp. 1-29.
- Cantelo, W. W. (1985). Control of *Megaselia halterata*, a phorid fly pest of commercial mushroom production, by insecticidal treatment of the compost or casing material. Journal of Entomological Science, 20(1), 50-54.
- Edwards, C. A., Arancon, N. Q., Vasko-Bennett, M., Askar, A., Keeney, G., & Little, B. (2010). Suppression of green peach aphid (*Myzus persicae*) (Sulz.), citrus mealybug (*Planococcus citri*) (Risso), and two spotted spider mite (*Tetranychus urticae*) (Koch.) attacks on tomatoes and cucumbers by aqueous extracts from vermicomposts. Crop Protection, 29(1), 80-93.
- González-Hernández, A.I., Suárez-Fernández, M.B., Pérez-Sánchez, R., Gómez-Sánchez, M.A., Morales-Corts, M.R. (2021). Compost tea induces growth and resistance against *Rhizoctonia solani* and *Phytophthora capsici* in pepper. Agronomy, 11, 781.
- González-Hernández, A.I., Pérez-Sánchez, R., Plaza, J., Morales-Corts, M.R. (2022) Compost tea as a sustainable alternative to promote plant growth and resistance against *Rhizoctonia solani* in potato plants. Scientia Horticulturae, 300, 111090.
- Ingham, E.R. (1999). What is compost tea? Part I. Biocycle, 40, 74–75.
- IRAC (2022) Mode of action classification scheme: version 10.4. CropLife, 41 p.
- Krumrine, J., Raubacher, F., Brooijmans, N. & Kuntz, I. (2003) Principles and methods of docking and ligand design.
  In: P.E. Bourne & H. Weissig (Eds.), Methods of Biochemical Analysis,

John Wiley & Sons, Inc. New Jersey, pp. 443-476.

- Kumar, S., Mahapatro, G.K., Yadav, D.K., Tripathi, K., Koli, P., Kaushik, P., Sharma, K. & Nebapure, S. (2022). Essential oils as green pesticides: An overview. Indian Journal of Agricultural Sciences, 92(11), 1298– 1305.
- Lebbal, A., Rahal, K. & Lebbal, S. (2021). In silico assessment of insecticidal activity of *Thymus algeriensis* and *T. numidicus*. In: Mediterranean Forum for PhD Students and Young Researchers, 2021/07/06-07, Online -CIHEAM Montpellier, France. Proceedings, pp 98-99.
- López-Martín, J. J., Morales-Corts, M. R., Pérez-Sánchez, R., & Gómez-Sánchez, M. Á. (2018). Efficiency of garden waste compost teas on potato growth and its suppressiveness against *Rhizoctonia*. Poljoprivreda i Sumarstvo, 64(4), 7-14.
- Malak, N., Niaz, S., Wadood, A., Nasreen, N., Ali, I., Iqbal, J., Swelum, A.A., Alkahtani, M.A., Zając, Z., & Khan, A., 2022. In silico approaches to develop herbal acaricides against *R*. (*Boophilus*) *microplus* and in vitro anti-tick activities of selected medicinal plants. Saudi Journal of Biological Sciences, 29(6), p.103302.
- Martín, C.C.G., Dorinvil, W., Brathwaite, R.A.I., Ramsubhag, A. (2012). Effects and relationships of compost type, aeration and brewing time on compost tea properties, efficacy against *Pythium ultimum*, phytotoxicity and potential as a nutrient amendment for seedling production. Biological Agriculture & Horticulture, 28, 185– 205.
- Martín, C.C.G. (2014) Potential of compost tea for suppressing plant diseases. CAB Review, 9, 1–38.

- McDonald, L.L., Guy, R.H. & Speirs, R.D. (1970). Preliminary evaluation of new candidate materials as toxicants, repellents, and attractants against stored-product insects. Marketing Research Report No. 882. US Agricultural Research Service: USA. 8 p.
- Morales-Corts, M. R., Pérez-Sánchez, R., & Gómez-Sánchez, M. Á. (2018). Efficiency of garden waste compost teas on tomato growth and its suppressiveness against soilborne pathogens. Scientia Agricola, 75, 400-409.
- Nur, T. P., Priatna, S. J., & Gofar, N. (2023). Assessing the Quality of Compost Tea Made from Swamp-Growing Lotus Plants. Journal of Smart Agriculture and Environmental Technology, 1(3), 78-83.
- Orosz, V., Tomócsik, A., Demeter, I., Aranyos, T. J., & Makádi, M. (2021). Control of plant pathogen Fusarium spp. with compost, compost tea application–A review. COLUMELLA–Journal of Agricultural and Environmental Sciences, 8(2), 55-71.
- Pane, C., Celano, G., Villecco, D., & Zaccardelli, M. (2012). Control of *Botrytis cinerea*, Alternaria alternata and Pyrenochaeta lycopersici on tomato with whey compost-tea applications. Crop Protection, 38, 80-86.
- Pant, A.P., Radovich, T.J., hue, N.V., & Paull, R.E. (2012). Biochemical properties of compost tea associated with compost quality and effects on pak choi growth. Scientia Horticulturae, 148, 138-146.
- Pavela, R., Vrchotová, N., & Šerá, B. (2009). Repellency and toxicity of three Impatiens species (Balsaminaceae) extracts on *Myzus*

*persicae* Sulzer (Homoptera: Aphididae). Journal of Biopesticides, 2(1), 48-51.

- Pilla, N., Tranchida-Lombardo, V., Gabrielli, P., Aguzzi, A., Caputo, M., Lucarini, M., Durazzo, A. & Zaccardelli, M. (2023). Effect of compost tea in horticulture. Horticulturae, 9(9), 984.
- Ponti, L., Altieri, M. A., & Gutierrez, A. P. (2007) Effects of crop diversification levels and fertilization regimes on abundance of *Brevicoryne brassicae* (L.) and its parasitization by *Diaeretiella rapae* (M'Intosh) in broccoli. Agricultural and Forest Entomology, 9(3), 209-214.
- Sanchis Solera, J. (1996). Comparación entre los diversos medios de cultivo comerciales para aislamiento de hongos (levaduras y mohos). In Proyecto Microkit 1999 Para Optimizar la Sensibilidad de los Parámetros de Muestras Microbiológicos del aire: Laboratorios Microkit, S.L.: Madrid, Spain.
- Saxena, H.O., Tripathi, Y.C., Pawar, G., Kakkar, K. & Mohammad, N. (2014). Botanicals as biopesticides: active chemical constituents and biocidal action. In: P.K., Khatri & P.B. Mesharm, (Eds.), Familiarizing with local biodiversity notes on systematics of plants and insects. Tropical Forest Research Institute: India, pp 219-240.
- Scheuerell, S., Mahaffee, W. (2002). Compost tea: principles and prospects for plant disease control. Compost Science & Utilization, 10, 313–338.
- Shalaby, G. A., El-Gizawy, E. S., & El-Magd, A. (2012). Effect of mineral nitrogenous fertilization and compost tea on insect infestation of sugar beet and yield characteristics. Journal of

Plant Protection and Pathology, 3(8), 825-834.

- Singh, B. & Kaur, A. (2018). Control of insect pests in crop plants and stored food grains using plant saponins: A review. Lwt Food Science & Technology, 87, 93-101.
- Stafford, D. B., Tariq, M., Wright, D. J., Rossiter, J. T., Kazana, E., Leather, S. R., Ali, M., & Staley, J. T. (2012). Opposing effects of organic and conventional fertilizers on the performance of a generalist and a specialist aphid species. Agricultural and Forest Entomology, 14(3), 270-275.
- Stella, M., & Suhaimi, M. (2010). Selection of suitable growth medium for freeliving diazotrophs isolated from compost. Journal of Tropical Agriculture and Food Science, 38, 211–219.
- Surabhi, S., & Singh, B. K. (2018). Computer aided drug design: an overview. Journal of Drug delivery and Therapeutics, 8(5), 504-509.
- Suwandi, S., Irsan, C., Muslim, A., & Herlinda, S. (2020). Protection of chili pepper from mosaic virus disease and *Aphis gossypii* by a fermented water extract of compost. IOP Conference Series: Earth and Environmental Science, 468(1), p. 012043.

- Tlak Gajger, I., & Dar, S. A. (2021). Plant allelochemicals as sources of insecticides. Insects, 12(3), 189.
- Vargas-Gil, S.; Pastor, S.; March, G.J. (2006). Quantitative isolation of biocontrol agents *Trichoderma* spp., *Gliocladium* spp. and actinomycetes from soil with culture media. Research in Microbiology, 164, 196– 205.
- Weigand S. & Bishara S.I. (1991) Status of insect pests of faba bean in the Mediterranean region and methods of control. Options Méditerranéennes: Série A., 10, 67-74.
- Wójcicka, A. (2010). Cereal phenolic compounds as biopesticides of cereal aphids. Polish Journal of Environmental Studies, 19(6), 1337-1343.
- Wickerham, L.J. (1951). Taxonomy of yeasts. U.S. Dept. Agric. Technical Bulletin, 1029, 1–56.
- Yatoo, A. M., Ali, M. N., Baba, Z. A., & Hassan. B. (2021). Sustainable management of diseases and pests in vermicompost crops by and Α review. vermicompost tea. Agronomy for Sustainable Development, 41(1), 7.

## Εντομοκτόνος δράση του εκχυλίσματος κομπόστ απορριμμάτων κήπου κατά των αφίδων *Aphis craccivora* (Koch): *In vitro* και *in silico* μελέτη

## SALIM LEBBAL<sup>\*1</sup>, TOUFIK BENHIZIA<sup>1</sup>, RODRIGO PÉREZ-SÁNCHEZ<sup>2</sup> & MARIA REMEDIOS MORALES-CORTS<sup>2</sup>

<sup>1</sup>Abbas Laghrour University, Department of Agricultural Sciences, Laboratory of Management, Conservation and Valorization of Agricultural and Natural Resources (LMCVANR), Khenchela/Algeria

<sup>2</sup>University of Salamanca, Faculty of Agricultural and Environmental Sciences, Salamanca/Spain

#### ΠΕΡΙΛΗΨΗ

Οι αφίδες αποτελούν σημαντικούς εχθρούς πολλών καλλιεργειών. Η χρήση χημικών εντομοκτόνων έχει προκαλέσει σημαντικά περιβαλλοντικά προβλήματα και προβλήματα στην ανθρώπινη υγεία. Έτσι, η αποτίμηση των φυσικών προϊόντων αποτελεί μια ενδιαφέρουσα εναλλακτική. Τα εκχυλίσματα κομπόστ (τσάγια κομποστοποίησης) είναι φυσικά προϊόντα που θεωρούνται υποκατάστατα της χρήσης κοινών φυτοφαρμάκων. Η παρούσα μελέτη στογεύει να εξετάσει την πιθανή εντομοκτόνο δράση του εκχυλίσματος κομπόστ απορριμμάτων κήπου, μέσω προσεγγίσεων in vitro και in silico. Πέντε συγκεντρώσεις του εκχυλίσματος κομπόστ (20, 40, 60, 80 και 100%) δοκιμάστηκαν έναντι των αφίδων Aphis craccivora, σε σχέση με αρνητικούς και θετικούς μάρτυρες. Οι δοκιμές απώθησης και τοξικότητας πραγματοποιήθηκαν σε εργαστηριακές συνθήκες. Επιπλέον, πραγματοποιήθηκε αξιολόγηση της ικανότητας αναστολής ορισμένων ενώσεων κομποστοποίησης έναντι της ακετυλοχολινεστεράσης με χρήση μοριακής σύνδεσης. Τα αποτελέσματα αποκάλυψαν ότι το κομπόστ είχε πολύ ασθενή εντομοκτόνο δράση του A. craccivora (όπου η διορθωμένη θνησιμότητα δεν ξεπέρασε το 24%) σε σύγκριση με το γημικό εντομοκτόνο μάρτυρα. Επιπλέον, η δοκιμή απώθησης έδειξε ότι το κομπόστ είχε κάποια απωθητική δράση σε σύγκριση με το χημικό εντομοκτόνο μάρτυρα που φαίνεται να έχει ελκυστική δράση. Όσον αφορά τα αποτελέσματα της μοριακής δέσμευσης, το pirimicarb (δραστική ουσία του γημικού εντομοκτόνου) κατέγραψε καλύτερη βαθμολογία S από τις τρεις ενώσεις κομποστοποίησης.