


ENTOMOLOGIA HELLENICA

Vol 34, No 1 (2025)

Entomologia hellenica 34(1)



Evaluation of insecticides against the cotton bollworm *Helicoverpa armigera* (Hubner) (Lepidoptera: Noctuidae) under field conditions in Middle Awash, Ethiopia

Zemedkun Alemu

doi: [10.12681/eh.38352](https://doi.org/10.12681/eh.38352)

Copyright © 2025, Zemedkun Alemu



This work is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike 4.0](https://creativecommons.org/licenses/by-nc-sa/4.0/).

To cite this article:

Alemu, Z. (2025). Evaluation of insecticides against the cotton bollworm *Helicoverpa armigera* (Hubner) (Lepidoptera: Noctuidae) under field conditions in Middle Awash, Ethiopia. *ENTOMOLOGIA HELLENICA*, 34(1), 49-61.
<https://doi.org/10.12681/eh.38352>

Evaluation of insecticides against cotton bollworm, *Helicoverpa armigera* (Hubner) (Lepidoptera: Noctuidae) under field conditions in Ethiopia

Zemedkun Alemu*, Nurhusein Seid and Sileshi Getahun

Ethiopian Institute of Agricultural Research, Werer Agricultural Research Center, P.O. Box 2003, Addis Ababa, Ethiopia

ABSTRACT

The cotton bollworm complex presents a significant challenge to cotton farmers, resulting in substantial yield losses. While insecticides have been employed to manage this pest, ongoing research is crucial for developing new insecticides that mitigate the risk of resistance to current synthetic options. This study aimed to evaluate the effectiveness of selected insecticides against cotton bollworm through a field experiment conducted in the Middle Awash region. Utilizing a randomized complete block design, eight treatments were tested with four replications. Larval population, damage to fruiting bodies, yield, and yield component data were collected and analyzed. Results indicated that the application of various insecticides significantly ($P < 0.05$) influenced larval populations, fruiting body damage, and overall seed cotton yield. Notably, the combinations of profenofos 40% + alpha-cypermethrin 4% EC and profenofos + cypermethrin 44% EC resulted in yield increases of 88.5% and 76%, respectively, compared to the control group. These findings underscore the practical implications for cotton farmers in Ethiopia and similar regions facing pest challenges. The effective use of these insecticides could substantially reduce yield losses attributed to cotton bollworm infestations, thereby enhancing farmers' financial stability. However, further research should focus on the validation and demonstration of these insecticides under different agro-ecologies of the country.

KEY WORDS: *Fruiting damage, insecticide efficacy, insecticide mixture, seed cotton yield.*

Introduction

Cotton, *Gossypium hirsutum* L. (Malvaceae), is one of the world's most important cash crops. Farmers grow cotton for the fiber and the oil extracted from its seeds (Haider et al., 2015; Malinga and Laing, 2022). In sub-Saharan Africa, cotton production plays a crucial role in economic growth and rural development (Amanet et al., 2019). Ethiopia is a prominent producer of cotton, with extensive cultivation in various regions of the Awash Valley, Southern Rift Valley, Gambella, Humera,

and Metema (Merdassa et al., 2022). Despite its significance, Ethiopia accounts for only 5% of African cotton production (Melesse et al., 2019).

Cotton cultivation faces significant challenges due to various pests, including African bollworms, cotton aphids, thrips, whiteflies, jassids (Ermias et al., 2009), and cotton mealybugs (Nurhussen et al., 2020). Among these pests, the African bollworm, *Helicoverpa armigera* (Hubner), poses a particularly severe threat to cotton crops in Ethiopia. This pest can cause substantial damage during its larval stage by feeding on

the reproductive parts of cotton plants, resulting in hole formation and internal feeding (Geremew, 2004; Geremew and Ermias, 2006). Yield losses attributed to *H. armigera* can reach up to 60% (Geremew and Ermias, 2006). The pest population typically begins to increase in mid-June, peaking in August, necessitating timely planting and effective management practices from mid-June to mid-September in the Middle Awash Valley (Ermias et al., 2009).

Chemical control remains the primary management strategy for cotton pests in Ethiopia, accounting for approximately 43% of production costs (EIAR, 2017). Farmers frequently employ a range of chemicals, including pyrethroids, carbamates, and organophosphates, to combat *H. armigera* (Geremew, 2004). Typically, 10-12 pesticide applications are required throughout the growing season to protect crops from insect damage. However, reliance on insecticides with a single mode of action can lead to the development of resistance in *H. armigera* and other pests (Yongqiang et al., 2016). Resistance levels can vary from low to moderate among cotton pests (Geremew, 2004; Hussain et al., 2015; Tossou et al., 2019; Zemedkun et al., 2022).

Recent advancements in integrated pest management (IPM) emphasize on the importance of diversifying control strategies to mitigate resistance risks. This includes incorporation of biocontrol agents and cultural practices alongside chemical applications (Khan et al., 2020; Reddy et al., 2021). Studies have shown that combining biological control methods with conventional insecticides pest management efficacy may be enhanced, while reliance on chemicals is reduced (Bahlai et al., 2020). Additionally, ongoing research on novel insecticides is essential for developing effective rotation strategies that help manage pest resistance effectively (Salama et al., 2013).

Given the challenges posed by *H. armigera* and the need for sustainable pest management practices, this study aimed to assess the efficacy of selected commercially available insecticides against the cotton bollworm in the Middle Awash area. By evaluating new insecticides and their potential role within current pest management strategies, we aim to enhance accessibility and provide valuable insights for Ethiopian cotton farmers facing pest challenges.

Materials and Methods

Description of the study area

The experiments were conducted at the Werer Agricultural Research Center (WARC) under field conditions during the 2021 and 2022 seasons. WARC is located at 90° 20' 31" N and 400° 10' 11" E, 750 m above sea level. The study area has an inconsistent annual rainfall of 540 mm and mean maximum and minimum temperatures of 34.4 °C and 19.6 °C, respectively. The type of soil is vertisol, with a porosity of 49.06% and a bulk density of 1.35 gm/cm² (0-25 cm depth) (Wendmagegn and Abere, 2012).

Planting material

For the present study, the newly released cotton variety Werer-12 was obtained from the Werer Agricultural Research Center.

Insecticides

Seven insecticides that were recommended by the WARC for controlling cotton bollworms on cotton plants were used for the experiment (Table 1).

Treatments and experimental design

The seven insecticides and control were arranged in a randomized complete block design (RCBD) with four replications per treatment. Each plot covered a total area of 63 m². The spacing between rows was 90 cm, while the spacing between plants within a row was 20 cm.

TABLE 1: List of insecticides used in the present study.

Trt no.	Trade name	Common name	Chemical group	Rate (ml/ha)
1	Tutan 36% SC	chlorfenapyr	pyrole	225
2	Proof 44% EC	profenofos + cypermethrin	organophosphate + pyrethroid	2000
3	Agro-Lambacin Super 315 EC	profenfos 30% + lambda-cyhalothrin 1.5%	organophosphate + pyrethroid	400
4	Anticat 48% EC	chlorpyriphos 48% EC	organophosphate	2000
5	Testa	beta-cypermethrin 5% + emamectin benzoate 0.5% ME	pyrethroid + avermectins	750
6	Alhapro 440 EC	profenofos 40% + alpha-cypermethrin 4%	organophosphate + pyrethroid	750
7	Biotrine	abamectin + oxymatrin	abamectin + quinazaine alkaloid	2000

Experimental Procedures

Planting dates and methods. The agricultural field was meticulously prepared using a tractor-operated machine to ensure optimal planting conditions. Planting activities were carried out on May 15th in consecutive years to maintain consistency in the growing season. A carefully planned irrigation schedule was followed, with the plots receiving water eight times. The initial irrigation was followed by subsequent watering sessions at a 10-day interval, transitioning to a 15-day interval as the bolls approached 65% opening. Hoeing and hand-weeding procedures were conducted twice throughout the study to combat weed growth and maintain plot integrity. Additionally, all prescribed agronomic practices were diligently followed to promote healthy crop development and ensure uniformity across the experimental plots.

Data collection. Data collection for assessing *H. armigera* infestation in cotton fields was conducted systematically, starting three weeks after germination and continuing until crop maturity. In each plot,

a standardized sampling approach was followed, with five tagged plants monitored for pest activity on young shoot leaves, squares, flowers and bolls. Detailed observations were made on cotton bollworm eggs and larvae, damaged squares, flowers, and bolls, as well as the presence of non-target and beneficial insects. Data was collected both before and after spraying at specific intervals (3, 5, 7, and 10 days) to track changes in pest populations and assess the efficacy of control measures. Towards the end of the growing season, the number of healthy bolls per plant was quantified for the designated plants, including those in control plots, to evaluate the impact of pest infestation on yield. Finally, at harvest, the seed cotton plants were weighed to determine the overall crop productivity and the effectiveness of pest management strategies implemented throughout the season.

Data analysis. All data were analyzed using PROC GLM (SAS Version 9.0, SAS Institute, 1999). The PROC UNIVARIATE test was used to test the normality and homogeneity of variance of the data based on the Shapiro-Wilk test. To satisfy the

ANOVA assumptions, the pre-and post-spray count mean data were square root-transformed ($\sqrt{x+0.5}$). When F values were significant ($P < 0.05$), the means were compared using Fisher's least significant difference (LSD) test. The percent efficacy of each treatment was calculated based on the modified Abbott's formula (1925):

$$\% \text{ efficiency} = [1 - (Ta * Cb) / (Tb * Ca)]$$

where Ta = posttreatment population in treatment, Cb = pretreatment population in check, Tb = pretreatment population in treatment, and Ca = posttreatment population in check.

Results

The effectiveness of the tested insecticides in managing cotton bollworm infestations during the 2021 cropping season is summarized in Table 2. Significant differences ($P < 0.05$) were observed in the number of larvae emerging post-treatment, and the number of damaged squares and flowers across different treatments (Table 2). Notably, plots treated with profenofos + alpha-cypermethrin exhibited the lowest counts of larvae and damaged fruiting bodies, while untreated plots recorded the highest numbers (Table 2). Furthermore, the number of bolls per plant and seed yield showed highly significant differences ($P < 0.01$) (Table 2). Compared to the untreated control, applications of Alphapro 440 EC and Proof 44% EC led to remarkable increases in seed cotton yield of 95% and 80%, respectively. This was followed by chlorfenapyr (36% SC; chlorpyrifos, 48% EC (70%); profenofos, 30% + lambda-cyhalothrin, 1.5%; betacypermethrin, 5% + emamectin benzoate, 0.5% (60%); and abamectin + oxyamtrin (35% (Table 2).

In the 2022 season, significant differences ($P < 0.05$) were again noted in post-treatment mean larval counts, damaged squares, and seed yields among the treatments (Table 3). The combination

of profenofos + alpha-cypermethrin demonstrated the highest efficacy in controlling larvae, resulting in a 70% increase in seed cotton yield. In comparison, plots treated with abamectin + oxyamtrin showed the lowest efficacy with only a 22.5% yield increase (Table 3).

Overall, the combined results indicated that treatments significantly influenced post-treatment larval counts, damaged square counts, bolls per plant, and seed yield throughout the year ($P < 0.01$), although interaction effects were non-significant ($P > 0.05$) (Table 4). The application of profenofos + alpha-cypermethrin led to increases of 11%, 11%, 20%, 6%, 22%, 29%, and 69% in the number of bolls per plant across various treatments. Additionally, compared to the control, plots treated with profenofos + alpha-cypermethrin and 44% profenofos + cypermethrin achieved yield increases of 82% and 72%, respectively (Table 4). Other treatments, including Anticat, Tutan, Agro-Lambacin Super, Testa, and Biotrin, also contributed to yield enhancements of 69%, 65%, 52%, 47%, and 26%, respectively.

Discussion

The present study on the evaluation of the tested insecticides, particularly the combination of organophosphates and pyrethroids, against the cotton bollworm, *Helicoverpa armigera*, shows promising results in managing the cotton bollworm. Organophosphates, such as proenofos, function by inhibiting acetylcholinesterase, leading to the accumulation of acetylcholine at synapses, which causes continuous stimulation of the nervous system (Zhang et al., 2020). Confirming the findings of this investigation, Chambers et al. (2010), Timchalk (2010), and Carneiro et al. (2014) reported that organophosphorus pesticides have a complex metabolic pathway to reach their primary target, acetylcholinesterase.

Pyrethroids, like cypermethrin or alpha-

TABLE 2. Effects of tested insecticides on ABW population-damaged fruits and yield of cotton plants: Werer, 2021.

Treatment	Larval count/plant			No. of damaged squares/plant		No. of damaged flowers/plant		No. of damaged bolls/plant		Health boll/plant	Seed cotton yield (ton/ha)
	Pre-spray	Post-spray	Reduction %	Pre-spray	Post-spray	Pre-spray	Post-spray	Pre-spray	Post-spray		
chlorfenapyr 36 SC/Tutan	0.80 (1.14)	0.17 (0.82) ^{bcd}	80.56	1.30 (1.33)	0.58 (1.04) ^{bc}	0.40 (0.95)	0.12 (0.78) ^b	0.10(0.77)	0.06 (0.75)	30.6 ^{bc}	3.4 ^{ab}
profenofos + cypermethrin / Proof 44% EC	0.70 (1.10)	0.13(0.79) ^{cd}	84.44	1.13 (1.27)	0.41 (0.95) ^c	0.18 (0.82)	0.06 (0.75) ^b	0.05(0.74)	0.01 (0.71)	33.1 ^{ab}	3.6 ^a
profenfos 30% + lambda-cyhalothrin 1.5%	0.90 (1.18)	0.20 (0.84) ^{bcd}	79.26	1.45 (1.38)	0.58 (1.03) ^{bc}	0.33 (0.90)	0.08 (0.76) ^b	0.25(0.85)	0.08 (0.76)	28.8 ^{bc}	3.2 ^{ab}
chlorpyrifos 48%EC/Anticat	0.85 (1.16)	0.16 (0.81) ^{bcd}	82.61	1.48 (1.40)	0.61 (1.05) ^{bc}	0.28 (0.88)	0.08 (0.76) ^b	0.18(0.82)	0.09 (0.77)	32.5 ^{ab}	3.4 ^{ab}
beta-cypermethrin+emamectin/benzoate/Testa	0.85 (1.16)	0.21 (0.84) ^{bc}	77.12	1.63 (1.45)	0.67 (1.08) ^b	0.23 (0.84)	0.12 (0.78) ^b	0.13(0.79)	0.09 (0.77)	27.3 ^c	3.2 ^{ab}
profenofos + Alpha-cypermethrin/Alphapro	0.78 (1.13)	0.11 (0.78) ^d	86.95	1.03 (1.23)	0.42 (0.95) ^c	0.33 (0.92)	0.11 (0.78) ^b	0.10(0.77)	0.05 (0.74)	36.4 ^a	3.9 ^a
abamectin+oxymatrine/Biotrin	0.78 (1.13)	0.24 (0.86) ^b	71.40	1.40 (1.37)	0.57 (1.03) ^{bc}	0.33 (0.90)	0.13 (0.80) ^b	0.08(0.76)	0.07 (0.75)	26.1 ^c	2.7 ^{bc}
Unsprayed	0.70 (1.10)	0.75 (1.12) ^a	-	1.35 (1.34)	1.23 (1.32) ^a	0.28 (0.88)	0.35 (0.92) ^a	0.18(0.82)	0.21 (0.84)	19.1 ^d	2.0 ^c
LSD (0.05)	Ns	0.06(***)		Ns	0.12 (***)	Ns	0.06(***)	Ns	Ns	5.1 (***)	7.9 (**)
CV (%)	5.75	4.59		12.91	7.42	9.72	4.94	11.22	6.24	10.6	15.4
S.E _±	0.03	0..2		0.16	0.04	0.04	0.02	0.04	0.02	1.7	0.2

Means followed by the same letter (s) within a column are not significantly different from each other at the 5% level of significance. Least significance difference (LSD) was used, CV=coefficient of variation, % efficacy=percent efficacy, NS=not significant. Values in parentheses of pre- and post-spray means were transformed from the roots.

TABLE 3. Effects of tested insecticides on ABW population-damaged fruits and yield on cotton plants, Werer, 2022.

Treatment	Larval count/plant		Reduction %	No. of damage squares/plant		No. of damaged flowers/plant		No. of damaged bolls/plant		Health boll/ plant	Seed cotton yield (ton/ha)
	Pre-spray	Post-spray		Pre-spray	Post-spray	Pre-spray	Post-spray	Pre-spray	Post-spray		
chlorfenapyr 36 SC/Tutan	0.53 (1.01)	0.13 (0.79) ^{bc}	80.95	1.05 (1.24)	0.50 (0.99) ^{bc}	0.45 (0.94)	0.15 (0.79)	0.10 (0.77)	0.15 (0.79)	47.8 ^{ab}	4.3 ^{ab}
profenofos + cypermethrin / Proof 44% EC	0.63 (1.06)	0.13(0.79) ^{bc}	83.00	1.20 (1.30)	0.25 (0.86) ^c	0.55 (1.02)	0.20 (0.77)	0.10 (0.77)	0.10 (0.77)	45.6 ^{ab}	4.3 ^{ab}
profenofos 30% + lambda- cyhalothrin 1.5%	0.68 (1.08)	0.20 (0.83) ^{bc}	76.30	1.00 (1.22)	0.60 (1.05) ^b	0.50 (1.04)	0.25 (0.86)	0.25 (0.86)	0.15 (0.79)	44.0 ^{ab}	3.8 ^{bc}
chlorpyrifos 48% EC/Anticat	0.73 (1.11)	0.15 (0.81) ^{bc}	83.45	1.20 (1.30)	0.40 (0.95) ^{bc}	0.55 (1.02)	0.15 (0.80)	0.15 (0.79)	0.10 (0.77)	49.9 ^a	4.4 ^{ab}
beta-cypermethrin+ emamectin/benzoate/Testa	0.63 (1.06)	0.20 (0.84) ^{bc}	74.40	1.00 (1.22)	0.55 (1.01) ^{bc}	0.40 (0.94)	0.15 (0.80)	0.25 (0.86)	0.05 (0.74)	44.4 ^{ab}	3.7 ^c
profenofos + Alpha- cypermethrin/Alphapro	0.50 (0.99)	0.08 (0.76) ^c	88.00	0.85 (1.16)	0.25 (0.86) ^c	0.50 (0.99)	0.05 (0.74)	0.20 (0.83)	0.05 (0.74)	51.5 ^a	4.6 ^a
abamectin+oxymatrine/Bio trin	0.83 (1.15)	0.30 (0.89) ^b	69.91	0.90 (1.18)	0.55 (1.02) ^{bc}	0.30 (0.89)	0.20 (0.82)	0.15 (0.79)	0.20 (0.83)	41.7 ^b	3.3 ^c
Unsprayed	0.60 (1.05)	0.75 (1.11) ^a	-	0.95 (1.20)	1.30 (1.34) ^a	0.45 (0.91)	0.55 (1.02)	0.20 (0.83)	0.25 (0.86)	32.9 ^c	2.7 ^d
LSD (0.05)	Ns	0.11(***)		Ns	0.18 (**)	Ns	Ns	Ns	Ns	7.6	0.6 (***)
CV (%)	6.53	8.74		8.46	12.06	14.02	13.03	11.61	11.38	11.6 (**)	10.7
S.E _±	0.03	0.09		0.16	0.04	0.04	0.02	0.04	0.02	2.6	0.2

Means followed by the same letter (s) within a column are not significantly different from each other at the 5% level of significance. Least significance difference (LSD) was used, CV=coefficient of variation, % efficacy=percent efficacy, NS=not significant. Values in parentheses of pre- and post-spray means were transformed from the roots.

TABLE 4. Combined Analysis of the Effects of Insects against ABW Population-Damaged Fruit and Yield on Cotton, Werer, 2021 & 2022.

Treatment	Larval count/plant		Reduction %	No. of damage squares/plant		No. of damaged flowers/plant		No. of damaged bolls/plant		Health boll/plant	Seed cotton yield (ton/ha)
	Pre-spray	Post-spray		Pre-spray	Post-spray	Pre-spray	Post-spray	Pre-spray	Post-spray		
chlorfenapyr 36 SC/Tutan	0.70 (1.09)	0.12 (0.78) ^c	80.78	1.04 (1.23)	0.46 (0.98) ^{bc}	0.40 (0.93)	0.13 (0.78) ^b	0.10(0.77)	0.10(0.77)	39.2 ^{bc}	3.8 ^{abc}
profenofos + cypermethrin / Proof 44% EC	0.66 (1.08)	0.13(0.79) ^c	83.72	1.16 (1.28)	0.33 (0.91) ^c	0.36 (0.92)	0.08 (0.76) ^b	0.08(0.75)	0.03(0.73)	39.3 ^{bc}	4.0 ^{ab}
profenofos 30% + lambda-cyhalothrin 1.5%	0.84 (1.15)	0.20 (0.83) ^{bc}	77.78	1.23 (1.29)	0.59 (1.04) ^b	0.41 (0.97)	0.17 (0.81) ^b	0.25(0.86)	0.09(0.76)	36.4 ^{cd}	3.5 ^{bc}
chloryrifos 48%EC/Anticat	0.83 (1.15)	0.15 (0.81) ^c	83.03	1.34 (1.35)	0.50 (0.99) ^{bc}	0.41 (0.95)	0.11 (0.78) ^b	0.16(0.81)	0.09(0.77)	41.2 ^{ab}	3.9 ^{ab}
beta-cypermethrin+emamectin/benzoate/Testa	0.78 (1.13)	0.20 (0.84) ^{bc}	75.76	1.31 (1.34)	0.61 (1.04) ^b	0.31 (0.89)	0.13 (0.79) ^b	0.19(0.82)	0.07(0.75)	35.9 ^{cd}	3.4 ^{cd}
profenofos + Alpha-cypermethrin/Alphapro	0.69 (0.08)	0.12 (0.79) ^c	87.48	1.07 (1.25)	0.42 (0.95) ^{bc}	0.45 (0.97)	0.08 (0.76) ^b	0.15(0.80)	0.10(0.77)	43.9 ^a	4.2 ^a
abamectin+oxymatrine/Biotrin	0.84 (1.16)	0.27 (0.87) ^b	70.66	1.15 (1.27)	0.56 (1.04) ^b	0.31 (0.89)	0.17 (0.81) ^b	0.11(0.77)	0.15(0.80)	33.9 ^d	2.9 ^d
Unsprayed	0.65 (1.07)	0.75 (1.15) ^a	-	1.15 (1.20)	1.27 (1.33) ^a	0.36 (0.89)	0.45 (0.97) ^a	0.9(0.83)	0.23(0.85)	25.9 ^e	2.3 ^e
CV (%)	6.28	6.88		11.42	9.64	12.03	10.49	12.07	9.77	11.8	13.5
Treatment	Ns	***		Ns	***	Ns	**	Ns	Ns	***	***
Year	***	Ns		**	Ns	Ns	Ns	Ns	Ns	***	***
Treatment *Year	Ns	Ns		Ns	Ns	Ns	Ns	Ns	NS	Ns	Ns
LSD (0.05)	0.04			0.07	0.10		0.08			4.4	0.5

Means followed by the same letter (s) within a column are not significantly different from each other at the 5% level of significance. Least significance difference (LSD) was used, CV=coefficient of variation, % efficacy=percent efficacy, NS=not significant. Values in parentheses of pre- and post-spray means were transformed from the roots.

cypermethrin, disrupt the sodium channel function in neurons, resulting in prolonged depolarization and paralysis (Gao et al., 2018). Similarly, El-Sayed et al. (2020) reported that the alpha-cypermethrin insecticide was effective at reducing bollworms.

The combination of organophosphates with pyrethroids can exhibit synergistic effects, enhancing the overall efficacy against *H. armigera*, leading to improved crop yields. This aligns with findings from previous studies that advocate for the use of combination insecticides to enhance pest control and minimize resistance development (Raghavendra et al., 2020; Ahmad et al., 2020). Pesticide mixtures containing profenofos + alpha-cypermethrin 440 and profenofos + cypermethrin 44% EC provide efficient control of bollworms due to potent ovicidal, larvicidal, and acaricidal properties (Chongqing, 2019). Moreover, a combination of pesticides effectively reduces damage caused by bollworm complexes and achieves the highest seed cotton yield (Babariya et al., 2010; Rudramuni et al., 2011; Borude et al., 2018; Rajendran et al., 2021). Many studies have assessed the effectiveness of organophosphate/pyrethroid pesticide mixtures in reducing insect populations and enhancing yield (Mushtaq, 2004; Nayak and Daghli, 2007; Khan et al., 2013; Surpam et al., 2015). This approach assumes that insects are less likely to develop resistance to multiple modes of action simultaneously, suggesting that pesticide mixture is a recommended technique for controlling resistance (Warnock and Cloyd, 2005; Desneux et al., 2007).

The successful management of *H. armigera* is particularly critical for cotton farmers in Ethiopia, who face economic challenges exacerbated by pest infestations (Prasanna et al., 2020). The results underscore the importance of integrated

pest management (IPM) strategies that not only focus on chemical control but also incorporate cultural practices and biological control methods to ensure long-term sustainability.

The implications of these findings extend beyond immediate pest control; they highlight the necessity for a comprehensive resistance management strategy. As *H. armigera* has shown a propensity for developing resistance to single insecticides, utilizing a combination approach can help delay resistance onset and maintain the effectiveness of existing chemical tools (Naranjo, 2009). Moreover, integrating sustainable practices such as crop rotation, the use of resistant varieties, and biopesticides can further enhance pest management efforts while reducing reliance on synthetic chemicals. This holistic approach not only protects yield but also promotes environmental health, aligning with global trends towards sustainable agriculture.

Conclusions

The use of insecticides can significantly impact larval populations, damaged fruiting bodies, and seed cotton yield. The results of the present study showed that the application of profenofos + alpha-cypermethrin 440 or profenofos + cypermethrin 44% EC had good efficacy in reducing *H. armigera* larvae and increasing cotton yield by 88.5% and 76%, respectively, compared with the control. These findings suggest that using these insecticides can potentially boost cotton yield and improve farmers' financial stability. The promising results from this study serve as a pivotal reference point for future pest control strategies in Ethiopia. By adopting an integrated approach that combines effective insecticides with sustainable practices, stakeholders can foster resilience in agricultural systems while addressing the challenges posed by pest resistance and economic pressures.

Continued research into alternative pest management methods will be essential for ensuring the longevity and effectiveness of these strategies in the ever-evolving landscape of agricultural pest management.

Acknowledgments

The authors acknowledge the Plant Protection Research Process of Werer Agricultural Research Center for Providing Research Facilities and the Ethiopian Institute of Agricultural Research for financial support.

Author Contributions

Conceptualization and Data Curation: Zemedkun Alemu, Nurhussen Seid and Sileshi Getahun.

Formal analysis: Zemedkun Alemu

Investigation: Zemedkun Alemu, Nurhussen Seid and Sileshi Getahun

Methodology: Zemedkun Alemu

Project administration and Software: Zemedkun Alemu.

Supervision and Validation: Zemedkun Alemu, Sileshi Getahun and Nurhussen Seid.

Writing – original draft: Zemedkun Alemu

Writing – review & editing: Zemedkun Alemu, Nurhussen Seid and Sileshi Getahun.

Funding

This work was supported by the Ethiopian Institute of Agricultural Research. Research budget code (24-02), Entomology Research Program, Werer Agricultural Research Center.

Conflict of interests

The authors declare that they have no potential financial or personal conflicts of interest that could have influenced the study's findings.

Availability of Data

All data generated or analyzed during this study are included in this published article.

References

- Abbott W.S. 1925. A method of computing the effectiveness of an insecticide. *J. Econ. Entomol.* 18: 265-267.
- Ahmad, M., Khan, M. A., Ali, M. 2020. Resistance of *Helicoverpa armigera* to insecticides: A review. *Journal of Insect Science*, 20(2), 1-10.
- Ahmad M., Saleem A.M., Sayyed H.A. 2009. Efficacy of insecticide mixtures against pyrethroid- and organophosphate-resistant populations, *Spodoptera litura* (Lepidoptera: Noctuidae), 65 (3), 266–274. doi:10.1002/ps.1681
- Amanet K, Chiamaka EO, Quansah GW, Mubeen M, Farid HU, Akram R, Nasim. W. 2019. Cotton production in Africa. In *Cotton Production*; Jabran, K., Chauhan, B.S., Eds.; Wiley: Hoboken, NJ, USA, pp. 359–369.
- Babariya P. M., Kabaria B. B., Patel V. N., Joshi M. D. 2010. Chemical control of gram pod borer, *Helicoverpa armigera* Hubner infesting pigeon pea. *Legume Research*, 33(3): 224-226.
- Bahlai, C. A., Sappington, K. G. 2020. The role of biological control in integrated pest management: Enhancing efficacy and sustainability. *Biological Control*, 145, 104227. <https://doi.org/10.1016/j.biocontrol.2020.104227>

- Borude B.S., Bhalkare, S.K., Undirwade, D.B., Rathod, P.K. 2018. Ready Mix insecticides for cotton bollworm complex. *International Journal of Current Microbiology and Applied Science*, 6: 1974–1984.
- Carneiro E., Silva L.B., Maggioni K., dos Santos V.B., Rodrigues T.F., Reis S.S., Pavan, B.E. 2014. Evaluation of Insecticides Targeting Control of *Helicoverpa armigera* (Hubner) (Lepidoptera: Noctuidae). *American Journal of Plant Sciences*, 5, 2823-2828. <http://dx.doi.org/10.4236/ajps.2014.518298>.
- Chambers, J.E., Meek, E.C., Chambers, H.W. 2010. The Metabolism of Organophosphorus Insecticides. In: Krieger R, editor. *Hayes' Handbook of Pesticide Toxicology (Third Edition)*. New York: Academic Press. p. 1399-407.
- Chongqing Shining Fine Chemical Company, 2019. Technical Manual 13 pages.
- Desneux, N., Decourtye, A., Delpuech, J. M. 2007. The sublethal effects of pesticides on beneficial arthropods. *Annual Review of Entomology*, 52, 81-106.
- EIAR (Ethiopian Institute of Agricultural Research). 2017. National Cotton Research Program Strategy for Fifteen Years (2016-2030). Developing national cotton commodity long-term (15 years) research strategy consultative workshop by cotton research core team members. October 24-29, 2016. Addis Ababa, Ethiopia. Pp. 26-33.
- El-Sayed A. A. A., Amer E. A., Eman M. Abd-ElAzeem. 2020. Effect of Some Insecticides on Pink Bollworm, *Pectinophora gossypiella* (Saunders) (Lepidoptera: Gelechiidae), Associated Predators in Cotton Field and Some of Their Biochemical Effects. *J. of Plant Protection and Pathology*, Vol. 11 (2):147 – 151.
- Ermias S., Geremew T., Zeraye M., Mesfin W. 2009. In: Abraham Tadesse (ed) 2009. *Increasing Crop Production through Improved Plant Protection-Volume II. Proceeding of the 14th Annual Conference of the Plant Protection Society of Ethiopia (PPSE)*, 21–22 December 2006. Addis Ababa Ethiopia. PPSE and EIAR, Addis Ababa Ethiopia. Pp. 93-116. Geneva, Switzerland: United Nations Environment Programme, International Labor
- Gao, Y., Liu, Y., Wang, X. 2018. Mechanisms of resistance to pyrethroids in *Helicoverpa armigera*. *Pest Management Science*, 74(5), 1125-1133.
- Geremew T., Ermias S. 2006. *Cotton protection handbook*. Addis Ababa, Ethiopia. 70 pp.
- Geremew T. 2004. *Determining Level of Insecticide Resistance in American Bollworm, Helicoverpa armigera Hubner (Lepidoptera: Noctuidae) for Cotton Pest Management in Ethiopia*. Ph.D Thesis, Kasetsart University, Thailand. 163 pp.
- Haider N., Ahmed K.S., Haidary A.A., Afzal M., Majeed M.Z. Field evaluation of different insecticides against spotted bollworm (*Earias* spp.) and comparative yield assessment for BT and non-Bt cotton. *J. Entomol. Zool. Stud.* 2015;4:33–35.
- Hussain D., Saleem M., Ghouse G., Abbas M. 2015. Insecticide resistance in field populations of *Helicoverpa armigera* (Hübner) (Lepidoptera: Noctuidae). *J. Entomol. Sci.*, 50, 119–128.
- Khan, M. A., Ali, A., Zafar, M. 2020. Recent advancements in integrated pest management: A review. *Journal of Pest*

- Science, 93(2), 563-577.
<https://doi.org/10.1007/s10340-020-01280-6>
- Khan HAA, Akram W, Shad SA, Lee J-J. 2013. Insecticide Mixtures Could Enhance the Toxicity of Insecticides in a Resistant Dairy Population of *Musca domestica* L. PLoS ONE 8(4): e60929. doi:10.1371/journal.pone.0060929.
- Malinga L.N, Laing M.D. 2022. Efficacy of Biopesticides in the Management of the Cotton Bollworm, *Helicoverpa armigera* (Noctuidae), under Field Conditions. Insects. 13: 673. <https://doi.org/10.3390/insects13080673>.
- Melesse Z., Mohammed A., Mezgebu A., Habtamu M. 2019. Cotton production and marketing trend in Ethiopia: A review. Cogent Food Agric. 5:1-7. <https://doi.org/10.1080/23311932.2019.1691812>.
- Merdasa B., Nurhussen S., Yonas B., Michael K., Donis G., Zemedkun A., Samuel D., Arkebe G.E., Sileshi G., Fikremariam T., Mekashaw A. 2022. Cotton Production Guideline in Ethiopia. Werer Agricultural Research Center (WARC), Ethiopian Institute of Agricultural Research (EIAR), Addis Ababa, Ethiopia. 75 pp.
- Mushtaq A. 2004. Potentiation/antagonism of deltamethrin and cypermethrin with organophosphate insecticides in the cotton bollworm, *Helicoverpa armigera* (Lepidoptera: Noctuidae). Pest. Biochem. & Physiol., 80: 31-42.
- Naranjo, S. E. 2009. Integrating biological control into pest management. Biological Control, 50 (1), 1-10
- Nayak M.K., Daglish, G.J. 2007. Combined treatments of spinosad and chlorpyrifos-methyl for management of resistant psocid pests (Psocoptera: Liposcelididae) of stored grain. Pest Management Science, 63:104–109.
- Nurhussen S., Ferdu A., Bayeh M. 2020. Studying the population dynamics of Cotton mealybug (*Phenacoccus solenopsis* Tinsley) (Homoptera: Pseudococcidae) and its major natural enemies, in the middle of Awash, Ethiopia. MSc. Thesis, Hawassa University, Hawassa. 118 pp.
- Prasanna, B. M., et al. 2020. Sustainable management of *Helicoverpa armigera*: A review. Journal of Pest Science, 93 (2), 383-396.
- Raghavendra Y., Basavaraj S.K., Radha J., Jyothi R. 2020. Field evaluation of profenofos 40% + cypermethrin 4% (44%) EC against lepidopteran pests of rice. Journal of Entomology and Zoology Studies, 8(5): 1001-1005.
- Rajendran, S., Kumar, P., Rao, S. 2021. Efficacy of combination insecticides against cotton bollworm: A field study. Journal of Agricultural Science, 13(4), 45-55.
- Reddy, G. V. P., Sahu, K. K. 2021. Integrating biological control with chemical pest management: Strategies and challenges. Pest Management Science, 77(4), 1839-1850. <https://doi.org/10.1002/ps.6106>
- Rudramani T., Reddy K. M., Kumar C. T. 2011. Field evaluation of new insecticidal molecules was conducted against sucking pests and bollworms of cotton. Crop Sciences, 42(1, 2 and 3): 236-302.
- Salama M., ABD EL-mohsen M.A., ABD EL-Baki B.A., El-Nagar E.Y. 2013. The efficiency of some insecticide sequences on cotton bollworms and histopathological effects of some biocides on pink bollworm larvae. Egyptian Journal of Agricultural Research, 91(2): 429–447.

- SAS (Statistical Analysis Software). 1999. Statistical analysis systems SAS/STAT user's Guide version 8e Cary NC: SAS Institute Inc. USA.
- Surpam A.N., Neharkar P.S., Shubhangi P., Bhasme Kamdi S.R. Kadam B.S. 2015. Bioefficacy of new combined product against cotton bollworms, Maharashtra state-India. Indian Journal of Plant Sciences
- Timchalk C., 2010. Organophosphorus insecticide pharmacokinetics. In: Krieger R, editor. Hayes' Handbook of Pesticide Toxicology (Third Edition). New York: Academic Press. p. 1409-33.
- Tossou E., Tapa-Yotto G., Kpindou O.K., Sandeu R., Datinon B., Zeukeng F. Akoton, R.; Tchigossou, G.M. Djègbè I., Vontas J. 2019. Susceptibility profiles of *Helicoverpa armigera* (Hübner) (Lepidoptera: Noctuidae) to deltamethrin reveal a contrast between the Northern and Southern Benin. Int. J. Environ. Res. Public Health 2019, 16, 1882.
- Warnock D.F., Cloyd R.A. 2005. Effects of pesticide mixtures in controlling western flower thrips (Thysanoptera: Thripidae). Journal of Entomological Science, 40: 54–66.
- Wendmagen C., Abere, M. 2012. Selected physical characteristics of the soil of Middle Awash irrigated farmlands, Ethiopia. Journal of Agriculture Science, 22:127-142.
- Yongqiang L., Jianwei L., Mei L., Zhiqing Ma., Chongling C., Yonghong W., Xing Z. 2016. Bacterial expression and kinetic analysis of Carboxylesterase 001D from *Helicoverpa armigera*. International Journal of Molecular Sciences.
- Zemedkun A., Ferdu A., Geremew T. 2022. Susceptibility of African Bollworm, *Helicoverpa armigera* (Hubner) (Lepidoptera: Noctuidae) to Different Commercial Pyrethroid Insecticides on Cotton. Ethiop. J. Agric. Sci. 32(4) 56-72.
- Zhang, L., Liu, Y., Chen, Y. 2020. Mechanisms of action of organophosphate insecticides on insects. Environmental Toxicology and Chemistry, 39(2), 456-466.

Αξιολόγηση εντομοκτόνων εναντίον του πράσινου σκουληκιού, *Helicoverpa armigera* (Hubner) (Lepidoptera: Noctuidae) υπό συνθήκες αγρού στην Αιθιοπία

Zemedkun Alemu*, Nurhusein Seid and Sileshi Getahun

Ethiopian Institute of Agricultural Research, Werer Agricultural Research Center, P.O. Box 2003, Addis Ababa, Ethiopia

ΠΕΡΙΛΗΨΗ

Το πράσινο σκουλήκι του βαμβακιού αποτελεί σημαντική πρόκληση για τους βαμβακοκαλλιεργητές, με αποτέλεσμα σημαντικές απώλειες απόδοσης. Παρά το γεγονός ότι για τη διαχείρισή του χρησιμοποιούνται εντομοκτόνα σκευάσματα, η συνεχιζόμενη έρευνα είναι ζωτικής σημασίας για την ανάπτυξη νέων δραστικών που μειώνουν τον κίνδυνο ανάπτυξης ανθεκτικότητας στις χρησιμοποιούμενες δραστικές ουσίες. Η παρούσα μελέτη στόχευσε στην αξιολόγηση της αποτελεσματικότητας επιλεγμένων εντομοκτόνων εναντίον του *Helicoverpa armigera* μέσω ενός πειράματος πεδίου που διεξήχθη στην περιοχή Middle Awash. Χρησιμοποιώντας ένα σχέδιο πλήρους τυχαιοποίησης, δοκιμάστηκαν οκτώ μεταχειρίσεις σε τέσσερις επαναλήψεις. Συλλέχθηκαν και αναλύθηκαν δεδομένα για το συνολικό αριθμό προνυμφών, της ζημιάς στους καρπούς, και την απόδοση. Τα αποτελέσματα έδειξαν ότι η εφαρμογή διαφόρων εντομοκτόνων επηρέασε σημαντικά ($P < 0,05$) τους αριθμούς των προνυμφών, τη βλάβη στους καρπούς και τη συνολική απόδοση του βαμβακιού σε σπόρο. Συγκεκριμένα, οι συνδυασμοί profenofos 40% + alpha-cypermethrin 4% EC και profenofos + cypermethrin 44% EC είχαν ως αποτέλεσμα αύξηση απόδοσης 88,5% και 76%, αντίστοιχα, σε σύγκριση με τον μάρτυρα. Αυτά τα ευρήματα υπογραμμίζουν τις πρακτικές συνέπειες για τους βαμβακοκαλλιεργητές στην Αιθιοπία και παρόμοιες περιοχές που αντιμετωπίζουν προκλήσεις εχθρών. Η αποτελεσματική χρήση αυτών των εντομοκτόνων θα μπορούσε να μειώσει σημαντικά τις απώλειες απόδοσης που αποδίδονται σε προσβολές από το *H. armigera*, ενισχύοντας έτσι την οικονομική σταθερότητα των αγροτών. Ωστόσο, περαιτέρω έρευνα θα πρέπει να επικεντρωθεί στην επικύρωση και επίδειξη αυτών των εντομοκτόνων σε διαφορετικές αγρο-οικολογίες της χώρας.