


ENTOMOLOGIA HELLENICA

Vol 34, No 1 (2025)

Entomologia hellenica 34(1)



First report of *Hemiberlesia cyanophylli* (Signoret) (Hemiptera: Diaspididae) in Tunisia and North Africa

Monia Ben Halima Kamel, Zouari Sana, Ben Cheick Zahra, Kaydan Bora

doi: [10.12681/eh.39724](https://doi.org/10.12681/eh.39724)

Copyright © 2025, Monia Ben Halima Kamel, Zouari Sana, Ben Cheick Zahra, Kaydan Bora



This work is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike 4.0](https://creativecommons.org/licenses/by-nc-sa/4.0/).

To cite this article:

Ben Halima Kamel, M., Zouari, S., Ben Cheick, Z., & KAYDAN BORA, M. (2025). First report of *Hemiberlesia cyanophylli* (Signoret) (Hemiptera: Diaspididae) in Tunisia and North Africa. *ENTOMOLOGIA HELLENICA*, 34(1), 78–82.
<https://doi.org/10.12681/eh.39724>

First report of *Hemiberlesia cyanophylli* (Signoret) (Hemiptera: Diaspididae) in Tunisia and North Africa

M. Ben Halima Kamel^{1*}, S. Zouari¹, Z. Ben Cheik² & M. Kaydan Bora^{3,4}

¹LR21AGRO5: Gestion des productions horticoles en mode conventionnel et biologique, Institut Supérieur Agronomique Chott Mariem (ISA CM), Université de Sousse, 4042 Tunisie

²LR21AGRO2 : Agrobiodiversité et Ecotoxicologie, Institut Supérieur Agronomique Chott Mariem (ISA CM), Université de Sousse, 4042, Tunisie

³Biotechnology Application and Research Centre, Çukurova University, 01250, Adana, Türkiye

⁴Çukurova University, Imamoğlu Vocational School, 01330, Balcalı, Adana, Türkiye

ABSTRACT

Hemiberlesia cyanophylli (Signoret, 1869) (Hemiptera: Coccoomorpha: Diaspididae) was determined for the first time on *Opuntia triacanthos* Willd (Cactaceae) in a private garden at Akouda (Sousse, Tunisia) in 2022. The infestation was observed on the racket of *O. triacanthos*. This is the first record of *H. cyanophylli* in Tunisia, as well as for North Africa. Two other scale insects: *Parlatoria oleae* (Colvée, 1880) (Diaspididae) and *Planococcus citri* (Risso, 1813) (Hemiptera: Pseudococcidae) were also recorded on other ornamental plant species respectively, i.e. *Rosa indica* (Rosaceae) and *Kalanchoë* sp. (Crassulaceae).

KEY WORDS: *Coccoomorpha*, *Opuntia triacanthos*, Tunisia, North Africa.

The observation of a scale insect (Fig. 1), a honeydew and a sooty mould (Fig. 2), as well as unusual damage (Fig. 3) detected on an ornamental plant in a private garden in the region of Akouda (Sousse, Tunisia), led the authors to organize a survey for Coccoomorpha species presence and identification. The survey revealed the presence of three such species:

Planococcus citri (Risso, 1813) (Hemiptera: Pseudococcidae) (Fig. 1), *Parlatoria oleae* (Colvée, 1880) (Hemiptera: Diaspididae) (Fig. 3) and *Hemiberlesia cyanophylli* (Signoret, 1869) (Hemiptera: Diaspididae) (Fig. 5-6). The first two have previously been reported in Tunisia (Jarraya, 2003, Mansour et al., 2011a & 2011b).

Planococcus citri (Risso, 1813)

The citrus mealybug (Fig. 1) is one of the major pests of citrus and other tree crops. This pest is definitely recognized as a major citrus pest in Tunisia (Mahfoudhi & Dhoubi, 2009). The citrus mealybug is a common polyphagous species known from all zoogeographical regions (Ben-Dov et al., 2013).

Samples were collected during spring 2023 on *Kalanchoë* sp. (Crassulaceae) at Akouda (collected by the first author and identified as *P. citri* by the 3rd), after having observed an unusual damage, the presence of honeydew, sooty mould (Fig. 2) and cochineal colonies.

*Corresponding author: kamoniamn@yahoo.fr



FIG. 1: *P. citri* colony on *Kalanchoë*



FIG. 2: Damage of *P. citri* on *Kalanchoë*

***Parlatoria oleae* (Colvée, 1880)**

Insect samples were collected from *Rosa indica* (Rosaceae) during spring and summer 2023 during this study. Symptoms observed included unusual damage and stem drying. Different stages of armoured scale insects were detected (Fig. 3).

Interestingly, *P. oleae* is the least common armoured scale observed on olive trees in Tunisia comparatively to *Aspidiotus nerii* Bouché (Diaspididae) (Mansour et al, 2011a).



FIG. 3: Female and eggs of *P. oleae* on *Rosa indica*

***Hemiberlesia cyanophylli* (Signoret, 1869)**

During the present study, *H. cyanophylli* was determined for the first time for Tunisia, having been collected in a private garden at Akouda (Sousse, Tunisia) from *Opuntia triacanthos* Willd (Cactaceae) plants, during 2022. This is the first record of *H. cyanophylli* from Tunisia and North Africa. The species was collected from *O. triacanthos* (Fig. 4) by the first author of this paper and identified by the last. All stages of *H. cyanophylli* were presented (Fig. 4). Symptoms included discolouration and covering by scales of the host plant racket (Fig. 4). Samples were stored in 75% ethanol and mounted on glass slides according to the methodology described by Kosztrab and Kozar (1988). Species identification was performed using the characteristics given by Gill (1988) and Miller & Davidson (2005). Sample specimens have been deposited in Çukurova University, Biotechnology Application and Research Center, Balcali, Adana, Turkey.

Hemiberlesia cyanophylli (cyanophyllum scale) is a highly polyphagous species that has been recorded from hosts belonging to 176 plant genera from 77 plant families. It is widely distributed in the tropical and subtropical regions (Scale Net, 2016, accessed 1 December 2024). It is a pest of bananas, worldwide, a known pest of tea in Taiwan (Chua and Wood, 1990), a common scale insect for avocado trees (Takumasa & Jazmín 2016), a pest of olive trees in Peru (Canales Canales and Valdivieso, 1999) and of high economic significance for olive trees in Chile (Claps et al., 2001). Moreover, Foldi (2001) listed this species as an occasional pest in France. It is similar to the latania scale (*Hemiberlesia lataniae* (Signoret, 1869), which is found in Tunisia, infesting grapevines. The latania scale is a rare species for vineyards (Mansour et al. 2011b), but has also been detected on other hosts in Tunisia (Balachowsky 1948).



FIG. 3: Racket discolored of *O. triacanthos* and infested by *H. cyanophylli* (left), *H. cyanophylli* colony on *Opuntia triacanthos* (middle), *H. cyanophylli* female on dorsal view (right).

In conclusion, during the present study, *H. cyanophylli* was observed for the first time in Tunisia on *O. triacanthos* plants and a severe damage of *R. indica* by *P. oleae*

was observed. Other studies on economic crops such as olive trees or *Rosa* sp. are necessary to confirm the impact of the aforementioned species.

References

- Balachowsky, A. 1948. Les cochenilles de France, d'Europe, du Nord de l'Afrique et du Bassin Méditerranéen. IV. Monographie des Coccoidea Classification - Diaspidinae (Première partie). Actualités Scientifiques et Industrielles. Entomol. Appl. 1054: 243-394.
- Canales Canales, A. and L. Valdivieso Jara. 1999. Handbook for biological control in olives. (In Spanish.) Servicio Nacional de Sanidad Agraria, Jesus Maria, Peru. 37pp.
- Chua, T.H. and B.J. Wood. 1990. Other tropical fruit trees and shrubs. In: D. Rosen (ed.), Armoured scale insects, their biology, natural enemies and control. Vol. 4B. World Crop Pests. Elsevier, Amsterdam, the Netherlands. pp. 543-552.
- Claps, L.E., V.R.S. Wolff, and R.H. González. 2001. Catálogo de las Diaspididae (Hemiptera: Coccoidea) exóticas de la Argentina, Brasil y Chile. Revista de la Sociedad Entomológica Argentina. 60: 9-34.
- Davidson, J.A. and D.R. Miller. 1990. Ornamental plants. In Rosen, D (ed.) Armored scale insects, their biology, natural enemies and control. Elsevier, Amsterdam, the Netherlands, 4 (B). pp. 603-632.
- Foldi, I. 2001. Liste des cochenilles de France (Hemiptera, Coccoidea). Bulletin de la Société Entomologique de France. 106: 303-308.
- Gill, R.J. 1988. The scale insects of California: Part 1, The soft scales (Homoptera, Coccoidea: Coccidae). California Department of Food and Agriculture. Sacramento. California. USA. pp. 142.
- Jarraya, A. 2003. Principaux nuisibles des plantes cultivées et des denrées stockées en Afrique du Nord: leur biologie, leurs ennemis, leur dégâts et leur contrôle. Tunis Edition Climat Pub.
- Kosztrab, M. and E. Kozar. 1988. Scale Insects of Central Europe. Series Entomologica. 41. Akademiai Kiado. Budapest. pp. 456.
- Mahfoudhi, N. and M.H. Dhouibi. 2009. Survey of mealybugs (Hemiptera: Pseudococcidae) and their natural enemies in Tunisian vineyards. African Entomology. 17(2): 154-160.
- Mansour, R., K. Grissa Lebdi, P. Suma & M. Russo. 2011a. A survey of scale insects (Hemiptera: Coccoidea) occurring on olives in Tunisia. Journal of Entomological and Acarological Research, Ser. II, 43 (3): 315-322.
- Mansour, R., G. Mazzeo, A. La Pergola, K. Grissa Lebdi and A. Russo 2011b: A survey of scale insects (Hemiptera: Coccoidea) and tending ants in Tunisian vineyards. Journal of Plant Protection Research. 51: 197-203.
- Miller, D.R. and J.A. Davidson. 2005. Armoured Scale Insect Pests of Trees and Shrubs (Hemiptera: Diaspididae). Ithaca, UK: Cornell University Press. pp 442.
- Scale Net, 2016 <https://scalenet.info/catalogue/Hemiberlesia%20cyanophylli/>. Accessed 1 December 2024.
- Takumasa, K. and A.M. Jazmín. 2016. Scale insects (Hemiptera: Coccoidea) associated with avocado crop, *Persea americana* Mill. (Lauraceae) in Valle del Cauca and neighboring departments of Colombia. Insecta Mundi. 0465: 1-24.

Πρώτη αναφορά του *Hemiberlesia cyanophylli* (Signoret) (Hemiptera: Diaspididae) για την Τυνησία και τη Βόρειο Αφρική

M. Ben Halima Kamel^{1*}, S. Zouari¹, Z. Ben Cheik² & M. Kaydan Bora^{3,4}

¹LR21AGRO5: Gestion des productions horticoles en mode conventionnel et biologique, Institut Supérieur Agronomique Chott Mariem (ISA CM), Université de Sousse, 4042 Tunisie

²LR21AGRO2 : Agrobiodiversité et Ecotoxicologie, Institut Supérieur Agronomique Chott Mariem (ISA CM), Université de Sousse, 4042, Tunisie

³Biotechnology Application and Research Centre, Çukurova University, 01250, Adana, Türkiye

⁴Çukurova University, Imamoğlu Vocational School, 01330, Balcalı, Adana, Türkiye

ΠΕΡΙΛΗΨΗ

Το *Hemiberlesia cyanophylli* (Signoret, 1869) (Hemiptera: Coccoomorpha: Diaspididae) καταγράφηκε για πρώτη φορά σε φυτά του είδους *Opuntia triacanthos* Willd (Cactaceae), σε έναν ιδιωτικό κήπο στο Akouda (Sousse, Τυνησία), το 2022. Αυτή είναι η πρώτη καταγραφή του *H. cyanophylli* για την Τυνησία, καθώς και τη Βόρειο Αφρική γενικότερα. Δύο άλλα συγγενή είδη, το *Parlatoria oleae* (Colvée, 1880) (Diaspididae) και το *Planococcus citri* (Risso, 1813) (Hemiptera: Pseudococcidae) καταγράφηκαν επίσης σε άλλα είδη καλλωπιστικών φυτών, πιο συγκεκριμένα τα *Rosa indica* (Rosaceae) και *Kalanchoë* sp. (Crassulaceae).