

## DEPOSITIONAL ENVIRONMENTS AND HYDROCARBON POTENTIAL OF THE MIOCENE DEPOSITS OF ZAKYNTHOS ISLAND

Mpotziolis Ch.<sup>1</sup>, Kostopoulou S.<sup>2</sup>, Triantaphyllou M.<sup>2</sup>, Maravelis A.<sup>3</sup> and  
Zelilidis A.<sup>1</sup>

<sup>1</sup>University of Patras, Department of Geology, Laboratory of Sedimentology, 26504 Patras, Greece, A.Zelilidis@upatras.gr

<sup>2</sup>Faculty of Geology and Geoenvironment, University of Athens, Panepistimioupolis 15784, Athens, Greece,

<sup>3</sup>School of Environmental and Life Sciences, University of Newcastle, Callaghan 2308 NSW, Australia, Angelos.Maravelis@newcastle.edu.au

### Abstract

*Structure, sedimentary features, and biostratigraphy were taken into account to provide additional knowledge into the depositional conditions that influenced the sedimentary sequence along the southern coast of Zakynthos. This succession is exposed from Keri village to Ag. Sostis peninsula and appears to have been influenced by intense tectonic activity. The active tectonics resulted in the formation of at least four coarsening upward cycles. The co-occurrence of Bouma sequence subdivisions and slump horizons within the studied sediments suggest a deep-sea depositional environment. Grain size analysis indicates a regional swallowing upward trend. TOC and CaCO<sub>3</sub> contents are presented with both positive and negative correlations while the TOC values indicate that the study area contain samples with source rock potential.*

**Key words:** Sedimentology, source rocks, Bouma sequence, grain-size analysis.

### Περίληψη

*Λαμβάνοντας υπόψη τις ιζηματογενείς δομές, τα τεκτονικά χαρακτηριστικά και την βιοστρωματογραφία μπορούμε να αντλήσουμε γνώσεις για τις συνθήκες απόθεσης που επηρέασαν την ιζηματογενή ακολουθία κατά μήκος της νότιας ακτής της Ζακύνθου. Η ιζηματογενής ακολουθία από το χωριό Κερί μέχρι και τη χερσόνησο του Αγίου Σώστη φαίνεται να έχει επηρεαστεί από έντονη τεκτονική δραστηριότητα, παράγοντας τουλάχιστον τέσσερις κύκλους με αυξανόμενο κοκκομετρικό μέγεθος προς τα πάνω. Η παρουσία των υποδιαιρέσεων της ακολουθίας Bouma και των οριζόντων ολίσθησης εντός των υπό μελέτη ιζημάτων προτείνουν ένα βαθύ θαλάσσιο περιβάλλον απόθεσης. Η κοκκομετρική ανάλυση έδειξε μια σταδιακή πλήρωση του κάθε κύκλου ιζηματογένεσης, αλλά και ολόκληρης της περιοχής. Επιπροσθέτως, η ανάλυση του TOC και του CaCO<sub>3</sub> έδειξε ότι υπάρχει μια σχέση του ποσοστού τους με το περιβάλλον απόθεσης, ενώ οι τιμές TOC δείχνουν ότι η περιοχή μελέτης περιλαμβάνει δείγματα με δυνατότητα γένεσης υδρογονανθράκων.*

**Λέξεις κλειδιά:** Ζάκυνθος, ιζηματολογία, μητρικά πετρώματα, ακολουθία Bouma.

## 1. Introduction

Marine depositional environments with low depositional rate are typified by increasing carbonate content towards the coarse fractions (Saadallah & Kukal, 1969). The organic material is intimately related to sedimentary environments (Folk, 1968). The accumulation of organic matter requires: (1) high organic productivity, (2) high subsidence rates and (3) anoxic conditions. Such conditions occur either because of overproduction and deposition of organic material, or because of limited water circulations resulting in non-oxygen recycle.

In contrast, erosion results to the oxidation of the organic material of the rock and to the dissolution and precipitation of minerals in both sandstones and limestones. The dissolution of calcium carbonate ( $\text{CaCO}_3$ ) is affected by the temperature, pressure and the partial pressure of carbon dioxide ( $\text{CO}_2$ ). The dissolution of calcium carbonate ( $\text{CaCO}_3$ ) is controlled by the dissolution of carbon dioxide ( $\text{CO}_2$ ). In particular, the more carbon dioxide ( $\text{CO}_2$ ) dissolved in water, the more the calcium carbonate ( $\text{CaCO}_3$ ) will be dissolved. The  $\text{CO}_2$  is dissolved rapidly at high pressures and low temperatures and thus, the  $\text{CaCO}_3$  is dissolved more in the deep ocean waters of the, compared to shallow, surface water. The saturation of  $\text{CaCO}_3$  is also related to the pH. The higher the pH of the water is, the greater is the concentration of  $\text{CO}_2$ . Since more  $\text{CO}_2$  is incorporated to the deep ocean waters because of the respiration of organisms, the more easily the water erodes the calcareous shells. The organic matter and amount of oxygen are also affected by the level of sea level (Reolid et al. (2010)).

The decomposition of marine sediments from the oxygen and the oxidation of organic material, results in the release of  $\text{CO}_2$  into the water. Such process reduces the pH of the water and leads to a subsaturation of the  $\text{CaCO}_3$ . Through this procedure, the  $\text{CaCO}_3$  may be subjected to post-consequential dissolution (Martin & Sayles, 2004). The dissolution of  $\text{CaCO}_3$  by organisms can be independent of the saturation of water environments (Milliman et al., 1999). On the other hand,  $\text{CaCO}_3$  in pelagic sediments is controlled by several factors described by (Khim et al., 2011). The extent of carbon provided in the seabed at a constant volume of dilution factor (productivity), variations in the content of carbon in carbon feed rate constant (factor of dissolution), change in the proportions of calcareous/silicate particles in the flow and dilution of non biogenic material such as wind and volcanic suspensions are the principles factors that affect the  $\text{CaCO}_3$  content.

When there is an inverse correlation between  $\text{CaCO}_3$  and Corg in sediments, then the  $\text{CO}_2$  that is produced by the decomposition of Corg in the subsurface layer of water (where we have no circulations and  $\text{O}_2$  recycling) and sediments, resulted in a decrease in the pH of water, thus expediting the dissolution of  $\text{CaCO}_3$  produced in the surface layer of water. This happens when the sediment's Corg percentage is above 12% or there are seasonal anoxic bottoms (Dean, 1999; 2002). On the other hand, when there is a positive correlation between the two percentages, occurs due to removal of  $\text{CO}_2$ . At high temperatures, where the solubility of  $\text{CaCO}_3$  and the pH in the surface layer of water is reduced, thus accelerating the production of  $\text{CaCO}_3$  (Hodell et al., 1998). The interruption of this relation in stratigraphic column shows that the  $\text{CaCO}_3$  in sediments is mainly controlled by dissolution of the subsurface layer of water and sediment due to the decomposition of organic material.

For this purpose, five sections have been selected from Keri to Kalamaki area, where fifty samples have been selected in order to determine grain-size; total organic carbon (wt. % ToC) and  $\text{CaCO}_3$  contents (Figure 1).

## 2. Geological Setting

The study area is situated at the western part of Greece. Western Greece represents the southern edge of the Dinarides-Albanides-Hellenides active margin and has been the focus of oil/gas exploration over the last decade (e.g. Zelilidis et al., 2003; Karakitsios and Rigakis, 2007; Maravelis et al., 2012). It has been influenced by both a compressional and an extensional tectonic

regime, because of the motion of the Adriatic and the Apulia plates. Zakynthos Island is part of the parautochthonous Apulian lithospheric plate of Hellenic mountain belt and contains rocks of two different geotectonic zones, the western Pre-Apulian zone and the eastern Ionian zone (Underhill, 1985).

The region under study was influenced by the Ionian thrust during the early Pliocene (Zelilidis et al., 1998). As a result, the area was fragmented and separated into two discrete sedimentary basins, the westward Alikanas foreland basin and the eastward Gerakas piggy-back basin (Zelilidis et al., 1998). The Foreland basin hosts the oldest rocks seen in the area (Kontopoulos et al., 1997). The study area was served as a continental shelf during Tortonian and thus, mudstones that contain rare sandstone beds were accumulated (Keri area, Figure 1). The upward coarsening upward trend suggests a progressive regional swallowing upward. A large-scale NNW directed normal fault brings in contact the Tortonian shelf with the underlying Late-Cretaceous limestones (Figure 1). Messinian evaporites have been also identified (Agios Sostis area, Figure 1). Such salts evolve into a depositional sequence that contains evaporitic beds of turbiditic origin and further upslope, early Pliocene deep-sea sands. The sedimentation develops with the deposition of shelf deposits with hummocky cross-stratification. This depositional environment is represented by “trubi” marly limestone.




Figure 1 - Geological map of the study area (IGME, 1980, Zakynthos).

### 3. Tectonostratigraphy

The activity of the Ionian thrust during the early Pliocene is the most important tectonic event that affected the study area. This orogenic event separated the main basin into two discrete sub-basins, a westward foreland and an eastward piggy-back basin (Zelilidis et al., 1998). The foreland basin is typified by major NNW – SSE anticline, which is crossed-cut by both eastward and westward directed faults. Vrachionas anticline was developed during the Pliocene as part of the foreland-propagating fold and thrust system within Pre-Apulian zone (Underhill, 1989). The Pre-Apoulian zone is characterized by a compression regime, after Miocene time (Sorel, 1976; Underhill, 1989), followed by a Pliocene extension regime (Underhill, 1989). Nowadays, there is a shift in the tectonic regime and the study area is typified by a NNE – SSW directed regime. Finally, palaeomagnetic measurements indicate a late Pleistocene 25° clockwise rotation (Duermeijer et al., 1999).

#### 4. Materials and Methods

A suite of fifty samples were selected from five different sections (Figure. 2). The base of the depositional sequence is exposed at section one while, the younger sediments are exposed at sections four and five(Figure 2).

Section one: This section is cited at the southern end of Keri gulf. It is eight meters in total thickness and is composed of at least eight coarsening upward sedimentary cycles. Each cycle is up to 1m in total thickness and is represented by a basal mudstone that evolves upslope into coaly beds. Coals are with 30 cm in total thickness.



**Figure 2 - Map showing the study area and the sections of where the samples were taken. (Photo from Google Earth).**

Section two: This section corresponds to the laterally equivalent of section one, is located at Keri gulf and is presented with up to 6 meters total thickness. This outcrop consists of a basal gray/brown mudstone up to 25cm thick interbedded with thin-bedded sandstones. There is an upward increase in mudstone-sandstone ratio.

Section three: This section is located northeast of the section two between the Keri gulf and Agios Sostis peninsula. In this area a sedimentary succession with up to 65 meters in total thickness is exposed (Figure 3).This sequence is characterized by a thick slump horizon, placed at the top and with 25 meters in total thickness (figure 4). This slump horizon overlies sandstone beds, one meter thick with no evidence of deformation. The deformation unit underlies undisturbed mudstone strata. Throughout the sedimentary sequence Bouma subdivision are the dominant sedimentary features (Figure 5).



**Figure 3 - Panoramic view of Section three.**



Figure 4 - Slump horizon within section three.




Figure 5 - Bouma sequence within section three.

Section four: This section represents the upper parts of the studied stratigraphy (Kontopoulos et al., 1997). In the current research samples were collected at the base of the section. The upper parts of both sections four and five are represented by carbonates. The "Trubi" marly limestones are considered as the typical early Pliocene sedimentary facies of the Mediterranean. Such facies overlain the turbiditic sequence, which contains sandstone and re-sedimented evaporitic turbidites (Kontopoulos et al., 1997). Both the upper part of the studied section four (overlying the stratigraphic column of this work), and the upper part of the Kalamaki cross-section (Section five) consist of the typical for the Mediterranean environment during the Early Pliocene, "trubi" marly limestone (Figures 6 and 7), formed over Pliocene turbiditic sequence, with sandstone and re-sedimented evaporitic clasts turbidites (Kontopoulos et al., 1997).



Figure 6 - The "trubi" marly limestone on the upper part of section four, at Agios Sostis.



Figure 7 - The "trubi" marly limestone on the upper part of section five at Kalamaki.

According to the above description a synthetic stratigraphic column was produced (Figure 8). From the whole sequence a detailed calcareous nannofossil biostratigraphy showed that the age of section one and two sediments is Tortonian, and only from slumps we found many resedimented mostly Oligocene and Aquitanian to Burdigalian sediments (Figure 8). Samples of section three and section five (e.g. samples 30, 45, 52) display rare to common small *Gephyrocapsa* spp. coccoliths (2.5-3.0  $\mu\text{m}$ ), within a rich but redeposited from Eocene, Late Oligocene, Aquitanian-Burdigalian, Messinian, assemblage. *Gephyrocapsa* spp. presence along with the absence of *Discoaster* species (*D. tamalis*, *D. pentaradiatus*, *D. surculus*); only *Discoaster* cf. *brouweri* has been documented, enables tentative age assignment in Early Pliocene (subbottom of *Gephyrocapsa* spp. at 4.33 Ma, Lourens et al., 2004). Sample 36 from section three yields nannofossil flora (*Reticulofenestra pseudoumbilicus*, *Pseudoemiliana lacunosa*, small *Gephyrocapsa* spp.) of Early Pliocene (NN14-15 biozone).

As depicted on Figure 8, a synthetic stratigraphic column was constructed based on sedimentologic evidence. The detailed calcareous nannofossil biostratigraphy indicates a Late Miocene (Tortonian) age for the sections one and two. Samples from the slump horizon determined as Oligocene-early Miocene in age (Figure 8). Sections three and five (e.g. samples 30, 45, 52) display rare to common small *Gephyrocapsa* spp. coccoliths (2.5-3.0  $\mu\text{m}$ ), within a rich but re-deposited fossils from Eocene, Late Oligocene, Miocene (Aquitanian-Burdigalian, Messinian) sedimentary sequence. The occurrence of *Gephyrocapsa* spp. in association with the absence of *Discoaster* species (only *Discoaster* cf. *brouweri* has been documented), suggests an Early Pliocene age (subbottom of *Gephyrocapsa* spp. at 4.33 Ma, Lourens et al., 2004). Sample 36 from section three yields nannofossil flora (*Reticulofenestra pseudoumbilicus*, *Pseudoemiliana lacunosa*, and small *Gephyrocapsa* spp.) of Early Pliocene (NN14-15 biozone).

One cycle of section four at Agios Sostis and fourteen cycles of section five in Kalamaki display the influence of the Ionian thrust during the depositional evolution. The section five is intimately related to the activity of the Ionian thrust because of its proximity to the thrust front whereas; the remotely positioned section four is little influenced by the thrust activity.

## 5. Methodology - Materials and Methods

Sieve analysis for sediments sandstone (>63 $\mu\text{m}$ ) and pipette analysis for mudstones (<63 $\mu\text{m}$ ) were used in order to determine grain size and their parameters. The determination of organic carbon content was based on the method of titration (Gaudette et al., 1974). The content of calcium carbonate was measured with the complete dissolution of calcium carbonate ( $\text{CaCO}_3$ ) with acetic acid ( $\text{CH}_3\text{COOH}$ ) (Varnavas, 1979).

## 6. Results

The grain size analysis aims to determine the lithology and grain size parameters in order to describe the particle size distribution of sediments. The data exported from the grain size analysis were used for the construction of cumulative grading curve and the determination of the lithologic character of the sediments. The statistical parameters were calculated based on the Folk & Ward (1957) method.

The grain size analysis suggests that silt is the main lithology with minor presence of clay. Coarser material is totally absent. As portrayed on Figure 9, the depositional depth of the studied samples is variable. In Figure 11, it is depicted that the selected samples display a random scatter in terms of the stagnant water.

Generally, the main lithology is siltstone. The analysis of the statistical parameters suggests that studied sediments were mostly deposited in a deep basin depth, but a considerable number of samples were deposited in low depth too. Additional evidence for the deep basin depth is provided by the common presence of Bouma subdivisions in section three.




Figure 8 - Synthetic stratigraphic column of the studied area (age determination was according to Martini, 1971).




Figure 9 - Diagram of Valia and Cameron (1979), showing the depositional depth (Sk1=asymmetry, Mz=mean and σ1= deviation).




Figure 10 - Diagram of Steward (1958) for the determination of the depositional environment.





Figure 11 - Diagram of Passega (1957) for the determination of the depositional environment.

The amount of organic matter in sedimentary rocks is usually measured as the total organic carbon content (TOC) and expressed as a percentage of the dry rock. TOC is not a clear indicator of petroleum potential. as thick, organic-rich potential source rocks cannot become effective source rocks without sufficient burial and thermal maturation (Kenneth et al., (2007)).


Organic carbon content of the studied sediments exhibits variable contents, ranging from 0 to 3.19 wt. % (Figure 12).



**Figure 12 - The content of TOC in studied deposits, where the line depict the lower limit of 0,5% that differentiate the potential source rocks with those that have no potential.**

Studies of numerous global samples of different ages have led to the conclusion that the minimum TOC value required for the designation as an immature source rock is 0.5 wt. % (Hunt, 1979; Hedberg & Moody, 1979; Tissot & Welte, 1984). Thus, these values suggest that the studied sedimentary sequence contains samples with poor to good source rock potential.

The calcium carbonate of the studied samples ranges between 1.76 to 61.29 (Figure 13).



**Figure 13 - The content of CaCO<sub>3</sub> in studied deposits.**


In the study area the content of calcium carbonate is high among the fine material, suggesting high sedimentation rates. The combination of total organic carbon (TOC) content and calcium carbonate contents in sedimentary rocks can be employed to add knowledge into the depositional conditions during sedimentation (Figure 14). There is not a constant trend between these two parameters throughout the studied sedimentary succession. The contents of the total organic carbon and calcium carbonate display both positive and negative correlation. This fact indicates that the CaCO<sub>3</sub> content is mainly controlled by dissolution in the subsurface layer of water and sediment due to the decomposition of organic material.

The first stratigraphic cycle is typified by a positive correlation between the two parameters. The high percentage of CaCO<sub>3</sub> indicate high sedimentation rates while the variable TOC content can be explained by the existence of seasonal anoxic bottom (no circulation and recycling of O<sub>2</sub>) (Dean, 1999; 2002). Additionally, low organic productivity and high rate of burial of the organic matter resulted in no oxidization. In the other cycles, however, the lower content of CaCO<sub>3</sub> indicates lower sedimentation rates and so the TOC content is not influenced by the depositional environment.

The reduction in the content of CaCO<sub>3</sub> from cycle one to four indicates a gradual swallowing of the basin. Cold climatic conditions are also suggested and are in agreement with the Miocene regional icehouse conditions (Bertolani - Marchetti, (1985). Such conditions may potentially have resulted into an increase in solubility of CaCO<sub>3</sub>. However, the third cycle is characterized by the



presence of slump horizon and thus, the low rates of  $\text{CaCO}_3$  cannot be connected to low sedimentation rates. The slumps led to scarification of the subsurface layer and the production of suspensions, which probably concluded to the low content of  $\text{CaCO}_3$  caused by the dilution due to non-biogenic material (Khim et al., 2011). The high rates of TOC observed in a group of samples of this cycle possible can be related to quick of burial rate, where the organic matter does not have time to be oxidized.



**Figure 14 - Correlation of  $\text{CaCO}_3$  and TOC contents within studied deposits.**

## 7. Conclusions

Grain size analysis suggests a regional upward swallowing upward trend. Sedimentation took place in a deep basin with steep western margins during Tortonian to early Pliocene. The  $\text{CaCO}_3$  content indicates high deposition rates, while TOC contents indicate potential source rocks for hydrocarbon generation. The comparison of TOC and  $\text{CaCO}_3$  indicates that these two parameters are presented with both positive and negative correlation. This character is intimately related to the dissolution of the  $\text{CaCO}_3$  at the subsurface layer of water and sediment because of the decomposition of organic material, in oxic conditions.

## 8. References

- Bertolani–Marchetti D. 1985. Pollen palaeoclimatology in the Mediterranean since Messinian time, in: D. J. Stanley and F. C. Wezel (Editors), *Geological Evolution of the Mediterranean basin*. Springer, New York, pp. 525.
- Dean W.E. 1999. The carbon cycle and biogeochemical dynamics in lake sediments, *Journal of Paleolimnology*, v. 21, p. 375–393.
- Dean W.E. 2002. A 1500-year record of climatic and environmental change in Elk Lake, Minnesota II: Geochemistry, mineralogy, and stable isotopes, *Journal of Paleolimnology* v.27
- Duermeijer C. E., Krijgsman W., Langereis C. G., Meulenkamp J. E., Triantafyllou M.V. and Zachariasse W.J. 1999. A late Pleistocene clockwise rotation phase of Zakynthos (Greece) and implications for the evolution of the western Aegian arc, *Earth and Planetary Science Letters* 173 (1999), p. 315-331.
- Folk R.L. 1968. Petrology of Sedimentary Rocks, *Hemphill Publishing Co*, Austin, Texas, 170p
- Folk R. L. and Ward W. S. 1957. Brazos river bar. A study in the significance of grain size parameters. *Jour. Sed. Petr.*, 27, Nol, 3-26.
- Gaudette H.E., Flight W.R., Toner L. and Folger D.W. 1974. An inexpensive titration method for the determination of organic carbon in recent sediments, *Jour. Sed. Petr.*, 44, 249-253.
- Hedberg R.M. and Moody J.D. 1979. Petroleum prospects of the deep offshore, *Am. Assoc. Pe t. Geol. Bull.* 63:p. 286-300.
- Hodell D.A. and Schelske C.L. 1998. Production, sedimentation, and isotopic composition of organic matter in Lake Ontario, *American Society of Limnology and Oceanography*. 43(2), p 200-214.

- Hunt J.M. 1979. *Petroleum Geochemistry and Geology*, Freeman, San Francisco ISBN 0-7167-1005-6.
- Karakitsios V. and Rigakis N. 2007. Evolution and petroleum potential of Western Greece, *Journal of Petroleum Geology* 30 (3):197-218.
- Kenneth, E.P., Leslie B.M., Zenon C.V. and Paul G.L., 2007. Source-Rock Geochemistry of the San Joaquin Basin Province, California, Chapter 11, *Petroleum Systems and Geologic Assessment of Oil and Gas in the San Joaquin Basin Province, California*
- Khim B.K., Kim H.J., Cho Y.S., Chi S.B., Yoo C. M. 2011. Orbital Variations of Biogenic CaCO<sub>3</sub> and Opal Abundance in the Western and Central Equatorial Pacific Ocean During the Late Quaternary, *Terr. Atmos. Ocean. Sci.* 23 -1, 107-117.
- Kontopoulos N., Zeligidis A., Piper D. and Mudie P. 1997. Messinian evaporates in Zakynthos, Greece. *Palaeo*, 129, 361-367.
- Maravelis A., Makrodimitras G. and Zeligidis A. 2012. Hydrocarbon prospectivity in Western Greece, *Oil and Gas European Journal* 38 (2), 84-89.
- Martin W. and Sayles F. 2004. The recycling of biogenic material of the seafloor, in: Holland, H.D. and Turekian, K.K. (eds.), *Treatise on Geochemistry*, Vol. 7, Sediments, Diagenesis and Sedimentary Rocks (ed. F.T. Mackenzie), Elsevier. Amsterdam, 37-66 pp.
- Martini E. 1971. Standard Tertiary and Quaternary Calcareous Nannoplankton Zonation, *Proceedings of the II Planktonic Conference*, Roma, 1970, A. Farinacci, ed., Ed. Tecnoscienza, p. 739 - 785, Rome.
- Milliman J.D., Troy P.J., Balch W.M., Adams A.K., Li Y.H. and Mackenzie F.T. 1999. Biologically mediated dissolution of calcium carbonate above the chemical lysocline, *Deep Se Res.* I, 46, 1653-1669.
- Moila R.J. and Visser D. 1968. Textural parameters: an evaluation, *Jour. Sed. Petr.*, V260, p. 45-63.
- Passega R. 1957. Texture as characteristic of clastics deposition: *Am. Soc. Assoc. Petrol. Geol. Bull. Tulsa*, Okland, 41, 1952-1984.
- Passega R. 1964. Grain size representation by CM patterns as a geologic tool, *Jour. Sed. Petrol. Tulsa Ok.*, 34, No. 4, 830-847.
- Reolid M., Nagy J., Rodríguez-Tovar F.J. 2010. Ecostratigraphic trends of Jurassic agglutinated foraminiferal assemblages as a response to sea-level changes in shelf deposits of Svalbard (Norway), *Palaeogeography, Palaeoclimatology, Palaeoecology*, 293: 184-196.
- Saadallah A. and Kukal Z. 1969. Grain size and carbonate content in coastal sediment of Iraq, *Jour. Iraq G. Soc.*, 2, 3-10.
- Sorel D. 1976. Etude neotectonique dans l'arc egeen externe occidentale: les iles ioniennes de Ke fallinia et Zakinthos et l'Elide, *These 3me cycle*, Universite de Paris Sud, Orsay.
- Stewart H.B. 1958. Sedimentary reflections of depositional environments in San Miguel Lagoon, Baja California, Mexico, *Am. Assoc. Petr. Geol. Bull.*, V42, p. 2567-2618.
- Tissot B. and Welte D.H. 1984. *Petroleum Formation and Occurrence*, Springer Verlag.
- Underhill J. R. 1985. Neogene and Quaternary tectonics and sedimentation in Western Greece, *PhD thesis* Cardiff, Wales, University College, University of Wales.
- Underhill J.R. 1989. Late Cenozoic deformation of the Hellenides foreland, western Greece, *Bulletin of the Geological Society of America*, 101, p 613-634.
- Valia H.S. and Cameron B. 1977. Skewness as a Paleoenvironmental indicator, *Jour. Sed. Petr.*, 47, No2, 784-793.
- Varnavas S.P. 1979. Geochemistry of sediments from the eastern Pacific, *PhD Thesis* University London pp. 431.
- Zeligidis A., Piper D.J.W., Vakalas I., Avramidis P. and Getsos K.. 2003. Oil and gas plays in Albania: Do equivalent plays exist in Greece? *Journal of Petroleum Geology* 26 (1):29-48.
- Zeligidis A., Kontopoulos N., Avramidis P. and Piper D. 1998. Tectonic and sedimentological evolution of the Pliocene-Quaternary basins of Zakynthos Island, Greece: case study of the transitions from compressional to extensional tectonics, *Basin Research*, 10, 393-408.