Monitoring temporal changes of the surface water area of the Burullus and Manzala lagoons using automatic techniques applied to a Landsat satellite data series of the Nile Delta coast

KH. M. DEWIDAR

Faculty of Science, New Damietta City, Department of Environmental Science, Mansoura University, Box 34517, Egypt

Corresponding author: khdewidar@yahoo.com

Received: 1 February 2010; Accepted: 13 September 2011; Published on line: 21 November 2011

Abstract

This study introduces the automated shoreline techniques used to monitor the temporal change of the surface water area of the Burullus and Manzala lagoons. In this study, a series of Landsat image data are acquired at intermittent intervals between 1972 and 2006 for the Burullus lagoon and between 1972 and 2007 for the Manzala lagoon. All Landsat images were radiometrically calibrated and converted to reflectance values. The reflectance values of each date were atmospherically corrected using the 6S model. The automated shoreline technique was checked against field observations using GPS over the four seasons for each lagoon during reconnaissance for the shoreline boundary. The accuracy of the extracted shoreline boundary for each lagoon was validated by calculating the area of a large aquaculture farm in the study area from satellite imagery and the available topographic maps. The resulting accuracy of this technique was approximately 97.5%. From the spatial temporal analysis of the satellite data, the results indicate that the rate change of the aquatic surface area is -7.3 km²/yr and -2.7 km²/ yr for the Manzala and Burullus lagoons, respectively, during the approximately 35-year study period. The changes detected in this study indicate that over this time period, the surface water area of the Manzala and Burullus lagoons has decreased by 62.6% and 61.9, respectively, compared with their original size.

Keywords: Shoreline position extraction; Coastal lagoon; Landsat; Surface area changes; Nile Delta.

Introduction

Coastal lagoons and wetlands have many benefits: they support local economies based on fisheries, agriculture and timber production: they are a source of food and energy; they offer diverse opportunities for recreation and tourism (ALIAUME et al., 2007); and they provide protection to shorelines against storms and other meteorological perturbations. During the last few decades, the northern lagoons of the Nile delta have been rapidly impacted by eutrophication from wastewater drainage and have been altered by human activities. These activities include new transportation roads, drying processes, and urban encroachment. Population increases, together with rural and urban activities, not only directly affect the availability and quality of natural resources, but also induce secondary effects that must be evaluated from a regional viewpoint (MOUFADDAL, 2005).

Several studies have been conducted on the different aspects of the Burullus lagoon ecology (EL SABROUTI, 1984; SAAD, 1988; ABDEL MONEIM, et al., 1990; FA-TOUH, 1990; MAIYZA, et al., 1991; EL KARACHLLY, 1991; EL SHERIF, 1993; ABOUL EZZ, 1995; SHAKWEER, et al., 1998). These results show that the water quality of the lagoon has changed over time. These changes are related to human intervention and eutrophication processes (BELT-AGY, 1985; SAAD, 1990; KHEDR & LOVETT-DOUST, 2000; RADWAN & SHAKWEER, 2004). The bottom sediments of the lagoon are predominantly silty clay, with high organic content in some parts, and large areas of shelly to silty muddy sand (COUTLIER & STANLEY, 1987). Approximately 75 islands are scattered throughout the Burullus lagoon, with various surface areas, offering a variety of habitat types. Other studies have indicated that the size and shape of the Burullus lagoon has suffered from shrinking and, that many of the islands have increased in size (GURIGU-IS et al., 1996; KHEDR, 1999).

Studies on the hydro-chemical characteristics of the Manzala lagoon have been carried out by many authors (EL WAKEEL & WAHBY, 1970; WAHBY et al., 1972; BISHAI & YOUSEF, 1977; DOWIDAR & ABDEL MOATI, 1983; ABDEL BAKY & EL GHOBASHY, 1990; FAHMI et al., 1997). A United Nations Development Program in 1997 compiled background infor-

mation about the lagoon to be used for monitoring by remote sensing. This project reported that large amounts of particulate matter, nutrients, and bacteria are transported to Manzala lagoon via drains. BATTAGLIA & YORK, 2002 reported that dissolved oxygen differed significantly between sites, whereas conductivity, temperature, total suspended solids, ammonia and nitrate were shown to be significantly different between sampling days. In DEWIDAR et al., 2008, a Chlorophyll-a model derived from remote sensing data indicated that the concentration of chlorophyll is high at the eastern and southern boundaries of the lagoon. High Chlorophyll-a concentration were associated with high concentrations of total phosphorous and high turbidity. These high concentrations may be attributed to the effect of drains and eutrophication conditions. Their study also indicated that Manzala lagoon has suffered from different levels of eutrophication resulting from human activities and poor management. RAMDANI et al., 2009, stated that zooplankton communities in Manzala lagoon were generally dominated by rotifers and that the planktonic communities were comparatively highly diverse with essentially freshwater species.

Previous remote sensing studies have been carried out to extract shoreline position. . MCFEETERS (1996) used a Normalised Difference Water Index (NDWI) for the detection of water bodies. FRIHY et al., 1998, used a screen manual digitising method to detect the water body of the Manzala lagoon. The waterline can also be mapped through image classification including density slice analysis (RYAN et al. 1991; GORMAN et al. 1998; BRAUD & FENG 1998; MOORE, 2000; FRAZIER & PAGE 2000; PAJAK & LEATHERMAN 2002; STOCKDON et al., 2002; HORRITT et al., 2003). Time series remotely sensed

images with a medium spatial resolution are ideal data sources for mapping coastal land uses and monitoring their changes over a large area (SHI et al., 2001). BOSWORTH et al., 2003 used a segmentation technique for multispectral Landsat TM imagery achieved using multi-resolution combined with watershed pyramids with variational region growth. DI et al., 2003, investigated a novel approach to the automatic extraction of shorelines from high-resolution IKONOS satellite imagery using a mean shift segmentation algorithm as a first step, followed by a local refinement process. Monitoring changes in the coastline is an important task in some applications, such as cartography and the environmental management of the entire coastal zone (DELLEPIANE 2004; ALESHEIKH et al. 2004). FOODY et al. (2005) proposed a technique called a super-resolution mapping approach applied to soft classification. LIRA (2006) proposed a new methodology to segment open water bodies based on a variant of principal component analysis. Furthermore, the use of satellite remote sensing data was found to be a cost effective approach to quantifying changes over large geographic regions (SESLI et al. 2008). DEWIDAR & FRIHY (2008) used an automated waterline technique for detecting pre and post beach response to engineering hard structures at the northwestern part of the Nile delta from Landsat satellite data. AHMED et al., 2009, used ISODATA unsupervised techniques to classify Manzala lagoon into three main classes of open water, emergent vegetation and dry land. They indicated that Manzala lagoon has declined in area by approximately 50%.

The aim of the present work is to quantify the loss of the water body areas of the Manzala and Burullus lagoons over a period of 35 years (1972-2007) using automatic

techniques applied to Landsat time series data. The automatic technique developed in this study can be used to update north Egyptian coastal lagoon maps. The results of this research are also important for decision makers to manage the north coastal lagoons of Egypt for future generations and conserve it as a natural defence against expected climate changes.

Study areas

The coastal lagoons and the Nile River of Egypt are the largest wetlands in the north of Africa, representing approximately 25% of the total wetlands of the Mediterranean Sea (SAAD, 2003). These lagoons contribute no less than 50% of the total wild fish catch produced in Egypt (SAAD, 2003). Moreover, they are important sites for the passage and breeding of migratory birds. In the following two sections, the study areas are described in detail.

Manzala lagoon

Manzala lagoon is the most economically important fishing ground among the northern Delta lagoons in Egypt. This lagoon supplies the Egyptian people with approximately 50% of the total Egyptian fish catch (BISHAI & YOSEF, 1977). Manzala lagoon is situated at the eastern margin of the Damietta branch between 31° 45' - 32° 20' E longitude and 31° 00' - 31° 30' N latitude (Fig. 1). The area was reduced from 1647 km² to approximately 862 km² during the early 1990,s to approximately 777 km² in 1993 (ABDEL BAKY & EL GHOBASHY, 1990) as a result of extensive agricultural reclamation activities, especially on the southern and south western sides of the lagoon. Within the lagoon are hundreds of islands which vary in size and shape. Some islands in the north, which often have a long axis




Fig. 1: The main geomorphological features derived from Landsat-7 ETM+ image data dated 2007 for Manzala lagoon.

parallel with the coast, have been attributed to former coastal sand ridges, whereas islands further south, which are less elongated in a north-south direction, may reflect deposition along former branches of the Nile (ARBOUILLE & STANELY 1991; AY-ACHE, et al., 2009). The lagoon lies within the borders of five of Egypt's governorates (Dakahliya, Damietta, Port Said, Ismailiya and Sharkiva) and is connected to the Mediterranean via an artificial opening, the El-Gamil inlet (100 meters wide), approximately 10 km west of Port Said city. The northwestern side of the lagoon is connected with the Damietta estuary by the El Suffara and El Ratama canals north of Damietta City. The exchange of water and biota between the lagoon and the adjoining Mediterranean Sea is possible through these outlets. The northern part of the lagoon is affected by marine water invasion through the El Gamil inlet, while the southern and southwestern parts annually receive approximately 6695×10³ m³ of sewage effluents and agricultural and industrial drainage waters without treatment from several drains and canals, namely Bahr El Bakar, Ramsis, Hadous, El Matariya, El Serw, Faraskour, Inaniya, and Port Said city. The major part of the drainage water (78%)is discharged by the Bahr El-Bakar, Ramsis and Hadous drains. The latter drains are heavily contaminated by sewage and industrial wastes. The construction of the international coastal road and the fast growing new human settlements on either side of the high way are also among the serious threats to biodiversity in the lagoon (Fig. 3A).

The lagoon's average depth is approx-

imately 1.25 m (DEWIDAR, *et al.*, 2008), and it is traversed by several sandy and clayey islets that divide the lagoon into several more or less isolated small basins known locally as "Bahour". To the north of the lagoon is a recently protected area called Ashtoum El Gamil. This area was established by the Egyptian Prime Ministers Decision No.459/1988 & Modify Deicer No.2780/1988. The area is approximately 175 km² and is completely located inside the lagoon (Fig. 1). The main purpose for creating this protected area was the protection of many resident and migratory species of birds, saline and fresh water fish, natural plants and historical sites scattered throughout the lagoon. However, there is a proposal to increase the size of this protected area to encompass larger, more important parts of Manzala lagoon.

Burullus lagoon

Burullus lagoon lies on the eastern side of the Rosetta branch of the Nile River, occupying a central position along the Mediterranean Nile delta coast of Egypt. The lagoon extends between longitude 30° 30' and 31° 10' E and latitudes 31° 21' and 31° 35' N (Fig. 2). Burullus is the second largest of the Nile delta coastal lagoons at approximately 53 km long and 13 km wide, and it has water depths ranging from 0.5 to 2.5 m (FRIHY & DEWIDAR, 1993). GUIR-GUIS et al., (1996) stated that the area of the Burullus lagoon has been reduced to 452 km². TORAB and AZAB (2007) mention that the area of the Burullus lagoon has been reduced during the last century from 8688.5 km² (1925) to 553.4 km² (1984) to 489.4 km² (2001) as a result of human activities and land reclamation projects, especially after the construction of the High Aswan Dam. MOUFADDAL et al. (2008) stated that the Burullus lagoon surface area has deceased to 460 km² as a result of extensive agricultural reclamation activities, especially on the southern and southeast-




Fig. 2: The main geomorphological features derived from Landsat-7 ETM+ image data dated 2006 for Burullus lagoon.

ern sides of the lagoon. Burullus lagoon is connected to the Mediterranean Sea at its northeastern edge through the Burullus inlet (Fig. 3B), which is approximately 250 m wide and 5 m deep. The northern border is separated from the Mediterranean Sea by a strip of land covered with sand bars and dunes. Seven drains and fresh water canals are connected to its eastern, southern and western shores. The lagoon barriers are sandy and range from 0.4 to 5.5 km in width. These barriers are generally < 1.5 m above mean sea level, with beach face slopes ranging between 5° and 13°. Low relief backshores and foredunes characterise the western barrier. The eastern barrier is narrow and backed by barchans coastal dunes. These dunes encroach landward onto a cultivated coastal flat. Water is discharged into the lagoon through a series of seven land drains. The estuarine water of the Rosetta mouth of the Nile River is also mixed with the lagoon water through the Brimbal Canal. These water sources and others (rainfall and seepage of underground water) cause

a rise in the lagoon level inducing a lagoon sea current (SAAD, 1990). Burullus lagoon is designated as a wetland nature reserve under the International Ramsar Convention of 1988 and is one of the significant wetlands along the Mediterranean coast. The lagoon has been a major area for human activities, including urban and rural development, agriculture reclamation projects, and recreation, navigation and industrial activities.

Material and Methods

In this study, a series of image data were acquired at periodic intervals between 1972 and 2006 for Burullus lagoon and between 1972 and 2007 for Manzala lagoon, i.e., covering a time span of approximately 35 years (Table 1). The images were acquired at unequal intervals at a good quality level, with no effective clouds or sensor defects such as striping. The study area is visible in the TM and ETM+ scenes (Path / Row 176/38 and 177/39) and in MSS (Path / Row 190/38,

Table 1
Acquired dates, sensor type and spatial resolution of remote sensing data used in this study.

	Bı	urullus lago	oon	Manzala lagoon			
Serial	Acquired	Sensor	Spatial	Acquired	Sensor	Spatial	
#	date	type	Resolution	date	type	Resolution	
			(meters)			(meters)	
1	31/08/1972	MSS	57.00	31/08/1972	MSS	57.00	
2	11/09/1984	TM	28.50	03/01/1973	MSS	57.00	
3	10/08/1990	TM	28.50	20/09/1984	TM	28.50	
4	17/06/1993	TM	28.50	11/04/1995	TM	28.50	
5	18/04/1997	TM	28.50	07/06/1998	TM	28.50	
6	12/02/2000	TM	28.50	11/11/2000	ETM+	14.25	
7	22/02/2001	ETM+	14.25	24/08/2003	ETM+	14.25	
8	17/06/2002	ETM+	14.25	18/06/2005	ETM+	14.25	
9	09/08/2004	ETM+	14.25	04/03/2007	ETM+	14.25	
10	12/06/2006	ETM+	14.25				

190/39, 191/38 and 191/39). All image scenes were subjected to image processing using ERDAS Imagine software version 9.1 (ERDAS, 2006). In this study, automated shoreline positions were formed for each date with image data. The satellite image data were processed according to the following steps.

Geometric corrections

The study data were geometrically corrected to the Universal Transverse Mercator (UTM) map projection system zone number 35 and number 36 north. A layer stack was formed for all 8 bands of the ETM+ sensor for each date (ERDAS, 2006). Recent Landsat-7 ETM+ image data were rectified to a topographic map scale of 1:50000 with more than 35 ground control points evenly distributed within 2006 for Burullus lagoon and 2007 for Manzala lagoon. A second-degree polynomial and nearest neighbour interpolation were used for resampling the image to the UTM reference system. The image rectification accuracy was < 0.5pixel for each date. Other dates of satellite images scenes were registered to the rectified image of 2006 for Burullus lagoon and 2007 for Manzala lagoon. The image registration accuracy is < 0.4 pixels between each date and the 2006 and 2007 imagery.

Atmospheric corrections

In this study, all Landsat images were radiometrically calibrated and converted to reflectance values. Radiometric calibration can greatly improve our ability to compare landcover change at one location and is required for studies involving multiple temporal uses of Landsat images in two or more geographically district locations. ERDAS Imagine Model Maker was used to convert the DN of the image values to apparent at sensor reflectance values. The re-

flectance values of each date were atmospherically corrected using the 6S model (VERMOTE et al., 1997). The input parameters of the 6S model describe the atmospheric conditions, aerosol model and concentration; target and sensor altitude, band definitions; definitions of the target and environment reflectance and latitude and longitude of the target; and azimuth and zenith angles of the Sun and sensor. Continental type aerosol was assumed and a locally measured visibility value for each date was taken from the stations of the Egyptian Meteorological Authority. The atmospheric corrected data were checked with the standard spectral reflectance curves of four materials: sand, mud, vegetation and water (LILLESAND, et al., 2008). The method of checking the atmospheric data included the creation of reflectance profiles for each material type (sand, mud, vegetation and water) from the corrected image using the spectral profile module of the ERDAS Imagine software. Using this module, the behaviour of each spectral band curve was graphically checked against the standard.

Shoreline position extraction

To extract the shoreline position an image threshold was determined using band 4 (0.8-1.1μm) for MSS and band 7 (2.0-2.35µm) for TM/ETM+ (shortwave infrared) to create a binary image or image mask (zero value for water and one value for land) for each date. The selections of these bands depend on the strong capability of water to absorb the incident energy at particular wavelengths. Thus, the water appears black in colour in these bands and with a sharp edge detected between water and land. To ensure the waterline mapping accuracy in the case of the MSS image data 3 x 3 edge enhancement filters were used to sharpen the boundary between the water and land

classes. Binary images (masked images) are used as input layers in the unsupervised classification ISODATA algorithm to form a complete separation between the land class and the water class and to remove the effect of submerged vegetation (macrophytes) and scattered islands inside the lagoon. Horizontal and vertical Sobel filters (ERDAS, 2006) were used on each unsupervised classified image of each date to enhance edge detection. Some editing to remove small objects and fill holes was carried out for each filtered image on each date. The filtered images for each date were converted to vector layers using the Raster to Vector module (ERDAS, 2006).

Validation of shoreline extraction

To validate the resultant water body vectors, each vector dataset was superimposed over the false colour composite 543 RGB for TM/ETM+ for each date. This false color composite enhanced the sharp edges between vegetation, wetland and surface water bodies. During this check, some net lines were removed from the boundary vectors due to the ridges between basins (Fig. 3C). The fieldwork took place during the four sea-

sons to carry out reconnaissance for the water boundary of each lagoon using the Garmin Etrex Summit GPS. To detect the accuracy of shoreline extraction in this study, a large aquaculture farm of El Matariya was selected. The farm area measured from a topographic map (1:50,000) was 4.0 km². Vectors extracted from this map were then superimposed over the extracted vectors from satellite data. Additionally, the resultant calculated area value was checked through personal communication with the Ministry of Water Resources and Irrigation. The area value calculated by this approach was approximately 3.9 km² yielding an accuracy of ~ 97.5%.

Image differencing

Image differencing is a technique whereby changes in brightness values or other datasets are determined by the cell-by-cell subtraction of co-registered image datasets (SINGH 1989, GREEN et al. 1994). To quantify the changes between the resultant water body areas, the image differencing technique was used between each vector dataset (Table 2 & 3). The change of water body surface area between each two dates may be

Table 2
Some numerical indicators of Manzala lagoon surface area loss.

Year	Boundary area	Islands	Time interval	Changed	Unchanged
	(km²)	(km ²)	(year)	(km²)	(km²)
1972	1079.4514	137.8	1972-1973	22.7188	1056.8354
1973	1072.5638	137.8	1973-1984	109.1993	970.3549
1984	992.8424	135.4	1984-1995	205.0346	874.5196
1995	923.401	126.4	1995-1998	196.5536	883.0006
1998	929.9031	132.4	1998-2000	232.8163	846.7379
2000	861.6439	129.1	2000-2003	235.6176	843.9366
2003	860.693	133.9	2003-2005	259.5957	819.9842
2005	844.0394	114.6	2005-2006	290.5385	789.0157
2007	800.9148	125.1			

Table 3
Some numerical indicators of Burullus lagoon surface area loss.

Year	Boundary area	Islands	Time interval	Changed	Unchanged
	(km²)	(km ²)	(year)	(km²)	(km ²)
1972	682.6	30.1	1972-1984	70.2638	612.3796
1984	626.7	32.6	1984-1990	213.6698	468.9736
1990	474.5	30.7	1990-1993	218.0388	464.6046
1993	476.3	34.2	1993-1997	218.7327	463.9107
1997	470.9	32.8	1997-2000	229.8094	452.834
2000	464.6	30.6	2000-2001	228.5244	454.1447
2001	461.2	33.0	2001-2002	227.3936	455.2498
2002	462.3	33.6	2002-2004	228.5244	454.1447
2004	459.4	33.8	2004-2006	229.9893	452.6541
2006	457.0	34.7			



Fig. 3: Photographs of the study areas showing, (A) the international coastal road crossing Manzala lagoon, (B) the Burullus inlet, (C) drying processes inside the southern part of Manzala lagoon, (D) the height of reeds inside Burullus lagoon.

attributed to an increase in the size of the islands due to siltation and drying processes.

Results and Discussions

Manzala lagoon

By observing the spatial distribution of surface water area for Manzala lagoon (Fig. 4 and Table 2), it can be concluded that the lagoon has lost much of its water surface area due to drying processes, fish farms and land reclamation. The changes between 1972 and 1973 are insignificant compared with those between 1973 and 1984. These changes may be attributed to faster drying processes in the north of the lagoon instigated by building an alternative coastal road due to

the coastal erosion of the Damietta headland during this period. This coastal road connects the cities of Damietta and Port Said. Additionally, large areas in the northwest have been converted to fish ponds for intensive aquaculture. During the second decade, between 1984 and 1995, the water surface body of the Manzala lagoon suffered from greater shrinkage due to artificial drying to build a wastewater treatment plant on the eastern side of the lagoon. The changes during the third decade are attributed to the increase of human interventions, such as urban encroachment and land reclamation activities on the eastern and southwestern sides of the lagoon. The result of this study agrees with the study of ABDEL BAKY & EL GHOBASHY (1990), which used a differ-




Fig. 4: Vector layers of Manzala lagoon derived from satellite time series data.

ent approach. The results of this study also indicate that the rate of Manzala's aquatic surface area loss over 35 years is –7.3 km²/yr (Fig. 5A). This result means that the Manzala lagoon has been heavily impacted by eutrophication caused by wastewater from the drainage system and by the drying processes of land reclamation and urban encroachment due to the expansion of the Dameitta governorate to the east of the lagoon. By comparing the results of this study with other studies (AHMED, et al., 2009), there is general agreement in the trend and pattern of shrinkage for the water surface area of Manzala lagoon.

The above situation justifies the creation of improvement works to maintain the present area of the lagoon water body without

further area loss. These improvement works could include dredging to create three parallel radially submerged channels that are 4 m deep and approximately 17 km in total length. This dredging plan can lead to a decrease of reed areas in the lagoon and contribute to the establishment of equilibrium between the reed areas and the free water surfaces of the lagoon system. These channels would facilitate the movement of water and sediment in the lagoon, thus preventing siltation processes. Second, a built up boundary buffer zone around the lagoon could be constructed to prevent random access and drying processes by fishermen or others. These suggestions have been discussed with the stakeholders in the study area to keep Manzala lagoon a healthy ecosys-




Fig. 5: Rate of Manzala (A) and Burullus (B) aquatic surface area change.

tem. Finally, some stakeholders have suggested using new technologies for waste management to decrease pollution loads.

Burullus lagoon

By comparing the change of water surface area between 1972 and 1984, the surface water area has decreased by 55.9 km² (Fig. 6 and Table 3) in Burullus lagoon. During the period between 1984 and 1997, the water surface body of the Burullus lagoon suffered from more shrinkage due to drying processes and land reclamation at the eastern side and the southern part of the lagoon (Table 3). By comparing the results of this study with other studies of area changes of the Burullus lagoon (GUIRGUIS et al., 1996; MOUFADDAL et al., 2008), there is gen-

eral agreement in the trend and pattern of shrinkage of the water surface area. During the period between 1997 and 2006, the rate of shrinkage decreased due to the Burullus lagoon becoming a conserved area under the International Ramsar convention of 1988. The rate of surface water area shrinkage is -2.7 km²/ yr (Fig. 5B) during the entire period of approximately 35 years. The islands inside the Burullus lagoon have only been slightly changed, which may be attributed to seasonal changes and changes in the density of reeds. Presently, the main problem the Burullus lagoon faces is the increase in the tallest reeds inside the lagoon (Fig. 3D), which may be attributed to the harvesting of reeds by fishermen and an increase in the eutrophication of the




Fig. 6: Vector layers of Burullus lagoon derived from satellite time series data.

southern part of the lagoon. To overcome this problem, harvesting and dredging processes should be alternated.

Conclusions

The results of the present study show that automated shoreline techniques can be used as a tool for monitoring long-term changes of surface water areas of the northern Nile Delta lagoons. The results of this study may also be important for decision makers to manage the north coastal lagoons for future generations and to conserve them as a natural defence against expected climate change. The accuracy of this technique is more than 95% for the detection of the surface water areas across the study area. The changes detected in this study indicate that the surface water area of the Manzala and Burullus lagoons has decreased by 62.6% and 61.9%, respectively, from their original size in 1972 over 35 years. The above situation justifies the creation of improvement works to maintain the area of the Manzala lagoon water body without further loss.

The results of this study may allow the decision makers to visualise the temporal changes of the Burullus and Manzala lagoons. Furthermore, the study may be an alarms for all stakeholders who are concerned with north Egyptian coastal lagoons. The resource value of the Nile Delta coastal lagoons has brought about the need to protect and conserve them. These lagoons should be continuously monitored so that temporal changes in their environment can be analysed. The importance of determining the cause, extent, and spatial distribution of these changes can be used in different aspects of environmental studies, land suitability analyses and environmental management in Egypt.

Acknowledgment

The author thanks Dr Paul Alpine, Chairman of Remote Sensing and Photogrammetry Society, School of Geography, University of Nottingham, UK for reading this manuscript and for his comments about this work. The author also thanks the postgraduate students of the Environmental Science Department of Mansoura University for their help in the field survey.

References

ABDEL BAKY, T.E. & EL GHOBASHY A.E., 1990. Some Ecological aspects of the western region of Manzala lagoon, Egypt. I-Physico-chemical characteristics. *Mansoura Science Bulletin*, 17: 166-184.

ABDEL MONEIM, M.A., EL SARRAF W.M. & BELTAGY A.I., 1990. Distribution of some heavy metals in lake Burullus, Egypt. *Bulletin of National Institute of Oceanography & Fisheries*, 16: 203-221.

ABOUL EZZ, S.M., 1995. Zooplankton of lake Burullus, Egypt. *Bulletin of National Institute of Oceanography & Fisheries*, 21: 233-261.

AHMED, M., EL LEITHY, B., THOMP-SON, J., FLOWER, R., RAMDANI, M., AYACHE, F. & HASSAN, S., 2009. Application of remote sensing to site characterization and environmental change analysis of North Africa coastal lagoons. *Hydrobiologia*, 622: 147-171.

ALESHEIKH, A.A., GHORBANALI, A. & TALEBZADEH, A., 2004. Generation the coastline change map for Uremia Lagoon by TM and ETM+ imagery. In: *Map Asia Conference. Beijing, China*.

ALIAUME, C., DO CHI, T., VIAROLI, P. & ZALDIVAR, J.M., 2007. Coastal lagoons of Southern Europe: recent changes

- and future scenarios. *Transitional Waters Monographs*, 1 (1): 1-12.
- ARBOUILLE, D. & STANLEY, D.J., 1991.

 Late Quaternary evolution of the Burullus lagoon region, north central Nile delta, Egypt. *Marine Geology*, 99 (1-2): 45-66. AYACHE, F., THOMPSON, J.R., FLOWER, R.J., BOUJARRA, A., ROUATBI, F. & MAKINA, H., 2009. Environmental characteristics landscape history and pressures on three coastal lagoons in the Southern Mediterranean Region: Merja Zerga (Morocco), Ghar El Melh (Tunisia) and Lake Manzala (Egypt). *Hydrobiologia*, 622: 15-43.
- BATTAGLIA, M. & YORK, P., 2002. Water quality changes through the Adrel pond. In: *UTS Freshwater Ecology Report 2002*. Department of Environmental Sciences, University of Technology, Sydney.
- BELTAGY, A.I., 1985. Sequences and consequences of pollution in northern Egyptian lakes. Lake Burullus. *Bulletin of National Institute of Oceanography & Fisheries*, 11: 73-97.
- BISHAI, H.M. & YOSEF, S.F., 1977. Some aspects of the hydrography, physicochemical characteristics and fisheries of Manzala Lake. *Bulletin of National Institute of Oceanography & Fisheries*, 7: 32-58.
- BOSWORTH, J., KOSHIMIZU, T. & ACTON, S.T., 2003. Multi-resolution segmentation of soil moisture imagery by watershed pyramids with region growing merging. *International Journal of Remote Sensing*, 24: 741-760.
- BRAUD, D.H. & FENG, W., 1998. Semiautomated construction of the Louisiana coastline digital land/water boundary using Landsat Thematic Mapper satellite imagery. In: Louisiana Applied Oil Spill Research and Development Program, OSRAPD Technical Report Series 97-002. Department of Geography & Anthro-

- pology, Louisiana State University.
- COUTLIER, V. & STANLEY, D.J., 1987. Late quaternary stratigraphy and paleogeography of the eastern Nile delta, Egypt. *International Journal of Marine Geology*, 27: 257-339.
- DAVIS, J.A. & FROEND, R., 1999. Loss and degradation of wetlands in southwestern Australia. Underlying cause, consequences and solutions. *Wetlands Ecology & Management*, 7: 13-23.
- DELLEPIANE, S., DE LAURENTIIS, R. & GIORDANO, F.G., 2004. Coastline extraction from SAR images and a method for the evaluation of coastline precision. *Pattern Recognition Letters*, 25 (13): 1461-1470.
- DI, K., WANG, J., MA, R. & LI, R., 2003. Automated shoreline extraction from high resolution IKONOS satellite imagery. In: ASPRS 2003 Annual Conference Proceedings, May 2003, Anchorage, Alaska.
- DEWIDAR, KH., EL EZABY, K., ABDEL DAYM, H. & IBRAHIM, M., 2008. Mapping some water quality parameters by using Landsat-7 ETM+ for Manzala lagoon, Egypt. In: *The 3rd International Conference, Environmental Sciences & Technology, Egypt 2008 April 1-3, 2008*. Environmental Research Division, National Research Centre, Cairo, Egypt.
- DEWIDAR, KH. & FRIHY, O., 2007. Pre and post beach response to engineering hard structures using landsat time-series at the northwestern part of the Nile delta, Egypt. *Journal of Coastal Conservation*, 11 (2): 133-142.
- DOWIDAR, M. & ABDEL MOATI, R., 1983. Distribution of nutrients salts in Lake Manzala (Egypt). International Mediterranean, 28: 185-188.
- EL KARACHLLY, A.G., 1991. Production function analysis in Lake Burullus fish-

- eries. Bulletin of National Institute of Oceanography & Fisheries, 17: 37-43.
- EL SABROUTI, M.A., 1984. Mineralogy and sources of bottom sediments of Lake Burullus, Egypt. *Journal of Africa Earth Science*, 2: 151-153.
- EL SHERIF, Z.M., 1993. Phytoplankton standing crop, diversity and statistical multispecies analysis in lake Burullus, Egypt. *Bulletin of National Institute of Oceanography & Fisheries*, 19: 213-233.
- EL WAKEEL, S.K. & WAHBY, S.D., 1970. Hydrography and chemistry of Manzala Lake, Egypt. *Archiv für Hydrobiologie*, 67: 173-200.
- ERDAS, 2006. Leica Geosystems Geospatial Imaging, LLC, 5051 Peachtree Corners Circle, Suite 100, Norcross, GA, 30092, USA.
- FAHMI, M.A., ABBAS, M.M. & EMARA, H.I., 1997. Chemical studies on the impact of pollution in lake Manzala, Egypt. Bulletin of High Institute of Public Health, 24: 31-44.
- FRIHY, E., DEWIDAR, KH., NASR, S. & EL RAEY, M., 1998. Change detection of the northeastern Nile Delta of Egypt: shoreline changes, Spit evolution, margin changes of Manzala lagoon and its islands. *International Journal of Remote Sensing*, 19 (10): 1901-1912.
- FATOUH, S.A., 1990. Analysis of fishery exploitation in Lake Burullus. *Bulletin of National Institute of Oceanography & Fisheries*, 16: 109-118.
- FOODY, G.M., MUSLIM, A.M. & ATKIN-SON, P.M., 2005. Super-resolution mapping of the waterline from remotely sensed data. *International Journal of Remote Sensing*, 26 (24): 5381-5392.
- FRAZIER, P.S. & PAGE, K.L., 2000. Water body detection and delineation with Landsat TM data. *Photogrammetric Engineer*ing & Remote Sensing, 66 (12): 1467-1467.

- FRIHY, O. & DEWIDAR, KH., 1993. Influence of shoreline erosion and accretion on texture and heavy mineral compositions of beach sands of the Burullus coast, north-central Nile delta, Egypt. *Journal of Marine Geology*, 114 (1-2): 91-104.
- GORMAN, L., MORANG, A. & LARSON, R., 1988. Monitoring the coastal environment. IV. Mapping, shoreline changes, and bathymetric analysis. *Journal of Coastal Research*, 14 (1): 61-92.
- GREEN, K., KEMPKA, D. & LACKEY, L., 1994. Using remote sensing to detect and monitor landcover and landuse change. Photogrammetric *Engineering* & *Remote Sensing*, 60 (3): 331-337.
- GUIRGUIS, S.K., HASSAN, H.M., EL RAEY, M.E. & HUSSAIN, M.M., 1996. Multi-temporal change of Lake Burullus, Egypt, from 1983 to 1991. *International Journal of Remote Sensing*, 17: 2915-2921.
- HORRITT, M.S., MASON, D.C., COBBY, D.M., DAVENPORT, I.J. & BATES, P.D., 2003. Waterline mapping in flooded vegetation from airborne SAR imagery. *Remote Sensing of Environment*, 85 (3): 271-281.
- KHEDR, A.A., 1999. Floristic composition and phytogeography in a Mediterranean deltaic lake (Lake Burullus), Egypt. *Ecologia Mediterranea*, 25 (1): 1-11.
- KHEDR, A.A. & LOVETT-DOUST, J., 2000. Determinants of floristic diversity and vegetation composition on the islands of Lake Burullus, Egypt. *Applied Vegetation Science*, 3: 147-156.
- LILLESAND, T.M., KIEFER, R.W. & CHIPMAN, J.W., 2008. Remote sensing and image interpretation. Wiley, Hoboken, NJ.
- LIRA, J., 2006. Segmentation and morphology of open water bodies from multispectral images. *International Journal*

- of Remote Sensing, 27: 4015-4038.
- MAIYZA, I.A., BELTAGY, A.I. & EL MA-MONEY, M.H., 1991. Heat balance of Lake Burullus, Egypt. *Bulletin of National Institute of Oceanography & Fisheries*, 17: 45-55.
- MCFEETERS, S.K., 1996. The use of the Normalized Difference Water Index (NDWI) in the delineation of open water features. *International Journal of Remote Sensing*, 17: 1425-1432.
- MOORE, L.J., 2000. Shoreline mapping techniques. *Journal of Coastal Research*, 16 (1): 11-124.
- MOUFADDAL, W., 2005. State of the coastal water along the Nile Delta front, Egypt, as revealed by satellite imagery. *International Journal of Remote Sensing*, 26: 4877-4883.
- MOUFADDAL, W., EL SAYED, E. & DAGHADY, E., 2008. Updating morphometric and idaphic information of lakes Edku and Burullus (NW Nile delta, Egypt) with the aid of satellite remote sensing. *Egyptian Journal of Aquatic Research*, 34: 291-310.
- PAJAK, M.J. & LEATHERMAN, S., 2002. The high water line as shoreline indicator. *Journal of Coastal Research*, 18 (2): 329-337.
- RADWAN, A.M. & SHAKWEER, L.M., 2004. On the present status of the environmental land fishery of lake Burullus. *Egyptian Journal of Aquatic Research*, 30: 43-98.
- SAAD, A.M., 1988. Studies on the bottom deposits of the Egyptian lakes. *Bulletin of National Institute of Oceanography & Fisheries*, 14: 123-132.
- SAAD, A.M., 1990. State of the Egyptian delta lakes, with particular reference to pollution problems. In: *Regional Symposium Environmental Studies (UNARC)*, Alexandria.

- SAAD, A.M., 2003. Impact of diffuse pollution on the socio-economic development opportunities in the coastal Nile Delta lakes. In: *Diffuse Pollution Conference, Dublin.*
- SESLI, A.F., KARSLI, F., COLKESEN, I. & AKYOL, N., 2008. Monitoring the changing position of coastlines using aerial and satellite image data: an example from the eastern coast of Trabzon, Turkey. *Environmental Monitoring & Assessment*, 153 (1-4): 391-403.
- SHAKWEER, L.M., EL EBIARY, E.H. & ZAKI, M.A., 1998. Comparative study on the major biochemical constituents in the muscles of Mugil cephalous inhabiting the Mediterranean water, the northern Delta lakes and fish farms of Egypt. *Bulletin of National Institute of Oceanography & Fisheries*, 24: 29-101.
- SHI, Y., PING, D. & SHEN, C., 2001. Changes in progradation rate of the tidal flats at the mouth of the Changjing (Yangtze) River, China. *Geomorphology*, 38: 167-180.
- SINGH, A., 1989. Digital change detection techniques using remote sensing data. *International Journal of Remote Sensing*, 10: 989-1003.
- STOCKDON, H.F., SALLENGER, A.H., LIST, J.H. & HOLMAN, R.A., 2002. Estimation of shoreline position and change using airborne topographic Lidar data. *Journal of Coastal Research*, 18 (3): 502-513.
- TORAB, M. & AZAB, M., 2007. Modern shoreline changes along the Nile Delta coast as an impact of construction of the Aswan high dam. *Geographia Technica*, 2: 69-76.
- UNDP, 1997. *Project Document. Lake Manzala Engineered wetland*. United Nation Development Program, Project Number: EGY/93/G31, March 1997.

- VERMOTE, E., TANRE, D., DEUZE, J., HERMAN, M. & MORCRETTE, J., 1997. Second simulation of the satellite signal in the Solar spectrum, 6S: An overview. *IEEE Transactions on Geosciences & Remote Sensing*, 35: 675-686.
- WAHBY, S.D., YOUSSEF, S.F. & BISHARA, N.F., 1972. Further studies on the hydrography and chemistry of Lake Manzala. *Bulletin of National Institute of Oceanography & Fisheries*, 2: 401-422.