

Mediterranean Marine Science

Vol 12, No 2 (2011)



First record of Pontian Monkey Goby, *Neogobius fluviatilis* (Pallas, 1814) in the Evros River (Greece); Is it an alien species?

S. ZOGARIS, A. APOSTOLOU

doi: [10.12681/mms.47](https://doi.org/10.12681/mms.47)

To cite this article:

ZOGARIS, S., & APOSTOLOU, A. (2011). First record of Pontian Monkey Goby, *Neogobius fluviatilis* (Pallas, 1814) in the Evros River (Greece); Is it an alien species?. *Mediterranean Marine Science*, 12(2), 454-461.
<https://doi.org/10.12681/mms.47>

First record of Pontian Monkey Goby, *Neogobius fluviatilis* (Pallas, 1814) in the Evros River (Greece); Is it an alien species?

S. ZOGARIS¹ and A. APOSTOLOU²

¹ Hellenic Center for Marine Research, Institute of Inland Waters, P.O. Box 712,
19013 Anavissos, Hellas

² The Institute of Biodiversity and Ecosystem Research, Bulgarian Academy of Sciences,
2 Gagarin Street 1113 Sofia, Bulgaria

Corresponding author: zogaris@ath.hcmr.gr

Received: 7 September 2011; Accepted: 7 October 2011; Published on line: 2 November 2011

Abstract

The Pontian Monkey Goby, *Neogobius fluviatilis* (Pallas, 1814), was recorded for the first time in Greece in August 2011. Eight specimens were collected in the Greek-Turkish section of the Evros River, 65 kilometers upstream of its river-mouth. Although the species has been recently discovered in the Tundza, a tributary of the Evros in Bulgaria, it has never before been found in the Evros' main stem. Although the lower Evros has been poorly researched by ichthyologists, it is unlikely that a conspicuous medium-sized fish would go unnoticed in this river; and, it is therefore suggested to be a probable alien. However, since the Evros basin has had geological connections to the Marmara Sea and Black Sea in the past and it is immediately adjacent to native populations of *N. fluviatilis*, the species status is categorized as questionable until genetic and morphological studies are completed.

Keywords: *Neogobius fluviatilis*; Greece; Bulgaria; Turkey; Evros; Maritsa; Meriç; Alien.

Introduction

A recent electrofishing expedition revealed the existence of the Pontian Monkey Goby *Neogobius fluviatilis* (Pallas, 1814) on the lower main stem of the river Evros-Maritsa-Meriç (hereafter referred as Evros, the river's classical Greek name). This is the first record of the species in Greece and its second published documentation in lotic environments in this river basin. Although many

river reaches have been recently surveyed in the Bulgarian section of the Evros, the species was recently found only in one section of the Bulgarian Tundza tributary (STEFANOV *et al.*, 2008), where it is well established and provisionally considered as non-native.

The native distribution of *N. fluviatilis* is known to encompass many Black Sea river basins and parts of the Caspian basin including the lower parts of associated lot-

ic systems (Dnieper, Dniester, Danube, Don and Volga rivers), and the Sea of Marmara basins (KOTTELAT & FREYHOF, 2007). According to NEILSON & STEPIEN (2011) certain Caspian populations are currently considered a different species, *Neogobius pallasii*. Moreover, *N. fluviatilis* is notorious for its remarkable East-to-West invasions within Europe in recent decades. Assisted by river navigation channel creation and accidental introductions the species began a rapid expansion in the rivers of the northern Black Sea since 1970 (COP *et al.*, 2005). By the first years of this century alien populations of the species have been found far up the Danube in Germany, throughout the main stem of the Vistula in Poland, and more recently as far west as the Rhine in the Netherlands (VAN KESSEL *et al.*, 2009).

The new distributional locality and specimens

The Evros river remains one of the poorest studied large river basins in the Balkans in terms of its ichthyofauna, and this is especially true of its lower section where several new discoveries have recently been made (KOTTELAT & ECONOMIDIS, 2006; FREYHOF *et al.*, 2008; ZOGARIS *et al.*, 2009a; KOUTRAKIS *et al.*, 2011). Although the Bulgarian section was rather well explored by the 1960's (RUSSEV, 1966), the Greek tributaries of the Evros were first surveyed for fish in 1970-72 by ECONOMIDIS (1974). After that, the wider border area was heavily militarized following worsening Greek-Turkish relations since the 1974 Cyprus conflict. Access to the 200 kilometer border zone of the Evros river in Greece and in Turkey is strictly restricted and very few fish surveys have been conducted. Therefore, it is possible that even native fish species or long-established aliens may have gone unnoticed in the lower Evros.

Members of the Institute of Inland

Waters-HCMR (IIW-HCMR) recently begun electrofishing surveys and have now widely sampled the Evros' Greek tributaries. Between 2009-2011, fifty-five river sites have been surveyed for their ichthyofauna within Greece's section of the Evros river basin, following the rapid quantitative sampling protocol (described in ECONOMOU *et al.*, 2007). During one of these surveys, on the 25th of August 2011, eight specimens of *N. fluviatilis* were caught on a one-bank 500 meter longitudinal transect of the main stem of the Evros using a Smith-Root 24 volt back-pack electrofisher (battery output voltage: 50 to 990V; 400W maximum continuous). The location of the sampling site (40° 58' 35 N, 26°19'40 E) is 65 river kilometers from the Evros' river-mouth between the villages of Tychemo and Gemisti at an elevation of about 7 meters above sea level. Eighteen fish species were caught at this site including the following aliens in descending order of abundance: *Carassius gibelio* (68 individuals), *Pseudorasbora parva* (13), *Lepomis gibbosus* (6), *Gambusia holbrooki* (5). The only other goby species present was the native *Proterorhinus semilunaris* (25 individuals). *Neogobius fluviatilis* inhabited sand and fine gravel substrates in the main channel of the river near its shoreline at depths ranging from 30 to 60 cm. The length of the collected specimens ranged between 58 – 71 mm *SL*. Five of the collected specimens were preserved in alcohol (80% solution) and three were transported alive to an aquarium (Fig.1).

N. fluviatilis is easily identified from other large freshwater gobiid species by the following attributes (PINCHUK *et al.*, 2003; KOTTELAT & FREYHOF, 2007): (1) the second dorsal fin is uniformly lowering down from the first rays to the last ones; (2) the first branched ray of the second dorsal fin is about twice as long as the penultimate ray;



Fig. 1: First *Neogobius fluviatilis* (Pallas, 1814) caught in Greece (Photo: S. Zogaris, 27.08.2011; ex situ at HCMR-Anavissos).

and (3) the lack of a black spot in the posterior part of the first dorsal fin. Also, the width of its fairly large head is equal to or a bit greater than the height of the head, and terminates in a leaf-shaped snout. Morphologically the specimens of the Evros are immediately identified as *N. fluviatilis*, as they show all the above characters. However no complete morphological or comparative genetic work has ever been published on samples from the Evros.

Distribution in adjacent basins of Bulgaria and Turkey

Regular sampling by the Institute of Biodiversity and Ecosystem Research (IBER) in Bulgaria has verified that the species is

fairly widespread in the whole Bulgarian Danube section, the lower section of some Danube tributaries, and several Black Sea coastal lakes and rivers of the country (Fig 2). It has also been recorded in the Black Sea shore in sandy bottoms and shallow waters (1-2m depth) such as in Butamia Bay (42° 03' 24 N, 27° 59' 16 E) near Sinemorets village. Despite this, the species is presumably absent from several sites, such as Bulgaria's Northern Black Sea lakes Shabla-Ezherits complex, Durankulak and near Varna. The Bulgarian populations outside the Danube/Black Sea area are provisionally considered as non-native (POLACIK *et al.*, 2008). The only known populations within the Evros basin are in the upper section of



Fig. 2: Recently documented *Neogobius fluviatilis* distribution in Bulgaria and NW Turkey (data from IBER surveys and available bibliography). Where specific location information is not known, but presence has been confirmed in the basin, a question mark symbol (?) is shown. The new site in the Evros is shown with a five-point star symbol.

the Tundja River (42° 03' 48 N, 26° 30' 18 E), where the species was first collected in 2006; and an unconfirmed population in a reservoir within the Sazliika tributary of the Evros (42° 18' 12 N, 25° 53' 45 E) (STEFANOV *et al.*, 2008).

Our observations of *N. fluviatilis* nearly on the Greek-Turkish border and its establishment in a large section of the Bulgarian Tundza (180 river kilometers upstream of our new find), suggest that the species is most likely established in Eastern Thrace, within Turkish territory. However, we have found no published records of the species in the Turkish tributaries of the Evros (ERKAKAN, 1983; BALIK, 1985; ILHAN, A., *pers. com.*). Furthermore although the species is known to occupy a wide area, including rivers and wetlands on the south-

ern shores of the Sea of Marmara (KOTTELAT & FREYHOF, 2007), specific geographic documentation is presumably scarce in the published literature (SARI *et al.*, 2006; ÖZULUĞ *et al.*, 2007; SASI & BERBER, 2010) (Fig. 2).

N. fluviatilis is often one of the most abundant fish species in its native range (KOTTELAT & FREYHOF, 2007) and this has been confirmed in the Bulgarian Black Sea basins. Survey data from recent IBER surveys in Bulgaria always showed higher biomass of *N. fluviatilis* in comparison with other gobies, including the invasive *Neogobius melanostomus* when more than one species are established at the same site. Only in the case of Tundza and Evros does *N. fluviatilis* show lower abundance in comparison with *Proterorhinus semillunaris*. The

abundance of *N. fluviatilis* in the Tundza is 4-10 specimens per 100 m river transect using electrofishing (perhaps similar to the situation in the new site on the lower Evros). The fact that *N. fluviatilis* tends to inhabit primarily the lower courses of rivers including the main stem of the Danube (unlike *Neogobius gymnotrachelus* which is also met in upland cool-water locations), indicates the species' preference for larger lotic waters and even slightly brackish water.

Are the Evros' Pontian Monkey Gobies alien?

In our opinion, the newly recorded population of *N. fluviatilis* is probably not native to the Evros basin, but there is currently no reliable evidence to prove that it was definitely absent from this river in the past. Based on the species' conspicuous behavior and rather large size, it is unlikely that it would go unnoticed by past researchers, if it was native. The most probable hypothesized vector for invasion is that it may have been transferred accidentally through fish stocking practices to the Tundza or to Bulgarian artificial reservoirs, and has now established localized river populations. Fish stocking and associated translocations are frequent in the Bulgarian part of the basin and several freshwater organisms have infiltrated to Greece through trans-boundary rivers, especially from Bulgaria (ECONOMIDIS *et al.*, 2000; ZENETOS *et al.*, 2009).

However, caution is needed in accepting an alien status for *N. fluviatilis* in the Evros due to the close proximity of this river to the species' native distribution range (i.e. Sea of Marmara). Moreover, if the Evros and the Sea of Marmara basins are located within a common biogeographical ecoregional area should the alleged non-native be called an alien or a translocated species? Although there is still some controversy regarding regional biogeographical bound-

aries (ZOGARIS *et al.*, 2009b), the 'Thrace Freshwater Ecoregion' as defined in ABELL *et al.* (2008) includes the entire Marmara basin and the wider northeastern Aegean basins (Evros, Nestos and Strymon). In fact, this broad freshwater biogeographical ecoregion delineation seems to be supported by evidence from regional geological history, creating conditions that may challenge the alien status of the species within the Evros.

The Evros basin has been affected several times by influences of Black Sea/Marmara waters and associated biota since the time it is postulated to have had its river-mouth in the Sea of Marmara, 1.5 million years ago (OKAY & OKAY, 2002). Furthermore, in the Late Pliocene, the Tundza tributary is considered to have been independently flowing directly into the Black Sea and later became a branch of the Evros due to a river-capture event (ANGELOVA, 2003; STEFANOV *et al.*, 2008). During the Pleistocene, the biogeographic relationship between the Evros and the Black Sea/Sea of Marmara biota is said to have had a direct connection due to the glacially-affected outflow of Black Sea water through the Dardanelles to the northern Aegean land-mass. This hypothesis, known as the Aegeopotamos, proposes that the Black Sea freshwater flood outflow had direct confluences with the Thracian rivers during this geological event (KOSSWIG & BATTALGIL, 1943; BANARESCU, 2004). This unique geological history may explain the native status of several Pontic and Anatolian species in the Evros, including the native population of Ponto-Caspian *Proterorhinus semillunaris* and the presence of the rare cobitid, *Cobitis puncticulata*, formerly considered unique to the Sea of Marmara wetlands, but recently also discovered in the Evros tributary at Lira in Greece (FREYHOF *et al.*, 2008).

All the above suggest that it is best to

provisionally consider *N. fluviatilis* in the Evros as “questionable” with respect to its alien status, since the origin of the Evros specimens have not been genetically inspected. A questionable status (as *per* ZENETOS *et al.*, 2010) may be characterized by insufficient or unverified information to confirm alien status. It is well known that labeling a species as ‘alien’ requires extreme caution (CARLTON, 1996) and must include specific biogeographical considerations (COP *et al.*, 2005). The difficulty of interpretation is greater when the alleged alien is discovered immediately next to its native range, within a common ecoregion, and where historic species distribution data are scarce or non-existent. A genetic and morphologic comparison among *N. fluviatilis* from the Evros, Danube, Marmara and Black Sea basins would be appropriate for the clarification of the Evros specimens’ phylogeny and population origins. Further surveys and monitoring are urgently needed within the Evros river basin, in order to assess potential “invasive” behavior of the species; including range alterations, population trends, and habitat-use in both lotic and transitional waters.

Acknowledgements

The EcoDOM project partially funded surveys in the Evros in 2011 and we are grateful to Elias Dimitriou and all members of the Institute of Inland Waters-HCMR for assistance. The World Wildlife Fund-Greece provided local support at Dadia National Park and volunteers in field work, including: Diogenis Kiziridis, Markos Tsigganis, Yann Lalaoui, Julia Meis, Marisa Weber and Özgül Yahyaoglu. We are especially indebted for local assistance to Theodora Skartsi (WWF Greece), Andreas Athanasiades (Evros Delta Management Body) and Gian-

na Arabatzi (Soufli Forest Service). We thank Radek Šanda, Marcelo Kovačić, Jörg Freyhof, Ali İlhan and Müfit Özuluğ for relevant discussions and distributional information. Aris Vidalis (Biodiversity East) assisted with cartography and Emily K. Green reviewed the manuscript for English. Finally, we are greatly indebted to Panos S. Economidis for assistance within an early draft of this manuscript.

References

- ABELL, R., THIEME, M.L., REVENGA, C., BRYER, M., KOTTELAT, M. *et al.*, 2008. Freshwater ecoregions of the world: a new map of biogeographic units for freshwater biodiversity conservation. *BioScience*, 58: 403-41.
- ANGELOVA, D., 2003. Endodynamic determination and rearrangement of the valley systems in Bulgaria at the Pliocene – Quaternary boundary. *Geografia Fisica e Dinamica Quaternaria*, 6: 19-24.
- BALIK, S., 1985. Taxonomical revision and present situations of the inland water fishes from Turkish Thrace. *Tubitak Doğa Bilim Dergisi*, 9(2): 147-160.
- BANARESCU, P.M., 2004. Distribution pattern of the aquatic fauna of the Balkan Peninsula. P. 203–217. In: *Balkan biodiversity: pattern and process in the European hotspot*. H.I. Grifóths, Krystufek B., & Reed J.M. (Eds). Kluwer Academic Publishers, Dordrecht.
- CARLTON, J.T., 1996. Biological invasions and cryptogenic species. *Ecology*, 77 (6): 1653-1655.
- COPP, G.H., BIANCO, P.G., BOGUTSKAYA, N.G., ERŐS, T., FALKA, I. *et al.*, 2005. To be or not to be a non-native freshwater fish? *Journal of Applied Ichthyology*, 21: 242-262.
- ECONOMIDIS, P.S., 1974. *Morphological*,

- taxonomical and zoogeographical study of the freshwater fish of E. Macedonia and W. Thrace*. PhD Thesis. Aristotle University of Thessaloniki, Thessaloniki, Greece, 179 pp. (In Greek).
- ECONOMIDIS, P. S., DIMITRIOU, E., PAGONI, R., MICHALOUDI, E. & NATSIS, L., 2000. Introduced and translocated fish species in the inland waters of Greece. *Fisheries Management & Ecology*, 7(3): 239-250.
- ECONOMOU, A.N., ZOGARIS, S., CHATZINIKOLAOU, Y., TACHOS, V., GIAKOUMI, S. *et al.*, 2007. Development of an ichthyological multimetric index for ecological status assessment of Greek mountain streams and rivers. Technical report. Hellenic Center for Marine Research – Institute of Inland Waters / Hellenic Ministry for Development, 189 pp. (in Greek)
- ERKAKAN, F., 1983. The fishes of the Thrace Region. *Hecettepe Bulletin of Natural Sciences & Engineering* (Ankara), 12: 39-48.
- FREYHOF, J., STELBRINK, B., ÖZULUĞ, M. & ECONOMIDIS, P.S., 2008. First record of *Cobitis punctulata* from Europe with comments on its conservation status (Teleostei: Cobitidae). *Folia Zoologica*, 57 (1-2): 16-19.
- KOSSWIG, C. & BATTALGIL, F., 1943. Zoogeographie der türkischen Süßwasserfische. *Review of the Faculty of Science University of İstanbul*, Série B 7:145-165.
- KOTTELAT, M. & ECONOMIDIS, P.S., 2006. *Squalius orpheus*, a new species of cyprinid fish from Evros drainage, Greece (Teleostei: Cyprinidae). *Ichthyological Exploration of Freshwaters*, 17(2): 181-186.
- KOTTELAT, M., & FREYHOF, J., 2007. *Handbook of European Freshwater Fishes*. Publications Kottelat, Cornol, Switzerland, 646 pp.
- KOUTRAKIS, E., SAPOUNIDIS, A., FAVRE-KREY, L., KREY, G. & ECONOMIDIS, P.S., 2011. Incidental catches of Acipenseridae in the estuary of the River Evros, Greece. *Journal of Applied Ichthyology*, 27: 366-368.
- NEILSON, M.E. & STEPIEN C.A., 2011. Historic speciation and recent colonization of Eurasian monkey gobies (*Neogobius fluviatilis* and *N. pallasii*) revealed by DNA sequences, microsatellites, and morphology. *Diversity & Distributions*, 17 (4): 688-702.
- OKAY, N. & OKAY, A., 2002. Tectonically induced Quaternary drainage diversion in the northeastern Aegean. *Journal of the Geological Society*, 159 (4): 393-399.
- ÖZULUĞ, M., TARKAN A.S., GAYGUSUZ Ö. & GÜRSOY Ç., 2007. Two new records for the fish fauna of lake Sapanca Basin (Sakarya, Turkey). *Journal of Fisheries Sciences*, 1 (3): 152-159.
- PINCHUK, V.I., VASIL'eva, E.D., VASIL'EV, V.P., & MILLER, P., 2003. *Neogobius fluviatilis* (Pallas, 1814). p.223-252. In: *The freshwater Fishes of Europe*. Mugilidae, Atherinidae, Atherinopsidae, Blenniidae, Odontobutidae, Gobiidae 1. Miller, P. (Ed). Aula-Verlag, Wiebelsheim.
- POLACIK, M., JANAC, M., TRICHKOVA, T., VASSILEV, M., KECKEIS *et al.*, 2008. The distribution and abundance of the *Neogobius* fishes in their native range (Bulgaria) with notes on the non-native range in the Danube River. *Archiv für Hydrobiologie*, 18: 193-208.
- RUSSEV, B., 1966. Hydrobiologische Untersuchungen der Marica.I. p.231-291. – In: *Die Fauna Thrakiens*. Sammelwerk. Band III. Sofia. Verlag der Bulgarischen

- Akademie der Wissenschaften, Sofia.
(in Bulgarian)
- SARI, H.M., BALIK, S., USTAOĞLU, M.R. & ILHAN, A., 2006. Distribution and Ecology of Freshwater Ichthyofauna of the Biga Peninsula, North-western Anatolia, Turkey. *Turkish Journal of Zoology*, 30: 35-45.
- SASI, H. & BERBER, S., 2010. Some biological characteristics of the Monkey Goby in Anatolia. *Asian Journal of Animal & Veterinary Advances*, 5(3): 229-233.
- STEFANOV, T., VASSILEV, M., & DOBREV, D., 2008. First occurrence of *Neogobius fluviatilis* (Pallas, 1814) (Pisces: Gobiidae) in the Aegean Sea Watershed. *Acta Zoologica Bulgarica*, 60 (3): 331-334.
- VAN KESSEL, N., DORENBOSCH, M. & SPIKMANS, F., 2009. First record of Pontian monkey goby, *Neogobius fluviatilis* (Pallas, 1814), in the Dutch Rhine. *Aquatic Invasions*, 4(2): 421-424.
- ZENETOS, A., PANCUCCI-PAPADOPOULOU, M.A., ZOGARIS, S., PAPA-STERGIADOU E., VARDAKAS, A., *et al.*, 2009. Aquatic alien species in Greece (2009): Tracking sources, patterns and effects on the ecosystem. *Journal of Biological Research –Thessaloniki*, 12: 135 - 172.
- ZENETOS, A., GOFAS, S., VERLAQUE, M., ÇINAR, M.E., GARCÍA RASO, E. *et al.*, 2010. Alien species in the Mediterranean Sea by 2010. A contribution to the application of European Union's Marine Strategy Framework Directive (MSFD). Part I. Spatial distribution. *Mediterranean Marine Science*, 11 (2): 381-493.
- ZOGARIS, S., ECONOMOU, A.N. & DIMOPOULOS, P., 2009a. Ecoregions in the Southern Balkans: should they be revised? *Environmental Management*, 43: 682-697.
- ZOGARIS, S., VARDAKAS, L., ECONOMOU, A.N. & ECONOMIDIS, P.S., 2009b. A preliminary list of fishes of the Dardia – Lefkimi – Soufli Forest National Park. p. 135-137. In: *The Dardia-Lefkimi-Soufli Forest National Park, Greece: Biodiversity, Management, and Conservation*. G. Catsadorakis & H. Kallander (Eds). WWF-Greece, Athens.