



Mediterranean Marine Science

Vol 11, No 2 (2010)



Catch and size selectivity of small-scale fishing gear for the smooth-hound shark Mustelus mustelus (Linnaeus, 1758) (Chondrichthyes: Triakidae) from the Aegean Turkish coast

T. CEYHAN, O. HEPKAFADAR, Z. TOSUNOGLU

doi: <u>10.12681/mms.73</u>

To cite this article:

CEYHAN, T., HEPKAFADAR, O., & TOSUNOGLU, Z. (2010). Catch and size selectivity of small-scale fishing gear for the smooth-hound shark Mustelus mustelus (Linnaeus, 1758) (Chondrichthyes: Triakidae) from the Aegean Turkish coast. *Mediterranean Marine Science*, *11*(2), 213–224. https://doi.org/10.12681/mms.73

Catch and size selectivity of small-scale fishing gear for the smooth-hound shark *Mustelus mustelus* (Linnaeus, 1758) (Chondrichthyes: Triakidae) from the Aegean Turkish coast

T. CEYHAN, O. HEPKAFADAR and Z. TOSUNOĞLU

Ege University, Faculty of Fisheries, 35100 Bornova, Izmir, Turkey

Corresponding author: tevfik.ceyhan@ege.edu.tr

Received: 29 September 2009; Accepted: 28 June 2010; Published on line: 8 October 2010

Abstract

Catch rate, CPUE, biomass ratios and size selectivity from traditional longline and trammel nets of Turkish coastal small-scale fisheries were investigated in order to describe the smooth-hound shark (*Mustelus mustelus*) fishery. The SELECT method was used to estimate the selectivity parameters of a variety of models for the trammel nets inner panel of 150 and 170 mm mesh sizes. Catch composition and proportion of the species were significantly different in longline and trammel nets. While mean CPUE of longline was 119.2 ± 14.3 kg/1000 hooks, these values for 150 and 170 mm trammel nets were 5.3 ± 1.2 kg/1000 m of net and 12.7 ± 3.9 kg/1000 m of net, respectively. Biomass ratios of the by catch to smooth-hound catch were found to be 1:0.32 for 150 mm trammel net, 1:0.65 for longline and 1:0.73 for 170 mm trammel net. The estimated modal lengths and spreads were found to be 91.1 and 16.2 cm for 150 mm and 103.2 and 18.4 cm for 170 mm, respectively. The modal lengths of the species as well as the spread values increased with mesh size.

Keywords: Smooth-hound shark; Mustelus mustelus; CPUE; Biomass ratio; Selectivity.

Introduction

Because of their K-selected life-history strategy (characterized by slow growth, late attainment of sexual maturity, long life spans, low fecundity and natural mortality, and a close relationship between the number of young produced and the size of the breeding biomass), sharks and rays appear to be particularly vulnerable to overexploitation (STEVENS *et al.*, 2000). This results in a low level of recruitment seldom capable of keeping pace with modern fishing technology (HOLTS *et al.*, 1998).

Globally, shark catches are divided into targeted and by-catch fisheries. Few elasmobranchs are subject to directed fisheries in the Mediterranean, with local fisheries mostly landing elasmobranchs as by-catch (FOWLER *et al.*, 2005). A similar situation occurs in Turkey. Targeted fisheries for the sharks have developed, due to the increase in domestic consumption around tourist areas, and export activity. Elasmobranch fisheries have globally been common and traditional with lesser importance around the world (BONFIL, 1994). In Turkish Seas, 64 elasmobranches species were reported (BILE-CENOĞ LU *et al.*, 2002). However, 38 species of these fish have a commercial value (FILIZ and TOĞ ULGA, 2002). In 2006, the harvested amount of chondrichthyans was 1532 t in Turkey, which is just 0.2% of the total elasmobranch catch in the world (FAO, 2000).

Despite the major socio-economic importance of small-scale fisheries in the whole eastern Mediterranean Sea, many aspects have not been studied comprehensively. In particular there is a lack of information on catch composition, catch rates and size selectivity of the mesh size in trammel net shark fishery. The main objectives of this study were to describe and compare elasmobranch catches, size selectivity for smooth-hound shark (*Mustelus mustelus*) in small-scale coastal fisheries, using trammel nets and longline in the Izmir Bay, Aegean Sea.

Material and Methods

A total of 22 fishing trials with trammel nets were carried out in the same fishing grounds visited by local fishermen from September 2006 to May 2007 in the Izmir Bay, Aegean Sea. In these trials, fishing depths varied from 28 to 55 m. Normal fishing practices were followed, with the setting of the gear during the daytime or afternoon and hauling after sunset. The stretched mesh sizes of the trammel nets were 150 and 170 mm and the total length of the nets were 2000 m and 5000 m, respectively. The hanging ratio was 0.5 for both mesh sizes of nets.

Experimental fishing trials with the longline were carried out by 6 trials in 2007, from July to August. Fishing was carried out by a commercial fishing vessel and took place on traditional fishing grounds. Fishing depths varied from 50 to 60 m. The longline used consisted of a 1 mm diameter monofilament main line with 0.7 mm diameter monofilament at intervals of approximately 9.2 m. These longlines were stored in two baskets each containing 200 hooks. The hook number used was 7, manufactured by Mustad. The total length of the longlines was 4 km.

All captured fishes were sorted, identified and measured (total length and weight) as they came aboard. The specimens of *M. mustelus* were separated by sex. Test for significance (p < 0.05) between total length and weight between males and females were performed by using Student t-test. Comparisons of differences between sex ratios of smooth-hound shark according to trials of each fishing gear were tested by Kruskal-Wallis H test.

Scientific names for each species were checked and confirmed using Fishbase (FROESE and PAULY, 2007).

Chi-square (χ^2) tests were performed to test variations in actual catches (kg) of species by type of fishing gear.

Fishing effort (f) and CPUE were calculated using following formula, modified from DE METRIO and MEGALOFONOU (1988): f = (a'/10) x g. where (a'/10) represents the average length of the nets and average number of hooks in longline, placed daily in the sea divided by the 10 net units. Therefore, a 10 net unit is equal to 10 x 100 = 1000 m for gillnets and $10 \times 100 = 1000$ hooks for longline. "g" is the number of fishing days. The CPUE was computed in biomass with the formula, CPUE = kg/f. Means were given with standard error $(\pm SE)$. Comparisons of differences between CPUEs of the three types of gear were tested by Kruskal-Wallis H test.

The SELECT (share each length class

catch total) method (MILLAR, 1992) was used to estimate the selectivity of the trammel nets. For a given length class, l, the numbers of fish, nlj, that encounter trammel net j are assumed to be observations of independent Poisson random variables,

 $n_{ij} \approx \text{Pois} (p_j \lambda_l r_j (l))$ where the expected count, $p_j \lambda_l$, is the product of the abundance of length class l fish, λ_l , and the relative fishing intensity of trammel net j, p_j . Relative fishing intensity of a trammel net is a combined measure of fishing effort and fishing power.

The log-likelihood of n_{lj} is

$$\sum_{l} \sum_{j} \{ n_l \log_e [p_j \lambda_l r_j (1)] - p_j \lambda_l r_j (1) \}$$

An appropriate software (Gillnet, ConS-

tat-DK) was used here for selectivity estimation (equal power over mesh sizes was assumed). How good the fit was was evaluated by comparison of deviances, the lowest deviance value corresponding to the best fitting model (DOS SANTOS *et al.*, 2003; ERZINI *et al.*, 2003), and the analysis of residual plots as in MILLAR and HOLST (1997). A collection of the most commonly used selection curves (HOLST *et al.*, 1996), estimated by this software, is given in Table 1.

Results

A total of 190 specimens of smooth-hound, 110 males and 80 females, were examined. The smallest specimen was 34 cm TL and weighed 350 g. The largest specimen was 141.1

Table	1
-------	---

Normal (fixed spread), normal (proportional spread), gamma, and log-normal selection curves (*l*= fish length, =dispersion/variance, m=mean, k= optimum catch length).

Model	Selection Curve
Normal location	$\exp\left(-\frac{(1-k_1\cdot m_j)^2}{2\sigma^2}\right)$
Normal scale	$\exp\left(-\frac{(1-k_1\cdot m_j)^2}{2k_2^2\cdot m_j^2}\right)$
Log-normal	$\frac{m_j}{l \cdot m_l} \exp\left(\mu - \frac{\sigma^2}{2} - \frac{\left(\log(l) - \mu - \log\left(\frac{m_j}{m_j}\right)\right)^2}{2k_2^2 \cdot m_j^2}\right)$
Gamma	$\left(\frac{l}{(a-1)\cdot k\cdot m_j}\right)^{a-1} \exp\left(a-1-\frac{l}{k\cdot m_j}\right)$
Bi-normal	$\exp\left(-\frac{(l-k_l\cdot m_j)^2}{2k_2^2\cdot m_j^2}\right) + c\cdot \exp\left(-\frac{(l-k_3\cdot m_j)^2}{2k_4^2\cdot m_j^2}\right)$

cm TL and weighed 7.5 kg (Table 2). There was no significant relationship between total mass and TL between both sexes (p>0.05).

Sex ratios of smooth-hound were found to be 1:0.59 for longline, 1:0.67 for 170 mm trammel net and 1:0.97 for 150 mm trammel net. (Table 3). No significant differences were identified between the sex ratios of smooth-hound shark according to trials of each fishing gear (KW=1.415, p>0.05).

In the longline fishery, three different elasmobranch species were captured constituting 72% of the total catch. These species were *Mustelus mustelus* (comprising 84% of elasmobranchs), *Myliobatis aquila* (10%) and *Raja clavata* (6%). In the trammel net fishery of 150 mm mesh size, 7 elasmobranch species were caught and their catches accounted for 92% of the total catch. 97% of the total catch was elasmobranchs in the second mesh size of trammel net (170 mm) and smooth-hound shark comprised 58% of the total catch (Table 4). Catch compositions were significantly different among fishing gears ($\chi^2 = 653.573$, p<0.001).

Fishing effort (f) was calculated to be 0.4, 5 and 2 for longline, 150 and 170 mm mesh size trammel net, respectively. CPUEs relative to fishing gears are shown in Table 5. Mean CPUE for the longline was 119.2 ± 14.3 kg/1000 hooks and mean CPUE for the 150 and 170 mm trammel nets were 5.3 ± 1.2 kg/1000 m and 12.6 ± 3.9 kg/1000 m, respectively. Median CPUE values differed significantly among gear types (KW=16.45, p < 0.05).

Biomass ratios of the by-catch to smoothhound were found to be 1:0.32 for 150 mm trammel net, 1:0.65 for longline and 1:0.73 for 170 mm trammel net (Table 6). No significant differences were identified between weight of by-catch species and weight of the smooth-hound shark by fishing gear (KW=2, p>0.05).

The estimated results of the SELECT model for the two mesh sizes of trammel nets are given in Table 7. The normal scale mod-

		TL (mm)				W (g)	
Sex	Ν	Range	Mean	SE	Range	Mean	SE
Male	110	38.85-141.1	106.21	2.31	350-7500	4724.35	190.45
Female	80	34-138.1	102.33	2.63	450-7300	4245.26	211.07
Male+ Female	190	34-141.1	104.58	1.73	350-7500	4522.63	142.29

 Table 2

 The composition of total length and weight of *M. mustelus*.

 Table 3

 Sex ratio of the smooth-hound by the different gear trials.

	Number o		
Gear	Male	Female	Sex ratio
Longline	32	19	1:0.59
Trammel net 150 mm	29	28	1:0.97
Trammel net 170 mm	49	33	1:0.67
Σ	110	80	

 Table 4

 List of all species, total weight (kg) and weight % of each species captured in all fishing trials.

	Family	Species	Total Weight (kg)	%W
	Triakidae	Mustelus mustelus Linnaeus, 1758	286.1	60.4
Muraenidae		Muraena helena Linnaeus, 1758	70.5	14.9
	Congridae	Conger conger Linnaeus, 1758	53.5	11.3
Longling	Myliobatidae	Myliobatis aquila Linnaeus, 1758	33.6	7.1
Longinic	Rajidae	Raja clavata Linnaeus, 1758	19.9	4.2
	Sparidaa	Dentex dentex Linnaeus, 1758	4.2	0.9
	Spandae	Dentex gibbosus Rafinesque, 1810	3.2	0.7
	Triglidae	Trigla lyra Linnaeus, 1758	2.6	0.5
		$\Sigma =$	473.6	100
	Triakidae	Mustelus mustelus Linnaeus, 1758	176.8	75.5
	Dasyatidae	Dasyatis pastinaca Linnaeus, 1758	20.2	8.6
	Lophiidae	Lophius piscatorius Linnaeus, 1758	9.0	3.9
	Zeidae	Zeus Faber Linnaeus, 1758	6.1	2.6
	Myliobatidae	Myliobatis aquila Linnaeus, 1758	5.0	2.1
Trammel	Scyliorhinidae	Scyliorhinus stellaris Linnaeus, 1758	4.0	1.7
net 150 mm Uranoscopidae Rajidae	Scyliorhinus canicula Linnaeus, 1758	3.5	1.5	
	Uranoscopidae	Uranoscopus scaber, Linnaeus, 1758	3.6	1.6
	Rajidae	Dipturus oxyrinchus Linnaeus, 1758	1.9	0.8
		Raja clavata Linnaeus, 1758	1.9	0.8
		Rostroraja alba Lacepède, 1803	1.1	0.5
	Pleuronectidae	Pleuronectes platessa Linnaeus, 1758	0.9	0.4
		$\Sigma =$	234.0	100.0
	Triakidae	Mustelus mustelus Linnaeus, 1758	396.4	8.0
	Squalidae	Squalus acanthias Linnaeus, 1758	156.2	3.2
	Dasyatidae	Dasyatis pastinaca Linnaeus, 1758	54.9	2.8
	Scyliorhinidae	Scyliorhinus stellaris Linnaeus, 1758	22.2	2.1
T	Seynorminade	Scyliorhinus canicula Linnaeus, 1758	19.3	0.4
I rammei	Rajidae	Rostroraja alba Lacepède, 1803	14.4	1.6
net 170 mm	Rajidae	Raja asterias Delaroche, 1809	2.7	1.3
	Lophiidae	Lophius piscatorius Linnaeus, 1758	10.8	0.4
	Zeidae	Zeus faber Linnaeus, 1758	8.8	0.0
	Uranoscopidae	Uranoscopus scaber Linnaeus, 1758	3.0	0.4
	Pleuronectidae	Pleuronectes platessa Linnaeus, 1758	0.3	0.0
		$\Sigma =$	688.9	20.2

Gear	Total weight of catches (kg)	CPUE
	54.1	135.25
	36.8	92
,	36.9	92.25
Longline	57.7	144.25
	33.2	83
	67.4	168.5
	7	3.5
	18.8	9.4
	8.4	4.2
	17.1	8.55
net 150 mm	40.2	20.1
	65.1	32.55
	20.2	10.1
	0.6	0.12
	16.7	3.34
	7.3	1.46
	32.1	6.42
	7.3	1.46
	19.9	3.98
Trammel	23	4.6
net 170 mm	52	10.4
	28.1	5.62
	19.4	3.88
	43.2	8.64
	23.1	4.62
	10.9	2.18
	16.4	3.28
	96.4	19.28

Table 5Fishing effort and CPUE by gear type.

el had the lowest model deviance as bi-normal model. In fact, the bi-normal model provided the best fit, based on the residual diagnostic plots. The fitted selectivity curves of the two trammel net (150 and 170 mm) are shown in Figure 1, as well as the corresponding deviance residuals for smoothhound shark. The estimated modal lengths and spreads for the two trammel net mesh sizes for the best model are shown in Table 8. The modal lengths and spread values increased with mesh size.



Fig. 1: Selectivity curves of trammel net for the smooth-hound shark and deviance residual plots. Full circle indicates a positive residual and an open circle a negative residual.

Gear	Total	Total catch	Smooth-hound shark	By-catch	By catch ratio
	landings	(kg)	(kg)	(kg)	
Longline	6	473.6	286.1	187.5	1:0.65
Trammel net 150 mm	7	234.0	176.8	57.2	1:0.32
Trammel net 170 mm	15	688.9	396.4	292.5	1:0.73

 Table 6

 Summary of total catch and by-catch ratios by gear.

	Table	7		
Summary of total	catch and	by-catch	ratios b	y gear.

	Equal Fishing Power			
Model	Parameters	Model Deviance	d.f	P value
Normal Location	(k. s)= (0.559. 1817)	14.37	10	0.1570
(Fixed spread)				
Normal scale	(k1.k2) = (0.607. 0.108)	12.17	10	0.2740
(Spread α mj)				
Gamma				
(Spread α mj)	$(\alpha. k) = (0.023. 25.245)$	14.37	10	0.1570
Log normal				
(Spread α mj)	$(\mu 1. s) = (4.419. 0.219)$	17.08	10	0.0726
Bi-Normal				
(Spread α mj)	$(a1. b1. a2. b2. w) = (0.607 \ 0.108. \ 1.648. \ 0.082. \ 0.552)$	12.17	7	0.0952

Table 8 Modal length and spread values for the best-fitting model of gill net selectivity model curves.

Trammel net	Model Length	Spread
150 mm	91.1	16.2
170 mm	103.2	18.4

Discussion

Analysis of the catch compositions and catch rates of both types of fishing gear showed that catches of smooth-hound shark in longline are very high. COELHO *et al.* (2005) reported that catches of elasmobranchs were high in the longline fishery of the coasts of Southern Portugal, owing to setting in deeper waters than the trammel nets. This results from the fact that longlines with baited hooks attract fish from considerable distances (BJORDAL and LØKKEBORG, 1996), whereas trammel nets depend on the normal movements of fish.

In this study, mean CPUE of the longline was 119.2 ± 14.3 kg/1000 hooks and mean CPUEs of the trammel nets 150 and 170 mm were 5.3 ± 1.2 kg/1000m and 12.6±3.9kg/1000m, respectively. MEGALO-FONOU et al. (2005), studying large pelagic sharks in the surface drifting longline fishery, emphasized that CPUEs were 3.8 fish/1000 hooks in the Alboran Sea and 1 fish/1000 hooks in the Adriatic Sea. They also reported that shark CPUE peaked during late spring and summer. Furthermore, thresher shark CPUE in the driftnet fishery ranged from 0.13 to 1.92 fish/fishing set (HOLTS et al. 1998). These results are not directly comparable and can only be regarded as informative data. To begin with, the catch rates are expressed in different units. Secondly, these types of fishing gear operate in different types of habitats (different depths), and lastly the shark species are different. SERENA and VACCHI (1997) reported that large elasmobranchs are often caught incidentally as by-catch in artisanal fisheries, especially in longline fisheries and trammel nets set near the bottom. However, there is lack of data about the CPUE of deep longlines and trammel nets for deep water fisheries.

Despite the importance of gillnet selectivity in fisheries assessment and management, there are few estimates for sharks. Gillnet selectivity models were also estimated for Mustelus antarcticus (KIRKWOOD and WALKER, 1986), Carcharhinus tilstoni and Carcharhinus sorrah (MCLOUGHLIN and STEVENS, 1994), Carcharhinus ob-(SIMPFENDORFER scurus and UNSWORTH, 1998), Carcharhinus plumbeus (MCAULEY et al. 2007) and Scyliorhinus canicula (FONSECA et al., 2005). Furthermore, gillnet selectivity parameters for the Atlantic sharpnose Rhizoprionodon terraenovae, blacknose Carcharhinus acronotus, finetooth Carcharhinus isodon, and bonnethead Sphyrna tiburo, sharks were estimated in multi-panel gillnets off the southeastern United States (CARLSON and CORTÉS, 2003). No selectivity estimates are available for smooth-hound shark caught by trammel nets. In this study, the estimated modal lengths and spreads of the trammel nets increased with mesh size. Our results are higher than the estimated first maturity size of 80 cm for female and 70-74 cm for male (COMPAGNO, 1984; GOOSEN and SMALE, 1987). However, first maturity size for the Mediterranean was 971 and 1172 mm for males and females, respectively (SAÏDI et al., 2008). Because of that, mesh size for trammel nets targeting M. mustelus should be larger than 170 mm since there is good evidence that selective fishing mortality can lead to changes in growth and juvenile survival for both sharks and batoids, leading to changes in population dynamics (STEVENS et al., 2000). Furthermore, JENNINGS and KAISER (1998) note that fishing acts as a selective force and life-history traits such as growth that are at least partly inheritable may be expected to evolve under sustained exploitation.

CAPAPEÉ et al. (2006) reported male specimens were ranging between 39 cm and 139 cm TL and weighing 195 g and 11 kg. The female smooth-hounds fluctuated within 39 cm and 150 cm TL, weighing between 201g and 8.1 kg for the coast of Senegal. In this study, male smooth-hound examined ranged between 38.85 cm and 141.1 cm TL and weighed between 350g and 7.5 kg. The females examined ranged between 34 cm and 138.1 cm TL and weighed between 450g and 7.3 kg. Because of the environment differences, the disagreements between minimum and maximum measurements in CAPAPEÉ et al. (2006) and this study are more or less an expected result.

Many coastal fishers are facing the dilemma of fisheries' collapse, the search for income, and the difficulty in sustaining fishing livelihoods (BERKES, 2001). BONFIL (1994) emphasized that fisheries for sharks and rays were common throughout the world and differ in both the species taken and in the type of gear and vessels used. This diversity has contributed to the difficulty in studying the fisheries and to the problems of collecting accurate data on yields and fishing effort. Statistics for elasmobranchs around the world need to be improved. Much data compilation and reviewing must be done on a country and regional basis to enable appraisal of exploitation levels and to make assessments of the status of elasmobranch stocks. In this study, we have put forth the fisheries and the selectivity data for the first time to develop management strategies for the smooth-hound shark. Further studies on animal removals (landing/discards as well as some key factors of the species biology (age, growth, reproductive biology) of smooth-hound and other sharks in the region are essential.

Acknowledgements

We thank two anonymous reviewers for their insightful comments which led to a much improved manuscript.

References

- BERKES, F., MAHON, R., MCKINNEY, P., POLLNAC, R. & POMEROY, R., 2001. *Managing small-scale fisheries; alternative directions and methods*. Ottawa, Canada, International Development Research Centre (IDRC), 320 pp.
- BILECENOĞLU, M., TAŞKAVAK, E., MATER, S. & KAYA, M., 2002. Checklist of the marine fishes of Turkey. *Zootaxa*, 113: 1-194.
- BJORDAL, Å. & LØKKEBORG, S., 1996. Longlining. Oxford, Fishing News Books, 156 pp.
- BONFIL, R., 1994. Overview of world elasmobranch fisheries. FAO Fisheries Technical Paper. No.341, Rome, FAO, 119 pp.
- CAPAPEÉ, C., DIATTA, Y., DIOP, M., VERGNE, Y. & GUÉLORGET, O.,

2006. Reproductive biology of the smoothhound, *Mustelus mustelus* (Chondricthyes: Triakidae) from the coast of Senegal (eastern tropical Atlantic). *Cybium*, 30 (3): 273-282.

- CARLSON, J.K. & CORTES, E., 2003. Gillnet selectivity of small coastal sharks off the southeastern United States. *Fisheries Research*, 60 (2): 405-414.
- COELHO, R., ERZINI, K., BENTES, L., CORREIA, C., LINO, P.G., MONTEI-RO, P., RIBEIRO, J. & GONÇALVES, J.M.S., 2005. Semi-pelagic longline and trammel net elasmobranch catches in Southern Portugal: catch composition, catch rates and discards. *Journal of Northwest Atlantic Fishery Science*, 35: 531-537.
- COMPAGNO, L.J.V., 1984. FAO species catalogue. Vol. 4. Sharks of the world. An annotated and illustrated catalogue of shark species known to date. Part 2
 Carcharhiniformes. FAO Fisheries Synopsis, 125 (4): 251-655.
- DE METRIO, G. & MEGALOFONOU, P., 1988. Catch, size distribution, growth and sex ratio of swordfish (*Xiphias gladius* L.) in the Gulf of Taranto. *FAO Fisheries Report*, No. 394: 91-102.
- DOS SANTOS, M.N., GASPAR, M., MONTEIRO, C.C. & ERZINI, K., 2003. Gillnet selectivity for European hake *Merluccius merluccius* from southern Portugal: implications for fishery management. *Fisheries Science*, 69 (5): 873-882.
- ERZINI, K., GONÇALVES, J.M.S., BENTES, L., LINO, P.G., RIBEIRO, J. & STERGIOU, K. I., 2003. Quantifying the roles of competing static gears: comparative selectivity of longlines and monofilament gillnets in a multi-species fishery of the Algarve (southern Portugal). Scientia Marina, 67 (3): 341-352.
- FAO, 2000. Fishstat Plus: Universal Software for Fishery Statistical Time Series.

Ver. 2.3.2 Rome, FAO Fish. Dept., Data and Statistics Unit.

- FİLİZ, H. & TOĞULGA, M., 2002. Commercial Elasmobranch species in Turkey's waters, their fisheries and management (in Turkish). P. 717-727. In: *Türkiye'nin Kıyı ve Deniz Alanları IV. Ulusal Konferansı Bildiriler Kitabı*. E. Özhan ve N. Alpaslan (Eds.) 5-8 Kasım 2002, İzmir, Türkiye.
- FONSECA, P., MARTINS, R., CAMPOS, A. & SOBRAL, P., 2005. Gill-net selectivity off the Portuguese western coast. *Fisheries Research*, 73 (3): 323-339.
- FOWLER, S.L., CAVANAGH, R.D., CAMHI, M., BURGESS, G.H., CAILLIET, G.M., FORDHAM, S.V., SIMPFENDORFER, C.A. & MUSICK, J.A., 2005. Sharks, Rays and Chimaeras: The Status of the Chondrichthyan Fishes. Status Survey. IUCN/SSC Shark Specialist Group, Gland, Switzerland, 461 pp.
- FROESE, R. & PAULY, D., 2007. *FishBase*. World Wide Web electronic publication. www.fishbase.org, version (04/2008).
- GOOSEN, A.J.J. & SMALE, M.J., 1997. A preliminary study of age and growth of the smooth-hound shark *Mustelus mustelus* (Triakidae). *South African Journal of Marine Science*, 18 (1):85-91.
- HOLST, R., MADSEN, N., MOTH-POULSEN, T., FONSECA, P. & CAMPOS, A., 1996. *Manual for gillnet selectivity*. Denmark, ConStat, 43 pp.
- HOLTS, D.B., JULIAN, A., SOSA-NISHIZAKI, O. & BARTOO, N.W., 1998. Pelagic shark fisheries along the west coast of the United States and Baja California, Mexico. *Fisheries Research*, 39 (2): 115-125.
- JENNINGS, S. & KAISER, M.J., 1998. The effects of fishing on marine ecosystems. p. 201-352. In: *Advances in Marine Bi*ology. J.H.S. Blaxter, A.J. Southward,

and P.A. Tyler (Eds.). London, Academic Press.

- KIRKWOOD, G.P. & WALKER, T.I., 1986. Gill-net mesh selectivities for gummy shark, *Mustelus antarcticus* Gunther, taken in southeastern Australian waters. *Australian Journal of Marine and Freshwater Research*, 37 (6): 689-697.
- MCAULEY, R.B., SIMPFENDORFER, C.A. & WRIGHT, I.W., 2007. Gillnet mesh selectivity of the sandbar shark (*Carcharhinus plumbeus*): implications for fisheries management. *ICES Journal of Marine Science*, 64 (9): 1702-1709.
- MCLOUGHLIN, K.J. & STEVENS, J.D., 1994. Gill-net mesh selectivities for two species of commercial carcharhinid shark taken in northern Australia. *Australian Journal of Marine and Freshwater Research*, 45: 521-534.
- MEGALOFONOU, P., YANNOPOULOS, C., DAMALAS, D., DE METRIO, G., DELFLORIO, M., de LA SERNA J.M. & MACIAS, D., (2005) Incidental catch and estimated discards of pelagic sharks from the swordfish and tuna fisheries in the Mediterranean Sea. *Fishery Bulletin*, 103: 620-634.
- MILLAR, R.B., 1992. Estimating the sizeselectivity of fishing gear by conditioning on the total catch. *Journal of the Amer*-

ican Statistical Association, 87: 962-968.

- MILLAR, R.B. & HOLST, R., 1997. Estimation of gillnet and hook selectivity using log-linear models. *ICES Journal* of Marine Science, 54: 471-477.
- SAÏDI, B., BRADAÏ, M.N. & BOUAÏN, A., 2008. Reproductive biology of the smooth-hound shark *Mustelus mustelus* (L.) in the Gulf of Gabés (south-central Mediterranean Sea). *Journal of Fish Biology*, 72: 1343-1354
- SERENA, F. & VACCHI M., 1997. Attività di studio sui grandi pesci cartilaginei dell'Alto Tirreno e del mar Ligure nell'ambito del programma LEM (Large Elasmobranchs Monitoring). Quaderni della Civica Stazione Idrobiologica di Milano, 22: 121-123.
- SIMPFENDORFER, C.A. & UNSWORTH, P., 1998. Gill-net mesh selectivity of dusky sharks (*Carcharhinus obscurus*) and whiskery sharks (*Furgaleus macki*) from south-western Australia. *Marine & Freshwater Research*, 49 (7): 713-718.
- STEVENS, J.D., BONFIL, R., DULVY, N.K. & WALKER, P., 2000. The effects of fishing on sharks, rays and chimaeras (chondrichthyans), and the implications on marine ecosystems. *ICES Journal of Marine Science*, 57: 476-494.