

## Mediterranean Marine Science

---

Vol 11, No 2 (2010)

---



### First record of the broad-banded cardinal fish *Apogon fasciatus* (White, 1790) from Turkey

C. TURAN, D. YAGLIOGLU, D. ERGUDEN, M.  
GURLEK, B. SONMEZ

doi: [10.12681/mms.85](https://doi.org/10.12681/mms.85)

---

#### To cite this article:

TURAN, C., YAGLIOGLU, D., ERGUDEN, D., GURLEK, M., & SONMEZ, B. (2010). First record of the broad-banded cardinal fish *Apogon fasciatus* (White, 1790) from Turkey. *Mediterranean Marine Science*, 11(2), 369–372.  
<https://doi.org/10.12681/mms.85>

## First record of the broad-banded cardinal fish *Apogon fasciatus* (White, 1790) from Turkey

C. TURAN, D. YAGLIOGLU, D. ERGUDEN, M. GURLEK and B. SONMEZ

Fisheries Genetics Laboratory, Faculty of Fisheries, Mustafa Kemal University, 31220 Iskenderun, Hatay, Turkey

Corresponding author: [cturan@ymail.com](mailto:cturan@ymail.com)

Received: 29 March 2010; Accepted: 16 September 2010; Published on line: 9 November 2010

---

### Abstract

Two specimens of the alien cardinal fish *Apogon fasciatus* (White, 1790) are recorded for the first time in Turkey and second time in the Mediterranean Sea. This is the fourth Indo-Pacific apogonid species documented in the Mediterranean Sea, and the introduction of this species to the eastern Mediterranean is due to migration from the Red Sea via the Suez Canal.

**Keywords:** *Apogon fasciatus*; Alien species; Eastern Mediterranean; Turkey.

---

The number of Red Sea fish species, introduced into the Mediterranean through the Suez Canal has reached 77 (ZENETOS *et al.*, 2008; GOLANI *et al.*, 2008; GOREN *et al.*, 2008; GOREN *et al.*, 2009; GOLANI *et al.*, 2010). Recently, GOREN *et al.* (2009) and GOREN *et al.* (2010) have added two additional alien species.

To date, forty nine alien fish species, originally from the Red Sea and the Indo-Pacific Ocean have been reported in Turkish waters (BILECENOGLU *et al.*, 2010). Recently one species was added to the Turkish ichthyofauna: *Pomadasys stridens* was recorded from Iskenderun Bay (BILECENOGLU *et al.*, 2009). An additional new species, *Apogon fasciatus* (White, 1790) is reported here.

The Apogonidae family has been represented in the Turkish waters up until recently by two species: the native species *Apogon imberbis* L. and the Lessepsian immigrant *Apogon pharonis* Bellotti, 1874 until 2005. However, in recent years two new Indo-Pacific cardinal fish have been recorded: *Apogon queketti* Gilchrist, 1903 (ERYILMAZ & DALYAN, 2006) and *Apogon smithi* (Kotthaus, 1970) (GOREN *et al.*, 2009) from the Turkish coasts of the Eastern Mediterranean.

Two specimens of broad-banded cardinal fish, 62 and 52 mm in total length (TL), 50 and 41 mm in standard length (SL), were collected by a commercial purse seiner (F/V Koca Reis) from Iskenderun Bay (36° 10' N - 35° 51' E), at a depth of 36 m on 25

March 2010 (Fig. 1). The specimens were preserved in 4% formalin. Samples were deposited at the Museum of the Faculty of Fisheries, Mustafa Kemal University, Iskenderun-Hatay (collection numbers: MSM-PIS/2010-1).

Descriptions of the examined specimens are as follows: first dorsal fin rays VII; the first spine is very short, the third spine is longest in the first dorsal fin and very robust. Second dorsal fin ray I + 9, anal fin ray 2 + 9, pectoral fin rays 15-16, caudal fin rays 22, developed gill rakes on first gill arch 16, three-four large scales on the opercula, a lateral line with pored scales 25-26 and 2 scales without pores on the base of the caudal fin. Head length 3.5-3.7 body depth 3.2-3.3, caudal peduncle depth 1.2-1.3 all at standard length (SL). Eye diameter 2.6-2.9, snout length 1.9-2.0,

postorbital length 0.8-0.9, pre-opercular length 0.7-0.9, interorbital space 1.5-1.7 all at head length. The body color is silver on the sides and whitish on the belly, with a broad, blackish mid-lateral stripe which extends to the end caudal fin. The dorsal and anal fins are in part whitish, and there is a dark bar on the rear margin of the fin (Fig 2). The morphometric measurements, meristic characters and color agree with descriptions of *Apogon fasciatus* given by FRAUSER (2005) and GOREN *et al.* (2009).

The broad-banded cardinal fish has a widespread distribution throughout the Red Sea, Mozambique, Fiji, New Caledonia, the Persian Gulf and east to the western Pacific (FRASER, 2005). The broad-banded cardinal fish inhabits coastal reefs and sandy or weedy areas, which can be found within




Fig. 1: Sampling location (●) of *Apogon fasciatus*.



**Fig. 2:** *Apogon fasciatus* (White, 1790) from Iskenderun Bay, Turkey.

the depth range of 2-128 m (LIESKE & MYERS, 1994; FROESE & PAULY, 2010). *A. fasciatus* reaches the size of 10.3 cm and feeds on zooplankton (RANDALL, 2005; GOREN *et al.*, 2009).

*A. fasciatus* is clearly distinguished from the other four Apogonids by a blackish mid-lateral stripe which extends to the end caudal fin on the body. *Apogon pharonis* has three dark, vertical bars on the flanks between the anterior of the dorsal and the caudal fin, and a dark ocellus within the first bar. *Apogon queketti* has dark spots on the scales, from 6-7 longitudinal stripes on the body and a black stripe on the margin of the anal fin. There is a black spot on the upper part of the first dorsal fin from the third spine to the seventh spine. The native species *Apogon imberbis* can easily be dis-

tinguished from lessepsian apogonid species by its red body colour and six spines in the first dorsal fin.

The first record of *A. fasciatus* from the Mediterranean Sea was given by GOREN *et al.* (2009) on the coast of Ashdod in Israel, collected by a trawler at a depth of 40 m. Our finding from Iskenderun Bay is the second record of the species from the Mediterranean Sea and indicates that *A. fasciatus* has extended its range northerly to the eastern Levant.

#### **Acknowledgements**

Thanks for financially support to TAGEM (TAGEM-09/AR-GE/11 coordinated by CT) and Captain Nevzat YESIL.

## References

- BILECENOGLU, M., 2010. Alien marine fish of Turkey - an updated review. p.189-217. In: *Fish invasions of the Mediterranean-change and renewal*. D. Golani & B. Appelbaum-Golani (Eds). Sofia, Pensoft Publishers.
- BILECENOGLU, M., KAYA, M. & ERYIGIT, A., 2009. New data on the occurrence of two alien fishes, *Pisodonophis semicinctus* and *Pomadasys stridens*, from the Eastern Mediterranean Sea. *Mediterranean Marine Science*, 10 (2): 151-155.
- ERYILMAZ, L. & DALYAN, C., 2006. First record of *Apogon queketti* Gilchrist (Osteichthyes: Apogonidae) in the Mediterranean Sea. *Journal of Fish Biology*, 69 (4): 1251-1254.
- FRASER, T.H., 2005. A review of the species in the *Apogon fasciatus* group with a description of a new species of cardinalfish from the Indo-West Pacific (Perciformes: Apogonidae). *Zootaxa*, 924: 1-30.
- FROESE, R. & PAULY, D., 2010. FishBase. World Wide Web electronic publication. <http://www.fishbase.org>, version (03/2010).
- GOLANI, D., APPELBAUM-GOLANI, B. & GON, O., 2008. *Apogon smithi* (Kotthaus, 1970) (Teleostei: Apogonidae), a Red Sea cardinalfish colonizing the Mediterranean Sea. *Journal of Fish Biology*, 72 (6): 1534-1538.
- GOLANI, D., SALAMAH, B. & SONIN, O., 2010. First record of the Emperor Angelfish, *Pomacanthus imperator* (Teleostei: Pomacanthidae) and the second record of the Spotbase Burrfish *Cylichthys spilostylus* (Teleostei: Diodontidae) in the Mediterranean. *Aquatic Invasions*, 5 (Suppl. 1): S41-S43.
- GOREN, M., YOKES, M.B., GALIL, B.S. & DIAMANT, A., 2008. Indo-Pacific cardinal fishes in the Mediterranean Sea - new records of *Apogon smithi* from Turkey and *A. queketti* from Israel. *JMBA2 - Biodiversity Records*.
- GOREN, M., GALIL, B.S., DIAMANT, A., GAYER, K. & STERN, N., 2009. First record of the Indo-Pacific cardinal fish *Apogon fasciatus* (White, 1790) in the Mediterranean Sea. *Aquatic Invasions*, 4 (2): 409-411.
- GOREN, M., GAYER, K., & LAZARUS, N., 2009. First record of the Far East chameleon goby *Tridentiger trigonocephalus* (Gill, 1859) in the Mediterranean Sea. *Aquatic Invasions*, 4 (2): 413-415.
- GOREN, M., STERN, N., GALIL, B.S. & DIAMANT, A., 2010. First record of the Indo-Pacific Arrow bulleye *Priacanthus sagittarius* Starnes, 1988 in the Mediterranean Sea. *Aquatic Invasions*, 5 (Suppl. 1): S45-S47.
- LIESKE, E. & MYERS, R., 1994. *Collins Pocket Guide. Coral reef fishes. Indo-Pacific & Caribbean including the Red Sea*. Haper Collins Publishers, 400 pp.
- RANDALL, J.E., 1995. *Coastal fishes of Oman*. Honolulu, University of Hawaii Press, 439 pp.
- ZENETOS, A., MERIÇ, E., VERLAQUE, M., GALLI, P., BOUDOURESQUE, C.F., GIANGRANDE, A., ÇINAR, M. & BILECENOGLU, M., 2008. Additions to the annotated list of marine alien biota in the Mediterranean with special emphasis on Foraminifera and Parasites. *Mediterranean Marine Science*, 9 (1): 119-165.