Mediterranean Marine Science

Vol 11, No 1 (2010)

The response of benthic foraminifer, ostracod and mollusc assemblages to environmental conditions: a case study from the Camalti Saltpan (Izmir-Western Turkey)

E. MERIC, N. AVSAR, A. NAZIK, F. BARUT, F. BERGIN, N. BALKIS, M.S. ONCEL, S. KAPAN-YESILYURT

doi: [10.12681/mms.88](https://doi.org/10.12681/mms.88)

To cite this article:

MERIC, E., AVSAR, N., NAZIK, A., BARUT, F., BERGIN, F., BALKIS, N., ONCEL, M., & KAPAN-YESILYURT, S. (2010). The response of benthic foraminifer, ostracod and mollusc assemblages to environmental conditions: a case study from the Camalti Saltpan (Izmir-Western Turkey). *Mediterranean Marine Science*, 11(1), 5–32.
<https://doi.org/10.12681/mms.88>

The response of benthic foraminifer, ostracod and mollusc assemblages to environmental conditions: a case study from the Camalti Saltpan (Izmir-Western Turkey)

E. MERIC¹, N. AVŞAR², A. NAZİK³, İ. F. BARUT⁴, F. BERGİN⁵,
N. BALKIS⁴, M.S. ÖNCEL⁶ and S. KAPAN-YEŞİLYURT⁷

¹ Moda Hüseyin Bey Sokak, 15/4, 34710, Kadıköy, İstanbul, Turkey

² Çukurova University, Dept. of Geology Engineering, 01330 Balcalı-Adana, Turkey

³ Adiyaman University, Faculty of Vocational and Technical Education, 02040, Adiyaman, Turkey

⁴ İstanbul University, Institute of Marine Sciences and Management, 34116 Vefa-İstanbul, Turkey

⁵ Boğaziçi University, Cultural Heritage Museum, 34134 Bebek İstanbul, Turkey

⁶ Gebze Institute of Technology, Dept. of Environmental Engineering, 41400 Kocaeli, Turkey

⁷ Canakkale Onsekiz Mart University, Dept. of Geological Engineering, 17100 Çanakkale, Turkey

Corresponding author: barutif@istanbul.edu.tr

Received: 19 December 2008; Accepted: 27 November 2009 Published on line: 4 March 2010

Abstract

The subject of this report is benthic foraminifer populations preserved in the saltpan of Camalti in the Province of Izmir. High salinity in certain habitats of *Ammonia tepida* Cushman may be the primary cause of the high rate of twins and triplets as well as other morphological abnormalities recorded within this species (50 % as compared to an anomaly rate of 1 % in normal marine waters). Thicker cyst membrane developing in extremely saline environments may encourage twins and other morphological deformities by denying free movement of the offspring. Ecological factors such as heavy metal contamination of ambient waters as well as contamination by other wastes are also not ruled out as leading to such developmental anomalies. Of the 27 collected samples, Number 5 (that is closest to the sea) includes the typical marine foraminifers. *Nonion depressulum* (Walker and Jacob), *Ammonia tepida* Cushman and *Porosonion subgranosum* (Egger) are the dominant species in other samples. A total of 63 abnormal individuals (8 triplets, 24 twins, and 31 morphological anomalies) was found within seven of the 27 samples collected. Ten samples contained freshwater ostracods: *Darwinula stevensoni* (Brady and Robertson), *Leptocythere lacertosa* Hirschmann, *Cyprideis torosa* (Jones), *Cyprideis* (C.) *anatolica* Bassiouni, and *Loxochoncha elliptica* Brady. Among these samples (some of which contained only a few species of ostracods - and those limited in number of offspring), one had an unusually high ratio of healthy foraminifers those with anomalies. Worthy of note in another sample was a high abundance of molluscs. Among pelecypods, were found *Ostrea edulis* Linné, *Lucinella divaricata* (Linné), *Pseudocama gryphina* Lamarck, *Cerastoderma edule* (Linné), and *Scrobicularia plana* da Costa; and among gastropods were identified *Hydrobia* (*Hydrobia*) *acuta* (Draparnaud), *Rissoa labiosa* (Montagu), *R. parva* (da Costa), *R. violacea* Desmarest, *Pirenella conica* (Blainville), *Bittium desayesi* (Cerulli and Irelli), *B. lacteum* Philippi and *B. reticulatum* Philippi.

Keywords: Benthic foraminiferal deformities; Gastropods; Molluscs; Geochemistry; Camalti Saltpan; Izmir-Turkey.

Introduction

Because of their short life cycles and special habitat requirements, benthic foraminifers are affected by environmental changes and can be used as bio-indicators of pollution in coastal environments (FERRARO *et al.*, 2006). Morphological abnormalities in foraminifer tests are observed in polluted environments and may represent a useful bio-marker for evaluating long-term environmental impacts (VILELA *et al.*, 2004). However, strong natural stress caused by hypersalinity or salinity variations is found to be more effective on abnormal test development, disturbing the test construction (GESLIN *et al.*, 1998 and 2002). Twins, triplets and other morphological aberrations have been reported worldwide from different locations with hypersaline environments, such as the Araruma Lagoon in Rio de Janeiro (STOUFF *et al.*, 1999a), several small inland pools in Israel (ALMOGI-LABIN *et al.*, 1992) and coastal environments of middle Aegean Sea affected by anthropogenic activities (TRANTAPHYLLOU *et al.*, 2005). Morphologically abnormal foraminifer specimens were observed in some of the locations in the Camalti Saltpan (İzmir, Turkey) where the salinity ranges between 40 and 273 ‰ in August (Table 1). The purpose of this study is to verify the twins, triplets and other morphological aberrations in foraminifers from the Camalti Saltpan and the possible causes of their origin.

Despite the observed relation between shell deformities in bivalves and tributyltin (TBT) pollution (MINCHIN *et al.*, 1996; ALZIEU, 2000; COELHO *et al.*, 2006), little if any studies have focused on the effect of TBT on gastropod shell morphology. TBT is one of the pollutants of concern in the North Sea (VAN DEN BROECK,

et al., 2009). This organometal has been used as a biocide in antifouling paints on ships, boats and off-shore installations and as fungicide in agriculture and wood preservation (BAUER *et al.*, 1997; OEHLMANN *et al.*, 1998a). High levels of TBT may affect the ecosystems and cause hazards to

Table 1
Annual averages salinity in sampling stations.

Sample No.	Salinity ‰ S		
	February	April	August
1	--	35,0	52,5
2	--	35,0	52,5
3	--	35,0	88,7
4	--	45,0	88,7
5	--	--	44,8
6	--	35,0	267,7
7	--	--	267,7
8	202,7	192,0	273,0
9	202,7	202,7	257,2
10	202,7	202,7	257,2
11	130,0	130,0	250,0
12	130,0	120,0	144,0
13	120,2	120,0	177,7
14	--	110,0	215,0
15	82,0	70,0	70,0
16	82,0	60,0	70,0
17	--	33,0	40,0
18	--	33,0	40,0
19	--	33,0	40,0
20	--	33,0	44,8
21	--	35,0	44,8
22	--	35,0	44,8
23	--	35,0	44,8
24	--	35,0	52,5
25	--	--	88,5
26	--	--	45,0
27	--	--	45,0

organisms, such as imposex, reproduction, deformities, etc. (ISMAIL, 2006). In bivalves TBT has the potential to induce shell deformities and may reduce reproductive success (BEAUMONT *et al.*, 1992; MINCHIN *et al.*, 1996; ALZIEU, 2000; COELHO *et al.*, 2006), while in female gastropods TBT can cause masculinization (SMITH, 1981; BAUER *et al.*, 1997; OEHLMANN *et al.*, 1998b; RILOV *et al.*, 2000; STRAND and JACOBSEN, 2002; TEWARI *et al.*, 2002). Thus, some molluscs can help in the monitoring and management of heavy metals in the coastal ecosystems.

In Turkey salt is produced from four different sources in seas, lakes, outcroppings and springs. The only saltpan utilizing seawater is located in the Camalti region in the Province of İzmir. The Camalti Saltpan, north of the city on the eastern reaches of the Gulf of İzmir, was established by the Italians in 1906. After the founding of the Republic of Turkey, it continued to operate under the name of 'Tuz Inhisari' (the 'Salt Monopoly') until it was incorporated into the Turkish State Monopolies (Tekel Genel Müdürlüğü) in 1932.

At the Camalti Saltpan, production is based on the very old and slow locomotive system: seawater is pumped into salt pools and the evaporation procedure is very slow. The crystallization pools have a soft substrate and need special modernization techniques. There are two by-products of salt production. The first one concerns the waste water discharge into the sea, which contains salt derivatives such as $MgCl_2$, Br, KCl and also the economically important $Mg(OH)_2$. Modernization of the system will make it economically feasible to enlarge the capacity of the saltpan. Turkey does not have solid potassium sources and the KCl used in the fertilization industry is imported.

The other by-product at Camalti Saltpan is *Artemia salina* living within high salinity pools and channels. It is used at aquarium and cultural fisheries as fish food and is produced and marketed at the Camalti Saltpan. Because this species is vital for salt formation, it is forbidden to collect and dispose of *Artemia salina* eggs (UCAL AND ERGEN, 1994).

The Camalti Saltpan (Fig. 1) covers an area of approximately 10 km², a long narrow coastal strip cordoned off from the Gulf southward of the mouth of the Gediz River (sections İzmir L 17 b2 and Kütahya K 1 on the official maps at 1/25000). There are 36 evaporation pools with a total surface area of 20000 m², 15 collection pools with a total surface of 600 m², 21 cold freshwater reservoirs with a total surface of 3000000 m², and 74 crystallization pools with a total surface of 1195000 m². In March and April the saltpan is rinsed with water from the sea to clear the flats of seaweed and other unwanted components; the rinse-water is subsequently diverted back into the sea. Then in May the seawater that has reached the Baumé temperature of 2.5-5.0 (25.5-52.25 ‰) is let into the collection pools and left to reach Baumé 12 (12 ‰), after which it is conducted into the freshwater reservoirs where $CaSO_4$ is deposited when it reaches Baumé 23 (25 ‰) at the cold water pools Fe_3O is precipitated with CO_3 . Brine (18 Baumé) is transferred to other pools for evaporation until it reaches 23 Baumé. From here the precipitate is taken into the hot-water crystallization pools, where it reaches Baumé 27.5 (28.5 ‰) within 3-4 days. A 25-30 cm layer of salt is precipitated here. At this stage the sodium chloride content of the precipitate is 94-95 % and its color is pinkish due to its plankton content. It is then stored in heaps over the winter. With drainage and winter

Fig. 1: Location map of study area.

precipitation, the sodium chloride content rises to 98%.

Materials and Methods

Investigation of foraminifer, ostracod and mollusc populations was performed on dead individuals. The research was carried out on sediment samples from 27 different points in the saltpan (Fig. 2). The samples were hand collected using a spoon, a total of 20 gr of each wet sample was soaked in 7% hydrogen peroxide (H_2O_2) solution for 24 hours. Then, the samples were dried after flushing under pressurized water in a sieve with a 0.063 mm mesh size. Carbon

tetrachloride was added to the dried samples to bring foraminifer and ostracod populations in the material to the surface. Both the foraminifer and ostracod specimens were collected on filter paper and separated by size using different sieves with 2.00, 1.00, 0.500, 0.250 and 0.125 mm mesh size. The mollusc populations were directly collected from the 0.063 mm mesh size sieve. The samples were analysed under a binocular microscope.

The heavy metals (Fe, Mn, Pb, Cu, Zn, Ni, Co and Cr) analyses on sediment samples were performed in the laboratory of the Institute of Marine Sciences and Management at Istanbul University. The samples

Fig. 2: Sampling stations of Camalti Saltpan.

that were taken during each extraction phase (both total and selective extraction) were analyzed on Shimadzu AA-6701-F Atomic Absorption Spectrophotometry and on the air-acetylene fire (LANDING AND LEWIS 1991; BOUGHRIET *et al.*, 1994). Aluminium was analyzed on nitrus oxide (N₂O) acetylene fire. In five replicate analyses precisions were: Al $\pm 8\%$, Fe $\pm 8\%$, Mn $\pm 4\%$, Pb $\pm 2\%$, Cu $\pm 8\%$, Zn $\pm 4\%$, Ni $\pm 14\%$, Co $\pm 18\%$ and Cr $\pm 4\%$ with 95% confidence interval. Hg analysis was performed on AAS with the cold vapor method on the Hydride Unit. The precision of five replicate analyses on Hg is 2% with a 95% confidence interval.

At the Gebze Institute of Technology laboratories, the sediment samples were dried at room temperature for 24 hours without any prewashing step and mollusc tests were separated. Nitric and hydrochloric

acids were added to 0.35 grams of test samples in a teflon container (BELZUNCE *et al.*, 2001). The material was dissolved using a Milestone Ethos 1600 microwave digestion apparatus. Chemical analyses were performed on Perkin Elmer SIMAA 6000 Simultaneous Atomic Absorption Spectrophotometry with the graphite oven technique.

We encountered diverse and abundant foraminifer fauna at stations 21, 22 and 23 which required different chemical analyses. Thus, geochemical analyses of major and minor elements were done in two different laboratories with different capabilities: Istanbul University, Institute of Marine Sciences and Management (Pb, Cu, Ni, Cd, Hg, Zn, Al, Cr, Fe, Na, Mg and Si), and Gebze Institute of Tecnology, Department of Environmental Engineering (As, Cd, Cu, Fe, Hg, Mn, Pb, Se, Si, Sn and V).

Results

Faunal assemblage composition

Benthic foraminifer assemblage

Eight genera and 13 species of benthic foraminifers were recognized in the 27 sediment samples. Sample 5, which was nearest to the sea, contained marine foraminifer populations of *Textularia bocki* Höglund, *Adelosina cliarensis* (Heron-Allen and Earland), *A. mediterraneensis* (Le Calvez and Le Calvez), *Quinqueloculina disparilis* d'Orbigny, *Q. seminulum* (Linné), *Nonion depressulum* (Walker and Jacob), *Ammonia compacta* Hofker, *A. tepida* (Cushman), *Elphidium complanatum* (d'Orbigny), and *E. crispum* (Linné). In the other 26 samples, however, *Nonion depressulum* (Walker and Jacob), *Ammonia tepida* Cushman, and *Porosonion subgronosum* (Egger) were dominant (CIMERMAN & LANGER, 1991; HOTTINGER *et al.*, 1993; SGARRELLA & MONCHARMONT-ZEI 1993; MERIC *et al.* 2004). In most of the samples, the foraminiferal assemblages were found to be composed of only two or three different genera and species, which are represented by few individuals; whereas in Samples 1 and 21, large populations were represented. In Sample 24, young individuals of *Quinqueloculina seminulum* (Linné) were abundant, but *Ammonia tepida* Cushman and *Nonion depressulum* (Walker and Jacob) were very rare.

Adelosina carinata striata Wiesner was previously found only in one of the samples collected from nearby Dikili Bay, during another study of benthic foraminifers in the Eastern Aegean coasts (MERIC *et al.* 2004). Its presence in samples 12, 19, 21 and 22, suggests that this species displays a high tolerance of salinity. Also, *A. cliarensis* (Heron-Allen and Earland) in Samples 18 and 19;

Spiroloculina ornata (d'Orbigny) in Sample 21, *Ammonia compacta* Hofker in Samples 9, 21 and 25, *Elphidium complanatum* (d'Orbigny) in Samples 2, 12 and 25, and *E. crispum* (Linné) in Sample 25 were found in the present study (Table 2; Plate 1-5).

Twins and triplets among foraminifers

In Samples 1, 2, 4, 6, 19, 21, 22, 23 and 27 (on the north-northeast and south-southeast of the saltpan; Fig. 2) an abundance of aberrant specimens, including twins, triplets and other morphological abnormalities were observed. In nine of the samples, a total of 55 individuals showing aberrant test morphology were determined; seven triplets (Pl. 1, Figs. 1-8), 20 twins (Pl. 1, Figs. 9-16; Pl. 2, Figs. 1-16), and 28 other examples of abnormal morphology (Pl. 3, Figs. 1-15; Pl. 4, Figs. 1-13). Sample 1 revealed a remarkably large population, as it was also observed in Sample 21. The salinities of Stations 1 and 21 in August of 2002 were 52.5‰ and 44.8‰, respectively (Table 1). Four triplets and four twins were counted in Sample 1, whereas two triplets and 10 twins were found in Sample 21. Another 12 specimens with abnormal chamber development were detected in these two samples. In addition, a total of 28 twins or triplets were observed in Sample 2 (eight individuals), Sample 4 (three individuals), Sample 6 (four individuals), Sample 19 (one individual), Sample 22 (three individuals), Sample 23 (one individual) and Sample 27 (three individuals). The percentage of abnormal individuals was found to be 1.50 %, however it was found that only 0.075 % of the population showed morphological abnormalities in marine sediment samples (MERIC, 1996). *Ammonia tepida* was found to be the dominant species with aberrant individuals in all the analysed samples, thus most of the twins, triplets and other morphologically

12

Medit. Mar. Sci., 11/1 2010, 05-32

abnormal individuals were *A. tepida* (Table 3).

The ostracod assemblages

In 10 of the 12 samples analyzed were found the following ostracod species: *Darwinula stevensoni* (Brady and Robertson), *Leptocythere lacertosa* Hirschmann, *Cyprideis*

torosa (Jones), *Cyprideis* (C.) *anatolica* Bassiouni, and *Loxoconcha elliptica* Brady. While some samples contained only one or two species with few specimens, Sample 21 displayed a relatively high individual abundance of ostracods as well as foraminifers (Table 2; Plate 5, figs. 10-13).

Table 3
Distribution of normal and abnormal benthic foraminiferal individuals.

Sample no.	Abnormal individuals	Twins	Triplets	Other abnormalities	Number of individuals	Abnormality %
1	15	4	4	7	457	3.28
2	8	1	1	6	186	4.3
3					153	
4	3			3	247	1.22
5					65	
6	4	2		2	147	2.72
8					15	
9					31	
10					11	
11					52	
12					158	
13					15	
14					148	
15					10	
16					1	
18					7	
19	1	1			86	1.16
20					2	
21	17	10	2	5	1534	1.11
22	3	1		2	127	2.36
23	1	1			109	0.92
25					23	
26					10	
27	3			3	62	4.84
Total	55	20	7	28	3656	1.50

Of the above, *Darwinula stevensoni* is characterized as thriving in sandy beds under fresh or slightly saline waters (oligohaline) with a salt content of 0.5-12 ‰, and generally in muddy sediments rich in organic matter (WAGNER 1957; KEYSER 1976; GUILLAUME *et al.*, 1985; BESONEN 1997; MISCHKE 2001). *Leptocythere lacertosa* is typical of rather brackish (mesohaline) littoral regions in tidal zones (VON MORKHOVEN 1963; BONADUCE *et al.*, 1975; GUILLAUME *et al.*, 1985). *Cyprideis torosa* is a frequent and ubiquitous species found in brackish coastal waters (mesohaline) with a salt content of 8-40 ‰. Preferring a rather high salt content (25-40 ‰), it is thus often found in lagoons, gulfs, and estuaries as well as at river mouths and along tidal plains (WAGNER 1957; BARBEITO-GONZALES 1971; BREMAN 1975; CARBONEL *et al.*, 1975; GUILLAUME *et al.*, 1985; WITTE 1993; BESONEN 1997; MISCHKE 2001). *Cyprideis (C.) anatolica* was described typically from river mouths (BASSIOUNI 1979). *Loxiconcha elliptica* is another frequent and ubiquitous species found in brackish coastal waters (mesohaline) with a salt content of 18-30 ‰; it thrives at relatively shallow depths (20 m or less) in lagoons, gulfs, estuaries, river mouths and tidal plains (BONADUCE *et al.*, 1975; CARBONEL *et al.*, 1975; PASCUAL & CARBONEL 1992; BESONEN 1997).

These genera and species have been recovered from waters with different salt content from 45 to 273 ‰. Most probably they were transported from the sea and trapped in the evaporation pools. The oligohaline species *Darwinula stevensoni* might have been transported from a nearby river branch or cold water reservoir. Samples 21 and 23, in which the greatest number of ostracods were found, come from areas with a salt

content of 45 ‰ in August (when collected), which is one of the lowest salinity values measured, indicating that low salinity might have enhanced the survival rate of ostracods.

The mollusc assemblages

A total of 10 of the 27 collected samples from the saltpan are rich in mollusc fauna. The pelecypods are represented by *Ostrea edulis* Linné, *Lucinella divaricata* (Linné), *Pseudochama gryphina* Lamarck, *Cerastoderma edule* (Linné), and *Scrobicularia plana* da Costa. The gastropods are represented by *Hydrobia (Hydrobia) acuta* (Draparnaud), *Rissoa labiosa* (Montagu), *R. parva* (da Costa), *R. violacea* Desmarest, *Pirenella conica* (Blainville), *Bittium desayesi* Cerulli and Irelli, *B. lacteum* Philippi, and *B. reticulatum* Philippi (Table 2; Plate 6). Although *Ostrea edulis* Linné and *Lucinella divaricata* (Linné) are found in the Black Sea as well, all of the species are Aegean Sea populations.

The habitats for these pelecypoda are inferred from the work of POPPE & GOTO (1991, 1993). *Ostrea edulis* generally lives in shallow waters at depths of up to 90 m, thriving in waters of 25 and 33 ‰ salinity. *Lucinella divaricata* is found from the tidal line to depths of 60 m in waters with 25 ‰ salinity and *Pseudochama gryphina* at depths from 10 to 60 m in salt content of about 30 ‰. *Cerastoderma edule* is found at depths of only a few meters along tidal plains, in sandy bays and also in fresh water. *Scrobicularia plana* thrives along tidal plains, at up to 30 m water depth between 25 and 30 ‰ salinity.

Gastropod habitats are also based on the work of POPPE & GOTO (1991, 1993). *Hydrobia (Hydrobia) acuta* lives in freshwater, saline mineral waters and seawater at depths varying from 3 to 40 meters. It can

tolerate up to 33 ‰ salinity. *Rissoa labiosa* lives in very shallow waters at depths up to 25 m, favoring a salt content between 20 and 30 ‰. *Rissoa parva* is found off the coastline at depths of 30 m in waters with a salt content of around 30 ‰. *Rissoa violacea* thrives at depths of 0-25 m in salty waters (25-33 ‰), while *Pirenella conica* inhabits shallow waters with low salinity. *Bittium desayesi*, *B. lacteum* and *B. reticulatum* are all found in seawater with a salt content as 25-30‰ from tidal plains down to 250 m.

The geochemistry of sediment and mollusc tests

Among the analyzed heavy metals, Hg is found to have the lowest value in ppb (Table 4).

Figure 3 illustrates the distribution of heavy metals. Fe is measured at highest ppm value compared to other heavy metals (Si, Al, Zn, Hg, Pb, Cd, Cr, Cu, Ni, Mg and Na).

Si, Al, Mg and Na values are considered as important. Zn values are close to each other in Samples 5, 9, 25. Cr has almost the same values in Samples 12 and 24, while Pb, Cu, Ni and Cd values are close to each other in all samples. Cd is < 0,01 in all samples. The Al, Mg, Na values show parallel curves in Samples 3, 4, 5, 6, 7, 8, 9, 10, 11, 14, 15, 16, 21, 22 and 23. When the locations of the stations are taken into consideration, we observe no significant difference between the saltpan and the pools. The Si value is highest in Samples 5, 13, 20, 22 and it is found to be lowest in Sample 19. Although the locations of Stations 19 and 20 are juxtaposed, their Si values are very different. This discrepancy can be explained by the fact that Station 19 is located on the point of discharge of a river branch into the saltpan.

Geochemical analysis of both sediment samples and mollusc tests from Samples 21 and 23 are given in Figure 4. The tests collected from Samples 21 and 23, where the

Fig. 3: The distribution of heavy metal and major cation concentrations in the sediment samples from Camalti Saltpan (%Fe, the others ppm) (by Istanbul University, Institute of Marine Sciences and Management).

Table 4

The values of heavy metal and major cation concentrations in *Cerastoderma edule* (Linné) and *Rissoa violacea* shells (Fe%, the others ppm) and in the sediment samples in Camalti Saltpan (Fe %, the others ppm)
(by Istanbul University, Institute of Marine Sciences and Management)

Station No.	Pb (ppm)	Cu (ppm)	Ni(ppm)	Cd (ppm)	Hg (ppb)	Zn(ppm)	Al(ppm)	Cr (ppm)	Fe (ppm)	Fe (%)	Na(ppm)	Mg(ppm)	Si (ppm)
1	18	17	23	<0.01	42	22	661.82	36.95	26838.5	2.68385	1053	813	5141.43
2	15	10	10	<0.01	26	4	501.95	18.605	21687	2.1687	1393.5	979.5	4911.57
3	13	11	14	<0.01	25	10	856.715	28.69	22313	2.2313	492	564	4839
4	16	9	10	<0.01	68	8	493.32	22.68	19287.8	1.92878	162	364.5	4415.58
5	45*	24	7	<0.01	66	79	211.135	30.11	26820.5	2.68205	156.93	102	5431.77
6	29*	31	5	<0.01	243	5	100.49	12.975	13803.8	1.38038	78	217.5	4209.93
7	26*	7	16	<0.01	141	4	433.43	26.055	21306	2.1306	1108.5	729	3871.2
8	17	1	7	<0.01	67	<0.01	298.43	16.435	9656.25	0.96563	477	387	3689.73
9	32*	22	26	<0.01	34	51	453.225	41.455	28069.5	2.80695	1192.5	786	2504.16
10	23*	6	12	<0.01	27	0.02	307.055	24.41	17506.3	1.75063	319.5	264	1693.65
11	22*	13	18	<0.01	40	17	490.275	29.315	23420.5	2.34205	1822.5	691.5	3145.35
12	23*	16	31	<0.01	77	27	829.815	75.235	36178.3	3.61783	418.5	352.5	2504.16
13	23*	9	23	<0.01	51	10	844.535	45.71	28195	2.8195	699	442.5	5443.86
14	12	<0.01	4	<0.01	26	<0.01	81.205	10.88	5348.25	0.53483	483	286.5	2552.55
15	22*	12	17	<0.01	49	8	567.93	47.165	26520	2.652	1474.5	817.5	2189.64
16	34*	30	37	<0.01	24	92*	567.93	60.185	32208.3	3.22083	541.5	381	2479.98

(continued)

Table 4 (continued)

Station No.	Pb (ppm)	Cu (ppm)	Ni(ppm)	Cd (ppm)	Hg (ppb)	Zn(ppm)	Al(ppm)	Cr (ppm)	Fe (ppm)	Fe (%)	Na (ppm)	Mg(ppm)	Si (ppm)
17	30*	14	21	<0.01	29	38	766.375	54.94	36043.8	3.60438	481.5	306	2310.63
18	31*	20	33	<0.01	10	41	556.255	46.445	33266.8	3.32668	832.5	531	1149.27
19	20	8	16	<0.01	29	10	838.44	45.255	26618.5	2.66185	774	565.5	508.08
20	23*	6	16	<0.01	52	7	592.795	28.48	19090.5	1.90905	493.5	615	5056.74
21	33*	14	10	<0.01	38	18	267.47	25.535	32068.5	3.20685	982.5	819	2637.24
22	29*	22	33	<0.01	1	36	787.69	51.345	38477.8	3.84778	1192.5	1737	5286.6
23	19	17	10	<0.01	14	24	558.79	28.205	34949.5	3.49495	21.45	403.5	3786.51
24	28*	23	36	<0.01	15	39	1300.295	63.9	41679.3	4.16793	516	754.5	4137.33
25	225*	7	9	<0.01	17	77	159.365	15.83	13901.8	1.39018	1024.5	625.5	4851.09
21 <i>C.edule</i>	160*	8	8	1.66*	30	<0.01	3.0249	15.64	1319.7	0.13197	276	538.5	5093.04
21 <i>C.edule</i>	73*	5	8	0.56*	71	<0.01	2.4565	13.72	1378.25	0.13783	73.5	229.5	5008.35
23 <i>R.violacea</i>	41*	35	14	2*	65	<0.01	2.1012	12.45	516.22	0.05162	328.5	412.5	6605.22
23 <i>R.violacea</i>	147*	14	21	1.2*	119	<0.01	4.0501	6.65	304.84	0.03048	86.19	580.5	5601.12

Fig. 4: a. The distribution of heavy metal and major cation concentrations (ppm); **b.** The distribution of heavy metal and major cation concentrations (% Fe, the others ppm) in in *Cerastoderma edule* (Linné) and *Rissoa violacea* Desmarest for samples 21 and 23 in Camalti Saltpan (by Istanbul University, Institute of Marine Sciences and Management).

fauna was found to be diverse, were analyzed and the values were compared with the sediment analyses. Si, Mg, Na and Al

(in ppm) values are significant. Cr and Zn values show a slight tendency to increase. In Figures 4a and b, Fe values are given in %, (in ppm) values are significant.

and the relation of values of other metals in sediment and mollusc tests to each other can be shown as .

The heavy metal analyses of the sediment samples and tests from Samples 21 and 23 were performed in Gebze at the Institute of High Technology and they are comparatively shown in Table 5 and Figure 5 . Fe, Hg, Mn, Pb, Si, V and As values tend to increase the sediments and test values

are found to be parallel. Sediment samples from Samples 21 and 22 $Si > Pb > Fe > Mn$, from Samples 23 $Pb > Si > Fe > Mn$.

The As, Cd, and V values are close to each other and Se and Sn values are very small.

In sediment from Sample 21; $Al < Mg < Na < Si$ and on the test $Na < Mg < Si$, from Sample 23; $Na < Al < Mg < Si$ and on the test $Na < Mg < Si$ (Table 6).

Fig. 5: The distribution of of heavy metals and major cation concentrations (ppm) in shells of *Cerastoderma edule* (Linné) and *Rissoa violacea* Desmarest for samples 21 and 23 in Camalti Saltpan (by Gebze Institute of Technology, Dept.of Environmental Eng.)

Table 5

The values of heavy metal and trace elements concentrations (ppm) in *Cerastoderma edule* (Linné) and *Rissoa violacea* Desmarest shells in samples 21, 22 and 23 in Camalti Saltpan (by Gebze Institute of Technology, Dept. of Environmental Eng.)

Sample location	As	Cd	Cu	Fe	Hg	Mn	Pb	Se	Si	Sn	V
Çamaltı 21	0,29	0,32	0,14	10,34	3,42	5,09	51,40	0,06	58,98	0,01	0,66
Çamaltı 22	0,27	0,09	0,26	10,15	2,30	9,57	32,80	0,03	107,59	0,01	1,85
Çamaltı 23	0,52	0,10	0,21	10,48	1,70	2,60	72,69	0,04	19,55	0,01	0,32
21 (<i>C.edule</i>)	1,12	0,07	0,06	<0.01	0,68	0,24	81,85	0,05	34,80	0,02	0,15
23 (<i>R.violacea</i>)	0,08	0,10	0,25	3,53	1,10	3,16	9,24	0,04	29,45	<0.01	0,47

Table 6
Correlation of Fe, Si, Al, Mg, Na (ppm) between their values in the shells of *Cerastoderma edule* (Linné) and *Rissoa violacea* Desmarest and their values in sediment samples.

	Sediment	Shell
Fe	High	Low
Mn	Low High	<i>Rissoa violacea</i> high in the shell <i>Cerastoderma edule</i> low in the shell
Pb	High Low	<i>Rissoa violacea</i> low in the shell <i>Cerastoderma edule</i> high in the shell
Si	High Low	<i>Cerastoderma edule</i> low in the shell <i>Rissoa violacea</i> high in the shell
Hg	High	Low

The chemical and physical parameters have been routinely measured by the saltpan management. Pump water data from the new area of the second saltpan which is near to Sample 27 and pool water data from crystalization pool are given in Table 7 and Table 8, respectively. Ca and Mg values vary between the pool water and the water from the pumping system:

Pool water Ca^{2+} : 0.4008- 0.4810 g/l
 Mg^{+2} : 30.6432-34.4979 g/l
 Pump water Ca^{2+} : 0.5010 g/l
 Mg^{+2} : 14.0448 g/l

Discussion and Conclusions

Foraminifer twins and triplets are rarely observed in natural environments (about 1% of the population) However, we found that in laboratory conditions, exposing *Ammonia tepida* to a hypersaline (50‰) environment greatly facilitates abnormal development, mainly of twins (STOUFF *et al.*, 1999b). (In this report the researchers also estimated that the maximum salinity tolerated by *Elphidium crispum* individuals was 42‰). Furthermore, twins have been ob-

served in a dominant *Ammonia* population in salt pools near the Dead Sea in Israel (ALMOGI-LABIN *et al.*, 1992). COLE (1960) determined that a delay in the disintegration of the cyst enclosing the microspherical individuals produced by schizogonic reproduction can lead to a fusing of the individual offspring. Thicker cyst walls grown in hypersaline environments disintegrate more slowly and bring about twins and triplets as well as many other abnormal developments. This delay may keep the young offspring locked within close contact for a considerably longer time. If two or more offspring develop at the same rate, their tests will have the same size and shape. Likewise, the first chambers of the tests may develop on the same or different planes. One can sometimes exhibit more development than the other (STOUFF *et al.*, 1999a). Thus the offspring trapped longer within the same limited space may become fused to each other, hence, will be destined to continue living in this state after the disintegration of the cyst wall.

An interesting characteristic among twins and triplets is that there is striking dif-

Table 7
The result of salt waters analysis from the new area of second saltpan
(near of the sample 27) (by Management Camalti Saltpan).

Sample Name	The date of sample collection	(Ca ⁺ ²⁺) (g/l)	(Mg ⁺ ²⁺) (g/l)	BOME (Be°) Values	(°C)	Baumé (Be°) Calculation with the factor of correction
New area A-1 1(West) Poolwater	05.08.2003	0,4008	30,6432	27,9	27,0	28,30
New area A-1 1(East) Poolwater	05.08.2003	0,4609	34,4979	28,4	28,0	28,85
New area A-2 2(West) Poolwater	05.08.2003	0,4810	31,8106	28,0	28,5	28,50
Saltwater of pump which feed the new area	07.08.2003	0,5010	14,0448	25,8	25,0	26,10
Saltwater from reserv feeding New area	07.08.2003	0,4609	14,3123	25,9	25,5	26,2
Feeding water from New area (2-A)	08.08.2003	0,5411	14,0205	24,5	34,0	25,3
2 Poolwater new area feeding water	11.08.2003	0,6012	14,3488	24,8	28,0	25,25
Feeding water from reserv number 36	09.06.2003	1,0020	11,5520	24,2	26,0	24,5
Feeding water to the new area	13.06.2003	0,6012	12,7680	24,25	32,5	25,0
Feeding water for the pools 1A-2A	06.06.2003	0,8016	11,4304	24,5	27,0	24,9
First salt water for A-1; A-2 New area	30.05.2003	0,9018	9,5456	22,1	23,0	22,5

ference in the formation of the twins or triplets. Sometimes they are attached from the very first chambers (Pl. 1, Fig. 9), but mostly they show great differences in the community, which then contradicts the mechanism of development proposed by STOUFF *et al.* (1999a). Therefore, the twin or triplet formation is caused by entrapment within cysts which have not disintegrated in time

in hypersaline environments; it appears to be a matter of chance, as the specimens illustrated on Plates 1 and 2 would seem to confirm.

Still another subject is the morphological changes that occur in the tests of foraminifers living in hypersaline environments. Test formation depends on the chemical composition of the water, its turbidity,

Table 8
The analysis of the salt precipitated in the crystallisation pools in the new area of Camalti Saltpan Management.

Sample Name	The date of sample collection	Quantity of insoluble matter (%gr)	Ca ²⁺ (%gr)	Mg ²⁺ (%gr)	SO ₄ ²⁻ (%gr)	CaSO ₄ (%gr)	MgSO ₄ (%gr)	MgCl ₂ (%gr)	NaCl (Dry) (%gr)
Salt of channel in New area	29.07.2003	0,0100	0,1603	0,4134	0,8374	0,5446	0,5679	1,1698	97,5078
10.Vinc A-1 Salt of pool	06.08.2003	0,0170	0,0601	0,3101	0,6378	0,2042	0,6187	0,7248	98,2353

pH, salt content and temperature, as well as the presence of heavy metals and environmental pollution (ALVE, 1991; SHARIFI *et al.*, 1991; ALMOGI-LABIN *et al.*, 1992; YANKO *et al.*, 1998). This may explain the morphological abnormality that occurs in foraminifer tests in hypersaline environments. Increasing concentrations of heavy metals in the evaporation pools might have been the cause of morphological abnormalities observed in the foraminifer specimens.

Adelosina carinata striata was previously found only in one of the samples collected from nearby Dikili Bay during another study of benthic foraminifers in the Eastern Aegean coasts. (MERIC *et al.* 2004). Its presence in four of our samples (No. 12, 19, 21, 22) suggests that this species displays a high tolerance of enhanced salinities. Also, *Adelosina cliarensis* in Samples 18 and 19; *Spiroloculina ornata* in Sample 21, *Ammonia compacta* in Samples 9, 21 and 25, *Elphidium complanatum* in Samples 2, 12 and 25, and *E. crispum* in Sample 25 were found in the present study.

Although the salinity generally tolerated by *Ammonia tepida* was accepted as 52.5‰, we show that in the Camalti Saltpan it can

survive to a limited number in hypersaline environments with 89‰ salinity. The study confirms that the twins and triplets, rarely seen in environments with normal salinity (the western Black Sea, the Gulf of İzmit, and the Aegean Sea) (MERIC 1996; MERIC *et al.*, 2001, 2003; TRIANTAPHYLLOU *et al.*, 2005), are considerably more abundant in hypersaline environments.

The ostracods were represented in the saltpan by 4 genera and 5 species. These genera and species have been recovered from waters with different salt content from 45 to 273 ‰. Most probably they were transported from the sea and trapped in the evaporation pools. The oligohaline species *Darwinula stevensoni* might have been transported from a nearby river branch or cold water reservoir. Samples 21 and 23, in which the greatest number of ostracods were found, come from areas with a salt content of 45 ‰ in August (when collected), which is one of the least salinity values measured, indicating that low salinity might have enhanced the survival rate of ostracods.

The mollusc fauna was also found not to be very rich; pelecypods were represented by 6 genera and 6 species and gastropods

were represented by 4 genera and 8 species. The habitats for these pelecypoda are inferred from the work of POPPE & GOTO (1991, 1993). *Ostrea edulis* generally lives in shallow waters at depths of up to 90 m, thriving in waters of 25 and 33 ‰ salinity. *Lucinella divaricata* is found from the tidal line to depths of 60 m in waters with 25 ‰ salinity and *Pseudochama gryphina* at depths from 10 to 60 m in salt content of about 30 ‰. *Cerastoderma edule* is found at depths of only a few meters along tidal plains, in sandy bays and also in fresh water. *Scrobicularia plana* thrives along tidal plains, but at up to 30 m water depth between 25 and 30 ‰ salinity.

Gastropod habitats are also based on the work of POPPE AND GOTO (1991, 1993). *Hydrobia* (*Hydrobia*) *acuta*, lives in freshwater, saline mineral waters and seawater at depths varying from 3 to 40 meters. It can tolerate up to 33 ‰ salinity. *Rissoa labiosa* lives in very shallow waters at depths up to 25 m, favoring a salt content between 20 and 30 ‰. *Rissoa parva* is found off the coastline at depths of 30 m in waters with a salt content of around 30 ‰. *Rissoa violacea* thrives at depths of 0-25 m in salty waters (25-33 ‰), while *Pirenella conica* inhabits shallow waters with low salinity. *Bitium desayesi*, *B. lacteum* and *B. reticulatum* are all found in seawater with a salt content of 25-30‰ from tidal plains down to 250 m.

The heavy metal analyses on mollusc tests showed that the Fe concentration is very much less compared to sediment values, indicating that accumulation in tests is not the same for every metal. So mollusc tests cannot be used for monitoring every kind of heavy metal pollution.

Acknowledgements

The authors would like to express their sincere thanks to Dr. Nuray Önoğlu (Izmir

Regional Directorate of the MTA) for the countless ways in which she facilitated our efforts and to Dr. Nerdin Kubanç (Istanbul University Faculty of Natural Sciences) for her participation, as well as to Mr. Murat Dundar (General Director of ASSAN Ar-Ge) and Technician Hüsnü Öztürk for the photographs taken with the Jeol JSM electron microscope. Baki Yokes (Haliç University) is gratefully acknowledged for revising the article.

References

- ALMOGI-LABIN, A., PERELIS-GROSSOVICZ, L. & RAAB, M., 1992. Living Ammonia from a hypersaline inland pool, Dead Sea area, Israel. *Journal of Foraminiferal Research*, 22 (3): 257-266.
- ALVE, E., 1991. Benthic foraminifera in sediment cores reflecting heavy metal pollution in Solfjord. Western Norway. *Journal of Foraminiferal Research*, 21 (1): 1-19.
- ALZIEU, C., 2000. Impact of tributyltin on marine invertebrates. *Ecotoxicology*, 9 (1-2): 71-76.
- BARBEITO-GONZALES, J.P., 1971. Die Ostracoden des Küstenbereiches von Naxos (Griechenland) und ihre Lebensbereiche. *Mitteilungen aus dem Hamburgischen Zoologischen Museum und Institut*, 67: 255-326.
- BASSIOUNI, M.A., 1979. Brackische und marine Ostracoden (Cytherideinae, Hemicytherinae, Trachyleberidinae) aus dem Oligozan und Neogen der Türkei. *Geologisches Jahrbuch Reihe B*, 31: 1-200.
- BAUER, B., FIORONI, P., SCHULTE-OEHLMANN, U., OEHLMANN, J. & KALBFUS, W., 1997. The use of *Littorina littorea* for tributyltin (TBT) effect monitoring - results from the Ger-

- man TBT survey 1994/1995 and laboratory experiments. *Environmental Pollution*, 96 (3): 299-309.
- BEAUMONT, A.R., NEWMAN, P.B. & SMITH, J. 1992. Some effects of tributyltin from anti-fouling paints on early development and veliger larvae of the mussel *Mytilus edulis*. In: *Proc. Ninth Int. Malacol. Congress*, Eds. E. Gittenberger and J. Goud, 55-63, Unitas Malacologia, Leiden, The Netherlands.
- BESONEN, M.R., 1997. *The Middle and Late Holocene Geology and Landscape Evolution of the Lower Acheron River Valley, Epirus, Greece*. MSc Thesis, University of Minnesota, Faculty of the Graduate School, 161 pp.
- BELZUNCE, M.J., SOLAUN O., FRANCO, J., VALENCIA, V. & BORJA, A., 2001. Accumulation of organic matter, heavy metals and organic compounds in surface sediments along the Nervión Estuary (Northern Spain). *Marine Pollution Bulletin*, 42 (12): 1407-1411.
- BONADUCE, G., CIAMPO, G. & MASOLI, M., 1975. Distribution of Ostracoda in the Adriatic Sea. *Pubblicazioni Della Stazione Zoologica di Napoli*, 40: 1-154.
- BOUGHRIET, A., DERAM, L. & WARTEL, M., 1994. Determination of dissolved chromium (III) and chromium (VI) in sea-water by atomic absorption spectrometry. *Journal of Analytical Atomic Spectrometry*, 9: 1135-1142.
- BREMAN, E., 1975. *Distribution of ostracodes in the bottom sediments of the Adriatic Sea*. PhD thesis. Vrije Universiteit te Amsterdam, 165 pp.
- CARBONEL, P., MOYES, J. & PEYPOUQUET, J.P., 1975. Utilisation des Ostracodes pour la mise en évidence et l'évolution d'une lagune holocene à l'Ouest de la Gironde, Golfe de Biscay. *Bulletins of American Paleontology*, 65 (282): 445-462.
- CIMERMAN, F. & LANGER, M.R., 1991. *Mediterranean foraminifera*. Academia Scientiarum et Artium Slovenica. Ljubljana. Classis IV 30: 118 p.
- COELHO, M.R., LANGSTON, W.J. & BEBIANNO, M.J., 2006. Effect of TBT on *Ruditapes decussatus* juveniles. *Chemosphere*, 63 (9): 1499-1505.
- COLE, W.S., 1960. Variability in embryonic chambers of *Lepidocyclus*. *Micropaleontology*, 6 (2): 133-144.
- FERRARO, L., SPROVIERI, M., ALBERICO, I., LIRER, F., PREVEDELLO, L. & MARSELLA, E., 2006. Benthic foraminifera and heavy metals distribution: A case study from the Naples harbour (Tyrrhenian Sea, Southern Italy). *Environmental Pollution*, 142 (2): 274-287.
- GESLIN, E., DEBENAY, J.P. & LESOURD, M., 1998. Abnormal wall textures and test deformation in *Ammonia* (hyaline foraminifer). *Journal of Foraminiferal Research*, 28 (2): 148-156.
- GESLIN, E., DEBENAY, J.-P., DULEBA, W. & BONETTI, C., 2002. Morphological abnormalities of foraminiferal tests in Brazilian environments: comparison between polluted and non-polluted areas. *Marine Micropaleontology*, 45 (2): 151-168.
- GUILLAUME, M.C., PEYPOUQUET, J.P. & ET TETART, J., 1985. Quaternaire et actuel. In: *Atlas des Ostracodes de France*. Edited by H.J. Oertli. *Bulletin des Centres de Recherches Exploration-Production Elf-Aquitaine - Memoire*, 9: 337-377.
- HOTTINGER, L., HALICZ, E. & REISS, Z. 1993. *Recent foraminifera from the Gulf of Aqaba, Red Sea*. Slovenska Akademija Znanosti in Umetnosti, Academia Scientiarum et Artium Slovenica: Ljubljana. 179 pp.

- ISMAIL, A., 2006. The use of intertidal molluscs in the monitoring of heavy metals and organotin compounds in the west coast of Peninsular Malaysia. *Coastal Marine Science* 30 (1): 401-406.
- KEYSER, D., 1976. *Zur Kenntnis der brackigen mangrouebewachsenen weichböden Südwest-Floridas unter besonderer Berücksichtigung ihrer ostracoden fauna*. Dissertation zur Erlangung des Fachbereichs Biologie der Universität Hamburg, 142 pp.
- LANDING, W.M. & LEWIS B.L. 1991. Thermodynamic modeling of trace metal speciation in the Black Sea. p. 125-160. In: *NATO-ASI Symposium Series "Black Sea Oceanography"*, E. Izdar and J. W. Murray (Eds.).
- MERİÇ, E., 1996. Twin forms in foraminifera from the Quaternary sediments of Gulf of İzmit (Turkey). *Revue Paléobiologie*, 15 (2): 461-467.
- MERİÇ, E., AVŞAR, N. & GÖRMÜŞ, M., 2001. Twin forms in recent benthic foraminifera from the northern Aegean Sea and western Black Sea regions (Turkey). *Revue Paléobiologie*, 20 (1): 69-75.
- MERİÇ, E., AVŞAR, N. & GÖRMÜŞ, M. & BERGIN F., 2003. Twin and triplet forms of Recent benthic foraminifera from the eastern Aegean Sea, Turkish Coast. *Micropaleontology*, 50 (3): 297-300.
- MERİÇ, E., AVŞAR, N. & BERGIN, F. 2004. *Benthic foraminifera of Eastern Aegean Sea (Turkey): Systematics and Autoecology*. Turkish Marine Research Foundation. Publication No.18: 306 pp.
- MINCHIN, D., STROBEN, E., OEHLMANN, J., BAUER, B., DUGGAN, C.B. & KEATINGE, M., 1996. Biological indicators used to map organotin contamination in Cork Harbour, Ireland. *Marine Pollution Bulletin*, 32 (2): 188-195.
- MISCHKE, S., 2001. Mid and Late Holocene palaeoenvironment of the lakes Eastern Juyan and Sogur Nur in NW China, based on ostracod species assemblages and shell chemistry. *Berliner Geowissenschaftliche Abhandlungen, Reihe E*, 35: 131 pp.
- OEHLMANN, J., BAUER, B., MINCHIN, D., SCHULTE-OEHLMANN, U., FIORONI, P. & MARKERT, B., 1998a. Imposed in *Nucella lapillus* and intersex in *Littorina littorea*: interspecific comparison of two TBT-induced effects and their geographical uniformity. *Hydrobiologia*, 378 (1-3): 199-213.
- OEHLMANN, J., STROBEN, E., SCHULTE-OEHLMANN, U. & BAUER, B., 1998b. Imposed development in response to TBT pollution in *Hinia incrassata* (Ström, 1768) (Prosobranchia, Stenoglossa). *Aquatic Toxicology*, 43 (4): 239-260.
- PASCUAL, A. & CARBONEL, P., 1992. Distribution and annual variations of *Loxiconcha elliptica* in the Gernika Estuary (Bay of Biscay). *Geobios*, 25 (4): 495-503.
- POPPE, G.T. & GOTO, Y., 1991. *European Seashells Volume I (Polyplacophora, Caudofoveata, Solenogaster, Gastropoda)*. Verlag Christa Hemmen. 352 pp.
- POPPE, G.T. & GOTO, Y., 1993. *European Seashells Volume II (Scaphopoda, Bivalvia, Cephalopoda)*. Verlag Christa Hemmen. 221 pp.
- RILOV, C., GASITH, A., EVANS, S.M. & BENAYAHU, Y., 2000. Unregulated use of TBT-based antifouling paints in Israel (eastern Mediterranean): high contamination and imposed levels in two species of marine gastropods. *Marine Ecology Progress Series*, 192: 229-238.
- SHARIFI, A.R.L., CROUDACE, W. & AUSTIN, R. L., 1991. Benthic

- foraminiferids as pollution indicators in Southampton Water, southern England. United Kingdom. *Journal of Micropaleontology*, 10 (1): 109-113.
- SGARRELLA, F. & MONCHARMONT-ZEI, M., 1993. Benthic foraminifera of the Gulf of Naples (Italy), systematic and autoecology. *Bollettino della Società Paleontologica Italiana*, 32 (2): 145-264.
- SMITH, B.S., 1981. Tributyltin compounds induce male characteristics on female mud snails *Nassarius obsoletus* = *Ilyanassa obsoleta*. *Journal of Applied Toxicology*, 1 (3): 141-144.
- STRAND, J. & JACOBSEN, J.A., 2002. Imposex in two sublittoral neogastropods from the Kattegat and Skagerrak: the common whelk *Buccinum undatum* and the red whelk *Neptunea antiqua*. *Marine Ecology Progress Series*, 244: 171-177.
- STOUFF, V., DEBENAY, J.-P. & LESOURD, M., 1999a. Origin of double and multiple tests in benthic foraminifera: observations in laboratory cultures, environments. *Marine Micropaleontology*, 36 (4): 189-204.
- STOUFF, V., GESLIN, E., DEBENAY, J.-P. & LESOURD, M., 1999b. Origin of morphological abnormalities in *Ammonia* (Foraminifera): Studies in laboratory and natural environments. *Journal of Foraminiferal Research*, 29 (2): 152-170.
- TEWARI, A., RAGHUNATHAN, C., JOSHI, H.V. & KHAMBHATY, Y., 2002. Imposex in rock whelks *Thais* and *Ocenebra* species (Mollusca, Neogastropoda, Muricidae) from Gujarat coast. *Indian Journal of Marine Science*, 31 (4): 321-328.
- TRIANAPHYLLOU, M.V., TSOUROU, T., KOUKOUSIOURA, O. & DERMITZAKIS, M.D., 2005. Foraminiferal and ostracod ecological patterns in coastal environments of SE Andros Island (Middle Aegean Sea, Greece). *Revue de Micropaléontologie*, 48 (4): 279-302.
- UCAL, K. & ERGEN, Z., 1994. Comparative study of *Artemia* egg and larval growth from Camalti (Izmir) and Ayvalık (Balıkesir) salterns. *Turkish Journal of Zoology*, (TUBITAK), 18: 19-24.
- WAGNER, C.W., 1957. *Sur les ostracodes du Quaternaire récent des Pays-bas et leur utilisation dans l'étude géologique des dépôts Holocènes*. The Hauge, Mouton and Company, 259 p.
- WITTE, L., 1993. *Taxonomy and biogeography of West African beach ostracods*, Verhandelingen der Koninklijke Nederlandse Akademie van Wetenschappen, Afdeling Natuurkunde, Eerste Reeks, 39, 84 p., Amsterdam, North-Holland.
- VAN DEN BROECK, H., DE WOLF, H., BACKELJAU, T. & BLUST, R., 2009. Comparative assessment of reproductive impairment in the gastropod mollusc *Littorina littorea* along the Belgian North Sea coast. *Science of the Total Environment*, 407 (8): 3063-3069.
- VILELA, C.G., BATISTA, D.S., BATISTANETO, J.A., CRAPEZ, M. & MCALLISTER, J.J., 2004. Benthic foraminifera distribution in high polluted sediments from Niterói Harbor (Guanabara Bay), Rio de Janeiro, Brazil. *Anais da Academia Brasileira de Ciências*, 76 (1): 161-171.
- VAN MORKHOVEN, F.P.C.M., 1963. *Post-Palaeozoic Ostracoda*. Volume II, generic descriptions. Elsevier, Amsterdam, 478 p.
- YANKO, V., AHMAD, M. & KAMINSKI, M., 1998. Morphological deformities of benthic foraminiferal tests in response to pollution by heavy metals; implications for pollution monitoring. *Journal of Foraminiferal Research*, 28 (3): 177-200.

PLATE 1

1-16. *Ammonia tepida* Cushman. Camalti Saltpan, İzmir, Turkey.

1. Triplet forms , x 66, Sample 1.	9. Twin forms , x 112, Sample 1.
2. Triplet forms , x 73, Sample 1.	10. Twin forms , x 73, Sample 1.
3. Triplet forms , x 98, Sample 1.	11. Twin forms , x 98, Sample 1.
4. Triplet forms , x 98, Sample 1.	12. Twin forms , x 98, Sample 1.
5. Triplet forms , x 56, Sample 2.	13. Twin forms , x 90, Sample 2.
6. Triplet forms , x 84, Sample 21.	14. Twin forms , x 84, Sample 6.
7. Triplet forms , x 112, Sample 21.	15. Twin forms , x 73, Sample 6.
8. Triplet forms , x 73, Sample 27.	16. Twin forms , x 140, Sample 21.

PLATE 2

1-12. *Ammonia tepida* Cushman. Camalti Saltpan, İzmir, Turkey.

1. Twin forms , x 140, Sample 21.	5. Twin forms , x 98, Sample 21.	9. Twin forms , x 90, Sample 21.
2. Twin forms , x 140, Sample 21.	6. Twin forms , x 112, Sample 21.	10. Twin forms , x 84, Sample 22.
3. Twin forms , x 165, Sample 21.	7. Twin forms , x 112, Sample 21.	11. Twin forms , x 98, Sample 23.
4. Twin forms , x 90, Sample 21.	8. Twin forms , x 140, Sample 21.	12. Twin forms , x 73, Sample 27.

13-14. *Quinqueloculina seminula* (Linné), Camalti Saltpan, İzmir, Turkey.

13. Twin forms , x 73, Sample 19.	14. Twin forms , x 90, Sample 23.

15. *Nonion depressulum* (Walker and Jacob), Camalti Saltpan, İzmir, Turkey.

Twin forms , x 98, Sample 1.

16. *Porosonion subgronosum* (Egger), Camalti Saltpan, İzmir, Turkey.

Twin forms , x 98, Sample 21.

PLATE 3

1-15. *Ammonia tepida* Cushman. Specimens have abnormal chamber development, Camalti Saltpan, İzmir, Turkey.

1. Spiral side, x 98, Sample 1.	9. Spiral side, x 140, Sample 2.
2. Spiral side, x 73, Sample 1.	10. Spiral side, x 90, Sample 2.
3. Spiral side, x 98, Sample 1.	11. Spiral side, x 112, Sample 4.
4. Spiral side, x 112, Sample 1.	12. Spiral side, x 125, Sample 4.
5. Umblical side, x 98, Sample 1.	13. Spiral side, x 125, Sample 4.
6. Spiral side, x 84, Sample 1.	14. Spiral side, x 84, Sample 6.
7. Umblical side, x 112, Sample 2.	15. Spiral side, x 90, Sample 21.
8. Spiral side, x 98, Sample 2.	

PLATE 4

1-8. *Ammonia tepida* Cushman. Specimens have abnormal chamber development, Camalti Saltpan, İzmir, Turkey.

1. Spiral side, x 150, Sample 21.	4. Spiral side, x 98, Sample 21.	7. Spiral side, x 98, Sample 27.
2. Umbilical side, x 195, Sample 21.	5. Umbilical side, x 125, Sample 22.	8. Umbilical side, x 73, Sample 27.
3. Umbilical side, x 125, Sample 21.	6. Spiral side, x 70, Sample 22.	

9. *Quinqueloculina cf. seminula* (Linné).

External view, x 73, Sample 6.

10-11. *Nonion depressulum* (Walker and Jacob).

10. External view, x 98, Sample 2. 11. External view, x 112, Sample 2.

12-13. *Porosonion subgronosum* (Egger).

12. External view, x 112, Sample 1. 13. External view, x 98, Sample 27.

PLATE 5

1. *Adelosina carinata striata* Wiesner. a and b, external views, x 85, Camalti Saltpan, Sample 19.
2. *Adelosina cliarensis* (Heron-Allen and Earland). External view, x 45, Camalti Saltpan, Sample 5.
3. *Adelosina* cf. *mediterraneensis* (Le Calvez J. and Y.). External view, x 40, Camalti Saltpan, Sample 5.
4. *Quinqueloculina seminula* (Linné). External views; a, x 40 and b, x 60, Camalti Saltpan, Sample 5.
5. *Nonion depressulum* (Walker and Jacob). External views, x 73, Camalti Saltpan, Sample 1.
6. *Ammonia compacta* Hofker. External views; a, spiral side and b, umbilical side, x 40, Camalti Saltpan, Sample 5.
7. *Ammonia tepida* Cushman. External views; a, spiral side and b, umbilical side, x 70, Camalti Saltpan, Sample 5; c, spiral side and d, edge view, x 65; Camalti Saltpan, Sample 1.
8. *Porosonion subgranosum* (Egger). a and b, external views; a, x 73, Camalti Saltpan, Sample 1; b, x 85, Camalti Saltpan, Sample 21.
9. *Elphidium complanatum* (d'Orbigny). External view, x 30, Camalti Saltpan, Sample 2.
10. *Leptocythere lacertosa* (Hirschmann). External view, left valve, x 98, Camalti Saltpan, Sample 24.
11. *Cyprideis* (C.) *anatolica* Bassiouni. External view, x 63, left valve, Camalti Saltpan, Sample 21.
12. *Cyprideis torosa* (Jones). External view, x 38, left valve, Camalti Saltpan, Sample 21.
13. *Loxoconcha elliptica* Brady. External view, x 63, left valve, Camalti Saltpan, Sample 23.

PLATE 6

- 1a-b.** *Hydrobia (Hydrobia) acuta* (Draparnaud), x 4.5, 1a apertural view, 1b dorsal view, Sample 21.
- 2a-b.** *Hydrobia (Hydrobia) acuta* (Draparnaud), x 3.5, 2a apertural view, 2b dorsal view, Sample 23.
- 3a-b.** *Rissoa labiosa* (Montagu), x 3, 3a apertural view, 3b dorsal view, Sample 25.
- 4a-b.** *Rissoa parva* (da Costa), x 6, 4a apertural view, 4b dorsal view, Sample 23.
- 5a-b.** *Rissoa violacea* Desmarest, x 3, 5a apertural view, 5b dorsal view, Sample 22.
- 6a-b.** *Pirenella conica* (Blainville), x 3.5, 6a apertural view, 6b dorsal view, Sample 21.
- 7a-b.** *Bittium desayesi* Cerulli and Irelli, x 4, 7a apertural view, 7b dorsal view, Sample 25.
- 8a-b.** *Bittium lacteum* (Philippi), x 4, 8a apertural view, 8b dorsal view, Sample 9.
- 9a-b.** *Bittium lacteum* (Philippi), x 4, 9a apertural view, 9b dorsal view, Sample 9.
- 10a-b.** *Bittium lacteum* (Philippi), x 2.5, 10a apertural view, 10b dorsal view, Sample 21.
- 11a-b.** *Bittium reticulatum* Philippi, x 2.5, 11a apertural view, 11b dorsal view, Sample 25.
- 12a-b.** *Columbella* sp., x 3, 12a apertural view, 12b dorsal view, Sample 26.
- 13a-b.** *Ostrea edulis* Linné, x 5.5, 13a left valve, external view, 13b left valve, internal view, Sample 21.
- 14a-b.** *Ostrea edulis* Linné, x 5, 14a left valve, external view, 14b left valve, internal view, Sample 21.
- 15a-b.** *Lucinella divaricata* (Linné), x 3.5, 15a left valve, external view, 15b left valve, internal view, Sample 9.
- 16a-b.** *Pseudochama gryphina* Lamarck, x 3, 16a right valve, external view, 16b right valve, internal view, Sample 21.
- 17a-b.** *Pseudochama gryphina* Lamarck, x 5.5, 17a left valve, external view, 17b left valve, internal view, Sample 21.
- 18a-b.** *Cerastoderma edule* (Linné), x 1.5, 18a left valve, external view, 18b left valve, internal view, Sample 17.
- 19a-b.** *Scrobicularia plana* da Costa, x 2, 19a left valve, external view, 19b left valve, internal view, Sample no 21.