Mediterranean Marine Science

Vol 11, No 1 (2010)

Season and size effects: changes in the quality of gilthead sea bream (*Sparus aurata* L.)

R. RODRIGUEZ, E. FOUNTOULAKI, K. GRIGORAKIS,
M. ALEXIS, R. FLOS

doi: [10.12681/mms.95](https://doi.org/10.12681/mms.95)

To cite this article:

RODRIGUEZ, R., FOUNTOULAKI, E., GRIGORAKIS, K., ALEXIS, M., & FLOS, R. (2010). Season and size effects: changes in the quality of gilthead sea bream (*Sparus aurata* L.). *Mediterranean Marine Science*, 11(1), 117–132.
<https://doi.org/10.12681/mms.95>

Season and size effects: changes in the quality of gilthead sea bream (*Sparus aurata* L.)

R. HURTADO-RODRÍGUEZ^{1,2,3}, E. FOUNTOULAKI², K. GRIGORAKIS²,
M. ALEXIS² and R. FLOS^{1,3}

¹ Departamento de Ingeniería Agroalimentaria y Biotecnología. Universidad Politécnica de Cataluña (UPC). Av. Canal Olímpico s/n, 08860, Castelfell de la Roca, Barcelona, Spain

² Hellenic Centre for Marine Research, P.O. Box 712, 19013 Anavissos, Hellas

³ Xarxa de Referència de Recerca i Desenvolupament en Aquicultura (XRAq), Spain

Corresponding author: efoudo@ath.hcmr.gr

Received: 30 June 2009; Accepted: 2 March 2010; Published on line: 26 March 2010

Abstract

The aim of this study was to investigate the potential effects of the harvesting season and fish size on the final quality of cage-farmed gilthead sea bream (*Sparus aurata*). Morphometric analysis, fat store estimation, fillet composition analysis, fillet fatty acids profile determination and sensory analysis were carried out on a commercially-sized cage farmed gilthead sea bream (*Sparus aurata* L.) population in April, June, July and November.

Both season and size had a significant effect in morphometric characteristics. The most pronounced size effect was the change of fish shape, becoming rounder with fish growth, the increase of filleting yield and condition index. An increase of both peritoneal and perivisceral fat was observed in summer months. In general, size had little effect on muscle composition. A significant size effect was observed in the fillet lipid content, which increased with it. Fillet lipid content was unexpectedly lower in the warm period (June-July). This can be possibly attributed to feeding restrictions that may have occurred, due to anoxic incidents related to high temperatures.

Slight seasonal differentiations occurred in the muscle fatty acids. No significant differences were found in the organoleptic characteristics of the two size groups compared.

Keywords: Fat depots; Fillet composition; Sea bream; Season; Size; Quality.

Introduction

Fish consumption is strongly recommended by health authorities due to its high ω -3 polyunsaturated fatty acid (PUFA) content, playing an important role in the pre-

vention of cardiovascular diseases (SCHMIDT *et al.*, 2005). An increasing percentage of fish consumption is of fish of aquaculture origin, aquaculture having experienced a boom in the last decade (FAO, 2006). The latest efforts in aquaculture focus on the

farming of new species and on the improvement of product quality and safety (FAO, 2006).

Farmed fish quality in production terms is related not only to the achieved growth, but also to the final characteristics of the fish, including its flesh composition and nutritional value, its organoleptic characteristics and its general acceptance.

Most of the studies concerning the final quality of common Mediterranean species like the sea bream (*S. aurata*) are related to its freshness and shelf life (LOUGOVOIS *et al.*, 2003; CHOULIARA *et al.*, 2004), growth (COMPANY *et al.*, 1999), rigor mortis (BAGNI *et al.*, 2006) or nutritional characteristics (IZQUIERDO *et al.*, 2005; TESTI *et al.*, 2006; FERREIRA *et al.*, 2007) as affected by handling, feeding, killing and post-mortem storage conditions. However, very little knowledge exists of the effects of the season on the muscle fatty acid profile (SENSO *et al.*, 2007) and on its general quality and somatometric characteristics (GRIGORAKIS *et al.*, 2002). Furthermore, these studies refer to fish of the same size produced at different seasons, i.e. they refer to different batches of fish with the same dietary history or origin. No data exists on how the muscle quality, the somatometric characteristics and fat depots of the same population change seasonally. Therefore, the aim of this study was to investigate the quality changes of a commercially-sized cage-farmed, gilthead sea bream population, in order to outline the potential effects of season and size on the final product quality.

Materials and Methods

Rearing conditions and samplings

Sea bream (*S. aurata*) of 65 g initial body weight were reared in 12 x 12 x 10 m³ cages in a fish farm in the Saronic Gulf (Poros is-

land, Greece), with a growing density of 10-12 kg/m³ for a period of 14 months. Fish were fed with a commercial feed at a concrete ratio based on feeding tables provided by the aquafeed company. The feed gross composition and fatty acid profile is presented in Table 1.

Fish were sampled from various cages of the same unit at various commercial sizes, and in four different sampling seasons (April, June, July and November). Water temperature in sampling periods was 15, 20, 25 and 18° C in April, June, July and November respectively (Table 2).

In total, 112 sea bream were killed by immersion in ice-water slurry, packed with flaked ice into polystyrene boxes and carried to the laboratory within 3-4 hours of harvesting. Samples of fish were kept at 4° C in a domestic refrigerator while sampling was performing.

Morphometric analysis, fillet composition analysis, fillet fatty acids profile determination and sensory analysis were conducted. The number of the individuals analyzed in each case is presented in Table 2.

Morphometric analysis, fat depots and fillet proximate composition

Total body weight, standard length, measured from the mouth to the base of the caudal fin, height, taken at the pectoral fin vertical line and maximum width, measured at the height of the pectoral fin, were determined in every fish. After dissection, liver, gonads, perivisceral fat (the fat stored inside the peritoneal cavity) and peritoneal fat (the fat located around the peritoneal cavity) were weighed.

Condition index (CI) were determined according to the following formula $CI = 100 \times W \text{ (g)} / Lst^3 \text{ (mm)}^3$ where W is the fish total body weight and Lst its standard length.

Peritoneal fat index, perivisceral fat index, hepatosomatic index (HSI), gonadosomatic index (GSI) and fillet yield, were calculated as percentages of the total body weight.

Fillet proximate composition analyses took place according to the official methods of AOAC (1998).

Fillet fatty acids profile determination

Samples for muscle fatty acid analysis were extracted from the front dorsal part of the fillet in a chloroform methanol (2:1 v/v) solution, within 7-8 hours after slaughter. Pools of total 1g tissue each were made from 3-4 different fish.

The fatty acids were extracted according to FOLCH *et al.* (1957) and the methyl esters were obtained as described by CHRISTIE (1989). The methyl esters were separated and quantified by gas chromatography with flame detector, using helium as carrier gas as described by FOUNTOULAKI *et al.* (2003). A standard fatty acid mixture (Supelco 37 component FAME mix) was used as a pattern and the data were analyzed using the Millenium Software. The feed methyl esters were obtained directly from a 50-70 mg of ground sample as described by LEPAGE & ROY (1984).

Atherogenicity index (AI) and thrombogenicity index (TI) were determined by using the formulae: $AI = (12:0 + 4 \times 14:0 + 16:0) / (\sum MUFA + \sum \omega 6 + \sum \omega 3)$ and $TI = (14:0 + 16:0 + 18:0) / [0.5 \sum MUFA + 0.5 \sum \omega 6 + 3 \sum \omega 3 + (\sum \omega 3 / \sum \omega 6)]$, where $\sum MUFA$ is the sum of mono-unsaturated fatty acids, $\sum \omega 3$ the total n-3 fatty acids and $\sum \omega 6$ the total n-6 fatty acids respectively.

Sensory analysis

In July, 10 sea bream were additionally sampled in order to compare the sensory quality of different sizes. Two different

Table 1
Feed composition and fatty acids profile
as % of the total fatty acids.

Feed composition %	
Protein	45.7
Fat	18.9
Moisture	7.2
Ash	9.3

Fatty acid	%
14:00	5.05
16:00	14.54
18:00	2.52
$\sum SFA$	25.34
16:1 $\omega 7$	5.77
18:1 $\omega 7$	2.73
18:1 $\omega 9$	15.41
20:1 $\omega 9$	5.06
$\sum \omega 9$	24.02
22:1 $\omega 11$	4.9
$\sum MUFA$	39.27
18:3 $\omega 3$	2.17
18:4 $\omega 3$	2.29
20:5 $\omega 3$	7.41
22:5 $\omega 3$	1.04
22:6 $\omega 3$	11.09
$\sum \omega 3$	26.92
18:2 $\omega 6$	11.55
20:4 $\omega 6$	0.66
$\sum \omega 6$	7.47
$\sum PUFA$	35.39
$\sum \omega 3 / \omega 6$	3.60
EPA + DHA	19.74

size groups were examined for their possible organoleptic differences (fish of <250 g and fish of 350 g). Within 8-10 hours of harvesting, fish were gutted, filleted and whole fillets were wrapped in aluminium

Table 2
Size ranges, size averages, and total samples analysed in every sampling.

Analysis	April	June	July	November
Size range (g)	126.0-323.9	207.8-374.1	172.2-376.3	369.2-500.3
Size average (g)	210.3	281.7	271.5	431.5
Morphometry	30 fish	30 fish	30 fish	10 fish
Composition	30 fish	30 fish	30 fish	10 fish
Organoleptic	-	-	20 fillets	-
Fatty acids*	5 pools	-	5 pools	3 pools

*Pools for fatty acid analysis were made of 3-4 fishes.

foil and steam cooked for 20 min. Quantitative Descriptive Analysis (QDA) (MURRAY *et al.*, 2001) was used to evaluate flesh taste, flavour, colour, oiliness, elasticity, hardness, stickiness and general acceptance. Ten assessors previously having received the training required for QDA (MURRAY *et al.*, 2001) conducted the sensory test. A 0 to 5 scale was used for rating the intensity of the characteristics.

Statistical analysis

Regression analysis against body weight was performed for each of the studied parameters within each season, and slopes were compared. In cases of non-significant weight effect, one-way analysis of variance (ANOVA) was used to find the effects of season. In cases of significant weight effects and parallel regression lines (i.e. similar effect for all seasons), ANCOVA with weight as covariance was used for group comparisons. In cases of significant weight effect and non-parallel lines, ANCOVA comparisons were made only for the season groups that were parallel, since comparisons among non-parallel lines do not make sense (RAUBEHEIMER & SIMPSON 1992).

Regression analysis was done by Statgraphics and analysis of variance by SPSS

13.0 software. The Tukey test was used to determine the possible significant differences among values ($p < 0.05$). Data from sensory analysis were analyzed by a non parametric Kruskal-Wallis test.

Results

Both season and size had a significant effect on morphometric characteristics of the studied fish. Morphometric characteristics altered by fish size in the June and July samples, i.e. in the summer period (Table 3). Thus, length/height and length/width were found to decrease with fish size within summer samplings, while condition index increased.

Table 4 shows the morphometric characteristics of the studied population at the different sampling seasons. Regarding the season, there was no significant effect on condition index, while length/height and length/width were found significantly increased in the July sampling.

Filleting yield increased with fish size for July fish, but remained unaffected by size in the rest of the samplings. Significant differences were also found in HSI and GSI with respect to the fish size, increasing with it, in November and April, respectively.

Table 3
Characteristics of regression lines describing morphometric characteristics and quality indexes
in relation to body weight.

	Intercept	Slope	*R ²	Sig ¹	% explain of the variability
Length/Height					
April	2.8536	-0.0003	-0.082	ns	0.7
June	2.9687	-0.0011	-0.473	**	22.4
July	3.8914	-0.0020	-0.686	**	47.0
Nov	3.2663	-0.0011	-0.457	ns	20.9
Length/Width					
April	6.6303	0.0016 ^b	0.150	ns	2.3
June	8.5753	-0.0077 ^a	-0.489	**	23.9
July	14.1796	-0.0107 ^a	-0.476	**	22.7
Nov	7.5265	-0.0019 ^{ab}	-0.240	ns	5.8
Condition Index					
April	2.9020	-0.0005 ^a	0.094	ns	0.9
June	2.2019	0.0027 ^b	0.595	**	35.4
July	2.2403	0.0020 ^b	0.62	**	38.4
Nov	2.0780	0.0017 ^{ab}	0.495	ns	24.5
Fillet Yield					
April	27.9608	0.0172 ^{ab}	0.261	ns	6.8
June	34.0053	-0.0084 ^a	-0.166	ns	2.7
July	25.0833	0.0355 ^b	0.0552	**	30.5
Nov	29.8282	0.0190 ^{ab}	0.586	ns	34.4
Gonadosomatic Index					
April	-0.3361	0.0022 ^a	0.773	**	59.8
June	-0.0164	0.0002 ^b	0.155	ns	2.4
July	0.0309	0.0002 ^b	0.256	ns	6.6
Nov	0.1290	0.0001 ^{ab}	0.021	ns	0.0
Hepatosomatic Index					
April	2.6441	-0.0032 ^b	-0.352	ns	12.4
June	1.1112	0.0005 ^c	0.157	ns	2.5
July	1.0220	0.0003 ^{ac}	0.117	ns	1.4
Nov	-0.4951	0.004 ^{1a}	0.775	**	60.1
Peritoneal Fat Index					
April	0.5148	-0.0001 ^b	-0.018	ns	0.03
June	0.5806	0.0016 ^{ab}	0.136	ns	1.8
July	-0.5298	0.0051 ^a	0.615	**	37.9
Nov	0.7760	-0.0005 ^b	-0.112	ns	1.3

(continued)

Table 3 (continued)

	Intercept	Slope	*R ²	Sig ¹	% explain of the variability
Perivisceral Fat Index					
April	4.0946	-0.082b	-0.311	ns	9.7
June	-1.2654	0.0014a	0.545	**	29.7
July	0.1832	0.0112a	0.545	**	29.7
Nov	-2.1225	0.0109a	0.561	ns	31.5
Total Fat Index					
April	4.6080	-0.0083b	-0.2946	ns	8.7
June	-0.6803	0.0162a	0.548	**	30.1
July	-0.3403	0.0167a	0.652	**	42.5
Nov	-1.346	0.0104ab	0.500	ns	25.

Different letters (a, b, c) stand for statistically significant differences ($p < 0.05$) between sampling seasons
¹Significance level of the regression (ns = no significant; * $p < 0.05$; ** $p < 0.01$)

*Correlation coefficient

Table 4
Morphometric characteristics, quality indexes of the studied population
at the different sampling seasons.

	April	15° C	June	20° C	July	25° C	Nov	18° C
	Mean	SEM ¹	Mean	SEM ¹	Mean	SEM ¹	Mean	SEM ¹
Length/Height	2.70a	0.03	2.68a	0.03	3.35c	0.03	2.99b	0.07
Length/Width	6.56a	0.20	6.46a	0.17	11.28c	0.17	7.72b	0.41
Condition Index	2.80	0.05	2.97	0.04	2.79	0.04	2.79	0.05
Fillet Yield	32.75ab	0.7	31.45a	0.60	34.72b	0.6	35.02b	1.39
Gonadosomatic Index	0.17	0.02	0.02	0.02	0.08	0.02	0.07	0.05
Hepatosomatic Index	1.96	0.06	1.27	0.05	1.10	0.05	1.32	0.13
Peritoneal Fat Index	0.64b	0.11	1.02c	0.09	0.87	0.09	0.13a	0.22
Perivisceral Fat Index	2.84b	0.28	2.77b	0.24	3.33b	0.24	1.38a	0.56
Total Fat Index	3.48b	0.31	3.79b	0.27	4.20b	0.26	1.51a	0.61
N	30		30		30		10	

Different letters (a, b, c) stand for statistically significant differences ($p < 0.05$).

¹Standard error of the means. N = Number of samples.

Significant size effect for perivisceral and total fat depots, increasing with it, was found only in summer samplings and for peritoneal fat in the July sampling (Table 3).

Seasonal significant differences were found in most of the quality indexes. Periv-

isceral and total fat depots were higher during the summer (June-July) and peritoneal fat was also significantly higher in June. Average peritoneal fat content in July seems also to be higher than in the spring and autumn, but this sampling could not be in-

cluded, since a different size-effect was observed for this sampling (non-parallel regression line).

Size did not affect muscle protein (Table 5), but affected muscle fat, which increased with fish size in April and July. Muscle moisture and ash decreased with size in July but remained unaffected by size in the rest of the samplings (Table 5).

With respect to fillet composition, protein was significantly higher during the sum-

mer (June and July) whereas lipid percentage was significantly lower in the same season. Moisture and ash content were found higher in June than in the rest of the samplings (Table 6).

The fatty acid profiles of the flesh are shown in Table 7. In all cases, the predominant fatty acids among the saturated ones (SFA) were myristic acid (14:0), palmitic acid (16:0) and stearic acid (18:0). Palmitoleic acid (16:1 ω 7) and oleic acid (18:1 ω 9)

Table 5
Characteristics of regression lines describing fillet composition in relation to body weight.

	Intercept	Slope	R ²	Sig ¹	% explain of the variability
Muscle Fat					
April	3.4069	0.0158	0.439	**	19.3
June	2.0674	0.0059	0.284	ns	8.1
July	1.3684	0.0130	0.411	*	16.9
Nov	8.4160	-0.0016	-0.043	ns	0.2
Muscle Protein					
April	19.7389	0.0013	0.089	ns	0.8
June	21.8147	-0.0011	-0.107	ns	1.1
July	20.9919	0.0014	0.1488	ns	2.2
Nov	19.5390	0.0016	0.1470	ns	2.2
Muscle Moisture					
April	67.2890	-0.0040	-0.012	ns	0.0
June	74.3742	-0.0062	-0.313	ns	9.8
July	76.0066	-0.0168	-0.539	**	29.1
Nov	70.1536	0.0004	0.011	ns	0.0
Muscle Ash					
April	1.4228	0.0000	0.054	ns	0.3
June	1.5336	-0.0001	-0.125	ns	1.5
July	1.5216	-0.0004	-0.389	*	15.2
Nov	1.3730	0.0000	0.063	ns	0.4
Energy (Kcal/g)					
April	1.2123	0.0011	0.335	ns	11.3
June	1.0841	0.0005	0.278	ns	7.7
July	0.9792	0.0013	0.459	*	21.1
Nov	1.7206	-0.0005	-0.147	ns	2.1

¹Significance level of the regression (ns = no significant; * $p < 0.05$; ** $p < 0.01$)

Table 6
Muscle composition of the studied population at the different
sampling seasons $\pm$ standard error of the means.

	April 15° C	June 20° C	July 25° C	Nov 18° C
Fat	7.41 $\pm$ 0.36 ^c	3.64 $\pm$ 0.31	4.90 $\pm$ 0.31 ^b	6.00 $\pm$ 0.71 ^{bc}
Protein	20.07 $\pm$ 0.14	21.50 $\pm$ 0.12 ^b	21.36 $\pm$ 0.12 ^b	20.09 $\pm$ 0.27
Moisture	65.84 $\pm$ 1.93	72.73 $\pm$ 1.66 ^b	71.44 $\pm$ 1.65 ^{ab}	71.89 $\pm$ 3.84 ^{ab}
Ash	1.42 $\pm$ 0.012	1.50 $\pm$ 0.01 ^b	1.42 $\pm$ 0.01	1.42 $\pm$ 0.02
*N	30	30	30	10

Different letters (a, b, c) stand for statistically significant differences (p < 0.05) *N = Number of samples.

Table 7
Seasonal changes of fillet fatty acids (expressed as % of the total fatty acid
content $\pm$ standard error of the means).

	April 15° C	July 25° C	Nov 18° C
Fatty acid	Mean	Mean	Mean
14:0	3.41 $\pm$ 0.08 ^a	4.06 $\pm$ 0.08 ^b	3.43 $\pm$ 0.02 ^a
16:0	14.59 $\pm$ 0.04	15.43 $\pm$ 0.31	14.71 $\pm$ 0.22
18:0	3.82 $\pm$ 0.03 ^b	3.64 $\pm$ 0.16 ^{ab}	3.30 $\pm$ 0.11 ^a
Σ SFA	23.11 $\pm$ 0.08 ^a	24.45 $\pm$ 0.42 ^b	22.64 $\pm$ 0.34 ^a
16:1 ω 7	5.71 $\pm$ 0.12 ^a	6.58 $\pm$ 0.23 ^b	5.96 $\pm$ 0.11 ^{ab}
18:1 ω 7	2.92 $\pm$ 0.012	2.77 $\pm$ 0.12	2.93 $\pm$ 0.09
18:1 ω 9	16.80 $\pm$ 0.23 ^{ab}	16.20 $\pm$ 0.55 ^a	17.99 $\pm$ 0.36 ^b
20:1 ω 9	2.25 $\pm$ 0.13 ^a	1.99 $\pm$ 0.19 ^a	3.53 $\pm$ 0.04 ^b
$\Sigma\omega$ 9	21.17 $\pm$ 0.29 ^a	19.67 $\pm$ 0.69 ^a	23.60 $\pm$ 0.08 ^b
22:1 ω 11	1.62 $\pm$ 0.05 ^a	1.54 $\pm$ 0.11 ^a	3.11 $\pm$ 0.17 ^b
Σ MUFA	31.15 $\pm$ 0.31 ^a	30.46 $\pm$ 0.9 ^a	35.34 $\pm$ 0.18 ^b
18:3 ω 3	1.62 $\pm$ 0.03 ^b	1.47 $\pm$ 0.04 ^{ab}	1.44 $\pm$ 0.06 ^a
18:4 ω 3	1.20 $\pm$ 0.03	1.28 $\pm$ 0.04	1.17 $\pm$ 0.04
20:5 ω 3	7.83 $\pm$ 0.09 ^a	8.57 $\pm$ 0.24 ^b	7.34 $\pm$ 0.06 ^a
22:5 ω 3	4.09 $\pm$ 0.05	4.33 $\pm$ 0.15	4.25 $\pm$ 0.21
22:6 ω 3	14.35 $\pm$ 0.33	13.85 $\pm$ 0.9	13.67 $\pm$ 0.13
$\Sigma\omega$ 3	30.83 $\pm$ 0.34	31.31 $\pm$ 0.79	29.48 $\pm$ 0.22
18:2 ω 6	11.75 $\pm$ 0.17 ^b	10.57 $\pm$ 0.32 ^{ab}	9.44 $\pm$ 0.47 ^a
20:4 ω 6	0.85 $\pm$ 0.06	0.84 $\pm$ 0.08	0.72 $\pm$ 0.04
$\Sigma\omega$ 6	13.79 $\pm$ 0.19 ^b	12.65 $\pm$ 0.32 ^b	11.39 $\pm$ 0.43 ^a
Σ PUFA	45.81 $\pm$ 0.33 ^b	44.27 $\pm$ 0.81 ^{ab}	41.98 $\pm$ 0.38 ^a
$\Sigma\omega$ 3/ $\Sigma\omega$ 6	2.24 $\pm$ 0.04 ^a	2.48 $\pm$ 0.07 ^{ab}	2.59 $\pm$ 0.11 ^b
EPA + DHA	22.18 $\pm$ 0.36	22.42 $\pm$ 0.84	21.01 $\pm$ 0.14
AI	0.37 $\pm$ 0.004 ^a	0.43 $\pm$ 0.008 ^b	0.37 $\pm$ 0.003 ^a
TI	0.19 $\pm$ 0.002	0.20 $\pm$ 0.005	0.19 $\pm$ 0.002
N	5	5	3

were the predominant fatty acids among the mono-unsaturated ones (MUFA). Among poly-unsaturated fatty acids (PUFA), the most abundant were linoleic (18:2 ω 6), EPA (20:5 ω 3) and DHA (22:6 ω 3). Concerning the global percentages, the most abundant were PUFAs (with $\sum\omega$ 3 predominating among them), followed by MUFAs (mostly $\sum\omega$ 9). Table 7 shows the seasonal variations in the fillet fatty acid profile of the studied population. $\sum$ SFA was significantly higher during the summer, due to the significantly higher content of 14:0.

$\sum$ MUFA were found higher in November. The 18:1 ω 9 content was lower in summer than in autumn and spring. With regard to PUFAs, $\sum\omega$ 6 were found significantly lower in the November sampling, mostly due to the lower 18:2 ω 6 content, and ratio $\sum\omega$ 3/ $\sum\omega$ 6 was found higher at this

season. Among the ω 3 fatty acids, the EPA percentage was higher in the summer than in the rest of the samplings.

With regard to the atherogenicity and thrombogenicity indexes (AI and TI respectively), only AI was found significantly higher in summer.

Sensory analysis did not show significant differences between the two studied size groups (Fig. 1).

Discussion

For most of the studied parameters (length/height, length/width, condition index, fillet yield, peritoneal, perivisceral and total deposit fat, fillet fat, moisture, ash and energy) the size effects were most profound in summer, mostly in July and to a lesser degree in June, than in the other samplings (Tables 3 and 5). This could be possibly due

Fig. 1: Sensorial scores of the descriptors for the studied size groups.

to the higher size ranging within the summer samplings and especially in July (Table 2).

Morphometric characteristics in our experiment were within the normal values for that species, although rearing conditions, i.e. growing density and feeding, can affect the morphological characteristics (FLOS *et al.*, 2002).

The facts of significant length/height and length/width reduction, as well as the significant increase with size (in June and July) of the condition index (Table 3), constitute serious indications that in commercial sizes the main fish growth is due to a gain in weight, while fish become more round-shaped.

In the present study condition index did not show significant seasonality (Table 4). A seasonality in condition index has been mentioned for Atlantic Salmon *Salmo salar* (YOUNG *et al.*, 2006), with highest values observed at the end of the warm period (in September) and a reduction observed thereafter during winter. However, condition index seasonality has been found to be dietarily affected by the dietary energy content (SOLBERG, 2004; YOUNG *et al.*, 2006), while positive correlation with fish size (YOUNG *et al.*, 2006) and influence of other dietary treatments such as food deprivation (GRIGORAKIS & ALEXIS, 2005) can further complicate a possible seasonal effect in condition index.

The average filleting yield in the present study (33.2%) remained slightly lower than the range of 34.3 – 48 % mentioned in literature (GRIGORAKIS & ALEXIS, 2005; TESTI *et al.*, 2006). Filleting yield increased with size only in July samples. The higher size heterogeneity for July fish compared to the rest of the samplings (Table 2) gives a possible explanation for significant size effect only for this sampling. Size effect explains why the present average filleting yield was found slightly lower than that of

the literature, since the present study also included a large number of smaller fish (150 – 250 g). The significant increase of filleting yield with size, also poses the question of finding the most profitable sizes for gilt-head sea bream filleting.

Gonadosomatic index was lower during the summer (June, July) and increased in November, but in all cases remained very low. A similar seasonality without reaching maturity has been previously described for gilthead sea bream by GRIGORAKIS *et al.* (2002).

Hepatosomatic index had its maximum value in April and decreased in summer. Liver lipid depot increase with water temperature drop has been previously mentioned by IBARZ *et al.* (2005) as a mechanism of adaptation to the cold. However, the reason for HSI summer reduction can also be due to dietary factors, and this will further be analysed.

Total depot of fat, considered as the sum of peritoneal and perivisceral fat indexes, is comparable to that previously found (GRIGORAKIS *et al.*, 2002; SANTINHA *et al.*, 1999). The seasonality observed for fat depot accumulation, being higher in the warm months (June and July) partially agrees with what was found by GRIGORAKIS *et al.* (2002) who also found increased fat depots at the end of the summer, but also for fish sampled later in the autumn.

The presently found average values for muscle protein (20.8 %), fat (5.8 %), moisture (71.3 %) and ash (1.4 %) are within the common limits reported for this species (FLOS *et al.*, 2002; GRIGORAKIS *et al.*, 2002; ÖZYURT *et al.*, 2005; SENSO *et al.*, 2007).

Fillet protein content and fillet lipid content showed an opposite trend. A significant increase in fillet protein content was found during the summer together with

a significant fillet lipid content decrease. Protein content stability at commercial sizes, with no seasonal or dietary variations has been mentioned by previous studies (GRIGORAKIS *et al.*, 2002; ÖZYURT *et al.*, 2005; SENSO *et al.*, 2007). This contradicts the present results where a seasonal effect was observed with protein found to be higher in summer fish.

Also a seasonal change was observed for muscle fat, but contrary to the literature which found either muscle fat increase during summer (GRIGORAKIS *et al.*, 2002) or no specific muscle fat trend throughout the year (SENSO *et al.*, 2007). The fact of such different seasonal trends indicates the sensitivity of gilthead sea bream muscle fat to dietary and environmental conditions. A possible explanation of low fat in the summer period could be due to muscle fat mobilization because of high metabolic needs in summer not met by the dietary intake. The low HSI values in the summer months (Table 4) could possibly indicate the same thing. Liver and muscle fat have been mentioned as the first fat depots mobilizing in the gilthead sea bream (GRIGORAKIS & ALEXIS, 2005). Insufficient dietary energy intake could be probably explained due to a frequent summer phenomenon in Greek aquaculture, where high water temperatures in combination with low levels of dissolved oxygen, often demand restriction of feeding.

Muscle lipid content and muscle moisture content showed an opposite trend, which is in accordance with the literature (WEATHERLY & GILL, 1983).

The fatty acid profile of the present study is in general typical for farmed fish species. However, saturated fatty acids were found slightly lower than the usual range of 26 – 31 % mentioned in the literature (GRIGORAKIS *et al.*, 2002; SAGLIK *et al.*, 2003; IBARZ *et al.*, 2005; IZQUIERDO

et al., 2005; VASILIADOU *et al.*, 2005; TESTI *et al.*, 2006; MNARI *et al.*, 2007), but similar to the levels mentioned by SENSO *et al.* (2007). Mono-unsaturated fats are within the values of 30 – 37 % mentioned in the literature (GRIGORAKIS *et al.*, 2002; SAGLIK *et al.*, 2003; VASILIADOU *et al.*, 2005; TESTI *et al.*, 2006; MNARI *et al.*, 2007). As in our study, in all of the previously mentioned literature, palmitic acid (16:0) is the predominant saturated fatty acid, while oleic acid (18:1 ω 9) predominated among the mono-unsaturated ones. Regarding ω 3 poly-unsaturated fatty acids, their total levels were found similar to those of SENSO *et al.* (2007) and MNARI *et al.* (2007), but higher than those mentioned in previous studies (GRIGORAKIS *et al.*, 2002; SAGLIK *et al.*, 2003; VASILIADOU *et al.*, 2005; TESTI *et al.*, 2006). The literature showed a great variability in ω 3/ ω 6 ratios, ranging from 1.47 (IBARZ *et al.*, 2005) to 4.09 (MNARI *et al.*, 2007), and present results are within this range. Also the total EPA and DHA levels are in accordance with GRIGORAKIS *et al.*, 2002; SAGLIK *et al.*, 2003; IBARZ *et al.*, 2005; IZQUIERDO *et al.*, 2005; VASILIADOU *et al.*, 2005; TESTI *et al.*, 2006; MNARI *et al.*, 2007; SENSO *et al.*, 2007.

Seasonal effects on the fillet fatty acid profile of cultured gilthead sea bream has rarely been studied (SENSO *et al.*, 2007). In general fish fatty acid changes due to the season, mainly include an increase in the unsaturation level (primarily in the polar lipid fraction) as a response to the cold, in order to maintain the appropriate fluidity of biological membranes (HAZEL & PROSSER, 1974; DELGADO *et al.*, 1994). In gilthead sea bream, an increase in unsaturation in white muscle, with artificially applied temperature drop, has been mentioned (IBARZ *et al.*, 2005). An increase of DHA at low tem-

peratures has been mentioned in a seasonal study of wild gilthead sea bream (ÖZYURT *et al.*, 2005). Our study, however, failed to show such a seasonal impact similar to the other existing source in cultured gilthead sea bream (SENSO *et al.*, 2007). Higher SFA levels in summer were also observed in our study, but were not followed with a respective decrease in unsaturates (Table 7). The existing data for wild gilthead sea bream (ÖZYURT *et al.*, 2005) also failed to show an increase in the total unsaturated fatty acids in colder periods (winter). In any case, fatty acid mobilization in wild fish is expected to be different, since maturation and spawning also play an important part in this process (ÖZYURT *et al.*, 2005), unlike in their farmed counterparts.

The atherogenicity index is the ratio between the main saturate and the main unsaturated fatty acids, and the higher its value, the higher the tendency of lipids to adhere to cells of the immunological and circulatory systems. The average value (0.39) was within the range of 0.25 (SENSO *et al.*, 2007) and 0.49 (AMERIO *et al.*, 1996) found for the same species. The present study indicated that AI is seasonally variable. However, the only existing previous reference to sea bream did not find significant seasonal variations in AI (SENSO *et al.*, 2007).

The thrombogenicity index is an indicator of the clot forming tendency in the blood vessels. According to our results, TI does not appear to depend on the season. Similarly SENSO *et al.* (2007) did not show any seasonal effect on TI. Since limited respective data occur in literature regarding AI and TI season and size effect, further research would be useful.

One of the most important aspects for consumers is the sensory characteristics of the fish. Organoleptic characteristics mainly depend on the fillet composition and its

fatty acid profile and both of them are strongly affected by dietary factors (IZQUIERDO *et al.*, 2005). Results on sensory analysis, not exhibiting any organoleptic differences since no significant differences were found, correlated with size in the fillet composition and were minor in the fatty acid profile.

Conclusion

Results in the present study showed that the quality of the cage-farmed sea bream fluctuates with the harvesting season. The most profound seasonal effect was the increase of deposit fat (both perivisceral and peritoneal) during the summer period. A minor seasonal effect was observed for muscle fatty acids. On the other hand, the size did not affect fillet protein content or organoleptic quality of the fish but had some significant positive effects on fillet fat. Size mainly affected the morphometric characteristics of the fish. In particular there is an indication the fish become more round-shaped with growth. Some quality indexes which are important for fish consumers, such as the condition index, fillet yield or peritoneal fat and total deposit fat, increased with size.

Acknowledgements

This study has been supported by the Generalitat de Catalunya and the Xarxa de Referència de Recerca i Desenvolupament en Aqüicultura (XRAq). The authors would like to acknowledge Lysandros Kyriakakos, from Kaloni S.A. for providing the fish and Dr. Ioannis Nengas for his useful help.

References

- AMERIO, M., RUGGI, C. & BADINI, C., 1996. Meat quality of reared fish: nu-

- tritional aspects. *Italian Journal of Food Science*, 8 (3): 221-229.
- AOAC, 1998. 16th Edition CD-ROM 3rd revision.
- BAGNI, M., CIVITAREALE, C., PRIORI, A., BALLERINI, A., FINOIA, M., BRAMBILLA, G. & MARINO, G., 2007. Pre-slaughter crowding stress and killing procedures affecting quality and welfare in sea bass (*Dicentrarchus labrax*) and sea bream (*Sparus aurata*). *Aquaculture*, 263 (1-4): 52-60.
- CHOULIARA, I., SAVVAIDIS, I.N., PANAGIOTAKIS, N. & KONTOMINAS, M.G., 2004. Preservation of salted, vacuum-packaged, refrigerated sea bream (*Sparus aurata*) fillets by irradiation: microbiological, chemical and sensory attributes. *Food Microbiology*, 21 (3): 351-359.
- CHRISTIE, W.W., 1989. *Gas Chromatography and Lipids: a Practical Guide*. The Oily Press, Ayr, Scotland, 307 pp.
- COMPANY, R., CALDUCH-GINER, J.A., KAUSHIK, S. & PÉREZ-SÁNCHEZ, J., 1999. Growth performance and adiposity in gilthead sea bream (*Sparus aurata*): risks and benefits of high energy diets. *Aquaculture*, 171 (3-4): 279-292.
- DELGADO, A., ESTEVEZ, A., HORTELANO, P. & ALEJANDRE, M.J., 1994. Analyses of fatty acids from different lipids in liver and muscle of sea bass (*Dicentrarchus labrax* L.). influence of temperature and fasting. *Comparative Biochemistry & Physiology*, 108A: 673-680.
- FAO, 2006. State of world aquaculture. Available on: <ftp://ftp.fao.org/docrep/fao/009/a0874e/a0874e00.pdf>
- FERREIRA-PINTO, J., NUNES, M.L. & CARDOSO, C., 2007. Feeding interruption and quality of cultured gilthead sea bream. *Food Chemistry*, 100 (4): 1504-1510.
- FLOS, R., REIG, L., OCA, J. & GINOVART, M., 2002. Influence of marketing and different land-based systems on gilthead sea bream (*Sparus aurata*) quality. *Aquaculture International*, 10 (3): 189-206.
- FOLCH, J., LEES, M. & SLOANE STANLEY, G.H., 1957. A simple method for the isolation and purification of the total lipid from animal tissue. *Journal of Biological Chemistry*, 226 (1): 497-509.
- FOUNTOULAKI, E., ALEXIS, M.N., NENGAS, I. & VENOU, B., 2003. Effects of dietary arachidonic acid (20:4n-6), on growth, body composition, and tissue fatty acid profile of gilthead bream fingerlings (*Sparus aurata* L.). *Aquaculture*, 225 (1-4): 309-323.
- GRIGORAKIS, K., ALEXIS, M.N., TAYLOR, K.D.A. & HOLE, M., 2002. Comparison of wild and cultured gilthead sea bream (*Sparus aurata*): composition, appearance and seasonal variations. *International Journal of Food Science and Technology*, 37 (5): 477-484.
- GRIGORAKIS, K. & ALEXIS, M.N., 2005. Effects of fasting on the meat quality and fat deposition of commercial-size farmed gilthead sea bream (*Sparus aurata* L.) fed different dietary regimes. *Aquaculture Nutrition*, 11 (5): 341-344.
- HAZEL, J.R. & PROSSER, C.L., 1974. Molecular mechanisms of temperature compensation in poikilotherms. *Physiological Reviews*, 54 (3): 620-677.
- IBARZ, A., BLASCO, J., BELTRÁN, M., GALLARDO, M.A., SÁNCHEZ, J. & SALA FERNÁNDEZ-BORRÁS, J., 2005. Cold-induced alterations on proximate composition and fatty acid profiles of several tissues in gilthead sea bream (*Sparus aurata*). *Aquaculture*, 249 (1-4): 477-486.

- IZQUIERDO, M.S., MONTERO, D., ROBAINA, L., CABALLERO, M.J., ROSEN LUND, G. & GINÉS, R., 2005. Alterations in fillet fatty acid profile and flesh quality in gilthead seabream (*Sparus aurata*) fed vegetable oils for a long term period. Recovery of fatty acid profiles by fish oil feeding. *Aquaculture*, 250 (1-2): 431-444.
- LEPAGE, I. & ROY, M., 1984. Improved recovery of fatty acid through direct transesterification without prior extraction or purification. *Journal of Lipid Research*, 25: 1391-1396.
- LOUGOVOIS, V.P., KYRANAS, E.R. & KYRANA, V.R., 2003. Comparison of selected methods of assessing freshness quality and remaining storage life of iced gilthead sea bream (*Sparus aurata*). *Food Research International*, 36 (6): 551-560.
- MNARI, A., BOUHLEL, I., CHRAIEF, I., HAMMAMI, M., ROMDHANE, M.S., EL CAFSI, M. & CHAOUCH, A., 2007. Fatty acids in muscles and liver of Tunisian wild and farmed gilthead sea bream (*Sparus aurata*). *Food Chemistry*, 100 (4): 1393-1397.
- MURRAY, J.M., DELAHUNTY, C.M. & BAXTER, I.A., 2001. Descriptive sensory analysis: past, present and future. *Food Research International*, 34 (6): 461-471.
- ÖZYURT, G., POLAT, A. & ÖZKÜTÜK, S., 2005. Seasonal changes in the fatty acids of gilthead sea bream (*Sparus aurata*) and White Sea bream (*Diplodus sargus*) captured in Iskenderum Bay, eastern Mediterranean coast of Turkey. *European Food Research and Technology*, 220: 120-124.
- RAUBENHEIMER, D. & SIMPSON, S.J., 1992. Analysis of covariance: an alternative to nutritional indices. *Entomologia Experimentalis et Applicata*, 62 (3): 221-231.
- SAGLIK, S., ALPASLAN, M., GEZLIN, T., CETLINTURK, K., TEKINAY, A. & GUVEN, K.C., 2003. Fatty acid composition of wild and cultivated gilthead seabream (*Sparus aurata*) and sea bass (*Dicentrarchus labrax*). *European Journal of Lipid Science and Technology*, 105 (2): 104-107.
- SANTINHA, P.J.M., MEDALE, F., CORRAZE, G. & GOMEZ, E.F.S., 1999. Effects of the dietary protein : lipid ratio on growth and nutrient utilization in gilthead seabream (*Sparus aurata* L.). *Aquaculture Nutrition*, 5 (3): 147-156.
- SCHMIDT, E.B., ARNESEN, H., DECATERINA, R., RASMUSSEN, L.H. & KRISTENSEN, S.D., 2005. Marine n – 3 polyunsaturated fatty acids and coronary heart disease. Part I. Background, epidemiology, animal data, effects on risk factors and safety. *Thrombosis Research*, 115 (3): 163-170.
- SENSO, L., SUÁREZ, M.D., RUIZ-CARA, T. & GARCÍA-GALLEGO, M., 2007. On the possible effects of harvesting season and chilled storage on the fatty acid profile of the fillet of farmed gilthead sea bream (*Sparus aurata*). *Food Chemistry*, 101 (1): 298-307.
- SOLBERG, C., 2004. Influence of dietary oil content on the growth and chemical composition of Atlantic salmon (*Salmo salar*). *Aquaculture Nutrition*, 10 (1): 31-37.
- TESTI, S., BONALDO, A., GATTA, P.P. & BADIANI, A., 2006. Nutritional traits of dorsal and ventral fillets from three farmed fish species. *Food Chemistry*, 98 (1): 104-111.
- VASILIADOU, S., AMBROSIADIS, I., VARELTZIS, K., FLETOURIS, D. & GAVRIILIDOU, I., 2005. Effect of smoking on quality parameters of farmed gilthead sea bream (*Sparus aurata*) and

- sensory attributes of the smoked product. *European Food Research and Technology*, 221 (3-4): 232-236.
- WEATHERLEY, A.H. & GILL, H.S., 1983. Protein, lipid, water and caloric contents of immature rainbow trout, *Salmo gairdneri* Richardson, growing at different rates. *Journal of Fish Biology*, 23 (6): 653-673.
- YOUNG, A., MORRIS, P.C., HUNTINGFORD, F.A. & SINNOTT, R., 2006. Replacing fish oil with pre-extruded carbohydrate in diets for Atlantic salmon, *Salmo salar*, during their entire marine grow-out phase: Effects on growth composition and colour. *Aquaculture*, 253 (1-4): 531-546.