

Mediterranean Marine Science

Vol 7, No 2 (2006)



Spectroscopic studies of humic acids from subsurface sediment samples collected across the Aegean Sea

F. SAKELLARIADOU

doi: [10.12681/mms.166](https://doi.org/10.12681/mms.166)

To cite this article:

SAKELLARIADOU, F. (2006). Spectroscopic studies of humic acids from subsurface sediment samples collected across the Aegean Sea. *Mediterranean Marine Science*, 7(2), 11–18. <https://doi.org/10.12681/mms.166>

Mediterranean Marine Science
Volume 7/2, 2006, 11-17

Spectroscopic studies of humic acids from subsurface sediment samples collected across the Aegean Sea

F. SAKELLARIADOU

Laboratory of Oceanography and Marine Geochemistry
Dept of Maritime Studies, University of Piraeus, Greece

e-mail: fsakellar@gmail.com

Abstract

Natural humic acids are biogenic, structurally complex and heterogeneous, refractory, acidic, yellow-to black-coloured organic polyelectrolytes of relatively high molecular weight. They occur in all soils, sediments, fresh waters, and seawaters. Humic acids represent the largest portion of nonliving soil organic matter. In the present paper, humic substances were isolated from marine subsurface sediment samples collected across the Aegean sea (in Greece) and especially from a marine area extending northwards of the Samothraki plateau towards the north-eastern part of the island of Crete. In a following step, humic preparations were studied using infrared and fluorescence spectroscopy (emission, excitation and synchronous-scan excitation spectra were obtained). The infrared spectra suggested functional chemical groups such as OH^- , C-H aliphatic, C=C, C=O/COO⁻, salts of carboxylic acids, and also, in some cases, silicate anions or C-O from alcohols, esters and ethers. Fluorescence emission, excitation and synchronous scan excitation provided some valuable information concerning a probable origin (marine and/or terrestrial) for the isolated humics.

Keywords: Fluorescence spectroscopy; Functional group; Humic substances; Infrared spectroscopy; Marine sediments.

Introduction

Natural humic substances comprise a wide class of naturally occurring, biogenic, structurally complex and heterogeneous, refractory, acidic, yellow-to black-coloured organic polyelectrolytes of relatively high molecular weight that occur in all soils, sediments, fresh waters, and seawater (THURMAN, 1986; SCHNITZER, 1978; STEVENSON, 1982). Humic acids are polycarboxylic acids with phenolic, alcoholic and carbonyl groups, aromatic rings and free radicals (SCHNITZER, 1978; SCHNITZER

& KHAN, 1972). They are very important constituents of sediments which perform in many functions, such as: (a) metal scavenging, concentration and transport (MANSKAYA & DROZDOVA, 1968); (b) growth regulation; (c) toxic material concentration and transport; (d) sediment formation (MANSKAYA & DROZDOVA, 1968).

Marine humates are of a rather dubious origin (RICE & MACCARTHY, 1991). Marine origins (NISSEMBAUM & KAPLAN, 1972; NISSEMBAUM, 1974), continental origins (YEN & TANG, 1977), or both (ROGERS & KOONS, 1968) have been suggested.

Humic substances can be studied using spectroscopic methods such as infrared and fluorescence spectroscopy. The most interesting portion of the infrared spectrum for the structural and analytical study of molecules is the medium IR region, between 4000 and 400 cm^{-1} . Fluorescence spectroscopy includes the three conventional modes of emission, excitation and synchronous scan. The former two fluorescence methods show limited applicability in resolving the spectra of complex polymers satisfactorily while the latter is characterized as a non-separative, non-destructive, highly sensitive technique, with limited sample preparation and as offering the opportunity to select various $\Delta\lambda$ that allow the spectral separation of overlapping fluorescence signals into individual components (MIANO & SENESI, 1992).

Materials and Methods

Humic substances were isolated from relatively enriched in organic carbon marine subsurface sediment samples from the Aegean sea (Table 1, Fig. 1). Humic substances were isolated following the pre-treatment step (freeze-drying), extraction step (removal of the alkaline soluble fraction), acidification step (acidification of the alkali-soluble extract to $\text{pH} \leq 2$ with HCl and precipitation), then centrifugation and finally purification of humic acids (THURMAN, 1988).

Humic substances were studied using (a) infrared spectroscopy with an IR spectrophotometer Perkin Elmer model 399 B on KBr pellets (1mg sample and 400mg KBr, spectrometry grade), and (b) fluorescence spectroscopy, using a Perkin Elmer LS-5 luminescence spectrophotometer on aqueous solutions of 100mg/l sample adjusted to pH8 with 0.05N NaOH. Emission spectra were recorded over the range 300-550nm at a fixed excitation wavelength of 360nm at pH7. Excitation spectra were obtained over a scan range of 270-500nm, by measuring the emission radiation at a fixed wavelength

Table 1
Burial depth, and corresponding water depth of each sediment sample.

Sample Number	Burial Depth (cm)	Water Depth (m)
1	4-7.5	182
2	336-339	182
3	40-45.5	995
4	40.5-44	690
5	45-54	720
6	226-228.5	365
7	16.5-19.5	340
8	243.5-252	340
9	37-39.5	368
10	177.5-183	368
11	24-31	415
12	196-206	415
13	104-110	568




Fig. 1: Map showing the sampling sites.

of 520nm at pH 7. Synchronous-scan excitation spectra were obtained by measuring the fluorescence intensity while simultaneously


scanning over both the excitation and emission wavelengths, and keeping a constant, optimized wavelength difference, $\Delta\lambda = \lambda_{em} - \lambda_{exc}$ (18nm) between them (SENESI, 1990).

Results and Discussion


Infrared Spectroscopy

According to STEVENSON & GOH (1982), MACCARTHY & RICE (1985), VANDENBROUCKE *et al.* (1985), SENESI *et al.* (1989), and SENESI (1992), the absorption bands could be interpreted as the following: (a) the 3400 cm^{-1} band is the one where a strong absorption due to hydrogen

bonding hydroxyl groups occurs; (b) the 2920 and 2860 cm^{-1} absorption bands are attributed to the asymmetric and symmetric stretching vibrations, respectively, of aliphatic C-H bonds in methyl and/or methylene units; (c) the band at 1710 cm^{-1} , which in terrestrial humic acids appears as a deep shoulder, is assigned to C=O stretch of carboxyl and carbonyl groups; (d) the 1600-1650 cm^{-1} band is assigned to aromatic C=C double bonds conjugated with C=O and/or COO; (e) the absorption at 1540 cm^{-1} is the amide band due to peptidic linkages; (f) the band at 1385 cm^{-1} is assigned at O-H deformation and C-O- stretch of phenols, COO- group vibrations, C-H deformation of CH_3 ; (g) the



Sample 3



Sample 5

Fig.2.: Selected I.R. spectra.

absorption around 1220 cm^{-1} is assigned to the C-O stretching vibration and OH bending deformations, due mainly to carboxyl groups; (h) an intense band around 1035 cm^{-1} is assigned to alcoholic and polysaccharide CO stretch and OH deformation and, eventually, Si-OH bend in silicate impurities. In the present study, the IR spectra recorded show that significant bands occur. The IR absorption characteristics resemble each other significantly while the relative intensities of specific bands vary.

Thus, through a thorough study of the IR spectra, it appears that all samples show characteristic peaks for the following: hydroxyl, methyl, methylene, aromatic bond, carbonyl, carboxyl, phenol, alcohol, polysaccharide and silicate impurities. In addition, samples 4 and 10 show the characteristic peak for amide. Comparing the recorded spectra of the samples 7, 9 and 11 with those of 8, 10 and 12 respectively, it is found that hydroxyls, aromatic bonds, and phenols decrease with burial depth increase. Selected IR spectra are shown in Figure 2.

Fluorescence spectroscopy

Marine humates are of a rather dubious origin (RICE & MACCARTHY, 1991). Debris of vascular plants, commonly found in coastal sediments, can be a significant source of sedimentary humic substances (ERTEL & HEDGES, 1985). Marine humates could be of a marine origin, on the basis of their genetic relation with degraded planktonic material (NISSEMBAUM & KAPLAN, 1972; Nissebaum, 1974) or of a continental one via lignin formation on the continents (YEN & TANG, 1977). Both origins might also be valid (ROGERS & KOONS, 1968). Interpretation of fluorescence emission, excitation, and synchronous-scan excitation spectra suggests the probable origin of the isolated humic acids (SENESI, 1990; BOTO & ISDALE, 1985; SAAR & WEBER, 1980; RYAN & WEBER, 1982). In humic polymers, the most efficient fluorophores are var-

Table 2
Possible origin for humic acids isolated.

Sample Number	Possible origin
1	marine
2	marine
3	terrestrial
4	terrestrial
5	terrestrial
6	marine
7	marine
8	terrestrial
9	marine
10	terrestrial
11	marine
12	marine
13	marine

iously substituted, condensed aromatic rings, and/or highly unsaturated aliphatic chains (MIANO *et al.*, 1998; SENESI *et al.*, 1991a; SEITZ, 1981). Data from the literature (SENESI, 1990; VISSER, 1983; MIANO *et al.*, 1988, SAAR & WEBER, 1980; RYAN & WEBER, 1982) indicate that: (i) as far as emission spectra are concerned, humic acids isolated from various terrestrial sources typically show a maximum-intensity wavelength varying between 435 and 465nm, while humic acids originating from marine samples generally show a maximum emission intensity which varies within a lower wavelength range, 410-450nm; (ii) as far as excitation spectra are concerned, the major excitation peak for terrestrial humic acids commonly occurs at a wavelength of about 390nm, with a number of secondary peaks or shoulders occurring on either side of the major peak, although at lower (360-345nm) or higher wavelengths (455-470nm), while humic acids of marine origin are characterized by one main excitation peak at variable wavelength (from 385 to 320nm); (iii) as far as synchronous-scan excitation spectra are concerned, the spectra of soil humic acids exhibit two major peaks whereas the marine humic acids give the most structured spectra. Thus, a




Fig. 3.: Selected fluorescence spectra (a: emission, b: excitation, c: synchronous scan).

thorough study of the recorded fluorescence spectra allows us to suggest that samples 1, 2, 6, 7, 9, 11, 12 and 13 are most probably of a marine origin while 3, 4, 5, 8 and 10 are of a terrestrial origin (Table 2). Selected fluorescence spectra are shown in Figure 3.

However, the study of any fluorescence spectrum does not allow the identification of individual structural components responsi-

ble for the macromolecule fluorescence of humic acids. Each spectrum is the sum of individual spectra of several fluorophores present in the complex and heterogeneous humic acid compound. The only way to be able to suggest structural components of a potential fluorescence contribution would be by applying various other techniques, such as degradation means, etc, to the humic acid

macromolecule methods, (SCHNITZER & KHAN, 1972; SENESI, 1990; SENESI *et al.*, 1991).

Conclusion

The spectroscopic methods that have been applied to the humic acids isolated from marine subsurface sediment samples from the Aegean Sea, allowed us to suggest the presence of various chemical units as well as the marine or terrestrial origin of each sample.

Thus, in all humic acid preparations there is a presence of hydroxyl, methyl, methylene, carbonyl, carboxyl, phenol, alcohol, polysaccharide groups, aromatic bond and silicate impurities. Occasionally, there is amide as well.

In two of the cores studied (samples 7, 8 and 9, 10) a different humic acid composition and origin was found between the upper and lower part. This composition differentiation could be attributed to the variation in oxygen content, particle size, sedimentation environment and relative productivity of terrestrial and marine environments (VANDENBROUCKE *et al.*, 1985). In a third core (samples 11 and 12), humic acids from the top and bottom part show variable composition, although they are of the same origin. This shows the impact on humic acid composition of the three first previously mentioned parameters.

References

- BOTO, K. & ISDALE, P., 1985. Fluorescent bands in massive corals result from terrestrial fulvic acid inputs to nearshore zone. *Nature*. London, 31: 396-397.
- ERTEL, J.R. & HEDGES, J.I., 1985. Sources of sedimentary humic substances: vascular plant debris. *Geochimica et Cosmochimica Acta*, 49: 2097-2107.
- MACCARTHY, P. & RICE, J.A., 1985. Spectroscopic methods (other than NMR) for determining functionalities in humic substances. p. 527-559. In: *Soil, Sediment and Water: Geochemistry, Isolation, and Characterization*. G.R. Ailen, D.M. McKnight, R.L. Wershaw, & P.E. MacCarthy (Eds), New York, Wiley-Interscience.
- MANSKAYA, S.M. & DROZDOVA, T.V., 1968. *Geochemistry of Organic Substances*, vol. 28, L. Shapiro & I.A. Breger (Eds). International Series of Monographs in Earth Sciences, N.Y., Pergamon, 345p.
- MIANO, T.M. & SENESI, N., 1992. Synchronous excitation fluorescence spectroscopy applied to soil humic substances' chemistry. p. 41-51. In: *The Science of the Total Environment*. Amsterdam, Elsevier Science Publisher B.V.
- MIANO, T.M., SPOSITO, M. & MARTIN, JR., 1998. Fluorescence spectroscopy of humic substances. *Soil Science Society of America Journal*, 52: 1016-1019.
- NISSEMBAUM, A., 1974. The organic geochemistry of marine and terrestrial humic substances: implication of carbon and hydrogen isotope studies. p. 39-52. In: *Advances in Organic Geochemistry*. B.Tissot & F. Biennier (Eds), Amsterdam, Elsevier.
- NISSEMBAUM, A., & KAPLAN, I.R., 1972. Chemical and isotopic evidence for the in situ origin of marine humic substances. *Limnology And Ocean*, 17(4): 570-582.
- RICE, J.A. & MACCARTHY, P., 1991. Composition of humin in stream Sediments and peat. p. 35-46 In: *Organic Substances and Sediments in Water, Part I: Humic and Other Substances*. R. A. Baker (Ed.), Michigan, Lewis Publishers.
- ROGERS, M.A. & KOONS, C.B., 1968. Glacial-eustatic control of organic carbon delta C¹³ values from post-Pliocene through recent-marine sequences in the Gulf of Mexico. *Geol. Soc. Amer. Ann. Mtg.*, Mexico City.

- RYAN, D.K. & WEBER, J.H., 1982. Copper (II) Complexing capacities of natural waters by fluorescence quenching. *Environmental Science and Technology*, 16: 866-872.
- SAAR, R.A. & WEBER J.H., 1980. Comparison of spectrofluorometry and ion-selective electrode potentiometry for determination of complexes between fulvic acid and heavy-metal ions. *Analytical Chemistry*, 52: 2095-2100.
- SCHNITZER, M., 1978. Humic substances: chemistry and reactions In: *Soil Organic Matter*. p. 1-64. M. Schnitzer & S.U. Khan (Eds), Amsterdam, Elsevier.
- SCHNITZER, M. & KHAN, S.U., 1972. Humic substances in the environment. Marcel Dekker, New York, 327p.
- SEITZ, W.R., 1981. Fluorescence methods for studying speciation of pollutants in water. *Trends in Analytical Chemistry*, 1: 79-83.
- SENESI, N., 1990. Molecular and quantitative aspects of the chemistry of fulvic acid and its interactions with metal ions and organic chemicals. Part II. The fluorescence spectroscopy approach. *Analytica Chimica Acta*, 232: 77-106.
- SENESI, M., 1992. Metal-humic substance complexes in the environment. molecular and mechanistic aspects by multiple spectroscopic approach. p. 429-496. In: *Biogeochemistry of Trace Metals*. D.C. Adriano (Ed.), Boca Raton, Lewis Publishers.
- SENESI, N., MIANO, T.M. & PROVENZANO, M.R., 1991. Fluorescence spectroscopy as a means of distinguishing fulvic and humic acids from aquatic, dissolved and sedimentary, and terrestrial sources. *Lecture notes in Earth Sciences*, p. 63-73, Berlin, Springer-Verlag.
- SENESI, N., MIANO, T.M., PROVENZANO, M.R., & BRUNETTI, G., 1989. Spectroscopic and compositional comparative of I.H.S.S. Reference and standard fulvic and humic acids of various origin. *The Science of the Total Environment*, 81/82: 143-156.
- SENESI, N., MIANO, T.M., PROVENZANO, M.R., & BRUNETTI, G., 1991. Characterization, differentiation, and classification of humic substances by fluorescence spectroscopy. *Soil Science*, 152/4: 259-271.
- STEVENSON, F.J., 1982. Humus chemistry. Genesis, composition, reactions. Wiley-Interscience, New York, 443p.
- STEVENSON, F.J. & GOH, K.M., 1982. Infra-red spectra of humic acids and related substances. *Geochim. Cosochic. Acta*, 35: 471-483.
- THURMAN, E.M., 1986. Organic geochemistry of natural waters. Dordrecht, Netherlands, Martinus Nijhoff, 497p.
- THURMAN, E.M., 1988. Isolation of soil and aquatic humic substances. Group Report. p. 31-43 In: *Humic Substances and Their Role in the Environment*. F.H. Frimmel & F. Christman (Eds), Chichester, Wiley-Interscience.
- VANDENBROUCKE, M. PELET, R. & DEBYSER, Y., 1985. Geochemistry of humic substances in marine sediments. p. 250-273 In: *Aquatic Humic Substances: Influences on Fate and Treatment of Pollutants*. I.M. Suffet & P. MacCarthy (Eds), Washington, D.C., American Chemical Society.
- YEN, T.F. & TANG, J.I.S., 1977. Chemical aspects of marine sediments. p. 1-38. In: *Chemistry of Marine Sediment*. T.F. Yen (Ed) T.F., Ann Arbor Science.

Accepted in June 2007

