

Mediterranean Marine Science

Vol 6, No 2 (2005)



Alien species on the coasts of Turkey

M.E. CINAR, M. BILECENOGLU, B. OZTURK, T. KATAGAN, V. AYSEL

doi: [10.12681/mms.187](https://doi.org/10.12681/mms.187)

To cite this article:

CINAR, M., BILECENOGLU, M., OZTURK, B., KATAGAN, T., & AYSEL, V. (2005). Alien species on the coasts of Turkey. *Mediterranean Marine Science*, 6(2), 119–146. <https://doi.org/10.12681/mms.187>

Alien species on the coasts of Turkey

M. E. ÇINAR¹, M. BILECENOĞLU², B. ÖZTÜRK¹, T. KATAGAN¹ and V. AYSEL³

¹Ege University, Faculty of Fisheries, Department of Hydrobiology, 35100 Bornova, Izmir, Turkey

²Adnan Menderes University, Faculty of Arts & Sciences, Department of Biology, 09010 Aydin, Turkey

³Onsekiz Mart University, Faculty of Arts & Sciences, Department of Biology, Çanakkale, Turkey

e-mail: melih.cinar@ege.edu.tr

Abstract

The compilation of data on alien species reported from the Turkish coasts yielded a total of 263 species belonging to 11 systematic groups, of which Mollusca had the highest number of species (85 species), followed by Crustacea (51), fishes (43) and phytobenthos (39). The Black Sea is represented by a total of 20 alien species, the Sea of Marmara by 48 species, the Aegean Sea by 98 species and the Levantine Sea by 202 species. The majority of aliens found in the Black Sea and the Sea of Marmara were transported via shipping, whereas the Levantine coast is extensively subjected to Lessepsian migration. Benthic habitats (soft and hard substrata) comprise 76% of the total alien species and the pelagic environment is inhabited by thirty-nine species. Almost 50% of aliens collected from the Turkish coasts were found only at 0-10 m depth. Eight species occur at depths deeper than 100 m. The impacts of aliens on the benthic and pelagic ecosystems are presented.

Keywords: Alien species; Species list; Impact; Black Sea; Sea of Marmara; Aegean Sea; Levantine Sea; Turkey.

Introduction

Species introduction is one of the major factors adversely affecting biological diversity (ELTON, 1958). The impacts of alien species on their new environment include restructuring established food webs, importing new diseases and competition with native organisms for food and space. Other significant ecological changes may occur when the invading organisms reproduce with native species, altering the gene pool (OCCHIPINTI AMBROGI, 2001). Invaders may belong to any taxonomical group, and are

capable of colonizing every ecosystem on earth, changing the ecological relations within communities, altering evolutionary processes and causing dramatic changes in native populations, including extinctions (MACK *et al.*, 2000).

The negative effects of aliens experienced world-wide have triggered the social and scientific media to take some precautions against the dispersal of aliens among regions or among localities within a particular region. Thus, compiled data regarding all aliens or a target organism (i.e. *Caulerpa taxifolia* in the Mediterranean Sea) are urgently required by scientists

and decision makers. In addition, monitoring programmes on spatio-temporal structures of communities particularly in the hot spot areas for aliens such as harbours, brackish and polluted waters should be undertaken both on local and global scales.

Turkey is surrounded by four seas with different hydrographical characteristics. The maritime traffic taking place through the Dardanel and Bosphorus straits and among commercial harbours makes the Turkish coasts more susceptible to invasions by aliens. The proximity of Turkey to the Suez Canal has resulted in dense settlements of Lessepsian migrants, especially in habitats along the Levantine coast of Turkey. Alien species and their roles in the benthic and pelagic ecosystems is increasingly becoming a subject of study in the country.

This paper reviews the alien species reported from the Turkish coasts and constitutes the first comprehensive database for future studies.

Methods

Data on alien species along the Turkish coasts

In this study, only reported data on aliens along the Turkish coasts have been taken into account. All calculations are based on species' records up to the end of 2005. Cryptogenic species have not been included in our compiled list. Some of the species in Table 1, which represent uncertainty in their real taxonomic position and distributional pattern, are classified as questionable or excluded. The questionable species will turn into aliens if their real taxonomic and distributional identities are clarified. In all calculations, including number of species per sea, only the established and questionable species have been taken into account. The importance of aliens in the total biota was estimated, based on the authors' databases.

Results and Discussion

List of alien species from the Turkish coasts

Table 1 includes all alien species reported from the Turkish coasts. The species are categorized by their origins, the mode of introduction and their habitat and depth preferences. The establishment success is assigned for each species. The first reported years of the species together with the relevant publications are also given for each sea.

The present data show that a total of 263 alien species, which belong to 11 systematic groups, occur along the Turkish coasts. Among the groups, Mollusca had the highest number of species (85 species), followed by Crustacea (51), fishes (43) and phytobenthos (39). Cnidaria is represented by only two species (*Rhopilema nomadica* and *Cassiopea andromeda*), Ctenophora by 2 species (*Mnemiopsis leidyi* and *Beroe ovata*), Pantopoda by one species (*Anoplodactylus californicus*), Bryozoa by one species (*Rhynchozoon larreyi*) and Echinodermata by 2 species (*Ophiactis savignyi* and *Synaptula reciprocans*).

Thirty-two species are considered as questionable or excluded in Table 1, mainly due to the uncertainty in their real taxonomical status and distributional patterns. The polychaetes, *Branchiosyllis exilis*, *Opisthosyllis brunnea*, *Rhodine loveni* and *Monticellina dorsobranchialis*, which were previously considered as Lessepsian migrants, indeed occur also in the western Mediterranean and the Atlantic Ocean. These species were also previously proposed to be excluded from the list of Lessepsian species (ERGEN et al., 2002; ÇINAR, 2003; 2005). RULLIER (1963) found 11 polychaete species with Indo-Pacific affinity from the Sea of Marmara (Table 1). Of the species, *Nereis zonata persica* and *Timarete aencyclochaeta* were also reported along the Levantine coast and considered as Lessepsian migrants (BEN-ELIAH, 1995). The other species have not been subsequently reported from elsewhere in the Mediterranean Sea.

A total of 176 Lessepsian and 48 ship-transferred species were reported along the Turkish

Table 1.

The list of alien species and their first year of reports from the Turkish coasts. The habitat and depth preferences of aliens along the coasts together with their origins and establishment success are also given. The marked species in the list are questionable or excluded ones. BS: Black Sea, SM: Sea of Marmara, AS: Aegean Sea, LS: Levantine Sea, O: Origin (IP: Indopacific, RS: Red Sea, AT: Atlantic, WA: Western Atlantic, ST: Subtropical, IO: Indian Ocean, PG: Persian Gulf, PO: Pacific Ocean, TA: Tropical Atlantic, CT: Circumtropical, BA: Boreal Atlantic) Es: Establishment success (E: Established, Q: Questionable, Ex: Excluded), MI= Mode of Introduction (L: Lessepsian, S: Shipping, G: Gibraltar, Aq: Aquaculture), H: Habitat [Hs: hard substrata (including algae, sponge), Ss: soft substrata (including phanerogams), P: pelagic, Pz: parasite], D: Depth range (I: 0-10 m, II: 11-50m, III: 51-100 m, IV: 101-200 m, V: 201-400, VI: 401-500 m).

| | BS | SM | AS | LS | Es | O | MI | H | D |
|--|---------------------|---------------------|---------------------|---------------------|----|--------|----|-------|------|
| PHYTOPLANKTON | | | | | | | | | |
| <i>Alexandrium tamarensis</i> (Lebour) Balech 1992 | - | - | 1984 ¹²⁴ | - | Q | WA | S | P | I |
| <i>Chaetoceros coarctatus</i> Lauder 1864 | - | - | 1987 ¹²⁵ | - | E | AT,PO | S | P | I |
| <i>Gymnodinium cf. mikimotoi</i> Miyake et Kominami ex Oda 1935 | - | - | 2002 ⁴⁴ | - | Q | ? | S | P | I |
| <i>Heterosigma cf. akashiwo</i> (Hada) Hada, 1968 | - | - | 2002 ⁴⁴ | - | Q | ? | S | P | I |
| <i>Rhizosolenia calcar-avis</i> M. Schultze, 1858 | 1999 ⁷¹ | 1993 ¹²³ | - | - | E | AT | S | P | I |
| <i>Scrippsiella trochoidea</i> (Stein) Loeblich III, 1976 | 1999 ⁷¹ | - | - | - | E | AT | S | P | I |
| <i>Thalassiosira nordenskioeldii</i> Cleve 1873 | 2002 ¹⁶⁰ | - | - | - | E | AT | S | P | I |
| PHYTOBENTHOS | | | | | | | | | |
| Cyanophyta (=Cyanobacteria) | | | | | | | | | |
| <i>Trichodesmium erythraeum</i> Ehrenberg, 1830 | - | - | 1996 ⁶⁹ | - | E | IP,RS | L | Hs,Ss | I |
| Rhodophyta | | | | | | | | | |
| <i>Acanthophora nayadiformis</i> (Delile) Papenfuss, 1968 | - | 1973 ¹⁷⁷ | 1970 ¹⁰³ | 1997 ²⁰ | E | RS | L | Hs | I,II |
| <i>Acanthophora muscoides</i> Linnaeus, 1753 | - | - | 1986 ¹⁷⁸ | - | E | AT | S | Hs | I,II |
| <i>Acrochaetium codicolum</i> Børgesen, 1927 | 1996 ²⁸ | 1986 ¹⁷⁸ | 1990 ⁵⁵ | 1997 ¹⁹ | E | IP,AT | S | Hs,Ss | I |
| <i>Asparagopsis armata</i> Harvey, 1855 | 1973 ¹⁷⁶ | 1986 ¹⁷⁸ | 1973 ¹⁷⁶ | 1969 ¹⁷⁴ | E | IP,AT | S | Hs | I |
| <i>Asparagopsis taxiformis</i> (Delile) Trevisan de Saint-Léon, 1845 | - | - | 2001 ²⁷ | - | E | CT | L | Hs | I |
| <i>Bonnemaisonia hamifera</i> Hariot, 1891 | - | 1986 ¹⁷⁸ | - | 1997 ²⁰ | E | PO,AT | ?S | Hs | I |
| <i>Botryocladia madagascariensis</i> G.Feldmann, 1945 | - | - | - | 2000 ¹⁶¹ | E | IO | ?S | Hs | I |
| <i>Chondria collinsiana</i> Howe, 1920 | - | 1986 ¹⁷⁸ | - | - | E | AT, IO | ?L | Hs | II |
| <i>Chondrophycus papillosum</i> (C. Agardh) Garbary & Harper 1998 | 1973 ¹⁷⁵ | 1957 ¹⁴³ | 1969 ¹⁷⁴ | 1969 ¹⁷⁴ | E | RS | ?L | Hs | I |
| <i>Ganonema farinosum</i> (Lamouroux) Fan & Wang, 1974 | 1995 ²¹ | 1899 ⁹¹ | 1986 ²⁶ | 1997 ¹⁹ | Q | IP,RS | ?L | Hs,Ss | I |

(continued)

Table 1 (continued)

| | BS | SM | AS | LS | Es | O | MI | H | D |
|---|---------------------|---------------------|---------------------|---------------------|----|--------|----|-------|------|
| <i>Gracilaria arcuata</i> Zanardini 1858 | - | 1986 ¹⁷⁸ | - | - | E | RS,IP | L | Hs | I |
| <i>Griffithsia corallinooides</i> (Linnaeus) Trevisan, 1845 | - | 1993 ²³ | - | - | E | AT, IP | G | Hs | I |
| <i>Hypnea spinella</i> (C. Agardh) Kützing, 1847 | - | - | 1987 ¹⁸ | - | E | AT,IP | S | Hs | I |
| <i>Hypnea variabilis</i> Okamura, 1909 | - | 1986 ¹⁷⁸ | 1986 ¹⁷⁸ | - | E | PO | S | Hs | I |
| <i>Laurencia intermedia</i> Yamada, 1931 | 1986 ²⁶ | - | - | - | E | PO,RS | S | Hs | I,II |
| <i>Lophocladia lallemandii</i> (Montagne) Schmitz, 1893 | - | - | 1970 ¹⁰³ | 1986 ⁵³ | E | IP | L | Hs | I,II |
| <i>Polysiphonia fucoides</i> (Hudson) Greville, 1824 | 1973 ¹⁷⁶ | - | 1973 ¹⁷⁶ | - | E | AT, IO | S | Hs | I,II |
| <i>Polysiphonia paniculata</i> Montagne, 1842 | 1986 ²⁶ | - | - | - | E | AT | S | Hs | I |
| <i>Polysiphonia kampsaxii</i> Boergesen, 1939 | - | - | 1984 ¹⁷ | 1997 ²⁰ | E | IP | S | Hs | I |
| <i>Radicilingua thysanorhizans</i> (Holmes) Papenfuss, 1956 | - | 1986 ¹⁷⁸ | 1986 ²⁶ | 1986 ⁵³ | E | AT | S | Hs | I |
| <i>Rhodophysema georgii</i> Batters, 1900 | - | 1986 ¹⁷⁸ | - | - | E | AT,PO | ?S | Hs | I |
| <i>Spyridia hypnoides</i> (Bory de Saint-Vincent) Papenfuss, 1968 | - | - | 1987 ¹⁸ | 1997 ¹⁹ | E | IO | S | Hs | I |
| Heterokontophyta | | | | | | | | | |
| <i>Chorda filum</i> (Linnaeus) Stackhouse 1797 | - | 1986 ¹⁷⁸ | - | - | ? | AT,PO | ?S | Hs | I,II |
| <i>Ectocarpus siliculosus</i> (Dillwyn) Lyngbye, 1819 | 1973 ¹⁷⁵ | 1899 ⁹¹ | 1899 ⁹¹ | 1986 ¹⁴⁶ | E | AT | S | Hs | I |
| <i>Halothrix lumbricalis</i> (Kützing) Reinke, 1888 | 2004 ²⁹ | 1993 ²³ | 1986 ¹⁷⁸ | 1993 ¹⁴⁵ | E | AT,PO | S | Hs | I |
| <i>Pilayella littoralis</i> (Linnaeus) Kjellman, 1872 | 1998 ²⁴ | 1993 ²³ | 1985 ¹⁰⁵ | 1985 ¹⁰⁵ | E | AT | S | Hs | I |
| <i>Protectocarpus speciosus</i> Boergesen, 1902 | - | 1993 ²³ | - | - | E | AT | S | Hs | I |
| <i>Sargassum latifolium</i> (Turner) C. Agardh, 1820 | - | 1986 ¹⁷⁸ | 1986 ¹⁷⁸ | - | E | RS | ?L | Hs | I,II |
| <i>Sphaerotrichia divaricata</i> (Agardh) Kylin, 1940 | - | 1986 ¹⁷⁸ | 1970 ¹⁰³ | 1986 ¹⁷⁸ | E | AT | S | Hs | I |
| <i>Styropodium schimperi</i> (Buchinger ex Kützing) Verlaque & Boudouresque, 1991 | - | - | - | 1997 ²⁰ | E | RS,AT | ?L | Hs | I |
| Chlorophyta | | | | | | | | | |
| <i>Bryopsis pennata</i> Lamouroux, 1809 | - | 1986 ²⁶ | 1978 ⁵² | 1978 ⁵² | E | AT | S | Hs | I |
| <i>Caulerpa racemosa</i> (Forsskål) J. Agardh, 1873 | - | - | 1991 ⁵⁴ | 1986 ⁵³ | E | CT | ?L | Hs,Ss | I,II |
| <i>Caulerpa scalpelliformis</i> (R.Brown ex Turner) C. Agardh, 1817 | - | - | - | 1998 ⁸⁶ | E | AT,IP | ?L | Ss | I,II |
| <i>Cladophora boodleoides</i> Børgesen 1925 | - | - | 1996 ⁷⁰ | - | E | AT | G | Hs | I |

(continued)

Table 1 (continued)

| | BS | SM | AS | LS | Es | O | MI | H | D |
|--|---------------------|---------------------|---------------------|---------------------|----|-------|----|-------|--------|
| <i>Codium fragile</i> (Suringar) Hariot ssp. <i>tomentosoides</i> (Goor) P.C. Silva, 1955 | - | - | 1986 ¹⁷⁸ | - | E | AT,PO | ?S | Hs | I |
| <i>Ulva fasciata</i> Delile, 1813 | 1990 ²⁵ | 1986 ²⁶ | 1977 ¹⁰⁴ | 1996 ²² | E | ?RS | ?L | Hs | I |
| Phanerogame | | | | | | | | | |
| <i>Halophila stipulacea</i> (Forsskål) Ascherson | - | - | 1985 ¹⁰⁵ | 1985 ¹⁰⁵ | E | IO,RS | L | Hs | I |
| FORAMINIFERA (Benthic) | | | | | | | | | |
| <i>Astacolus insolitus</i> (Schwager, 1866) | - | - | 2004 ¹³² | - | E | IP,RS | L | Ss | III |
| <i>Astacolus sublegumen</i> (Parr, 1950) | - | - | 2004 ¹³² | - | E | IP,RS | L | Ss | III |
| <i>Planogypsina acervalis</i> (Brady, 1884) | - | - | 2004 ¹³² | - | E | IP,RS | L | Ss | II |
| <i>Planogypsina squamiformis</i> (Chapman, 1901) | - | - | 2004 ¹³² | - | E | IP,RS | L | Ss | II,III |
| <i>Amphistegina lobifera</i> Larsen, 1976 | - | - | 2004 ¹³² | 2002 ¹³³ | E | IP,RS | L | Ss | I,II |
| CNIDARIA | | | | | | | | | |
| <i>Cassiopea andromeda</i> (Forsskål, 1775) | - | - | - | 2002 ³⁹ | E | IP,RS | L | Ss | I |
| <i>Rhopilema nomadica</i> Galil, Spanier & Ferguson, 1990 | - | - | - | 1995 ¹¹³ | E | IP,RS | L | P | ? |
| CTENOPHORA | | | | | | | | | |
| <i>Mnemiopsis leidyi</i> (Agassiz, 1865) | 1994 ¹³⁷ | 1994 ¹¹⁴ | 1994 ¹¹⁴ | 1993 ¹⁶⁷ | E | WA | S | P | I,II |
| <i>Beroe ovata</i> Mayer 1912 | 2001 ¹¹⁵ | 2004 ¹⁰⁸ | - | - | E | WA | S | P | I,II |
| POLYCHAETA | | | | | | | | | |
| <i>Lepidonotus carinulatus</i> (Grube, 1870) | - | 1963 ¹⁴⁹ | - | - | Q | IP,RS | ?S | ? | ? |
| <i>Harmothoe boholensis</i> (Grube, 1878) | - | 1963 ¹⁴⁹ | - | - | Q | IP,RS | ?S | ? | ? |
| <i>Harmothoe minuta</i> (Potts, 1910) | - | 1963 ¹⁴⁹ | - | - | Q | IP,RS | ?S | ? | ? |
| <i>Ancistrosyllis rigida</i> Fauvel, 1919 | - | 1963 ¹⁴⁹ | - | - | Q | IP,RS | ?S | ? | ? |
| <i>Sigambra constricta</i> (Southern, 1921) | - | 1963 ¹⁴⁹ | - | - | Q | IP,RS | ?S | ? | ? |
| <i>Opisthosyllis brunnea</i> Langerhans, 1879 | - | - | 2002 ⁶³ | - | Ex | IP,RS | L | Hs | I |
| <i>Branchiosyllis exilis</i> (Gravier, 1900) | - | - | 2002 ⁶³ | - | Ex | IP,RS | L | Hs | I |
| <i>Leonnates persicus</i> Wesenberg-Lund, 1949 | - | - | 2002 ⁶⁵ | 2003 ⁸⁵ | E | IP,RS | L | Ss | II-IV |
| <i>Nereis zonata persica</i> Fauvel, 1911 | - | 1963 ¹⁴⁹ | - | - | Q | IP,RS | ?L | ? | ? |
| <i>Pseudonereis anomala</i> Gravier, 1900 | - | - | 2005 ⁶⁴ | 1989 ³³ | E | IP,RS | L | Hs | I |
| <i>Glycera alba adspersa</i> Fauvel, 1939 | - | 1963 ¹⁴⁹ | - | - | Ex | IP | ?S | ? | ? |
| <i>Eurythoe complanata</i> (Pallas, 1766) | - | - | - | 1997 ⁸¹ | Q | ?IP | L | Hs | I |
| <i>Linopherus acarunculata</i> Monro, 1937 | - | - | - | 1997 ⁸¹ | E | RS | L | Ss | I |
| <i>Lumbrineris debilis</i> Grube, 1878 | - | 1963 ¹⁴⁹ | - | - | Q | IP | ?S | ? | ? |
| <i>Lysidice collaris</i> Grube, 1870 | - | - | 1998 ⁶¹ | 1997 ⁸¹ | Q | IP,RS | L | Hs,Ss | I,II |
| <i>Prionospio saccifera</i> Mackie & Hartley, 1990 | - | - | | 1999 ⁶² | E | IP,RS | L | Ss | III |
| <i>Polydora cornuta</i> Bosc, 1802 | - | - | 2005 ⁶⁶ | - | E | ? | S | Ss | I,II |
| <i>Streblospio gynobranchiata</i> Rice & Levin, 1998 | - | - | 2005 ⁶⁶ | - | E | WA | S | Ss | I,II |
| <i>Dasybranchus carneus</i> Grube, 1870 | - | 1963 ¹⁴⁹ | - | - | Q | RS | ?S | ? | ? |
| <i>Notomastus aberans</i> Day, 1957 | - | - | 1983 ¹⁴¹ | - | E | IP | L | Ss | I |

(continued)

Table 1 (continued)

| | BS | SM | AS | LS | Es | O | MI | H | D |
|---|----|---------------------|---------------------|---------------------|----|-------|----|----|-------|
| <i>Monticellina dorsobranchialis</i> (Kirkegaard, 1959) | - | - | 2006 ⁸⁴ | - | Ex | ? | L | Ss | ? |
| <i>Timarete dasylophius</i> (Marenzeller, 1879) | - | 1963 ¹⁴⁹ | - | - | Q | IP | ?S | ? | ? |
| <i>Timarete anchylochaeta</i> (Schmarda, 1861) | - | 1963 ¹⁴⁹ | - | - | Q | IP | ?S | ? | ? |
| <i>Metasychis gotoi</i> (Izuka, 1902) | - | - | 2002 ⁸³ | - | E | IP,RS | ?L | Ss | ? |
| <i>Rhodine loveni</i> Malmgren, 1865 | - | - | 1992 ⁸⁰ | 1998 ⁸² | Ex | IP,AT | L | Ss | II |
| <i>Pista unibranchia</i> Day, 1963 | - | - | 1998 ¹⁴² | 1997 ⁸¹ | E | IP,RS | L | Ss | I,II |
| <i>Ficopomatus enigmaticus</i> (Fauvel, 1923) | - | 1952 ⁶⁷ | 1976 ⁷⁹ | - | E | ST | S | Hs | I |
| <i>Hydroides dianthus</i> (Verrill, 1873) | - | - | 1865 ¹⁴⁸ | - | E | WA | S | Hs | I |
| <i>Hydroides elegans</i> (Haswell, 1883) | - | - | 1976 ⁷⁹ | 1991 ³⁴ | E | A,P | S | Hs | I |
| <i>Spirorbis marioni</i> Caullery & Mesnil, 1897 | - | - | 1991 ¹¹⁷ | - | E | P | S | Hs | I |
| PANTOPODA | | | | | | | | | |
| <i>Anoplodactylus californicus</i> Hall, 1912 | - | - | - | 1962 ¹⁵³ | E | CT | L | Hs | I |
| CRUSTACEA | | | | | | | | | |
| Copepoda | | | | | | | | | |
| <i>Centropages furcatus</i> (Dana, 1846) | - | 2000 ¹⁶⁴ | - | 2002 ¹⁶⁸ | E | IP,RS | L | P | I-III |
| <i>Calanopia elliptica</i> (Dana, 1846) | - | - | - | 2003 ¹⁵⁷ | E | IP,RS | L | P | I |
| <i>Calanopia biloba</i> Bowman, 1957 | - | - | - | 2002 ¹⁶⁸ | E | IP,RS | L | P | I-III |
| <i>Calanopia minor</i> Scott, 1902 | - | - | - | 2002 ¹⁶⁸ | E | IP,RS | L | P | I-III |
| <i>Labidocera pavo</i> Giesbrecht, 1889 | - | - | - | 2003 ¹⁵⁷ | E | IP,RS | L | P | I |
| <i>Euchaeta concinna</i> Dana, 1846 | - | - | - | 2002 ¹⁶⁸ | E | IP,RS | L | P | I-III |
| <i>Eucalanus crassus</i> Giesbrecht, 1888 | - | - | - | 2002 ¹⁶⁸ | E | IP,RS | L | P | I-III |
| <i>Eucalanus subcrassus</i> Giesbrecht, 1888 | - | - | - | 2002 ¹⁶⁸ | E | IP,RS | L | P | I-III |
| <i>Parocalanus latus</i> Andronov, 1972 | - | 2000 ¹⁶⁴ | - | 2002 ¹⁶⁸ | E | IP,RS | L | P | I-III |
| <i>Parocalanus elegans</i> Andronov, 1972 | - | 2000 ¹⁶⁴ | - | 2002 ¹⁶⁸ | E | IP,RS | L | P | I-III |
| <i>Acartia tonsa</i> Dana, 1848 | - | 2000 ¹⁶⁴ | 2001 ¹⁵⁵ | - | E | WA | S | P | ? |
| <i>Acartia hasanii</i> Unal, Shmeleva & Kideys, 2002 | - | - | - | 2002 ¹⁶³ | Q | ?RS | ?L | P | I |
| <i>Paracartia ioannae</i> Unal, Shmeleva & Kideys, 2002 | - | - | - | 2002 ¹⁶³ | Q | ?RS | ?L | P | I |
| <i>Paracartia janetae</i> Unal, Shmeleva & Kideys, 2002 | - | - | - | 2002 ¹⁶³ | Q | ?RS | ?L | P | I |
| Cirripedia | | | | | | | | | |
| <i>Heterosaccus dolffusi</i> Boschma, 1960 | - | - | - | 1997 ¹⁴⁰ | E | RS | L | Pz | I,II |
| Amphipoda | | | | | | | | | |
| <i>Maera hamigera</i> Haswell, 1879 | - | - | 1978 ¹²¹ | 1978 ¹²¹ | E | IP | L | Ss | I,II |
| <i>Stenothoe gallensis</i> Walker, 1904 | - | - | 1978 ¹²¹ | 1978 ¹²¹ | E | IP | L | Ss | I,II |
| Isopoda | | | | | | | | | |
| <i>Sphaeroma walkeri</i> (Stebbing, 1905) | - | - | 1976 ¹¹⁸ | - | E | IP | L | Hs | I,II |

(continued)

Table 1 (continued)

| | BS | SM | AS | LS | Es | O | MI | H | D |
|---|----|---------------------|---------------------|---------------------|----|-------|----|-------|--------|
| Tanaidacea | | | | | | | | | |
| <i>Apseudes intermedius</i> Hansen, 1985 | - | - | 1978 ¹¹⁹ | - | E | AT | ?S | Hs | I,II |
| Cumacea | | | | | | | | | |
| <i>Eocuma sarsi</i> (Kossmann, 1880) | - | - | 1983 ¹⁰⁹ | - | E | IP | L | Ss | I,II |
| Decapoda | | | | | | | | | |
| <i>Marsupenaeus japonicus</i> (Bate, 1888) | - | 2001 ¹⁷³ | 2001 ¹⁷³ | 1930 ¹³⁶ | E | IP | L | Ss | I-III |
| <i>Melicertus hathom</i> (Burkenroad, 1959) | - | - | - | 2002 ¹²⁸ | E | RS,IO | L | Ss | I-III |
| <i>Metapenaeopsis aegyptia</i> Galil & Golani, 1990 | - | - | - | 2004 ¹⁷⁰ | E | IP,RS | L | Ss | I-III |
| <i>Metapenaeopsis mogiensis consobrina</i> (Nobili, 1904) | - | - | - | 2004 ¹⁷⁰ | E | IP,RS | L | Ss | I-III |
| <i>Metapenaeus monoceros</i> (Fabricius, 1798) | - | - | - | 1961 ¹⁰⁶ | E | IP,RS | L | Ss | I-III |
| <i>Penaeus semisulcatus</i> de Haan, 1844 | - | - | - | 1930 ¹⁰¹ | E | IP,RS | L | Ss | I-IV |
| <i>Metapenaeus stebbingi</i> (Nobili, 1904) | - | - | - | 1981 ¹²⁰ | E | RS,IO | L | Ss | I-III |
| <i>Trachysalambria palaestinensis</i> Steinitz, 1932 | - | - | - | 1968 ⁹⁵ | E | RS | L | Ss | II,III |
| <i>Leptocheila pugnax</i> de Man, 1916 | - | - | - | 1981 ¹²⁰ | E | IP,RS | L | Ss | I-IV |
| <i>Palaemonella rotumana</i> (Borradaile, 1898) | - | - | - | 2002 ⁹² | E | IP,RS | L | Ss | I-IV |
| <i>Alpheus audouini</i> (Coutière, 1905) | - | - | - | 2002 ⁹² | E | IP,RS | L | Hs | I,II |
| <i>Alpheus inopinatus</i> Holthuis & Gottlieb, 1958 | - | - | - | 1969 ⁹³ | E | IO,RS | L | Hs,Ss | I,II |
| <i>Alpheus migrans</i> Lewinsohn & Holthuis, 1978 | - | - | - | 1994 ¹²² | E | RS | L | Ss | II,III |
| <i>Alpheus rapacida</i> de Man, 1908 | - | - | - | 1981 ¹²⁰ | E | IP | L | Ss | I-III |
| <i>Urocaridella antonbrunii</i> (Bruce, 1967) | - | - | - | 2004 ¹⁷⁰ | Ex | IP | L | Ss | I,II |
| <i>Ixa monodi</i> Holthuis & Gottlieb, 1956 | - | - | - | 1956 ¹⁰⁷ | E | RS | L | Ss | II,III |
| <i>Leucosia signata</i> Paulson, 1875 | - | - | - | 1982 ⁹⁹ | E | IP | L | Ss | I,II |
| <i>Myra subgranulata</i> Kossmann, 1877 | - | - | - | 1930 ¹³⁶ | E | IP,RS | L | Ss | I-IV |
| <i>Micippa thalia</i> (Herbst, 1803) | - | - | - | 1995 ⁷⁴ | E | IP | L | Hs,Ss | I-III |
| <i>Callinectes sapidus</i> Rathbun, 1896 | - | - | 2001 ¹⁷³ | 1961 ¹⁰⁶ | E | WA | S | Ss | I-III |
| <i>Charybdis hellerii</i> (Milne Edwards, 1867) | - | - | - | 1981 ¹²⁰ | E | IP,RS | L | Hs | I,II |
| <i>Charybdis longicollis</i> Leene, 1938 | - | - | - | 1961 ¹⁰⁶ | E | IP,RS | L | Ss | I-III |
| <i>Portunus pelagicus</i> (Linnaeus, 1758) | - | - | - | 1928 ¹⁰⁰ | E | IP,RS | L | Ss | I-III |
| <i>Thalamita poissonii</i> (Audouin, 1826) | - | - | - | 1961 ¹⁰⁶ | E | IP,RS | L | Hs,Ss | I-III |
| <i>Carupa tenuipes</i> Dana, 1851 | - | - | - | 2004 ¹⁷⁰ | ? | IP,RS | L | Hs | I-III |
| <i>Pilumnopeus vauquelini</i> (Audouin, 1826) | - | - | - | 1981 ¹²⁰ | E | IP,RS | L | Hs | I |
| <i>Calappa hepatica</i> (Linnaeus, 1758) | - | - | - | 2003 ³⁰ | E | IP,RS | L | Ss | I-III |
| <i>Atergatis roseus</i> (Rüppell, 1830) | - | - | - | 1990 ⁷⁶ | E | IP,RS | L | Hs | I,II |

(continued)

Table 1 (continued)

| | BS | SM | AS | LS | Es | O | MI | H | D |
|--|--------------------|---------------------|---------------------|---------------------|----|-------|----|-------|-------|
| <i>Daira perlata</i> (Herbst, 1790) | - | - | - | 1995 ⁷⁴ | ? | IP | L | Hs,Ss | I |
| <i>Eucrate crenata</i> de Haan, 1835 | - | - | - | 1992 ⁷⁵ | E | IP,RS | L | Ss | I-III |
| <i>Macrophthalmus graeffei</i> Milne Edwards, 1873 | - | - | - | 1995 ⁷⁴ | ? | IP,RS | L | Ss | I-III |
| Stomatopoda | | | | | | | | | |
| <i>Erugosquilla massavensis</i> (Kossmann, 1880) | - | 2004 ¹¹⁰ | - | 1961 ¹⁰⁶ | E | IP,RS | L | Ss | I-III |
| MOLLUSCA | | | | | | | | | |
| Gastropoda | | | | | | | | | |
| <i>Diodora ruppellii</i> (Sowerby, 1834) | - | - | - | 1995 ⁷² | E | IP,RS | L | Hs | I |
| <i>Smaragdia souverbiiana</i> (Montrouzier, 1863) | - | - | - | 1994 ⁴⁸ | E | IP,RS | ? | Ss | I |
| <i>Trochus erythreus</i> Brocchi, 1821 | - | - | - | 1995 ⁷² | E | RS | L | Hs | I |
| <i>Pseudominolia nedyma</i> (Melville, 1897) | - | - | - | 1995 ⁷² | E | RS,IO | L | Ss | I,II |
| <i>Stomatella impertusa</i> (Burrow, 1815) | - | - | - | 2000 ¹⁵² | ? | IP,RS | ? | Hs | I |
| <i>Cerithium scabridum</i> Philippi, 1848 | - | - | 1990 ³ | 1987 ¹²⁹ | E | RS,IO | L | Hs,Ss | I |
| <i>Rhinoclavis kochi</i> (Philippi, 1848) | - | - | - | 1987 ¹²⁹ | E | IP,RS | ?L | Ss | I |
| <i>Gibborissoa virgata</i> (Philippi, 1849) | - | - | - | 2002 ¹ | E | IP | L | ? | ? |
| <i>Finella pupoides</i> Adams, 1860 | - | - | - | 1990 ¹⁵⁹ | E | IP | L | Ss | I |
| <i>Clathrofenella ferruginea</i> (Adams, 1860) | - | - | - | 1995 ⁷² | E | IP,RS | L | Ss | I,II |
| <i>Cerithiopsis pulvis</i> (Issel, 1869) | - | - | - | 1990 ¹⁵⁹ | E | RS | ?L | Ss | I |
| <i>Cerithiopsis tenthrenois</i> (Melvill, 1896) | - | - | - | 1990 ¹⁵⁹ | E | IO | ?L | Ss | I |
| <i>Metaxia bacillum</i> (Issel, 1869) | - | - | - | 1995 ⁷² | E | RS | ?L | ? | ? |
| <i>Rissoina ambigua</i> (Gould, 1849) | - | - | - | 2004 ¹³⁵ | ? | IP | ? | ? | II |
| <i>Rissoina bertholleti</i> Issel, 1869 | - | - | - | 1991 ¹³⁹ | E | RS,IO | ?L | Ss | I |
| <i>Strombus persicus</i> (Swainson, 1821) | - | - | 1991 ¹³⁹ | 1983 ¹³⁸ | E | PG | ? | Hs,Ss | I,II |
| <i>Purpuradusta gracilis notata</i> (Gill, 1858) | - | - | - | 1983 ⁴⁵ | E | RS,IO | ?L | Hs | I |
| <i>Cycloscalpa hyalina</i> (Sowerby, 1844) | - | - | - | 1999 ⁹⁶ | E | IP,RS | ? | ? | ? |
| <i>Sticteulima cf. lentiginosa</i> (Adams, 1861) | - | - | - | 1994 ¹⁵⁸ | ? | IP | ? | ? | ? |
| <i>Ergalatax obscura</i> Houart, 1996 | - | - | - | 1995 ⁵⁰ | E | RS | ? | Hs | I |
| <i>Thais lacera</i> (Born, 1778) | - | - | - | 1991 ¹³⁹ | E | PG,IO | S | Hs,Ss | I |
| <i>Rapana venosa</i> (Valenciennes, 1846) | 1960 ⁹⁰ | 1996 ¹² | 1995 ⁷² | - | E | PO | S | Ss | I |
| <i>Zafra savignyi</i> (Moazzo, 1939) | - | - | - | 1991 ¹³⁹ | E | RS | ?L | Ss | I,II |
| <i>Zafra selasphora</i> (Melvill & Standen, 1901) | - | - | - | 1993 ¹⁴⁷ | E | RS,IO | ?L | ? | ? |
| <i>Lienardia michelsi</i> Iredale & Tomlin, 1917 | - | - | 2003 ¹³⁵ | - | ? | IP | ? | ? | III |
| <i>Murchisonella columna</i> (Hedley, 1907) | - | - | - | 1995 ⁴⁶ | ? | IP | ? | Hs | I |
| <i>Chrysallida fischeri</i> (Hornung & Mermod, 1925) | - | - | - | 1992 ¹³⁴ | E | RS | ?L | ? | I |

(continued)

Table 1 (continued)

| | BS | SM | AS | LS | Es | O | MI | H | D |
|---|--------------------|--------------------|---------------------|---------------------|----|-------|----|-------|--------|
| <i>Chrysallida maiae</i> (Hornung & Mermod, 1924) | - | - | - | 1990 ¹⁵⁹ | E | RS | L | Hs,Ss | I |
| <i>Chrysallida pиринтхелла</i> (Melvill, 1910) | - | - | - | 1992 ¹³⁴ | E | RS | ?L | Ss | I |
| <i>Adelactaeon amoenus</i> (Adams, 1851) | - | - | - | 1992 ¹³⁴ | E | IP | ?L | Ss | I |
| <i>Adelactaeon fulvus</i> (Adams, A., 1851) | - | - | - | 1992 ¹³⁴ | E | IP | ?L | Ss | I,II |
| <i>Styloptygma beatrix</i> Melvill, 1896 | - | - | - | 1992 ¹³⁴ | E | PG | ?L | ? | I |
| <i>Cingulina isseli</i> (Tryon, 1886) | - | - | - | 1989 ⁵ | E | RS | ?L | ? | ? |
| <i>Turbonilla edgarii</i> (Melvill, 1896) | - | - | - | 1992 ¹³⁴ | E | IP | ?L | ? | ? |
| <i>Syrnola cinctella</i> Adams, A., 1860 | - | - | - | 1998 ⁴ | ? | IP,RS | ? | ? | I |
| <i>Syrnola fasciata</i> Jickeli, 1882 | - | - | - | 1989 ⁵ | E | IP | ?L | Ss | I |
| <i>Iolaea neofelixoides</i> (Nomura, 1936) | - | - | - | 1998 ⁴ | ? | PO | ? | Ss | I |
| <i>Hinemoa cylindrica</i> (de Folin, 1879) | - | - | - | 2001 ⁵¹ | ? | IP | ? | ? | ? |
| <i>Leucotina</i> cf. <i>eva</i> Thiele, 1925 | - | - | - | 2001 ⁹⁶ | ? | IP | ? | ? | ? |
| <i>Acteocina crithodes</i> Melvill & Standen, 1907 | - | - | - | 2004 ¹³⁵ | ? | IP | ? | Ss | II |
| <i>Acteocina mucronata</i> (Philippi, 1849) | - | - | 1990 ⁶ | 1990 ⁶ | E | RS | ?L | Ss | ? |
| <i>Cylichmina girardi</i> (Audouin, 1826) | - | - | 1996 ⁴⁹ | 1990 ¹⁵⁹ | E | IP | L | Ss | I |
| <i>Pyrunculus fourierii</i> (Audouin, 1826) | - | - | - | 1989 ⁵ | E | IP,RS | L | Ss | II,III |
| <i>Bulla ampulla</i> Linnaeus, 1758 | - | - | - | 2004 ¹⁷¹ | E | IP | ?L | Ss | II |
| <i>Haminoea cyanomarginata</i> Heller & Thomson, 1983 | - | - | 2004 ¹⁷¹ | 2004 ¹⁷¹ | ? | RS | ?L | Hs | II |
| <i>Chelidonura fulvipunctata</i> Baba, 1938 | - | - | - | 1961 ¹⁵⁴ | E | IP | ?L | Ss | I,II |
| <i>Oxynoe viridis</i> (Pease, 1861) | - | - | - | 2004 ¹⁷¹ | ? | IP | ?L | Ss | I |
| <i>Elysia grandifolia</i> Kelaart, 1858 | - | - | - | 2004 ¹⁷¹ | Q | ?IO | ?L | Hs | I |
| <i>Elysia tomentosa</i> Jensen, 1997 | - | - | - | 2004 ¹⁷¹ | ? | ?IP | ? | Ss | I,II |
| <i>Bursatella leachii</i> Blainville, 1817 | - | - | 1961 ¹⁵⁴ | 1961 ¹⁵⁴ | E | CT | ?L | Ss | I |
| <i>Syphonota geographica</i> (Adams & Reeve, 1850) | - | - | - | 2004 ¹⁷¹ | ? | IP | L | Ss | I,II |
| <i>Plocamopherus ocellatus</i> Rüppell & Leuckart, 1830 | - | - | - | 2004 ¹⁷¹ | E | RS | L | Hs | I |
| <i>Hypselodoris infucata</i> Rüppell & Leuckart, 1828 | - | - | - | 2001 ⁵⁶ | E | IP,RS | ?L | Hs | I |
| <i>Melibe viridis</i> Kelaart, 1858 | - | - | - | 2004 ¹⁷¹ | ? | IP | ? | Ss | I |
| <i>Flabellina rubrolineata</i> (O'Donoghue, 1929) | - | - | 2004 ¹⁷¹ | 2004 ¹⁷¹ | E | IP,RS | ?L | Hs | I,II |
| <i>Siphonaria belcheri</i> Hanley, 1858 | - | - | - | 2001 ¹³ | E | IP,RS | L | Hs | I |
| Bivalvia | | | | | | | | | |
| <i>Anadara demiri</i> (Piani, 1981) | - | - | 1977 ⁶⁸ | - | E | IO | ?S | Ss | I,II |
| <i>Anadara inflata</i> (Reeve, 1846) | - | - | - | 2002 ⁶⁰ | ? | IO | ?S | Ss | ? |
| <i>Anadara inaequivalvis</i> (Bruguière, 1789) | 2003 ¹⁰ | 1996 ¹¹ | 1995 ⁷² | - | E | IP | ?S | Ss | I,II |
| <i>Anadara natalensis</i> (Krauss, 1848) | - | - | - | 1991 ¹³⁹ | E | IO,RS | L | Ss | I,II |
| <i>Brachidontes pharaonis</i> (Fischer, 1870) | - | - | 1990 ³ | 1985 ¹¹⁶ | E | IO,RS | L | Hs | I |
| <i>Septifer forskali</i> Dunker, 1855 | - | - | - | 2001 ¹³ | ? | RS | ? | Hs | II |

(continued)

Table 1 (continued)

| | BS | SM | AS | LS | Es | O | Mi | H | D |
|--|----|--------------------|---------------------|---------------------|----|-------|----|-------|-------|
| <i>Crassostrea gigas</i> (Thunberg, 1793) | - | 2004 ¹⁵ | - | 2001 ⁵⁷ | E | PO | Aq | Hs | I |
| <i>Saccostrea cucullata</i> (Born, 1778) | - | - | - | 2001 ⁵⁷ | E | IP,RS | ?S | Hs | I,II |
| <i>Dendrostrea frons</i> (Linnaeus, 1758) | - | - | - | 2001 ⁵⁹ | E | IP | ? | Hs | ? |
| <i>Pinctada radiata</i> (Leach, 1814) | - | - | 1990 ¹⁵⁹ | 1985 ¹¹⁶ | E | IP,RS | ?L | Hs | I |
| <i>Electroma vexillum</i> (Reeve, 1857) | - | - | - | 2005 ⁵⁸ | ? | IP,RS | ?L | Hs | I |
| <i>Malvufundus regulus</i> (Forsskål, 1775) | - | - | - | 1974 ⁸⁸ | E | IP,RS | L | Hs | I |
| <i>Spondylus cf. multisetosus</i> Reeve, 1856 | - | - | - | 2001 ⁵⁹ | Q | IP | ? | Hs | ? |
| <i>Spondylus spinosus</i> Schreibers, 1793 | - | - | - | 1999 ⁷³ | E | IP,RS | L | Hs | I,II |
| <i>Chama pacifica</i> Broderip, 1834 | - | - | - | 2001 ⁵⁹ | E | IP,RS | L | Hs | I,II |
| <i>Fulvia fragilis</i> (Forsskål, 1775) | - | - | 2005 ¹⁴⁴ | 1987 ¹²⁹ | E | IO,RS | ?L | Ss | I |
| <i>Afrocardium richardi</i> (Audouin, 1826) | - | - | - | 2000 ² | E | RS | L | Ss | ? |
| <i>Tellina valtonis</i> Hanley, 1844 | - | - | - | 2001 ⁹⁶ | E | IO,RS | L | Ss | I |
| <i>Psammotreta praerupta</i> (Salisbury, 1934) | - | - | - | 1999 ⁷³ | ? | WA | ? | Ss | IV |
| <i>Gastrarium pectinatum</i> (Linnaeus, 1758) | - | - | - | 1987 ¹²⁹ | E | IP,RS | L | Ss | I |
| <i>Clementia papyracea</i> (Gray, 1825) | - | - | - | 1995 ⁷² | E | IP,RS | L | Ss | III |
| <i>Paphia textile</i> (Gmelin, 1791) | - | - | - | 1991 ¹³⁹ | E | IP,RS | L | Ss | ? |
| <i>Ruditapes philippinarum</i> (Adams & Reeve, 1850) | - | - | 2001 ¹⁴ | - | E | PO | Aq | Ss | I |
| <i>Antigona lamellaris</i> Schumacher, 1817 | - | - | - | 1999 ⁷³ | ? | IP,RS | ? | Ss | ? |
| <i>Mya arenaria</i> Linnaeus, 1758 | - | 1996 ¹¹ | - | - | E | WA | ?S | Ss | II |
| <i>Gastrochaena cymbium</i> Spengler, 1783 | - | - | - | 1991 ¹³⁹ | E | IP,RS | L | Ss | I-III |
| <i>Laternula anatina</i> (Linnaeus, 1758) | - | - | - | 1995 ⁷² | E | IP,RS | L | Ss | I-III |
| Cephalopoda | | | | | | | | | |
| <i>Octopus aegina</i> Gray, 1849 | - | - | - | 1999 ¹⁵¹ | E | IP | ?L | Ss | III |
| <i>Sepioteuthis lessoniana</i> Lesson, 1830 | - | - | - | 2002 ¹⁵⁰ | E | IP | L | P | I-III |
| BRYOZOA | | | | | | | | | |
| <i>Rhynchocoelium larreyi</i> (Audouin, 1826) | - | - | 1979 ¹⁶⁵ | - | E | IP,RS | L | Ss | I |
| ECHINODERMATA | | | | | | | | | |
| <i>Ophiactis savignyi</i> (Müller & Troschel, 1842) | - | - | 1998 ⁶¹ | - | E | IP,RS | L | Hs | I |
| <i>Synaptula reciprocans</i> (Forsskål, 1775) | - | - | 2001 ¹⁷³ | - | E | IP,RS | L | Hs,Ss | ? |
| PISCES | | | | | | IP,RS | | | |
| Chondrichthyes | | | | | | | | | |
| <i>Carcharhinus altimus</i> (Springer, 1950) | - | - | - | 2000 ³¹ | Q | TA | G | Ss | I-VI |
| <i>Himantura uarnak</i> (Forsskål, 1775) | - | - | - | 1966 ³⁷ | C | IP,RS | L | Ss | I,II |
| Osteichthyes | | | | | | | | | |
| <i>Dussumieria elopsoides</i> Bleeker, 1849 | - | - | - | 1953 ³⁶ | E | IP,RS | L | P | I,II |
| <i>Etrumeus teres</i> (DeKay, 1848) | - | - | - | 1997 ³² | E | IP,RS | L | P | I,II |
| <i>Herklotischthys punctatus</i> (Rüppell, 1837) | - | - | - | 1984 ¹⁶⁹ | E | RS | L | P | I,II |
| <i>Enchelycore anatina</i> (Lowe, 1839) | - | - | - | 2002 ¹⁷² | E | TA | G | Hs | I,II |

(continued)

Table 1 (continued)

| | BS | SM | AS | LS | Es | O | MI | H | D |
|--|---------------------|---------------------|---------------------|---------------------|----|-------|----|-------|-------|
| <i>Saurida undosquamis</i> (Richardson, 1848) | - | - | 1973 ³⁸ | 1966 ³⁷ | E | IP,RS | L | Ss | I-III |
| <i>Parexocoetus mento</i> (Valenciennes, 1846) | - | - | 1966 ³⁷ | 1966 ³⁷ | E | IP,RS | L | P | I |
| <i>Hemiramphus far</i> (Forsskål, 1775) | - | - | 1950 ¹²⁶ | 1950 ¹²⁶ | E | IP,RS | L | P | I |
| <i>Hyporhamphus affinis</i> (Günther, 1866) | - | - | - | 1954 ⁷ | Q | IP,RS | L | P | I |
| <i>Fistularia commersonii</i> (Rüppell, 1835) | - | - | 2002 ⁴³ | 2002 ⁴² | E | IP,IO | L | Hs,Ss | I,II |
| <i>Hippocampus fuscus</i> Rüppell 1838 | - | - | - | 2004 ⁹⁸ | ? | IO,RS | L | Ss | I |
| <i>Syngnathus rostellatus</i> Nilsson, 1855 | - | - | - | 2004 ⁹⁸ | Q | BA | G | Hs,Ss | I |
| <i>Atherinomorus lacunosus</i> (Forster, 1801) | - | - | 1969 ⁹⁴ | 1950 ¹²⁶ | E | IP,RS | L | P | I |
| <i>Sargocentron rubrum</i> (Forsskål, 1775) | - | - | 1950 ¹²⁶ | 1950 ¹²⁶ | E | IP,RS | L | Hs | I-III |
| <i>Pelates quadrilineatus</i> (Bloch, 1790) | - | - | - | 1987 ¹³¹ | E | IP,RS | L | Ss | I,II |
| <i>Apogon pharaonis</i> Bellotti, 1874 | - | - | - | 1987 ¹³¹ | E | IP,RS | L | Hs,Ss | I,II |
| <i>Sillago sihama</i> (Forsskål, 1775) | - | - | 2004 ⁴¹ | 1994 ¹⁰² | E | IP,RS | L | Ss | I,II |
| <i>Alepes djedaba</i> (Forsskål, 1775) | - | - | 1969 ⁹⁴ | 1957 ⁹ | E | IP,RS | L | P | I-III |
| <i>Leiognathus klunzingeri</i> (Steindachner, 1898) | - | - | 1966 ³⁷ | 1943 ⁷⁸ | E | RS | L | Ss | I-III |
| <i>Upeneus moluccensis</i> (Bleeker, 1855) | - | - | 1956 ¹²⁷ | 1950 ¹²⁶ | E | IP | L | Ss | I-IV |
| <i>Upeneus pori</i> Ben-Tuvia & Golani, 1989 | - | - | - | 1950 ¹²⁶ | E | IP,RS | L | Ss | I,II |
| <i>Pempheris vanicolensis</i> Cuvier, 1831 | - | - | 1999 ¹³⁰ | 1994 ¹⁰² | E | IP,RS | L | Hs | I,II |
| <i>Heniochus intermedius</i> Steindachner, 1893 | - | - | - | 2003 ⁹⁷ | Q | IP,RS | L | Hs,Ss | I |
| <i>Chelon carinata</i> (Valenciennes, 1836) [= <i>Liza carinata</i>] | - | - | - | 1956 ¹²⁷ | E | RS,IO | L | P | I |
| <i>Liza haematocheila</i> (Temminck & Schlegel, 1845) [= <i>Mugil soiuy</i> Basilewsky, 1855] | 1992 ¹⁶⁶ | 1998 ¹¹² | 1998 ¹¹² | - | E | PO | Aq | P | I |
| <i>Sphyraena pinguis</i> Doiuchi & Nakabo, 2005 [= <i>S. chrysotaenia</i> Klunzinger, 1884] | - | - | 1969 ⁹⁴ | 1957 ⁹ | E | IP,RS | L | P | I,II |
| <i>Sphyraena obtusata</i> Cuvier, 1829 [= <i>S. flavicauda</i> Rüppell, 1838] | - | - | - | 2002 ⁴² | E | IP,RS | L | P | I,II |
| <i>Pteragogus pelvus</i> Randall, 1981 | - | - | - | 2000 ¹⁵⁶ | E | IP,RS | L | Hs,Ss | I,II |
| <i>Petrosomites aenylodon</i> Rüppell, 1838 | - | - | - | 2000 ¹⁵⁶ | E | IP,RS | L | Hs,Ss | I |
| <i>Oxyurichthys petersi</i> (Klunzinger, 1871) | - | - | 1999 ³⁵ | 1992 ¹¹¹ | E | RS | L | Ss | I-III |
| <i>Callionymus filamentosus</i> Valenciennes, 1837 | - | - | - | 1994 ¹⁰² | E | IP,RS | L | Ss | I-III |
| <i>Siganus luridus</i> (Rüppell, 1829) | - | - | 1973 ³⁸ | 1973 ³⁸ | E | IP,RS | L | Hs | I,II |
| <i>Siganus rivulatus</i> Forsskål, 1775 | - | - | 1950 ¹²⁶ | 1950 ¹²⁶ | E | IP,RS | L | Hs | I,II |
| <i>Scomberomorus commerson</i> Lacepède, 1800 | - | - | 1997 ⁴⁷ | 1987 ⁸⁹ | E | IP,RS | L | P | I,II |
| <i>Solea senegalensis</i> Kaup, 1858 | - | 1942 ⁷⁷ | 1942 ⁷⁷ | 1942 ⁷⁷ | Q | TA | G | Hs | I-III |
| <i>Cynoglossus sinusarabici</i> (Chabanaud, 1931) | - | - | - | 1957 ⁹ | E | RS | L | Hs | I,II |

(continued)

Table 1 (continued)

| | BS | SM | AS | LS | Es | O | MI | H | D |
|--|----|----|---------------------|---------------------|----|-------|----|-------|------|
| <i>Stephanolepis diaspros</i> Fraser-Brunner, 1940 | - | - | 1950 ¹²⁶ | 1950 ¹²⁶ | E | IP,RS | L | Hs,Ss | I,II |
| <i>Lagocephalus sceleratus</i> (Gmelin, 1789) | - | - | 2005 ⁸ | - | E | IP,RS | L | Ss | I,II |
| <i>Lagocephalus spadiceus</i> (Richardson, 1844) | - | - | 1966 ³⁷ | 1950 ¹²⁶ | E | IP,RS | L | Ss | I,II |
| <i>Lagocephalus suezensis</i> Clark & Gohar, 1953 | - | - | 2002 ⁴² | 1999 ¹⁶ | E | RS | L | Ss | I,II |
| <i>Sphoeroides pachygaster</i> (Müller & Troschel, 1848) | - | - | 2003 ⁸⁷ | 1999 ¹³⁰ | E | TA | G | Hs,Ss | I-V |
| <i>Torquigener flavimaculosus</i> Hardy & Randall, 1983 | - | - | - | 2003 ⁴⁰ | E | IP,RS | L | Ss | I,II |

coast. Forty-eight alien Crustaceans (94% of the total aliens) belong to Lessepsian migrants. Most of the plants (29 species) were introduced via ships. The other modes of introduction of species were migration through the Gibraltar straits and Aquaculture.

Distribution of alien species on the Turkish coasts

The number of alien species found along the Turkish coasts and their modes of introduction are presented in Figure 1. The Turkish Black Sea coast possessed the lowest number of species (20 species), whereas the Turkish Levantine coast had the highest (202 species). The importance of shipping as a vector of introduction of species gradually diminishes from the Black Sea to the Levantine Sea. Lessepsian migrants exclusively dominate benthic habitats of the Levantine Sea. The Aegean Sea appears to be more influenced by Lessepsian migration than ship transportation.

The majority (70% of the species) of the marine animals introduced by ships in the Aegean Sea were reported from Izmir Bay; *Polydora cornuta*, *Streblospio gynobranchiata*, *Ficopomatus enigmaticus*, *Hydroides dianthus*, *H. elegans*, *Spirorbis marioni*, *Anadara demiri* and *A. inequivalvis*. This is mainly attributed to two facts; 1) Alsancak Harbour, which is one of the most important commercial harbours in Turkey, is located in the polluted part of Izmir Bay, 2) This bay has been relatively well studied and

continuously monitored since the 1970's. Species-poor communities such as polluted or physically degraded environments are known to be more vulnerable to invasion than are other communities (ZIBROWIUS, 1992). Izmir Bay has been subjected to various pollution discharges, intense marine transportation and increasing human populations. Ocean-going ships approaching Alsancak Harbour for loading or unloading processes empty their ballast tanks just before entering the harbour and the weakening in the ecosystem due to pollution greatly facilitates the settlement of alien species in the area.

Rate of invasion

The new additions to the inventory of the marine alien species on the Turkish coasts greatly increased after 1980 and the last five years alone a total of 92 new species were recorded (Fig. 2). A decreasing pattern is obvious in the yearly rate of introduction over the past years (Table 2). While one new alien species was introduced every 15.3 weeks over the period 1981-2000 in the Aegean Sea, the time span has decreased to 8.7 between 2001 and 2005. The special interest in aliens by Turkish and foreign scientists in the last decade has increased and thereby, the real diversity of aliens within the groups of phyto-benthos, fishes and crustaceans inhabiting the Turkish coasts seems to be well documented. Nowadays, a number of projects regarding the actual status of aliens along the Turkish coasts




Fig. 1: The number of alien species along the Turkish coasts and their mode of introduction. PLT: All plants, PP: Phytoplankton, PhB: Phytobenthos, CTN: Ctenophora, POL: Polychaeta, CRU: Crustacea, MOL: Mollusca, OTH: Others. ΣS indicates the total number of species reported from the sea.




Fig. 2: Rate of introduction of alien species along the Turkish coasts.

Table 2.
Time span (weeks) for alien species introduction along the Turkish coasts.

| | 1961-1980 | 1981-2000 | 2001-2005 |
|----------------|---------------------------|-----------|-----------|
| | Weeks per 1 alien species | | |
| Black Sea | 208 | 69.3 | 65 |
| Sea of Marmara | 80 | 31.5 | 65 |
| Aegean Sea | 40 | 15.3 | 8.7 |
| Levantine Sea | 52 | 6.4 | 4.7 |
| Total | 16.3 | 3.7 | 2.8 |

and their probable impacts on the native biota are undertaken.

Depth and habitat preferences of alien species

The distribution of alien species by depth and habitat is indicated in Figure 3. The majority of aliens collected from the Turkish coasts were found in shallow waters. A total of 106 species (41% of the total number of aliens) solely occurred at depths ranging from 0 to 10 m, and 180 species (68%) at the depth interval 0-50 m. Species with a wide distributional range in the area are *Leonnates persicus* (10-200 m), *Penaeus semisulcatus* (0-200 m), *Leptochela pugnax* (0-200 m), *Palaemonella rotumana* (0-200 m), *Myra subgranulata* (0-200 m), *Carcharhinus altimus* (0-500 m), *Upeneus moluccensis* (0-200 m) and *Sphoeroides pachygaster* (0-400 m).

Benthic habitats (soft and hard substrata) harboured 76% of the total number of alien species (Figure 3). Thirty-nine species inhabited the pelagic environment. In soft substrata, all major groups, except for algae, are represented by a higher number of species. The majority of hard bottom organisms were reported among algae.

The echinoderm *Ophiactis savignyii* was only reported in canals of the sponge *Sarcotragus muscarum* from the southern part of the Aegean Sea [ÇINAR & ERGEN, 1998 (as *Ophiactis virens* (M. Sars, 1857)); ÇINAR *et al.*, 2002]. The sweeper, *Pempheris vanicolensis*, prefers cavern habitats (BILECENOĞLU & TASKAVAK, 1999). The shrimp scad, *Alepes djedaba*, is generally associated with the venomous jellyfish *Rhopilema nomadica* (KIDEYS & GÜCÜ, 1995). ÖKSNEBJERG *et al.* (1997)




Fig. 3: The habitat (upper graphic) and depth (lower graphic) preferences of alien species along the Turkish coasts.

reported the parasitic cirriped *Heterosaccus dollfusi* on the Lessepsian crab *Charybdis longicollis*. The nudibranch *Flabellina rubrolineata* was found on hydroids (YOKES & RUDMAN, 2004). ALBAYRAK & ÇEVIKER (2001) reported *Septifer forskali* on the bivalve *Spondylus spinosus*. *Electroma vexillum* was encountered in discharge canals of the Iskenderun Iron and Steel Factory, where hot water is discharged into the sea (ÇEVIKER *et al.*, 2005).

Impacts of alien species

The phytoplankton species, *Alexandrium tamarense*, *Gymnodinium* cf. *mikimotoi* and *Heterosigma* cf. *akashiwae*, which were reported from Izmir Bay and classified as questionable in Table 1, have a potential to cause a toxic and/or harmful bloom along the Turkish coasts (KORAY, 1984; BIZSEL & BIZSEL, 2002).

The green alga *Caulerpa racemosa* invaded soft and hard bottom habitats of the Levantine and Aegean coasts but its impact on the native biota has not yet been documented in the area. However, its occurrence off the Dardanel (salinity: 27 psu) indicates its highly adaptive and invasive character (OKUDAN *et al.*, 2002).

The negative effect of *Mnemiopsis leidyi* on the Black Sea ecosystem was summarized by KIDEYS (2002). This zooplanktonic predator reached enormous biomass levels in the summer of 1989, devastating the food chain of the entire Black Sea basin, and subsequently led to a sharp decrease in the anchovy production along the Turkish coast. The other ship-mediated ctenophore, *Beroe ovata*, which largely feeds on ctenophores, appeared in 1997 in the Black Sea and has resulted in the decrease in the population of *M. leidyi*.

The outburst of the population of *Rhopilema nomadica* off the Levantine coast of Turkey was reported to have negative consequences on human health, tourism and fisheries (KIDEYS & GÜCÜ, 1995). Many swimmers were stung and sought medical treatment. The blockage of nets of fishermen by individuals of *R. nomadica* also created major economical losses.

Rapana venosa, which feeds exclusively on bivalves, is responsible for the decrease in stocks of the bivalves *Ostrea edulis* Linné, 1758 and *Mytilus galloprovincialis* (Lamarck, 1819) in the Black Sea (BILECIK, 1990). The fishery of *R. venosa* was intense along the Turkish Black Sea coast, reaching up to 1166 tons in 1986, yielding 3.415.884 US \$ (BILECIK, 1990).

Low faunal diversity found on sponge samples collected from the southern Aegean Sea was due to the invasion of the Lessepsian species *Ophiactis savignyi* (ÇINAR *et al.*, 2002).

The species number and density of other groups within the sponge samples were sharply declined as compared to those collected from the northern sites where this echinoderm was not present. A negative correlation was estimated between number of specimens of the ophiuroid and the total number of specimens of other taxa.

The soft bottom near Alsancak Harbour in Izmir Bay was exclusively dominated by the spionid polychaete *Streblospio gynobranchiata*, which originated from the western Atlantic (ÇINAR *et al.*, 2005). Its density reached almost 34 300 ind.m⁻² in the area and accounted for almost 100% of faunal populations at some stations. The soft bottom benthic community around Alsancak Harbour seems to be restructured by this species.

KOÇAK *et al.* (1999) found dense populations of two alien serpulid species, *Hydroides elegans* and *H. dianthus*, on panels submerged in a polluted marina in Izmir Bay. The species density of *H. elegans* reached up to 98000 ind.m⁻² and that of *H. dianthus* up to 2 000 ind.m⁻². The above authors also reported that an alien serpulid species of eastern Pacific origin, *Spirorbis marioni*, inhabited port environments subjected to relatively less pollution.

The density of *Anadara demiri* was found to be 300 ind.m⁻² on grey mud and 30 ind.m⁻² on black mud in Izmir Bay (DEMIR, 1977).

Aliens might bring their own parasites with them. ØKSNEBJERG *et al.* (1997) reported that 90% of the specimens of *Charybdis longicollis* collected between Karatas and Fener Burnu (Levantine coast of Turkey) were infected by the rhizocephalan *Heterosaccus dollfusi*. They also stated that the prevalence of this parasite was estimated at 50-60% in the east of Karatas, with its prevalence gradually declining towards the east into Iskenderun Bay. GALIL & LÜTZEN (1995) found that 55% of the males and 43% of the females of *C. longicollis* were visibly infected by this parasite along the Israeli coast in May 1994.

From a large number of aliens inhabiting the Turkish coasts, only a few possess commercial value, mostly belonging to fishes and crustaceans. Little is known concerning the impact of




Fig. 4: Percentages of alien species in total fauna along the Turkish coasts.

commercially important alien species on local fisheries, since there is no available catch statistic that distinguishes native and alien species' landings. Due to the great number of species caught during various fisheries activities, governmental statistics tend to present relevant data on a family basis. For example, the "mullids" category includes four species (native *Mullus* spp. and Lessepsian *Upeneus* spp.); a similar situation is valid for sphyraenids (two native species vs. two Lessepsian species). Lessepsian clupeids, especially *Etrumeus teres*, form a remarkable proportion of the pelagic catch along the Levantine coast. The siganids are generally sold together with the native *Sarpa salpa* due to their similar morphologies (BILECENOĞLU & KAYA, 2002). At least seven fish species (*Upeneus moluccensis*, *Saurida undosquamis*, *Scomberomorus commerson*, *E. teres*, *Sphyraena pinguis*, *Siganus* spp.) are captured in large amounts (mainly by bottom trawls), and consumed throughout the Levantine and southern Aegean Sea coasts. In terms of biomass, *S. undosquamis* is the most prominent species in the bottom trawl catch composition (31.9% of total CPUE in Iskenderun Bay), followed by *U. moluccensis* and *Siganus rivulatus* (GÜCÜ *et al.*, 1994).

Some species have relatively smaller catches and thus mostly consumed locally, i.e.

Alepes djedaba, *Sillago sihama*, *Dussumieria elopsoides*, *Atherinomorus lacunosus*. Among alien crustaceans, the highest annual production belongs to *Callinectes sapidus* (ca. 200 tonnes/year), which is captured especially in lagoons in amounts as much as 2 tonnes/day during summer periods (ÖZCAN *et al.*, 2003). *Marsupenaeus japonicus* is one of the most valuable shrimp species along the Levantine coast, which is also imported to various cities in Turkey. Since they are not captured in large quantities, other commercial shrimps (*Metapenaeus* spp., *Penaeus semisulcatus*) and brachyurans (*Portunus pelagicus*) are, in general, consumed locally.

Importance of alien species in total fauna

The relative percentages of aliens in total fauna are shown in Figure 4. Mollusca had the highest score in the Black (3.1%) and Levantine (16.1%) Seas, Polychaeta in the Sea of Marmara (4.4%) and fishes in the Aegean Sea (5.9%). The score for Polychaeta in the Sea of Marmara does not reflect its true number as 11 of 12 aliens are questionable. The score seems to be accurate for the better-studied group, fishes, which has more aliens on the southern coasts than the northern coasts due to the influence of Lessepsian migrants. The highest alien percentages for all groups examined were found on the Levantine

coast. This partly shows the impact of Lessepsian migration on the Levantine fauna and partly the lack of detailed works on some groups on the coast. The total number of species given for Mollusca and Polychaeta from the Levantine coast of Turkey is highly underestimated and undoubtedly would increase as further studies are carried out. It is a common assumption that Lessepsian migrants account for at least 10% of the species inventory of the Levantine Sea (POR, 1978). BEN-ELIAHU (1995) estimated that the proportion of migrant polychaetes is almost 9%; for fishes 13% (GOLANI, 1998) and for decapod crustaceans 20% (GALIL, 1986).

Similarity among seas in terms of alien species

The dendrogram and MDS configuration in Figure 5 show that there are two groups in terms of the presence and absence of alien species in the Turkish seas. The Black Sea and the Sea of Marmara constitute one group, in which 16 species (mainly algae) are common. The other group with a similarity of 40% is composed from the Aegean and Levantine Seas, where 63 species (mainly lessepsian fishes) are common. There is a very weak similarity (20%) between the groups, which are placed at different corners of the MDS plot. As pointed out earlier, the southern seas seem to be more affected by Lessepsian migrants than ship-borne species, whereas shipping is the main vector in the transportation of aliens to the northern seas. The difference in the way of introduction seems to be the main reason for similarity or dissimilarity among the seas.

Conclusions

The compiled data show that a total of 263 established or questionable alien species inhabit the coast of Turkey, with the Levantine Sea having the majority of them (202 species). Detecting new aliens depends totally on accurate taxonomic identifications and the knowledge of local biodiversity. This requires the availability of systematic experts on biotic components in the area and a close cooperation among scientists both on local and global scales. The wealth




Fig. 5: Dendrogram and MDS plot showing affinities of the seas in terms of alien species they included.

of data accumulated on aliens depicts the importance of monitoring studies on the understanding of ecological and functional roles of aliens in the ecosystems. Finally, results of studies and projects should not be concealed among pages of reports and papers. They should be presented to governmental establishments and politicians to force them to take immediate precautions on the dispersal of aliens, particularly via ballast water discharges.

Addendum

Before the publication of this contribution, a total of 14 new alien species were reported from the Levantine (13 species) and Aegean [1 species (*Platax teira*)] coasts of Turkey (BILECENOĞLU & KAYA, 2006; ÇEVİKER & ALBAYRAK, 2006; ÇINAR *et al.*, 2006; ÖZTÜRK & CAN, 2006). These species belong to Hydrozoa (*Macrorhynchia philippina* (Kirchenpauer, 1872), Anthozoa (*Oculina patagonica*

De Angelis, 1908), Polychaeta (*Branchiomma luctuosum* Grube, 1869), Gastropoda (*Amathina tricarinata* (Linnaeus, 1767), *Aplysia dactylomela* Rang, 1828, *Chromodoris quadricolor* (Rüppell & Leuckart 1830)], Bivalvia [*Cardites akabana* (Sturany, 1899) and *Petricola hemprichi* Issel, 1869], Echinodermata [*Synaptula reciprocans* (Forsskål, 1775)], Tunicata [*Phallusia nigra* Savignyi, 1816, *Pyura* (=*Herdmania momus* (Savignyi, 1816) and *Symplegma brakenhielmi* (Michaelsen, 1904)] and Osteichthyes [*Parupeneus forsskali* (Fourmanoir and Guézé, 1976) and *Platax teira* (Forsskål, 1775)]. These findings increased the number of alien species known from the coast of Turkey from 263 to 277, and from 202 to 216 on the Levantine coast of Turkey.

References

- ¹AARTSEN, J. J. van, 2002. Indo-Pacific migrants into the Mediterranean. 1. *Gibborissoa virgata* (Philippi, 1849). *La Conchiglia*, 303: 56-58.
- ²AARTSEN, J. J. van & GOUD, J., 2000. European marine Mollusca: Notes on less well-known species. XV. Notes on Lusitanian species of *Parvicardium* Monterosato, 1884, and *Afrocardium richardi* (Audouin, 1826) (Bivalvia:Heterodontida: Cardiidae). *Basteria*, 64: 171-186.
- ³AARTSEN, J. J. van & KINZELBACH, R., 1990. Marine molluscs from the Iztuzu beach near Dalyan (Mediterranean coast of Turkey). *Zoology in the Middle East*, 4: 103-112.
- ⁴AARTSEN, J. J. van & RECEVIK, M., 1998. Two more Indo-Pacific species found in the Mediterranean. *La Conchiglia*, 286: 13-15.
- ⁵AARTSEN, J. J. van, BARASH, A. & CARROZZA, F., 1989. Addition to the knowledge of the Mediterranean mollusca of Israel and Sinai. *Bollettino Malacologico*, 25: 63-76.
- ⁶AARTSEN, J. J. van, CARROZZA, F. & LINDNER, G., 1990. *Acteocina mucronata* (Philippi, 1849), a recent Red Sea immigrant species in the eastern Medi- terranean. *Bollettino Malacologico*, 25: 285-298.
- ⁷AKSIRAY, F., 1954. Türkiye deniz balıkları tayin anahtarı. *Istanbul Üniversitesi Fen Fakültesi Hidrobiologi Araştırma Enstitüsü Yayınları*, İstanbul, 277 p.
- ⁸AKYOL, O., ÜNAL, V., CEYHAN, T. & BILECENOĞLU, M., 2005: First record of the silverside blaasop, *Lagocephalus sceleratus* (Gmelin, 1789), in the Mediterranean Sea. *Journal of Fish Biology*, 66: 1183-1186.
- ⁹AKYUZ, E., 1957. Observations on the İskenderun red mullet (*Mullus barbatus*) and its environment. *GFCM Proceedings and Technical Papers*, 4: 305-326.
- ¹⁰ALBAYRAK, S., 2003. On the Mollusca Fauna of the Black Sea near Istanbul. *Zoology in the Middle East*, 30: 69-75.
- ¹¹ALBAYRAK, S. & BALKIS, N., 1996a. Bivalve Fauna of the Bosphorus. *Istanbul Üniversitesi Fen Fakültesi Biyoloji Dergisi*, 59: 1-15.
- ¹²ALBAYRAK, S. & BALKIS, N., 1996b. Benthic Prosobranch Gastropods of the Bosphorus. *Istanbul Üniversitesi Fen Fakültesi Biyoloji Dergisi*, 59: 17-26.
- ¹³ALBAYRAK, S. & ÇEVIKER, D., 2001. Two extra-Mediterranean molluscs from southeast Turkey: *Siphonaria belcheri* Hanley, 1858 (Gastropoda: Siphonariidae) and *Septifer bilocularis* (Linnaeus, 1758) (Bivalvia: Veneridae). *Israel Journal of Zoology*, 47: 297-298.
- ¹⁴ALBAYRAK, S., ASLAN, H. & BALKIS, H., 2001. A contribution to the Aegean Sea fauna: *Ruditapes philippinarum* (Adam & Reeve, 1850) [Bivalvia : Veneridae]. *Israel Journal of Zoology*, 47: 299-300.
- ¹⁵ALBAYRAK, S., BALKIS, H. & BALKIS, N., 2004. Bivalvia (Mollusca) fauna of the Sea of Marmara. *Acta Adriatica*, 45: 9-26.
- ¹⁶AVSAR, D. & ÇIÇEK, E., 1999. A new species record for the central and eastern Mediterranean; *Sphoeroides cutaneus* (Günther, 1870) (Pisces: Tetraodontidae). *Oebalia*, 25: 17-21.

- ¹⁷AYSEL, V., 1984. Türkiye'nin Ege Denizi'ndeki *Polysiphonia* Greville (*Rhodomelaceae, Ceramiales*) türleri 1-Bölüm *Oligosiphonia*. *Doga Bilim Dergisi*, 8: 29-42.
- ¹⁸AYSEL, V., 1987. Türkiye'nin Ege Denizi Florası II.Kırmızı Algler (*Rhodophyta*). *Doga Türk Botanik Dergisi*, 11: 1-21.
- ¹⁹AYSEL, V., 1997a. Marine flora of Turkish Mediterranean coast 1. Red algae (*Rhodophyta*). *Turkish Journal of Botany*, 21: 155-163.
- ²⁰AYSEL, V., 1997b. Marine flora of Turkish Mediterranean coast 2. Brown algae (*Fucophyceae = Phaeophyceae*). *Turkish Journal of Botany*, 21: 329-334.
- ²¹AYSEL, V. & ERDUGAN, H., 1995. Check-list of Blacksea seaweeds, Türkiye (1823-1994). *Turkish Journal of Botany*, 19: 545-554.
- ²²AYSEL, V. & GEZERLER-SIPAL, U., 1996. Türkiye'nin Akdeniz kıyılarının deniz florası, 3. *Cyanophyceae, Chlorophyceae, Charophyceae ve Angiospermae*. *Ege Üniversitesi Su Ürünleri Fakültesi Dergisi*, 13: 247-257.
- ²³AYSEL, GÜNER, H. & DURAL, B., 1993. Türkiye Marmara Denizi florası II. *Phaeophyta ve Rhodophyta*. *Ege Üniversitesi Su Ürünleri Fakültesi Dergisi*, 10: 115-168.
- ²⁴AYSEL, V., DURAL, B., GÖNÜZ, A. & OKUDAN, E. S., 1998. Kırklareli (Karadeniz, Trakya, Türkiye) deniz florası. *X Ulusal Biyoloji Kongresi*, Samsun, 2: 333-342.
- ²⁵AYSEL, V., KESERCIOGLU, T., GÜNER, H. & AKÇAY, H., 1990. Trabzon deniz algleri. *X Ulusal Biyoloji Kongresi*, Erzurum, 2: 183-192.
- ²⁶AYSEL, V., ZEYBEK, N., GÜNER, H. & SUKATAR, A., 1986. Türkiye'nin bazı derin deniz algleri III. *Rhodophyta* (Kırmızı Algler). *Doga Türk Biyoloji Dergisi*, 10: 8-29.
- ²⁷AYSEL, V., DURAL, B., OKUDAN, E. S., ALPASLAN, M. & UYSAL, I., 2001. Gökçeada (Ege Denizi, Çanakkale) Deniz Florası. *Ulusal Ege Adaları 2001 Toplantısı*, Gökçeada: 125-142.
- ²⁸AYSEL, V., DURAL, B., SUKATAR, A., GÜNER, H. & ERDUGAN, H., 1996a. Zonguldak Deniz Algleri, Karadeniz, Türkiye. *XIII Ulusal Biyoloji Kongresi*, İstanbul, 5: 311-321.
- ²⁹AYSEL, V., ERDUGAN, H., DURAL-TARAKÇI, B., OKUDAN, E. S., SENKARDESLER, A. & AYSEL, F. 2004. Marine Flora of Sinop (Black Sea, Turkey), *Ege University Journal of Fisheries & Aquatic Sciences*, 21: 59-68.
- ³⁰BALKIS, H. & ÇEVIKER, D., 2003. A new exotic crab species [*Calappa hepatica* (Linnaeus, 1758)] for the Mediterranean fauna. *Israel Journal of Zoology*, 49: 320-321.
- ³¹BASUSTA, N. & ERDEM, Ü., 2000. İskenderun Körfezi balıkları üzerine bir araştırma. *Turkish Journal of Zoology*, 24:1-19.
- ³²BASUSTA, N., ERDEM, Ü. & MATER, S., 1997. İskenderun körfezi'nde yeni bir Lesepsiyen göçmen balık türü; Kızılıgözlü Sardalya, *Etrumeus teres* (Dekay, 1842). *Mediterranean Fisheries Congress*, Izmir: 921-924.
- ³³BEN-ELIAHU, M. N., 1989. Lessepsian migration in Nereidae (Annelida: Polychaeta). Some case histories. p. 125-134. In: *Proceedings of Environmental Quality and Ecosystem Stability*, edited by E. Spanier, Y. Steinberger & M. Luria.
- ³⁴BEN-ELIAHU, M. N., 1991. Red Sea serpulids (Polychaeta) in the eastern Mediterranean. p. 515-528. In: *Proceedings of 2nd International Polychaeta Conference, Copenhagen, 1986. Systematics, Biology and Morphology of Word Polychaeta*, edited by M. E. Petersen & J. B. Kirkegaard, Ophelia Supplement, 5.
- BEN-ELIAHU, M. N., 1995. A list of Polychaeta along the Levant coast. *Haasiana*, 1: 78-93.
- ³⁵BENLİ, H. A., CIHANGIR, B. & BİZSEL, K. C., 1999. Ege Denizi'nde bazı demersal balıkçılık kaynakları üzerine

- arastirmalar. *Istanbul Üniversitesi Su Ürünleri Dergisi*, ozel sayi: 301-369.
- ³⁶BEN-TUVIA, A., 1953. Mediterranean fishes of Israel. *Bulletin of the Sea Fisheries Research Station*, 8:1-40.
- ³⁷BEN-TUVIA, A., 1966. Red Sea fishes recently found in the Mediterranean. *Copeia*, 2: 254-275.
- ³⁸BEN-TUVIA, A., 1973. Man-made changes in the eastern Mediterranean Sea and their effect on the fishery resources. *Marine Biology*, 19: 197-203.
- ³⁹BILECENOĞLU, M., 2002. Türkiye kıyılarında ilk kez rastlanan bir denizanası türü: *Cassiopea andromeda* (Forsskål, 1775) (Cassiopeidae, Scyphozoa). *Sualtı Dünyası Dergisi*, 72: 42-43.
- ⁴⁰BILECENOĞLU, M., 2003. Kızıldeniz Göçmeni Balon Balığı (*Torquigener flavimaculosus* Hardy & Randall, 1983), Türkiye kıyılarından ilk gözlemler. *Sualtı Dünyası Dergisi*, 74: 38-39.
- ⁴¹BILECENOĞLU, M., 2004. Occurrence of the Lessepsian migrant fish, *Sillago sihama* (Forsskål, 1775) (Osteichthyes: Sillaginidae), from the Aegean Sea. *Israel Journal of Zoology*, 50: 420-421.
- BILECENOĞLU, M. & KAYA, M., 2002. Growth of marbled spinefoot *Siganus rivulatus* Forsskål, 1775 (Teleostei: Siganidae) introduced to Antalya Bay, eastern Mediterranean Sea (Turkey). *Fisheries Research*, 54: 279-285.
- BILECENOĞLU, M. & TASKAVAK, E., 1999. Some observations on the habitat of Red Sea immigrant sweeper, *Pempheris vanicolensis*, at the Mediterranean coast of Turkey. *Zoology in the Middle East*, 17: 67- 70.
- ⁴²BILECENOĞLU, M., TASKAVAK, E. & KUNT, K. B., 2002. Range extension of three Lessepsian migrant fish (*Fistularia commersoni*, *Sphyraena flavicauda*, *Lagocephalus suezensis*) in the Mediterranean Sea. *Journal of the Marine Biological Association of the United Kingdom*, 82: 525-526.
- ⁴³BILECENOĞLU, M., TASKAVAK, E., MATER, S. & KAYA, M., 2002. Checklist of the marine fishes of Turkey. *Zootaxa*, 113: 1-194.
- BILECENOĞLU, M. & KAYA, M., 2006. A new alien fish in the Mediterranean Sea – *Platax teira* (Forsskål, 1775) (Osteichthyes: Ephippidae). *Aquatic Invasions*, 1: 80-83.
- BILECIK, N., 1990. Distribution of sea snail “*Rapana venosa* (V.)” in the Turkish coasts of Black Sea and its impact on Black Sea fisheries. *Tarım Orman ve Koyisleri Bakanlığı Su Ürünleri Arastirma Enstitüsü Müdürlüğü Yayınları*, seri B, 1, 34 pp.
- ⁴⁴BIZSEL, N. & BIZSEL, C., 2002. New records of toxic algae *Heterosigma cf. akashiwo* and *Gymnodinium cf. mikimotoi* in the hypereutrophic Izmir Bay (Aegean Sea): Coupling between organisms and water quality parameters. *Israel Journal of Plant Sciences*, 50: 33-44.
- ⁴⁵BLÖCHER, M., 1983. A dead specimen of *Palmadusta gracilis* (Gaskoin, 1849) found in the Mediterranean. *La Conchiglia*, 10: 166-167.
- ⁴⁶BOGI, C., BUZZURRO, G. & GREPPI, E., 1995. Presenza di *Murchisonella columna* (Hedley, 1907) Nel Mediterraneo Orientale. *Bollettino Malacologico*, 31: 69-76.
- ⁴⁷BUHAN, E., YILMAZ, H., MORKAN, Y., BÜKE, E. & YÜKSEK, A., 1997. Güllük ve Gökova Körfezleri için yeni bir av potansiyeli: *Scomberomorus commerson* (Lacepède, 1800) (Pisces-Teleostei). *Mediterranean Fisheries Congress*, Izmir: 937-944.
- ⁴⁸BUZZURRO, G. & GREPPI, E., 1994. Presenza di *Smaragdia (Smaragdella) souverbiana* (Montrouzier, 1836) nel Mediterraneo Orientale. *Bollettino Malacologico*, 29: 319-321.
- ⁴⁹BUZZURRO, G. & GREPPI, E., 1996. The lessepsian molluscs of Tasucu (South-East Turkey). *La Conchiglia*, 279 (supplement): 3-22.
- ⁵⁰BUZZURRO, G., ENGL, W. & TÜMTÜRK, I., 1995. Bivalven und Gastropoden

- der Europäischen Meer (4): *Ergalatax martensi* (Dall, 1923) (Muricidae), ein neuer lessepsischer einwanderer von der Türkischen Südküste. *Club Conchylia Informationen*, 27: 17-18.
- ⁵¹BUZZURRO, G., HOARAU, A., GREPPI, E. & PELORE, J., 2001. Prima segnalazione di *Hinemoa cylindrica* (de Folin, 1879) per il Mediterraneo. *Bollettino Malacologico*, 37: 23-26.
- ⁵²CIRIK, S., 1978. Recherches sur la végétation marine des côtes Turques de la mer Egée. Etude particulière des Peyssonneliacées de Turquie. *Thèse Doctora III, Cycle Biologie Végétale, Université Pierre et Marie Curie*, Paris: 172 pp.
- ⁵³CIRIK, S., 1986. A propos de la végétation marine de la baie d'Akkuyu, Mersin, Turquie. *5th OPTIMA Meeting*, Istanbul: 69-81.
- ⁵⁴CIRIK, S. & ÖZTÜRK, B., 1991. Notes sur la présence d'une forme rare du *Caulerpa racemosa*, en Méditerranée orientale. *Flora Mediterranea*, 1: 217-219.
- ⁵⁵CIRIK, S., ZEYBEK, N., AYSEL, V. & CIRIK, S., 1990. Note préliminaire sur la végétation marine de l'île Gökçeada (Mer Egée-Nord, Turquie). *Thalassographica*, 13: 33-37.
- ⁵⁶ÇEVİK, C. & ÖZTÜRK, B., 2001. A new lessepsian mollusc *Hypselodoris infucata* (Rüppell & Leuckart, 1828) (Gastropoda: Nudibranchia) for the coasts of Turkey. *Turkish Journal of Zoology*, 25: 27-30.
- ⁵⁷ÇEVİK, C., ÖZTÜRK, B. & BUZZURRO, G., 2001. The presence of *Crassostrea virginica* (Gmelin, 1791) and *Saccostrea commercialis* (Iredale & Roughley, 1933) in the eastern Mediterranean Sea. *La Conchiglia*, 298: 25-28.
- ⁵⁸ÇEVİK, C., DOGAN, A., ÖNEN, M. & ZENETOS, A., 2005. First record of the Indo-Pacific species *Electroma vexillum* (Mollusca: Bivalvia: Pterioida) in the eastern Mediterranean. *Journal of the Marine Biological Association of the United Kingdom*, 2, *Biodiversity Records*, 4996.
- ⁵⁹ÇEVİKER, D., 2001. Recent immigrant Bivalves in the northeastern Mediterranean off Iskenderun. *La Conchiglia*, 298: 39-46.
- ⁶⁰ÇEVİKER, D. & ALBAYRAK, S., 2002. New record of an exotic Bivalve from the southeastern coast of Turkey: *Scapharca inflata* (Reeve, 1844). *La Conchiglia*, 305: 56-60.
- ÇEVİKER, D. & ALBAYRAK, S., 2006. Three alien molluscs from Iskenderun Bay (SE Turkey). *Aquatic Invasions*, 1: 76-79.
- ÇINAR, M. E., 2003. Ecology of Syllidae (Annelida: Polychaeta) from northern Cyprus (Eastern Mediterranean Sea). *Bulletin of Marine Science*, 72: 795-811.
- ÇINAR, M. E., 2005. Polychaetes from the coast of northern Cyprus (eastern Mediterranean Sea), with two new records for the Mediterranean Sea. *Cahiers de Biologie Marine*, 46: 143-161.
- ⁶¹ÇINAR, M. E. & ERGEN, Z., 1998. Polychaetes associated with the sponge *Sarcotragus muscarum* Schmidt, 1864 from the Turkish Aegean Coast. *Ophelia*, 48: 167-183.
- ⁶²ÇINAR, M. E. & ERGEN, Z., 1999. Occurrence of *Prionospio saccifera* (Spionidae: Polychaeta) in the Mediterranean Sea. *Cahiers de Biologie Marine*, 40: 105-112.
- ⁶³ÇINAR, M. E. & ERGEN, Z., 2002. Faunistic analysis of Syllidae (Annelida: Polychaeta) from the Aegean Sea. *Cahiers de Biologie Marine*, 43: 171-178.
- ⁶⁴ÇINAR, M. E. & ERGEN, Z., 2005. Lessepsian migrants expanding their distributional ranges; *Pseudonereis anomala* (Polychaeta: Nereididae) in Izmir Bay (Aegean Sea). *Journal of the Marine Biological Associations of the United Kingdom*, 85: 313-321.
- ⁶⁵ÇINAR, M. E., ERGEN, Z. & DAGLI, E., 2002. Occurrence of the lessepsian species *Leonnatus persicus* (Polychaeta: Nereididae) in Izmir Bay, Aegean Sea. *Journal of the Marine Biological Associations of the United Kingdom*, 82: 811-815.

- ÇINAR, M. E., KATAGAN, T., ERGEN, Z. & SEZGIN, M., 2002. Zoobenthos inhabiting *Sarcotragus muscarum* (Porifera: Demospongia) from the Aegean Sea. *Hydrobiologia*, 482: 107-117.
- ⁶⁶ÇINAR, M. E., ERGEN, Z., DAGLI, E. & PETERSEN, M. E., 2005. Alien species of spionid polychaetes (*Streblospio gynobranchiata* and *Polydora cornuta*) in Izmir Bay, eastern Mediterranean. *Journal of the Marine Biological Associations of the United Kingdom*, 85: 821-827.
- ÇINAR, M. E., BILECENOĞLU, M., ÖZTÜRK, B. & CAN, A., 2006. New records of alien species on the Levantine coast of Turkey. *Aquatic Invasions*, 1: 84-90.
- ⁶⁷DEMİR, M., 1952-1954. Bogazlar ve adalar sahillerinin omurgasız dip hayvanları. *Istanbul Üniversitesi Fen Fakültesi Hidrobiyoloji Araştırma Enstitüsü Yayınlarından*, 3: 615 pp.
- ⁶⁸DEMİR, M., 1977. On the presence of *Arca (Scapharca) amygdalum* Philippi, 1847 in the harbour of Izmir, Turkey. *Istanbul Üniversitesi Fen Fakültesi Mecmuası*, Serie B, 42: 197-202.
- ⁶⁹DURAL, B., 1996. Ege Denizi Cyanophycaceae türleri. *Ege Üniversitesi Su Ürünleri Fakültesi Dergisi*, 12: 267-292.
- ⁷⁰DURAL, B. & AYSEL, V., 1996. A new record for the algal flora of Mediterranean 1. *Cladophora boodeoides* Börgesen. *Turkish Journal of Botany*, 20: 465-468.
- ⁷¹EKER, E., GEORGIEVA, L., SENICHKINA, L. & KIDEYS, A. E., 1999. Phytoplankton distribution in the western and eastern Black Sea in spring and autumn 1995. *ICES Journal of Marine Science*, 56 (Supplement): 15-22.
- ELTON, C. S., 1958. The ecology of invasions by animals and plants. The University of Chicago Press Chicago, 181 pp.
- ⁷²ENGL, W., 1995. Specie prevalentemente lessepsiane attestate lungo le coste Turche. *Bollettino Malacologico*, 31: 43-50.
- ⁷³ENGL, W. & ÇEVIKER, D., 1999. New migrant species from southeast Turkey *Psammotreta praerupta* (Salisbury, 1934) and *Antigona lamellaris* Schumacher, 1817. *La Conchiglia*, 290: 17-20.
- ⁷⁴ENZENROSS, R. & ENZENROSS, L., 1995. Erstnachweise indopazifischer Brachyura (Crustacea: Decapoda) von detürkischen Mittelmeerküste. *Stuttgarter Beiträge zur Naturkunde*, 521: 1-4.
- ⁷⁵ENZENROSS, R., ENZENROSS, L. & KOCATAŞ, A., 1992. Lessepsian migration: two Indo-Pacific crabs species (Crustacea, Decapoda) found on the Turkish Mediterranean coast. *Journal of the Faculty of Science Ege University*, 14: 3-10.
- ⁷⁶ENZENROSS, L., ENZENROSS, R. & NIEDERHÖFER, H. J., 1990. Wissenschaftlich interessante Funde aus der Sammlung Enzenross (Marine Invertebraten). *Jahreshefte der Gesellschaften fuer Naturkunde in Wuerttemberg*, 145: 283-294.
- ⁷⁷ERAZI, R. A. R., 1942. The heterosomata of the Bosphorus, the Golden Horn and the Sea of Marmara. *Revue de la Faculte des Sciences de l'universite d'Istanbul*, 7: 235-262.
- ⁷⁸ERAZI, R. A. R., 1943. *Leiognathus mediterraneus* nov. sp. *Compte Rendu Société Turquie des Sciences physiques et naturelles*, 10: 49-53.
- ⁷⁹ERGEN, Z., 1976. Investigations on the taxonomy and ecology of Polychaeta from Izmir Bay and its adjacent areas. *Scientific Reports of the Faculty of Science Ege University*, 209: 1-73.
- ⁸⁰ERGEN, Z., 1992. The latest status of Polychaeta in the soft substrate of Izmir Bay. *Rapport de la Commission Internationale Exploration de la Mer Méditerranée*, 33.
- ⁸¹ERGEN, Z. & ÇINAR, M. E., 1997. Polychaeta of Antalya Bay (Mediterranean coast of Turkey). *Israel Journal of Zoology*, 43: 229-241.
- ⁸²ERGEN, Z., ÇINAR, M. E. & UNSAL, M., 1998. Polychaetes from the Manavgat River Delta (Turkish Mediterranean coast). *Rapport de la Commission Internationale Exploration de la Mer Méditerranée*, 35: 536-537.

- ⁸³ERGEN, Z., ÇINAR, M. E., DAGLI, E. & KURT, G., 2002. Lessepsian polychaete species from the Turkish coasts. p. 50-55. In: *Proceedings of Workshop on Lessepsian Migration*, edited by B. Öztürk & N. Basusta, Turkish Marine Research Foundation, 9.
- ⁸⁴ERGEN, Z., ÇINAR, M. E., DAGLI, E. & KURT, G., 2006. Seasonal dynamics of soft-bottom polychaetes in Izmir Bay (Aegean Sea, eastern Mediterranean). *Scientia Marina*, in press.
- ⁸⁵ERGEV, M. B., ÇINAR, M. E., MUTLU, E. & ERGEN, Z., 2003. Ecological features of the lessepsian migrant *Leonnatus persicus* (Polychaeta: Nereididae) from the Levant coast of Turkey. *Journal of the Marine Biological Associations of the United Kingdom*, 83: 1225-1226.
- ⁸⁶ERTAN, Ö., TURNA, O. I. & CORMACI, M., 1998. A New Record for the Marine Algal flora of Turkey: *Caulerpa scalpelliformis* (Brown ex Turner) C. Agardh (Caulerpaceae, Chlorophyceae). *Turkish Journal of Botany*, 22: 285-287.
- ⁸⁷ERYILMAZ, L., ÖZULUG, M. & MERİÇ, N., 2003. The Smooth Pufferfish, *Sphoeroides pachygaster* (Müller & Troeschel, 1848) (Teleostei: Tetraodontidae), new to the Northern Aegean Sea. *Zoology in the Middle East*, 28: 125-126.
- ⁸⁸FALCHI, S., 1974. Molluschi di provenienza Indopacifica lungo le coste Turche. *Conchiglie*, Milano, 10: 89.
- ⁸⁹FISCHER, W., BAUCHOT, M.-L. & SCHNEIDER, M., 1987. Fiches FAO d'identification des espèces pour les besoins de la pêche. Méditerranée et mer Noire. Zone de pêche 37. FAO & EEC, Rome: 761-1530.
- ⁹⁰FISCHER-PIETTE, E., 1960. *Rapana bezoar* I. sur la côte Turquie de la mer Noire. *Hydrobiologie*, Serie B5, 1-2: 51.
- ⁹¹FRITSCH, K., 1899. Beitrag zur flora von Constantinopel. I. Kryptogamen. *Denkschriften der kaiserlichen Akademie der Wissenschaften zu Wien Math*, 68: 219-250.
- GALIL, B., 1986. Red Sea decapods along the Mediterranean coast of Israel, ecology and distribution. p. 179-183. In: *Environmental quality and Ecosystem stability*, edited by Z. Dubinsky & Y. Steinberg, Bar Ilan Pres.
- GALIL, B. S. & LÜTZEN, J., 1995. Biological observations on *Heterosaccus dollfusi* Boschma (Cirripedia: Rhizocephala), a parasite of *Charybdis longicollis* Leene (Decapoda: Brachyura), a Lessepsian migrant to the Mediterranean. *Journal of Crustacean Biology*, 15: 659-670.
- ⁹²GALIL, B. S., FROGLIA, C. & NOEL, P. Y., 2002. CIESM Atlas of Exotic Crustaceans in the Mediterranean. Volume 2: Crustacean Decapods and Stomatopods, F. Briand (ed). Monaco: CIESM Publishers.
- ⁹³GELDIAY, R., 1969a. A report on the collection of Natantia (Crustacea, Decapoda) along the coast of Turkey from the Eastern Mediterranean to the vicinity of Izmir. *Scientific Reports of Faculty of Science Ege University*, 74: 1-17.
- ⁹⁴GELDIAY, R., 1969b. Izmir Körfezinin baslica balıkları ve muhtemel invazionları. *Ege Üniversitesi Fen Fakültesi Monografileri*, Izmir, 135 pp.
- ⁹⁵GELDIAY, R. & KOCATAŞ, A., 1968. The species of Penaeidae family from the Mediterranean Coast of Turkey. *Scientific Reports of Faculty of Science, Ege University*, 58: 1-15.
- ⁹⁶GIUNCHI, L., MICALI, P. & TISSELLI, M., 2001. Report of two new lessepsian immigrants from Turkey. *La Conchiglia*, 298: 47-48.
- ⁹⁷GOKOGLU, M., BODUR, T. & KAYA, Y., 2003. First record of the Red Sea bannerfish (*Heniochus intermedius* Steindachner, 1893) from the Mediterranean Sea. *Israel Journal of Zoology*, 49: 324-325.
- ⁹⁸GOKOGLU, M., BODUR, T. & KAYA, Y., 2004. First records of *Hippocampus fuscus* and *Syngnathus rostellatus* (Osteichthyes: Syngnathidae) from the Anatolian coast (Mediterranean Sea). *Journal of*

- the Marine Biological Association of the United Kingdom*, 84: 1093-1094.
- GOLANI, D., 1998. Distribution of Lessepsian migrant fish in the Mediterranean. *Italian Journal of Zoology*, 65: 95-99.
- ⁹⁹GRIPPA G., 1982. First record of *Leucosia signata* Paulson, 1875 from the South-Western Turkish coast (Decapoda, Brachyura, Leucosidae). *Quaderni del Laboratorio di Tecnologia della Pesca. Ancona*, 3: 335-338.
- ¹⁰⁰GRUVEL, A., 1928. Répartition géographique de quelques crustacés comestibles sur les côtes d'Égypte et de Syrie. *Comptes rendus de la société de biogéographie*, 5: 45-46.
- ¹⁰¹GRUVEL, A., 1930. Les richesses marines et fluviales de la Syrie. *Exploration actuelle Avenir Revue Scientifique*, Paris, 68: 33-41.
- ¹⁰²GÜCÜ, A.C., BINGEL, F., AVSAR, D. & UYSAL, N., 1994. Distribution and occurrence of Red Sea fish at the Turkish Mediterranean coast - northern Cilician Basin. *Acta Adriatica*, 34: 103-113.
- ¹⁰³GÜNER, H., 1970. Ege Denizi sahil algları üzerinde taksonomik ve ekolojik araştırma. *Ege Üniversitesi Fen Fakültesi İlmi Raporlar Serisi*, 76: 1-77.
- ¹⁰⁴GÜNER, H. & AYSEL, V., 1977. Izmir Körfezi'nde bulunan bazı *Ulva* türleri (*Chlorophyta*) hakkında taksonomik araştırma. *Ege Üniversitesi Fen Fakültesi Dergisi*, Seri B, 1: 241-251.
- ¹⁰⁵GÜNER, H., AYSEL, V., SUKATAR, A. & ÖZTÜRK, M., 1985. Türkiye Ege Denizi Floraşı I. Mavi-Yesil, Yesil, Esmer Algler ve Kapalı Tohumlular. *Doga Bilim Dergisi*, A2, 9: 272-282.
- ¹⁰⁶HOLTHUIS, L. B., 1961. Report on a collection of crustacea decapoda and stomatopoda from Turkey and the Balkans. *Zoologische Verhandelingen*, 47: 1-67.
- ¹⁰⁷HOLTHUIS, L. B. & GOTTLIEB, E., 1956. Two interesting cabs (crustacea, decapoda, brachyura) from Mersin Bay, S.E. Turkey. *Zoologische Verhandelingen*, 21: 287-301.
- ¹⁰⁸ISINIBILIR, M., TARKAN A. N. & KIDEYS, A. E., 2004. Decreased levels of the invasive ctenophore *Mnemiopsis* in the Sea of Marmara in 2001. p. 155-165. In: *The Invasion of the Black, Mediterranean and Caspian Seas by the American Ctenophore, Mnemiopsis leidyi Agassiz: A Multidisciplinary Perspective*, edited by H. Dumont, T. Shiganova & U. Niermann, Kluwer Academic Publishers, Dordrecht, The Netherlands.
- ¹⁰⁹KATAGAN, T., 1983. Türkiye'nin Ege Denizi litoral kumase'leri üzerine sistematik ve ekolojik araştırmalar. *Ege University Faculty of Science Journal, Serie B*, 6: 9-18.
- ¹¹⁰KATAGAN, T., KOCATAŞ, A., ZENGİN, M. & ATES, A. S., 2004. An Indo-Pacific stomatopod from the Sea of Marmara: *Erugosquilla massavensis* (Kossmann, 1880). *Crustaceana*, 77: 381-383.
- ¹¹¹KAYA, M., MATER, S. & BENLİ, H. A., 1992. A new Indo-Pacific gobiid fish *Oxyurichthys papuensis* (Val., 1837) for eastern Mediterranean coasts of Turkey. *Rapport de la Commission Internationale Exploration de la Mer Méditerranée*, 33: 298.
- ¹¹²KAYA, M., MATER, S. & KORKUT, A. Y., 1998. A new grey mullet species "Mugil so-iuy Basilewsky" (Teleostei: Mugilidae) from the Aegean coast of Turkey. *Turkish Journal of Zoology*, 22: 303-306.
- KIDEYS, A. E., 2002. Fall and rise of the Black Sea ecosystem. *Science*, 297: 1482-1484.
- ¹¹³KIDEYS, A. E. & GÜCÜ, A. C., 1995. *Rhopilema nomadica*: A poisonous indo-pacific scyphomedusan new to the Mediterranean coast of Turkey. *Israel Journal of Zoology*, 41: 615-617.
- ¹¹⁴KIDEYS, A. E. & NIERMANN, U., 1994. Occurrence of *Mnemiopsis* along the Turkish coasts (from northeastern Mediterranean to Istanbul). *ICES Journal of Marine Science*, 51:423-427.
- ¹¹⁵KIDEYS, A. E., & ROMANOVA, Z.,

2001. Distribution of gelatinous macrozooplankton in the southern Black Sea during 1996-1999. *Marine Biology*, 139: 535-547.
- ¹¹⁶KINZELBACH, R., 1985. Lesseps'sche wanderung: neue stationen von muscheln (Bivalvia: Anisomyaria). *Archiv für Molluskenkunde*, 115: 273-278.
- ¹¹⁷KNIGHT-JONES, P., KNIGHT-JONES, W. & ERGEN, Z., 1991. Sabelliform polychaetes, mostly from Turkey's Aegean coast. *Journal of Natural History*, 25: 837-858.
- KOÇAK, F., ERGEN, Z. & ÇINAR, M. E., 1999. Fouling organisms and their developments in a polluted and an unpolluted marina in the Aegean Sea (Turkey). *Ophelia*, 50:1-20.
- ¹¹⁸KOCATAŞ, A., 1976. Tanaiidaces et Isopodes recueillis dans le golfe d'Izmir, Turquie. *Crustaceana*, 30: 68-72.
- ¹¹⁹KOCATAŞ, A., 1978. Contribution a l'étude des peuplements des horizons supérieurs de substrat rocheux du Golfe d'Izmir. *Scientific Monographs of the Faculty of Science, Ege University*, 12: 1-93.
- ¹²⁰KOCATAŞ, A., 1981. Liste préliminaire et répartition des Crustacea Decapodes des eaux Turques. *Rapport de la Commission Internationale Exploration de la Mer Méditerranée*, 27: 161-162.
- ¹²¹KOCATAŞ, A. & KATAGAN, T., 1978. Türkiye denizlerinin littoral bentik amphipod'lari ve yayilislari. *TÜBITAK Temel Bilimler Arastirma Grubu*, Proje No TBAG-223, 63 p.
- ¹²²KOCATAŞ, A. & KATAGAN, T., 1994. Impact of Lessepsian Decapod and Stomatopod species on the biodiversity of the Turkish Seas. *XII. Ulusal Biyoloji Kongresi*: 290-294.
- ¹²³KOCATAŞ, A., KORAY, T., KAYA, M. & KARA, O. F., 1993. Review of the fishery resources and their environment in the Sea of Marmara. *General Fisheries Council for The Mediterranean, Studies and Reviews*, 64: 87-143.
- ¹²⁴KORAY, T., 1984. The occurrence of red-tides and causative organisms in Izmir Bay. *Ege Üniversitesi Fen Fakültesi Dergisi*, Seri B, 1: 75-83.
- ¹²⁵KORAY, T., 1987. One-celled microplankton species of Izmir Bay (Aegean Sea): A species list and a comparison with the records of adjacent regions. *Doga, Turkish Journal of Biology*, 11: 130-146.
- ¹²⁶KOSSWIG, C., 1950. Erythraische fische im Mittelmeer und an der grenze der Agais. *Syllegomena Biologica, Festschrift Kleinschmidt*: 203 -212.
- ¹²⁷KOSSWIG, C., 1956. Facts and problems offered by the erythrean invaders into the Mediterranean. *Balık ve Balıkçılık*, 4: 31-37.
- ¹²⁸KUMLU, M., EROLDOGAN, O. T., AKTAS, M. & GÖÇER, M., 2002. New Records: A new shrimp record for the Turkish Seas: *Melicertus hator* (Burkenroad, 1959) (Penaeidae: Crustacea). *Israel Journal of Zoology*, 48: 246-247.
- ¹²⁹LINDNER, G., 1987. Interessante Schneckenfunde an der Südtürkischen Mittelmeerküste (Reisemitbringsel der Familie SCHMIDT, Feldkirchen). *Club Conchyliia Informationen*, 19 : 32-43.
- MACK, R. N., SIMBERLOFF, D., LONSDALE, W. M., EVANS, H., CLOUT, M. & BAZZAZ, F.A., 2000. Biotic invasions: causes, epidemiology, global consequences, and control. *Ecological Applications*, 10: 689-710.
- ¹³⁰MATER, S. & BILECENOĞLU, M., 1999. Türkiye deniz balıkları. p. 790-808. In: *Genel Zoocografya ve Türkiye Zoocografyası*, edited by A. Demirsoy, Meteksan Matbaası, Ankara.
- ¹³¹MATER, S. & KAYA, M., 1987. Türkiye'nin Akdeniz sularında yeni kaydedilen üç balık türü, *Sudis hyalina Rafinesque*, *Pelates quadrilineatus* (Bloch), *Apogon nigripinnis* Cuvier (Teleostei) hakkında. *Doga, Türk Zooloji Dergisi*, 11: 45-49.
- ¹³²MERIC, E., AVSAR, N. & BERGIN, F., 2004. Benthic foraminifera of eastern Aegean Sea (Turkey) Systematics and

- Autoecology. *Turkish Marine Research Foundation*, 18, 306 pp.
- ¹³³MERİÇ, E., AVSAR, N., BERGIN, F. & YOKES, B., 2002. The proliferation of *Amphistegina* (Lessepsian migrants) population at three-islands (Uçadalar, Antalya), a new observation from the Turkish Mediterranean coast. p. 27-34. In: *Proceedings of Workshop on Lessepsian Migration*, edited by B. Öztürk & N. Basusta, Turkish Marine Research Foundation, 9.
- ¹³⁴MICALI, P. & PALAZZI, S., 1992. Contributo Alla Conoscenza Dei Pyramidellidae Della Turchia, Con Segnalazione di Due Nuove Immigrazioni Dal Mar Rosso. *Bulletino Malacologico*, 28: 83-90.
- ¹³⁵MIENIS, H. K., 2004. New data concerning the presence of lessepsian and indo-pacific migrants among the molluscs in the Mediterranean Sea with emphasize on the situation in Israel. p. 117-131. In: *1st National Malacology Congress*, Izmir, edited by B. Öztürk & A. Salman, Turkish Marine Research Foundation Press.
- ¹³⁶MONOD, T., 1930. Über einige indopazifische Decapoden der Meeresfauna Syriens. *Zoologischer Anzeiger*, 92: 135-141.
- ¹³⁷MUTLU, E., BINGEL, F., GÜCÜ, A.C., MELNIKOV, V. V., NIERMANN, U., OSTR, N. A. & ZAIKA, V. E., 1994. Distribution of new invader *Mnemiopsis* sp. and the resident *Aurelia aurita* and *Pleurobranchia pileus* populations in the Black Sea in the years 1991-1993. *ICES Journal of Marine Science*, 51: 407-421.
- ¹³⁸NICOLAY, K. & ROMAGNA-MANOJA, E., 1983. *Strombus (Conomurex) decorus raybaudii* n. ssp. *La Conchiglia*, 174-175: 20-21.
- ¹³⁹NIEDERHÖFER, H., ENZENROSS, L & ENZENROSS, R, 1991. Neue Erkenntnisse über die Ausbreitung von "Lesseps'schen Einwanderern" (Mollusca) an der Türkischen Mittelmeerküste. *Club Conchylia Informationen*, 13: 94-108.
- OCCHIPINTI AMBROGI, A., 2001. Transfer of marine organisms: a challenge to the conservation of coastal biocoenoses. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 11: 243-251.
- ¹⁴⁰ÖKSNEBJERG, B., ENZENROSS, R. & ENZENROSS, L., 1997. First record of Rhizocephala (Crustacea: Cirripedia) from Turkish waters, with notes on Lessepsian migration. *Stuttgarter Beiträge zur Naturkunde, Serie A (Biologie)*, 55: 1-7.
- OKUDAN, E. S., YURDABAK, F., TUNCER, S., ERDEMIR, C. C. & AYSEL, V., 2002. The distribution of *Caulerpa racemosa* (Forskkål) J. Agard on the coast of Turkey. p. 46-49. In: *Proceedings of Workshop on Lessepsian Migration*, edited by B. Öztürk & N. Basusta, Turkish Marine Research Foundation, 9.
- ¹⁴¹ÖNEN, M., 1983. The qualitative and quantitative investigation of the macrobenthic fauna found in the soft substratum of the Urla Harbour. *Ege University Faculty of Science Journal*, 6: 29-39.
- ¹⁴²ÖNEN, M., ERGEN, Z., KATAGAN, T. & ÇINAR, M.E., 1998. Güllük Lagünü bentik organizmalarının mevsimsel dinamigi. *XIV Ulusal Biyoloji Kongresi*, Samsun, 2: 203-226.
- ÖZCAN, T., ÖZCAN, G. & AKYURT, I., 2003. İskenderun Körfezi kıyısındaki kentsel alanda yengeç tüketim ve satın alma davranışları üzerine bir araştırma. *XII. Ulusal Su Ürünleri Sempozyumu*: 546-549.
- ¹⁴³ÖZTIG, F., 1957. Erdek sahillerinin deniz vejetasyonu hakkında. *Türk Biyoloji Dergisi*, 7: 12-13.
- ¹⁴⁴ÖZTÜRK, B. & POUTIERS, J. M., 2005. *Fulvia fragilis* (Bivalvia: Cardiidae): a lessepsian mollusc species from Izmir Bay (Aegean Sea). *Journal of the Marine Biological Association of the United Kingdom*, 85: 351-356.
- ¹⁴⁵ÖZTÜRK, M., 1993. Türkiye'nin Ege ve Akdeniz Kıyılarındaki *Chordariales* ve *Sporochnales* (*Phaeophyta*) üyelerinin yayılımı ve taksonomisi. *Turkish Journal of Botany*, 17: 119-146.

- of Botany*, 17: 237-247.
- ¹⁴⁶ÖZTÜRK, M. & GÜNER, H., 1986. Türkiye'nin Ege ve Akdeniz Kıyılarındaki Ectocarpales (Phaeophyta) üyelerinin yayılımı ve taksonomisi. *Turkish Journal of Botany*, 10: 459-472.
- ÖZTÜRK, B. & CAN, A., 2006. Indo-Pacific Gastropod Species in the Levantine and Aegean Seas. *Aquatic Invasions*, 1: in press.
- ¹⁴⁷PALAZZI, S., 1993. Ci sono due *Zafra* in mediterraneo, ma come si chiamano? *Notiziario del CISMA*, 14: 13-18.
- POR, F. D., 1978. Lessepsian migration—the influx of Red Sea biota into the Mediterranean by way of the Suez Canal. Springer, Berlin, 228 pp.
- ¹⁴⁸QUATREFAGES, M. A., 1865. Histoire naturelle des Annélides marins et d'eau douce, Annélides et Géphyriens, 2. Sédentaire (Paris), 562 pp.
- ¹⁴⁹RULLIER, F., 1963. Les Annélides polychètes du Bosphore, de la Mer de Marmara et de la Mer Noire, en relation avec celles de la Méditerranée. *Extrait des Rapports et Procès-verbaux des réunions de la CIESM*, 17: 161-260.
- ¹⁵⁰SALMAN, A., 2002. New report of the loliginid squid *Sepioteuthis lessoniana* Lesson, 1830 in the Mediterranean. *Israel Journal of Zoology*, 48: 249-250.
- ¹⁵¹SALMAN, A., KATAGAN, T. & BOLETZKY, S. von, 1999. New Cephalopod molluscs in the eastern Mediterranean: previously unnoted species or recent migrants? *Vie et Milieu*, 49: 11-17.
- ¹⁵²SCHNIEBS, K., 2000. Erster Nachweis einer Stomatelliden-Art im Mittelmeer (Gastropoda: Prosobranchia). *Malakologische Abhandlungen*, 20: 99-100.
- ¹⁵³STOCK, J. H., 1962. Some Turkish pycnogonid records. *Entomologische Berichten*, 22: 218-219.
- ¹⁵⁴SWENNEN, C., 1961. On a collection of Opisthobranchia from Turkey. *Zoologische Mededelingen*, 38: 41-75.
- ¹⁵⁵TARKAN, A. N., ISINIBILIR, M. & ERK, M. H., 2001. Dominant zooplankton species in the waters of the northern GökÇeada. *Ulusal Ege Adaları 2001 Toplantısı Bildiriler Kitabı, Turkish Marine Research Foundation*, İstanbul: 59-70.
- ¹⁵⁶TASKAVAK, E., BILECENOĞLU, M., BASUSTA, N. & MATER, S., 2000. Occurrence of *Pteragogus pelycus* Randall, 1981 (Teleostei: Labridae) and *Petroscoirtes ancyloodon* Rüppell, 1838 (Teleostei: Blenniidae) on the eastern Mediterranean coast of Turkey. *Acta Adriatica*, 41: 53-58.
- ¹⁵⁷TOKLU, B. & SARIHAN, E., 2003. The Copepoda and Cladocera (Crustacea) Fauna along the Yumurtalık-Botas coastline in İskenderun Bay. *Ege University Journal of Fisheries and Aquatic Sciences*, 20: 63-68.
- ¹⁵⁸TRINGALI, L., 1994. Levant Sea: A new lessepsian host of the genus *Sticteulima* F. C. Laseron, 1955. *La Conchiglia*, 273: 57-58.
- ¹⁵⁹TRINGALI, L. & VILLA, R., 1990. Rinvenimenti malacologici dalle coste Turche (Gastropoda, Polyplacophora, Bivalvia). *Notiziario del CISMA*, 11: 33-41.
- ¹⁶⁰TURKOGLU, M. & KORAY, T., 2002. Phytoplankton species' succession and nutrients in the southern Black Sea (Bay of Sinop). *Turkish Journal of Botany*, 26: 235-252.
- ¹⁶¹TURNA, I., CORMACI, M. & FURNARI, G., 2000. First Record of *Botryocladia madagascariensis* G. Feldmann (Rhodymeniaceae, Rhodophyceae) from the Gulf of Antalya (Mediterranean coast of Turkey). *Plant Biosystems*, 134: 111-115.
- ¹⁶²ÜNAL, A., 1970. Türkiye sahilinde yetisen deniz alglerinin sistematigi, Ankara, 64 p.
- ¹⁶³ÜNAL, E., SHMELEVA, A. A. & KIDEYS, A. E., 2002. Three new species of *Acartia* (Copepoda, Calanoida) from the northeastern Levantine Basin. p. 35-45. In: *Proceedings of Workshop on Lessepsian Migration*, edited by B. Özturek & N. Basusta, Turkish Marine Research Foundation, 9.

- ¹⁶⁴ÜNAL, E., SHMELEVA, A. A., ZAGORODNYAYA, J. & KIDEYS, A. E., 2000. Zooplankton structure and copepod species of the Sea of Marmara in spring 1998. p. 450-460. In: *Proceedings book of symposium on Marmara Sea 2000*, edited by B. Öztürk, M. Kadiroglu & H. Öztürk, TUDAV, Istanbul, 5.
- ¹⁶⁵ÜNSAL, I. & d'HONDT, J. L., 1979. Contribution à la connaissance des bryozoaires marins de Turquie (Eurystomata et Cyclostomata). *Vie Milieu*, 28-29: 613-634.
- ¹⁶⁶ÜNSAL, S., 1992. Türkiye denizleri için yeni bir kefal balığı türü: *Mugil so-iuy* Basilewsky. *Doga, Turkish Journal of Veterinary and Animal Sciences*, 16: 427-432.
- ¹⁶⁷UYDAL, Z. & MUTLU, E., 1993. Preliminary note on the occurrence and biometry of Ctenophoran *Mnemiopsis leidyi* finally invaded Mersin Bay. *Doga, Turkish Journal of Zoology*, 12: 229-236.
- ¹⁶⁸UYDAL, Z., KIDEYS, A. E., SHMEL-EVA, A. A., ZAGORODNYAYA, J. A. & GUBANOVA, A. D. 2002. Checklist of copepods (Calonida and Podoplea) from the northern Levantine basin shelf waters. *Hydrobiologia*, 482: 15-21.
- ¹⁶⁹WHITEHEAD, P. J. P., 1984. Clupeidae. p. 268-281. In: *Fishes of the North-eastern Atlantic and the Mediterranean*, edited by P. J. P. Whitehead, M.-L. Bauchot, J.-C. Hureau, J. Nielsen & E. Tortonese, UNESCO, Paris.
- ¹⁷⁰YOKES, B & GALIL, B. S., 2004. New records of alien decapods from the south-western coast of Turkey. *Rapport de la Commission Internationale Exploration de la Mer Méditerranée*, 37: 556.
- ¹⁷¹YOKES, B. & RUDMAN, W. B., 2004. Lessepsian opisthobranchs from south-western coast of Turkey; five new records for Mediterranean. *Rapport de la Commission Internationale Exploration de la Mer Méditerranée*, 37: 557.
- ¹⁷²YOKES, B., DERVISOGLU, R. & KARACIK, B., 2002. Likya kiyilarında denizel biyolojik zenginlik arastirmasi. *Sualti Bilim ve Teknoloji Toplantisi Bildiriler Kitabi*, Istanbul, p.166-181.
- ¹⁷³ZAITSEV, Y. & ÖZTÜRK, B., 2001. Exotic species in the Aegean, Marmara, Black, Azov and Caspian Seas. *Turkish Marine Research Foundation*, Istanbul, 267 pp.
- ¹⁷⁴ZEYBEK, N., 1969. Türkiye'nin Akdeniz algleri 1. Bodrum-Finike Körfezi sahil boyu. 2. Ege Denizi-Edremit Saros Körfezi-Sile. TBAG-124 nolu proje.
- ¹⁷⁵ZEYBEK, N., 1973a. Meeresalgen aus der Turkei. 1. Die Buchten von Edremit und Saros am Aegaeischen meer, 2. Die küste von Igneada bis Sile am Schwarzen meer. *Verhandl der Schweizerischen naturforschenden Gesellschaft*: 95-100.
- ¹⁷⁶ZEYBEK, N., 1973b. Türkiye'nin deniz algleri. *IV Biyoloji Kongresi*, Ankara, 214: 1-3.
- ¹⁷⁷ZEYBEK, N. & GÜNER, H., 1973. Çanakkale Bogazi ve Bozcaada deniz algleri. *Ege Üniversitesi Fen Fakultesi İlmi Raporlar Serisi*, 145: 1-19.
- ¹⁷⁸ZEYBEK, N., GÜNER, H. & AYSEL, V., 1986. The marine algae of Türkiye *Proceed. 5th OPTIMA Meeting*, Istanbul: 169-197.
- ZIBROWIUS, H., 1992. Ongoing modification of the Mediterranean marine fauna & flora by the establishment of exotic species. *Mésogée*, 51: 83-107.