

Mediterranean Marine Science

Vol 6, No 2 (2005)



Update of marine alien species in Hellenic waters

M.A. PANCUCCI-PAPADOPOULOU, A. ZENETOS, M.
CORSINI-FOKA, C.Y. POLITOU

doi: [10.12681/mms.188](https://doi.org/10.12681/mms.188)

To cite this article:

PANCUCCI-PAPADOPOULOU, M., ZENETOS, A., CORSINI-FOKA, M., & POLITOU, C. (2005). Update of marine alien species in Hellenic waters. *Mediterranean Marine Science*, 6(2), 147-158. <https://doi.org/10.12681/mms.188>

Update of marine alien species in Hellenic waters

M.A. PANCUCCI - PAPADOPOULOU, A. ZENETOS,
M. CORSINI - FOKA and C.-Y. POLITOU

Hellenic Centre for Marine Research, Institute of Oceanography
P.O Box 712, 19013 Anavissos, Attica, Greece

e-mail: apan@ath.hcmr.gr

Abstract

The list of marine alien species in Hellenic waters is updated taking into account new findings (published and unpublished data). According to the present work, the number of these species has increased from 90 (known until end 2003) to 128. Most of them are zoobenthic species followed by fish and macroalgae. An interannual analysis revealed an important increase of alien species during the last years. The study of their geographic distribution showed that their majority is present in the southeastern Aegean. More than 55% of them are well established, while about 40% are casual records. Their main pathway of introduction seems to be the Suez Canal followed by shipping, whereas the Straits of Gibraltar, aquaculture and the Straits of Dardanelles appear to play a less important role in their invasion of Hellenic waters. These findings are discussed considering environmental and anthropogenic factors.

Keywords: Alien marine species; Greece; Ionian; Aegean; Mediterranean; Lessepsian migration.

Introduction

It is now accepted that during the last decades the marine biota of the Eastern Mediterranean Sea is experiencing severe changes, due to the introduction of alien species. Even though biotic invasion via ballast waters, fouling, aquaculture, accidental introduction etc. play a significant role in the Eastern Basin, the main pathway of alien species' spreading remains the Suez Canal. Despite impediments such as the canal's length, shallowness, current regime, temperature and salinity extremes, hundreds of species of Indo-Pacific origin (the so-called "Lessepsian immigrants") have traversed the canal and settled

in the Eastern Mediterranean, an ongoing phenomenon well documented by POR (1978), ZIBROWIUS (1992), STREFTARIS *et al.* (2005). After a series of scattered publications dealing with single species or groups (KEVREKIDES & GALIL, 2003; CORSINI *et al.*, 2005; ZENETOS *et al.*, 2005; SIMBOURA & ZENETOS, 2005), the first attempt to compile available information on alien species recorded in Hellenic waters is an inventory prepared by PANCUCCI-PAPADOPOULOU *et al.* (2005) which takes into account records up to 2003. Two years later, with a plethora of new findings and the contribution of more experts in the field, it is considered necessary to revise the previous inventory.

The present effort comes to elucidate any discrepancy with the previous inventory, adding new records and discussing questionable cases.

Methods

The basis of the present work is the checklist of alien species in Hellenic waters reported by PANCUCCI *et al.* (2005). However, this has been revised on the basis of the annotated list of ZENETOS *et al.* (this volume), according to which some species in the checklist have been classified by Mediterranean experts as “questionable” or “excluded”, while some others previously excluded, have been added. Additional records of species sighted in the period 2004-2005, published or recorded in the grey literature, as well as species belonging to taxonomic groups not considered previously and unpublished records are included. Grey literature includes HCMR and EU technical reports, 2005 Scientific Congresses, personal communication by specialists, website (seaslug forum, algae-base).

The distribution of each taxon in the Hellenic Seas was established from the scrutiny of more than 100 references. Only some references of interest plus the more recent are cited due to space limitations.

All calculations are based on species’ records up to December 2005. Alien species have been grouped into five broad categories covering ecofunctional groups: phytoplankton (PP), phytobenthos (PB), zooplankton (ZP), zoobenthos (ZB) and fish (F). Cryptogenic species are not included in our compiled list.

According to their establishment success alien species have been classified into four main categories, namely established, casual, questionable and excluded, in the sense described in ZENETOS *et al.* (this volume). Excluded species are not counted in calculations, while questionable which may turn out to be established, are taken into account.

Results

A. Diversity of Alien Taxa

The total number of aliens recorded/brought to our attention until December 2005 reaches 140 species (Table 1) of which 7 are questionable and 13 excluded, according to the screening procedure described in detail in ZENETOS *et al.* (this volume). The majority of aliens is represented by zoobenthos (63 species) followed by fish (32 species) and macroalgae (16 species) (Fig. 1).




Fig. 1: The contribution of main marine groups of aliens in Hellenic waters.

Table 1

List of aliens per group.

Asterisk denotes new entries

ES: Establishment success (E: Established, C: Casual, Q: Questionable, Ex: Excluded)

Fish	ES	<i>Asparagopsis armata</i>	E
* <i>Etrumeus teres</i>	E	* <i>Caulerpa racemosa</i>	E
* <i>Iniistius pavo</i>	C	* <i>Ceramium bisporum</i>	Ex
* <i>Lagocephalus sceleratus</i>	E	* <i>Chondria collinsiana</i>	C
* <i>Lagocephalus suezensis</i>	E	* <i>Chondria polyrhiza</i>	C
* <i>Petroscirtes ancyllodon</i>	C	<i>Codium fragile tomentosoides</i>	E
* <i>Seriola fasciata</i>	C	<i>Colpomenia peregrina</i>	C
* <i>Tylerius spinosissimus</i>	C	* <i>Grateloupia subpectinata</i> (= <i>G. filicina</i> var. <i>luxurians</i>)	C
* <i>Tylosyrus crocodilus</i>	C	<i>Halophila stipulacea</i>	E
* <i>Upeneus pori</i>	C	* <i>Hypnea esperi</i>	Ex
<i>Alepes djedaba</i>	C	* <i>Hypnea spinella</i> (= <i>H. cervicornis</i>)	C
<i>Apogon pharaonis</i>	E	* <i>Hypnea valentiae</i>	C
<i>Atherinomorus lacunosus</i>	C	* <i>Laurencia okamurae</i> (= <i>L. coronopus</i>)	Ex
<i>Callionymus filamentosus</i>	C	* <i>Lophocladia lallemandii</i>	C
<i>Enchelycore anatina</i>	C	* <i>Neosiphonia sphaerocarpa</i>	C
<i>Fistularia commersonii</i>	E	* <i>Polysiphonia fucoides</i> (= <i>P. nigrescens</i>)	Ex
<i>Gaidropsarus granti</i>	Q	* <i>Sarconema filiforme</i>	Ex
<i>Hemiramphus far</i>	E	<i>Sarconema scinaoides</i>	E
<i>Lagocephalus spadiceus</i>	E	<i>Styopodium schimperi</i>	E
<i>Leiognathus klunzingeri</i>	E	* <i>Womersleyella setacea</i>	E
<i>Mugil soiuy</i> = <i>Liza haematocheila</i>	C		
<i>Parexocoetus mento</i>	E		
<i>Pempheris vanicolensis</i>	E		
<i>Pteragogus pelycus</i>	E		
<i>Sargocentron rubrum</i>	E		
<i>Saurida undosquamis</i>	E		
<i>Siganus luridus</i>	E		
<i>Siganus rivulatus</i>	E		
<i>Sphoeroides pachygaster</i>	E		
<i>Sphyraena chrysotaenia</i>	E		
<i>Sphyraena flavicauda</i>	C		
<i>Stephanolepis diaspros</i>	E		
<i>Upeneus moluccensis</i>	E		
Phytobenthos (PB)		Phytoplankton (PP)	
* <i>Acanthophora nayadiformis</i>	C	* <i>Alexandrium insuetum</i>	E
* <i>Acrothamnion preissii</i>	Ex	* <i>Alexandrium taylori</i>	E
		<i>Alexandrium tamarense</i>	Ex
		* <i>Histioneis detonii</i>	C
		* <i>Gymnodinium mikimotoi</i>	E
		* <i>Gymnodinium</i> (= <i>Karenia</i>) <i>breve</i>	E
		* <i>Gymnodinium catenatum</i>	E
		* <i>Gymnodinium fusus</i>	Ex
		* <i>Phaeocystis pouchettii</i>	E
		* <i>Prorocentrum mexicanum</i>	E
		* <i>Rhizosolenia alata</i>	Ex

(continued)

(continued)

Zoobenthos (ZB)

Crustacea

* <i>Balanus trigonus</i>	E
* <i>Calappa pelii</i>	C
* <i>Leucosia signata</i>	C
* <i>Megabalanus tintinnabulum</i>	Q
* <i>Myra subgranulata</i>	C
* <i>Percnon gibbesi</i>	E
* <i>Stenothoe gallensis</i>	E
<i>Alpheus rapacida</i>	C
<i>Callinectes sapidus</i>	E
<i>Charybdis longicollis</i>	E
<i>Erugosquilla massavensis</i>	E
<i>Ixa monodi</i>	C
<i>Marsupenaeus japonicus</i>	E
<i>Metapenaeopsis aegyptia</i>	E
<i>Metapenaeopsis mogiensis consobrina</i>	E
<i>Portunus pelagicus</i>	E
<i>Thalamita poissonii</i>	E
<i>Trachysalambria palaestinensis</i>	E

Mollusca

* <i>Aplysia dactylomeda</i>	E
<i>Acteocina mucronata</i>	C
<i>Anadara demiri</i>	E
<i>Brachidontes pharaonis</i>	E
<i>Bulla ampulla</i>	E
<i>Bursatella leachi</i>	E
<i>Cellana rota</i>	C
<i>Crassostrea gigas</i>	E
<i>Crepidula fornicata</i>	E
<i>Cylichna girardi</i>	C
<i>Fulvia fragilis</i>	E
<i>Gastrochaena cymbium</i>	C
<i>Haminoea cyanomarginata</i>	C
<i>Malvufundus regulus</i>	C
<i>Melibe fimbriata</i>	E
<i>Murex forskoeihli</i>	Q
<i>Mya arenaria</i>	C
<i>Nerita sanguinolenta</i>	C

<i>Petricola pholadiformis</i>	C
<i>Pinctada radiata</i>	E
<i>Pleurobranchus forskali</i>	C
<i>Polycerella emertoni</i>	C
<i>Pseudochama corbieri</i>	C
<i>Rapana rapiformis</i>	Q
<i>Rapana venosa</i>	C
<i>Smaragdia souverbiana</i>	Q
<i>Strombus persicus</i>	E
<i>Trochus erythraeus</i>	E

Polychaeta

* <i>Desdemona ornata</i>	E
* <i>Ficopomatus enigmaticus</i>	E
* <i>Hydroides dianthus</i>	E
<i>Branchiosyllis exilis</i>	Ex
<i>Cossura coasta</i>	Q
<i>Hydroides elegans</i>	E
<i>Lysidice collaris</i>	Q
<i>Metasychis gotoi</i>	E
<i>Notomastus aberans</i>	C
<i>Paradyte cf. crinoidicola</i>	C
<i>Prionospio pulchra</i>	C

<i>Prionospio salzi</i>	Ex
<i>Scoloplos chevalieri candiensis</i>	Ex
<i>Spirobranchus tetracerus</i>	C
<i>Spirorbis marioni</i>	C
<i>Timarete anchylochaeta</i>	Ex

Miscellanea

* <i>Oculina patagonica</i>	E
* <i>Synaptula reciprocans</i>	C
<i>Aspidosiphon mexicanus</i>	C
<i>Hippopodina feegeensis</i>	C
<i>Ophiactis savignyi</i>	C

Zooplankton (ZP)

* <i>Paracartia grani</i>	E
* <i>Arietellus pavoninus</i>	E
<i>Calanopia elliptica</i>	C
<i>Centropages furcatus</i>	C
<i>Mnemiopsis leidyi</i>	E
<i>Pseudocalanus elongatus</i>	E
<i>Cassiopeia andromeda</i>	C

Table 2
Alien species sighted since 2003.

Group	Species latin name	Year of introduction	Area	Reference
Fish	<i>Etrumeus teres</i>	2003	Kyklades: Paros, Naxos, Ios, N.Kriti	CORSINI <i>et al.</i> , 2005 KALLIANIOTIS & LEKKAS, 2005
	<i>Lagocephalus suezensis</i>	2003	Rodos	CORSINI <i>et al.</i> , in press
	<i>Tylosyrus crocodilus</i>	2003	Chalkidiki	SINIS, 2005
	<i>Upeneus pori</i>	2003	Rodos	CORSINI <i>et al.</i> , 2005
	<i>Iniistius pavo</i>	2004	Rodos	CORSINI <i>et al.</i> , in press
	<i>Petroscirtes ancylodon</i>	2004	Rodos	CORSINI <i>et al.</i> , 2005
	<i>Seriola fasciata</i>	2004	Rodos	CORSINI <i>et al.</i> , in press
	<i>Tylerius spinosissimus</i>	2004	Rodos	CORSINI <i>et al.</i> , 2005
	<i>Lagocephalus sceleratus</i>	2005	Rodos, Symi, N. Kriti	CORSINI <i>et al.</i> , 2005
Zoobenthos				
Crustaceans	<i>Myra subgranulata</i>	2004	Rodos	CORSINI & KONDILATOS, in press
	<i>Percnon gibbesi</i>	2004	Messiniakos, Kriti, Rodos	KAMBOUROGLOU <i>et al.</i> , in press
	<i>Leucosia signata</i>	2005	Rodos	CORSINI <i>et al.</i> , in press
	<i>Calappa pelii</i>	2005	Saronikos, N. Ionian	POLITOU, unpublished
Molluscs	<i>Aplysia dactylomeda</i>	2005	Messiniakos	<i>Sea slug website</i>
Cnidaria	<i>Oculina patagonica</i>	2005	Saronikos	SALOMIDI <i>et al.</i> , in press
Echinoderms	<i>Synaptula reciprocans</i>	2005	Rodos	ZIBROWIUS, unpublished

Thirty eight new records are added to the inventory of PANCUCCI *et al.* (2005), 16 of which were sighted in the period 2003-2005 (Table 2) and published within 2005 or are as yet unpublished. Among the remaining 22 species, 14 are earlier records hidden in grey literature or previously excluded because of some uncertainties. These include: the copepods *Paracartia grani*, *Arietellus pavoninus*, the polychaetes *Desdemona ornata*, *Ficopomatus enigmaticus* and *Hydroides dianthus* and many macroalgae (phytobenthos). The

remaining 8 are species of phytoplankton, which was not treated in the previous inventory. These are: *Alexandrium insuetum*, *Alexandrium taylori*, *Gymnodinium (=Karenia) breve*, *Gymnodinium mikimotoi*, *Histioneis detonii*, *Phaeocystis pouchettii*, *Prorocentrum mexicanum* and *Gymnodinium catenatum*.

As seen in Table 2, the additional records belong mostly to fish (9) and zoobenthos (7) which indicate that they are the most studied groups, as opposed to phytobenthos and zooplankton that are studied less.




Fig. 2: Chronological trends of aliens in Hellenic waters (ZB: zoobenthos, ZP: zooplankton, PB: phytobenthos, PP: phytoplankton and F: fish).

B. Chronological trends

Compared to the 17 species reported by POR (1978) in Hellenic waters about 100 years after the opening of the Suez Canal, a vast increase of alien species during the last two decades is noted. An increasing trend in the occurrence of aliens (Fig. 2) that started in the decade 1980-1990 coincides with the revival of Hellenic taxonomists mainly in zooplanktonic and zoobenthic groups. This continues in the next decade, when not only Hellenic but the world scientific interest is increased in aliens, and culminates in the present decade. It is worth noticing that intensive research during the last three years has revealed 16 new entries (Table 2).

C. Geographic distribution

Figure 3 shows aliens' zoogeographical patterns in Hellenic waters. The bulk is present in the southeastern Aegean, their presence decreasing significantly northwards and westwards. Ninety eight out of 126 alien species nowadays accepted as casual or established have been reported from the South-Central Aegean Sea (south of 38° parallel). Indeed, 69 aliens are known from the eastern part of the South Aegean (Dodekanisos Islands), with a peak of

63 along the NW coast of Rodos Island, while only 34 have been recorded from the rest of the South Aegean. This testifies the importance of the area as the main pathway of lessepsian immigrants spreading within the Mediterranean. Particular notice should be given to the 34 species found in the restricted area of the Saronikos and Argolikos Gulfs, closely related to the route of ships towards Peiraias, the biggest Hellenic port. A marked decrease of the aliens appears northwards, with only 36 species recorded in the North Aegean, 14 of them present in the Bay of Thessaloniki, where the second major port of Hellas is located. In Thessaloniki Bay the majority of species are those coming from the Black Sea or transported in ballasts, as opposed to the dominance of lessepsian immigrants observed in the South Aegean. Distribution of alien species in the Ionian coasts is even more restricted (30 species), mostly introduced via ballasts or fouling, with only a few, mainly fishes, being of Indo-Pacific origin.

D. Establishment success

Twenty species are considered as "questionable" or "excluded" in Table 1, mainly due to the uncertainty of their real taxonomical status




Fig. 3: Patterns of alien species distribution in Hellenic waters.

and distributional patterns. Of the 126 species (including 7 questionable records), 69 (55%) are well established, being locally abundant and even exploited in Hellenic waters and 50 (40.3%) consist of casual records reported sporadically from one or two locations. The worst invasive species in terms of spread and impact are encountered among molluscs (*Pinctada radiata*, *Strombus persicus*), crustaceans (*Marsupenaeus japonicus*, *Charybdis longicollis*, *Erugosquilla massavensis*), fish (*Siganus luridus*, *Siganus rivulatus*, *Sphyrna chrysotaenia*, *Fistularia commersonii*) and macroalgae (*Asparagopsis armata*, *Caulerpa racemosa*, *Codium fragile*).

E. Pathways of introduction

Figure 4 shows the percentage of invasion pathways in Hellenic waters. The most important way appears to be the Suez Canal opening (60%), while the second is the accidental transportation via shipping (25%). The gradual introduction from the western Mediterranean is also significant (5%), while the introduction from the Black Sea via Dardanelles is very limited.




Fig. 4: Pathways of introduction of alien species into Hellenic waters.

E1. Introduction of alien species in ballasts/fouling

Thirtyfour species are assumed to have been transferred via shipping. Ship borne invaders are represented by phytoplankton (5 species), zooplankton (2 species), phytobenthos (7), and zoobenthos (19) [Cnidaria (1), Cirripedia (2),

Mollusca (8), Polychaeta (4), Amphipoda (1), Decapoda (2), Hydrozoa (1)].

A review of the alien biota recorded in the broader area of the major Hellenic ports (Peiraias, Thessaloniki) that is, in the inner Saronikos and Thermaikos Gulfs (HCMR technical reports, SIOKOU-FRANGOY, 1999; VARDA-LA-THEODOROU, 1999; ZENETOS, 1994; ZENETOS *et al.*, 2002) has revealed the presence of 32 and 14 species, respectively, of which only 5 are common. These are: the dinoflagellate *Gymnodinium mikimotoi*, the nanoflagellate *Phaeocystis pouchettii*, the copepod *Paracartia grani*, the polychaete *Metasychis gotoi* and the mollusc *Crepidula fornicata*.

The presence of *Strombus persicus*, which was earlier attributed to shipping in the eastern Mediterranean (Iskenderun) (ZENETOS *et al.*, 2003) and subsequently in Hellas, is now argued. Its wide expansion in Rodos and many Hellenic areas including the Saronikos Gulf (the Peiraias port broader area) is rather due to its invasive character. Therefore, Lessepsian migration appears to be its means of transportation. Similarly, molecular studies of *Pinctada radiata* populations in the Saronikos (ZENETOS *et al.*, 2004) have rather excluded shipping as mode of transportation.

E2. Introduction of alien species through fish/shellfish imports

It is unknown when the Pacific oyster *Crassostrea gigas* was imported for mariculture in Hellas. Although it is not currently cultured, self maintained populations have been recorded in the Ionian Sea. Similarly, the pearl oyster *Pinctada radiata* was imported as early as 1963, but its cultivation did not prove prosperous and was thus abandoned. However, the species was naturalised and expanded in the Hellenic Seas. (ZENETOS *et al.*, 2003).

The So-iuy mullet (*Mugil soiuy*) has been introduced for exploitation in the Northern Aegean Sea and is commercially exploited.

The following three species, which are en-

countered in mariculture areas, are considered accidental introductions: *Grateloupia subpectinata*, *Prionospio pulchra* and *Hydroides dianthus*.

Discussion

Intensive research and data mining the last years have increased the number of aliens in Hellenic waters from 90 (end 2003, PANCUCCI-PAPADOPOULOU *et al.*, 2005) to 126 (present work) and in particular from 15 records of ship transported species reported in 2002 (ZENETOS *et al.*, 2002), to 34. Intensive work in Hellenic ports, currently in progress, has brought to light more species carried among hull fouling (unpublished data).

Introductions of alien species are still going on and a global increase in the number of successful colonisation events is being observed (GRUSZKA, 1999). This is certainly true for the biological invasion in Hellenic waters. After a period of “gradual” spread of alien species via the Suez Canal to the eastern basin, it seems that during the last years a significant increase has been occurring, which reflects a new order of things in the area. This can be related to the adaptation time of “invaders” to their new habitat or, more simply, to the increased interest of the scientific community during the two last decades. It is, however, undeniable that Indo-Pacific biota records are continuously increasing along Hellenic coasts. The significant increase observed in the rate of introduction in the South Aegean could be related to the sampling depth, as it is known that settlement of Erythrean species is favoured in shallow areas with warm waters and the attention of Hellenic scientists has only recently been turned to very shallow coastal benthos. On the other hand, we cannot exclude the possibility of a gradual ‘colonisation’ of this ‘hotspot’ region due to climatic changes.

POR (1990) predicted that the distribution of Erythrean immigrants “will certainly expand or shrink according to the climatic evolution in the area” and that “immigrants will expand further

west if the climate warms up further". Moreover, the occurrence and spread of thermophilic species in the Mediterranean, either indigenous or alien, the so-called tropicalization of the Mediterranean has been attributed to the combination of four factors, namely: the Atlantic flux, lessepsian migration, human action and present climate warming (BIANCHI & MORRI, 2003).

Regarding the South Aegean, during 1988-1992 anomalies in meteorological parameters (very cold winters; dry years) were observed, in association with changes in the water mass pathways and a 1-4 fold increase in salt transport from the Levantine into the Aegean in the upper 200 m layer between 1987 and 1994 (THEOCHARIS *et al.*, 1999). The increased outflow of the newly formed denser water through the Cretan Arc straits into the eastern Mediterranean was compensated for by inflowing Levantine surface and intermediate waters (WU *et al.*, 2000). Significant changes resulted in the south Aegean water mass characteristics, which considerably influenced the thermohaline circulation of the eastern Mediterranean, termed the Eastern Mediterranean climatic Transient (EMT).

Although there is no documentation of direct competition between Erythrean and indigenous species, there are many instances of sudden changes in abundance where competition is suspected. Thus, if the present characteristics of the south Aegean hydrography persist, a modification of the composition and structure of the fauna is entirely possible.

The geographic distribution of alien species in the three major subareas of Hellenic waters (South, North Aegean, Ionian) clearly reflects the main hydrographic features of the subareas and shows that with the exception of those brought by shipping, the percentage of alien species in the North Aegean is almost negligible. This indicates that some natural "frontiers" still exist and may prevent the invasion/establishment of undesirable species. All that is needed is effective legislation considering all pathways associated. Because, although it is unrealistic to

stop a physical (or man-mediated phenomenon like Lessepsian migration), it is feasible to stop invasion from other sources.

In conclusion, it seems that, apart from the increasing interest of the scientific community in the phenomenon, a gradual warming of the area is occurring over the last years, resulting in more favourable climatic conditions for the establishment of new invaders.

Addendum

While this paper was accepted for publication two more species were recorded bringing the total number of marine aliens to 128. The new findings are: the polychaete *Pseudonereis anomala* (KAMBOUROGLOU & NICOLAIDOU, 2006) and the decapod *Charybdis helleri* (KIRMITZOGLOU *et al.*, 2006).

Acknowledgements

We would like to thank many colleagues who have contributed in various ways to this work. In particular, thanks are due to Dr. H. Zibrowius (Marseille, France) and Dr. A. Athanassiades (Gotenberg, Sweden) for providing information on unpublished species, and Dr. P. Kasapidis, Dr. A. Magoulas, Dr. I. Siokou-Frangou, Dr. O. Gotsis-Skrettas, Dr. K. Pagou, and Dr. P. Panayiotides (all of HCMR) for sharing their experience on the occurrence of fish, zooplankton, phytoplankton and phytobenthos, respectively.

References

- ALGAEBASE website <http://www.algaebase.org>.
- BIANCHI, C.N. & MORRI, C., 2003. Global Sea warming and tropicalization" of the Mediterranean Sea: biogeographic and ecological aspects. *Biogeographia*, 24: 319-328.
- CORSINI, M., MARGIES, P., KONDILATOS, G. & ECONOMIDIS, P. S., 2005. Lessepsian migration in Aegean Sea: *Tylerius spinosissimus* (Pisces, Tetraodontidae) new for the Mediterranean, and six more fish records

- from Rhodes. *Cybium*, 29 (4): 347-354.
- CORSINI, M. & KONDILATOS, G., in press. On the occurrence of two brachyurans, *Myra subgranulata* and *Herbstia kondyliata*, on Rhodes Island (SE Aegean Sea). *Crustaceana*, 2006.
- CORSINI-FOKA, M., MARGIES, P. & SANTORINIOS, E., in press. First record of the exotic brachyuran *Leucosia signata* from Rhodes. *Proceedings of the 8th Pan-Hellenic Symposium of Oceanography & Fisheries, Thessaloniki, May 2006*, 2006.
- CORSINI, M., MARGIES, P., KONDILATOS, G. & ECONOMIDIS, P. S., in press. Three new exotic fish records from the SE Aegean Greek waters. *Scientia Marina*, 2006.
- GRUSZKA, P., 1999. The River Odra Estuary as a Gateway for Alien Species Immigration to the Baltic Sea Basin. *Acta hydrochimica et Hydrobiologica*, 27: 374-82.
- KALLIANIOTIS, A. & LEKKAS, V., 2005. First documented report on the Lessepsian migrant *Etrumeus teres* De Kay, 1842 (Pisces: Clupeidae) in the Greek Seas. *Journal of Biological Research*, 4: 225-229.
- KAMBOUROGLOU, V., CORSINI-FOKA, M., KOURAKLIS, P., DOUNAS, K., NICOLAIDOU, A., THESSALOU-LEGAKI, M. & ZENETOS, A., CORSINI, M., MARGIES, P., KONDILATOS, G. & ECONOMIDIS, P. S., in press. Three new exotic fish records from the SE Aegean Greek waters. *Scientia Marina*, 2006. A new colonizer in the Hellenic seas: the brachyuran *Percnon gibbesi*. *Proceedings of the 8th Pan-Hellenic Symposium of Oceanography & Fisheries, Thessaloniki, May 2006*.
- KAMBOUROGLOU, V. & NICOLAIDOU A., 2006. A new alien species in Hellenic waters: *Pseudonereis anomala* (Polychaeta, Nereididae) invades harbors in the Eastern Mediterranean. *Aquatic Invasions*, 1(2): 59-60.
- KEVREKIDIS, K. & GALIL, B.S., 2003. Decapoda and Stomatopoda (Crustacea) of Rodos island (Greece) and the Erythrean expansion NW of the Levantine Sea. *Mediterranean Marine Science*, 4 (1): 57-66.
- KIRMITZOGLOU, I, KITSOS M.-S., THESSALOU-LEGAKI M., TSELEPIDES A. & KOUKOURAS A., 2006. Investigation of the progress and possible Expansion of the limits of the lessepsian migratory current regarding Decapoda (Crustacea). Poster, *10th International Congress on the Zoogeography and Ecology of Greece and adjacent regions*, Patras (Greece) June 26-30, 2006. (Abstract: www.upatras.gr/zoogeography).
- PANCUCCI-PAPADOPOULOU, M.A., KEVREKIDIS, K., CORSINI-FOKA, M. & SIMBOURA, N., 2005. Changes in species: invasion of exotic species. p. 336-342. In: *State of the Hellenic Marine Environment*, edited by E. Paphathanassiou and A. Zenetos, Athens, HCMR Publications.
- POR, F.D., 1978. *Lessepsian migration - the influx of Red Sea biota into the Mediterranean Sea by way of the Suez Canal*. viii + 228 p. Springer, Berlin.
- POR, F.D., 1990. Lessepsian migration. An appraisal and new data. *Bulletin de l'Institut Océanographique Monaco*, Special No 7: 1-10.
- SALOMIDI, M., BELLOU N., PANCUCCI-PAPADOPOULOU M.A. & ZIBROWIUS, in press. First observation of an invasive scleractinian coral in Greek waters. Poster 41st EMBS in Cork, Ireland on 4-8 September 2006.
- SEASLUG website <http://www.seaslug.com>
- SIMBOURA, N. & ZENETOS, A., 2005. Increasing Polychaete diversity as a consequence of increasing research effort in Greek waters: new records and exotic species. *Mediterranean Marine Science*, 6(1): 75-88.
- SINIS, A. I., 2005. First record of *Tylosurus crocodilus* (Prion & Lesueur 1821) (Pisces: Belonidae) in the Mediterranean (North Aegean Sea, Greece). *Journal of Biological Research*, 4: 221-224.
- SIOKOU-FRANGOU, I., 1999. Pelagic copepods of the Saronikos Gulf. In: *Contributions*

- to the Zoogeography and Ecology of the Eastern Mediterranean region, Proceedings 7th International congress on the zoogeography and ecology of Greece and adjacent regions, 1: 473-483.
- STREFTARIS, N., ZENETOS, A. & PAPANATHANASSIOU, E., 2005. Globalisation in marine ecosystems: the story of non-indigenous marine species across European seas. *Oceanography and Marine Biology: An Annual Review*, 43: 419-453.
- THEOCHARIS, A., NITTIS, KONTOYANNIS, H., PAPAGEORGIOU, E. & BALOPOULOS, E., 1999. Climatic changes in the Aegean Sea influence the thermohaline circulation of the Eastern Mediterranean (1986-1997). *Geophysical Research Letters*, 26 (11): 1617-1620.
- VARDALA-THEODOROU, G.E., 1999. The occurrence of the Indo-Pacific molluscan species *Fulvia fragilis* (Forsskal, 1775) and *Bulla ampulla* L., 1758 in Elefsis Bay. *Newsletters Hellenic Zoological Society*, Fasc. 31: 10-11.
- WU, P., HAINES, K. & PINARDI, N., 2000. Toward an understanding of deep-water renewal in the eastern Mediterranean. *Journal of Physical Oceanography*, 30: 443-458.
- ZENETOS, A., 1994. *Scapharca demiri* (Piani, 1981): First finding in the North Aegean Sea. *La Conchiglia*, 271: 37-38.
- ZENETOS, A., DOSI, A., ABATZOPOULOS, T.J., TRIANTAFYLIDIS, A., BEJAOU, N., SOUFI, E., AMMAR, I. A. & IBRAHIM, A., 2004. Study to investigate an invading bioindicator in the Mediterranean, *Pinctada radiata* (Leach, 1814). Communication presented at ICES/IOC/IMO Study group on Ballast and other ship vectors, Cesenatico, Italy, March 21-24, 2004.
- ZENETOS, A., GOFAS, S., RUSSO, G. & TEMPLADO, J., 2003. *CIESM Atlas of exotic species in the Mediterranean. Molluscs*. Vol. 3. 376 pp. (www.ciesm.org/atlas).
- ZENETOS A., KOUTSOUBAS, D. & VARDALA-THEODOROU, E., 2005. Origin and Vectors of Introduction of Exotic Molluscs in Greek Waters. *Belgian Journal of Zoology*, 135 (2): 279-286.
- ZENETOS, A., LANTZOUNI, M. & NICOLAIDOU, A., 2002. An account of our knowledge on fouling and ballast invaders in Greek Waters. Communication presented at ICES/IOC/IMO Study group on Ballast and other ship vectors 5th meeting, Gothenburg, Sweden, March 18-19, 2002.
- ZENETOS, A., ÇINAR, M.E., PANCUCCI-PAPADOPOULOU, M.A., HARMELIN, J.G., FURNARI, G., ANDALORO, F., BELLOU, N., STREFTARIS, N. & ZIBROWIUS, H., 2005. Annotated list of marine alien species in the Mediterranean with records of the worst invasive species, *Mediterranean Marine Science*, 6(2): 63-118.
- ZIBROWIUS, H., 1992. Ongoing modification of the Mediterranean marine fauna and flora by the establishment of exotic species. *Mésogée*, 51: 83-107.

*Accepted in December 2005.
Published in October 2006.*

