

Mediterranean Marine Science

Vol 3, No 1 (2002)



Use of algae for monitoring of heavy metals in the River Vardar, Macedonia

Z. LEVKOV, S. KRSTIC

doi: [10.12681/mms.262](https://doi.org/10.12681/mms.262)

To cite this article:

LEVKOV, Z., & KRSTIC, S. (2002). Use of algae for monitoring of heavy metals in the River Vardar, Macedonia. *Mediterranean Marine Science*, 3(1), 99-102. <https://doi.org/10.12681/mms.262>

Use of algae for monitoring of heavy metals in the River Vardar, Macedonia

Z. LEVKOV¹ and S. KRSTIC

¹ Institute of Biology, Faculty of Natural Sciences
Gazi Baba bb P.O. Box 162 1000 Skopje, FY Republic of Macedonia
e-mail: zlevkov@iunona.pmf.ukim.edu.mk

Abstract

Aiming to resolve some of the problems regarding monitoring of heavy metals in rivers using Cladophora glomerata and epilithic algal communities, a year's survey of Co, Cd, Cu, Fe, Mn, Pb and Zn has been conducted on the river Vardar, FY Republic of Macedonia. Obtained results and statistical analysis clearly point out the well documented possibility of using epilithon (basically diatom communities) as a monitoring tool, since correlation patterns for epilithon are either better or the same as those for Cladophora, while at the same time epilithon is much more reliable for monitoring, especially in cases when no other plant material can be obtained.

Keywords: Biomonitoring, Heavy metals, *Gladophora*, Epilithon, Vardar (Axios).

Introduction

A large number of investigations into the effect of heavy metals on water environments and their accumulation in various hydrobionts can be found in literature data (KELLY & WHITTON, 1989a; GENTER 1996; WANG & LEWIS, 1997). Submerged and emerged photosynthetic organisms are very important components of river ecosystems, but their application to monitoring systems has not reached level of their significance (WHITTON & KELLY, 1995). Organisms show an integrating response to their environment, as well as to fluctuations in water quality, which may be missed by intermittent chemical analysis of water (COX, 1991). Besides direct chemical analysis of river water which is spread out in

large amounts very frequently, the content of components and chemical substances in tissues, organs and the whole body, enable the obtaining of valid data for monitoring purposes. Taking into account that concentration factors for heavy metals have a range 10^2 - 10^5 (KEENEY *et al.*, 1976; OERTEL, 1991), this kind of analysis has great importance in the detection of the incident or discontinued inputs of toxic substances, especially chemicals with a low content in river water (concentrations at detection limit of the instruments). In many cases, contents of heavy metals in algal thali are much better indicators of the pollution load of the ecosystem. This is especially pointed for lead (FORSTER, 1982; LEVKOV, 2001). The idea of using organisms, or communities of organisms, to register and evaluate certain

characteristics of the environment is based on the ecological theorem of congruence between the prominence of environmental factors and the requirements of the species (MARKERT *et al.*, 1997).

Ecological investigations on the River Vardar

Interest in heavy metal pollution of River Vardar increased after the building of a metal industry in the city of Veles in the late seventies. Cadmium, one of most toxic heavy metals, was determined in the River Vardar in high concentrations (VELJANOVSKI, 1982). The concentrations have maximal values in the region of Veles, but it could be determined downstream from Veles to the border with Greece. GRIZO (1995) gave a review of heavy metals pollution during 1981-1995 and noticed that the values of concentration of particular heavy metal are decreasing. But this situation was a result mainly of decreased production and functioning of metal industry, but not from the measurement that was taken from industry (VELJANOVSKI, 1987).

More complex saprobiological investigations took place in the mid-nineties as a basis for establishing the biomonitoring based on diatoms (KRSTIC & MELOVSKI, 1994; KRSTIC, 1995; KRSTIC *et al.*, 1994, 1996, 1997). Presented data reveal physico-chemical parameters (concentration of main cations, and anions, oxygen regime, pH, etc) and biological features (qualitative composition of diatom flora, abundance of particular species and their saprobiological characteristics).

The data concerning the heavy metal accumulation in sediments are very few (MELOVSKI *et al.*, 1997) and incidental, but useful in determining the hot spots in the River Vardar and the ability of sediment to concentrate heavy metals.

Taking into account all data on the River Vardar, as well as characteristics of the river net, industry and lack of waste water treatment plants in FYR Macedonia, the hydrobiology

team at the Institute of Biology started a project entitled "*Biomonitoring of heavy metals in the River Vardar*" to determine the level of contamination of algae with heavy metals and their impact on the algal community in the River Vardar as well as to determine the most valid algal object for biomonitoring of heavy metals in the River Vardar.

Materials and Methods

The River Vardar is the main watercourse in FYR Macedonia, having a catchment area of 28.410 km², more than 15 large tributaries and is 388 km long, up to its mouth in the Aegean sea. The concentration of settlements and industry in the River Vardar valley (cities Skopje and Veles, chemical industry, smelters, and irrigation catchment, Fig.1) makes it a complex ecosystem with strong pollution pressure, primarily due to lack of any waste water treatment facility (KRSTIC *et al.*, 1997).

Field investigations on the River Vardar were realized in a period of one year (1998-1999) on ten sampling sites (Fig. 1). Water temperature, conductivity, pH, and concentration of oxygen were measured monthly in the field with appropriate instruments. Water for analysis of concentration of filtrable heavy metals was collected in 1 ½ L polyethylene bottles. After filtration of 500 mL water, it was concentrated 10 times with evaporation and the addition of 1 mL "H₂SO₄" and measured on an atomic spectrometer Varian 10BQ.

The sediment was also collected monthly from five different positions at sampling point. Interstitial water was removed with filtration of sediment. After 24 h drying on T=105 °C, 0.5 g of dried material was digested (wet digestion) using the nitric-perchloric-sulfuric acid method (10:2:1) for preparing the probes for AAS according to MOORE & CHAPMAN (1985). Analysis of concentration of heavy metals in total sediment was performed due to



Fig. 1: River Vardar water course in FYR Macedonia and 10 sampling sites.

obtaining the initial data of heavy metal load in the river bed on the sampling sites.

Samples of *Cladophora*, collected monthly from five different positions at sampling site (5 replicates), were vigorously washed in river water, and stored in plastic bottles in a refrigerator (WHITTON *et al.*, 1989). Samples were then transported to the laboratory and washed approximately 10 times in distilled water remove all attached algal cells (checked under the microscope). After 24 h drying on $T=105^{\circ}\text{C}$, 0.5 g of dried material was digested (wet digestion) using nitric-perchloric-sulfuric acid method (10:2:1) for preparing the probes for AAS according to MOORE & CHAPMAN (1985). Samples of epilithon were scraped from the surface of five bigger stones with a plastic knife, dried out and checked for presence of

Cladophora glomerata. The procedure followed for digestion was the same for sediment as well. After dissolving of the digested material in distilled water, the samples were measured on atomic spectrometer Varian 10BQ.

The statistical analysis were done with STATISTICA for Windows (StatSoft, Inc. 1995). Two multiple regressions were performed. In the first, the concentration of heavy metals in water as an independent variable, and the content of heavy metals in sediment, *Cladophora* and epilithon as dependent variable was taken, to predict the linkage of these two variables. The second correlation analysis was performed as an attempt to predict the influence of sediment on the accumulation of heavy metals in algae.

Results and Discussion

The most variable value for pH was obtained on sampling site T₈ (Table 1), due to the great impact of waste waters of the chemical industry of Veles (CIV). The waste waters show great variability of pH value from 12.9 to 1.1 (KRSTIC, 1995). Low pH values have an the impact on the dissolving or sedimentation of heavy metals (TACK *et al.*, 1996), as well as on the accumulation processes in algae (VYMAZL, 1995). Beside pH, other physico-chemical parameters have an impact on heavy metal concentrations in natural waters: suspended particles (BOTELHO *et al.*, 1994a; PELLETIER, 1996), redox potential (VERLOO & COTTENIE, 1985) sulfides and phosphates (RECZYNSKA-DUTKA, 1991), chemical interactions (RULE & ALDEN, 1996), organic substances (WHITTON & SAY, 1975; BOTELHO *et al.*, 1994b). The analysis of the concentration of heavy metals in water often gave only an approximate view of heavy metal load. The relativity of obtained data could be established in short-term or incidental (discontinued) load of heavy metals (VOGEL & CHOVANEC, 1992).

The communal waste waters from the capital Skopje have a great impact on the River Vardar (KRSTIC, 1995), but the concentration of filterable heavy metals is not as high as on the next sampling sites. This could be as a result of the huge load of organic substances, phosphates and sulfides with communal origin. The maximal values for the concentration of Cd, Pb and Zn are determined on sampling site T₇ due to the load of waste waters from the metal industry in Veles. The other three analysed elements, Cu, Mn and Fe, show maximal values on the next sampling site (T₈ CIV) as a result of the direct input of waste waters with extremely low pH.

Comparing the obtained results (Table 1) with analogous results from other authors, (WHITTON *et al.*, 1989; RECZYNSKA-DUTKA, 1991), it can be found that for some ions (Cd, Pb and Zn) values are much higher. But, compared with analogous data from Vardar in the past several decades the tendency towards decreasing of values could be noticed. This situation is a result of the decreased level of functioning and production of the metal industry in Veles and definitely not as a result

Table 1
Average, maximal and minimal values of measured parameters along the River Vardar (T1-T10) water course in FYR Macedonia.

Parametrs	T1	T2	T3	T4	T5	T6	T7	T8	T9	T10
T water (°C)	12.25 (19.1-9.6)	11.12 (17.2-6.7)	13.16 (20.5-6.1)	14.33 (19.6-7.9)	13.51 (20.5-6.9)	14.81 (29.1-5.4)	15.87 (25.9-5.7)	17.94 (28.7-9.5)	16.36 (28.8-5.5)	16.23 (28.8-6.1)
pH	7.61 (8.63-6.75)	7.62 (7.97-7.05)	7.72 (8.73-7.29)	7.78 (8.71-7.42)	7.65 (7.92-7.15)	7.57 (7.87-7.25)	7.51 (7.97-7.25)	6.13 (11.3-2.61)	7.51 (7.95-7.06)	7.80 (8.54-7.32)
Conductivity (µS/cm)	102.40 (200-84.7)	200.00 (300-100)	254.55 (500-200)	340.00 (400-40)	345.45 (400-200)	354.55 (400-300)	427.27 (600-300)	854.55 (3700-300)	427.27 (700-300)	418.18 (600-300)
Cd (µg/L)	0.02 (0.05-0.01)	0.11 (0.6-0.01)	0.38 (1.2-0.01)	0.33 (1.9-0.01)	0.61 (2.9-0.01)	1.07 (4.9-0.01)	14.40 (36-0.5)	9.32 (14.4-0.1)	1.54 (4.7-0.01)	1.14 (3.3-0.01)
Cu (µg/L)	5.93 (13-2.9)	12.58 (56-4.4)	13.27 (40-7.8)	23.03 (90-12.3)	18.87 (84-8.2)	17.15 (59-7.5)	20.27 (70-9.6)	26.67 (86-9.9)	18.62 (98-6.9)	18.26 (86-6.9)
Fe (µg/L)	19.25 (40-2)	38.50 (96-5)	50.75 (79-22)	72.42 (262-21)	110.08 (311-26)	120.91 (265-26)	130.50 (377-32)	495.25 (1904-43)	74.17 (140-22)	66.92 (142-14)
Mn (µg/L)	1.81 (8.5-0.1)	3.55 (6.4-0.1)	5.13 (10.3-0.2)	19.13 (56.2-2.3)	10.38 (29.8-1.1)	19.58 (69.8-2.8)	47.91 (249-10.4)	61.33 (236-8.6)	12.47 (27-5.8)	9.69 (20-0.01)
Pb (µg/L)	12.31 (37-1)	26.25 (90-4)	27.58 (103-4)	37.75 (104-4)	40.58 (104-11)	39.99 (104-9)	62.83 (138-16)	41.83 (130-14)	40.08 (101-6)	40.17 (104-4)
Zn (µg/L)	6.92 (17.9-3.8)	30.58 (220-4.8)	15.61 (44.6-6)	15.05 (26.7-7.4)	16.29 (24.5-7.4)	14.96 (25-5)	168.47 (397.2-39.7)	96.98 (472.5-12.6)	16.39 (20.6-11)	30.13 (213.5-5.3)

of measures undertaken for the treatment of waste waters.

The sediment is a depot where a large amount of heavy metals is accumulated (JOHNSON, 1998), but the high heavy metal content in sediment could be the result of high background concentration. In addition the content of heavy metals in sediment could not be correlated with concentration in river water (VERLOO & COTENIE, 1985) due to their binding in undissolved sulfides and phosphates.

According to VOGEL & CHOVANEC (1992), there are several advantages in using sediment to monitor heavy metals: (i) emission sources can be located and characterized by sediment analyses; (ii) data from single sediment sample represent the pollution of the river over a longer time period; (iii) due to the low concentration in water, some substances can be hardly detected, and substances which accumulate in sediments can be easily detected by sediment analyses.

When the content of heavy metals in sediment is analysed, two different approaches could be applied: (i) heavy metal in the whole sediment and (ii) heavy metals in different fractions of sediment. The fine sediment fraction (<0.2 mm) is particularly useful for determination of heavy metal load in rivers and

to distinguish between natural (background, geogenic) and anthropogenic sources (KRALIK, 1999). The size fraction of sediment could be divided into two groups: (1) coarse particles less than 0.2 mm and (2) fine particles less 0.02 mm. Nevertheless, there is experimental uncertainty associated with all presently available methods of metal speciations. In particular, solid phase fractionation schemes suffer from serious limitations: sample handling, reagent selectivity and specificity, interference, etc. (TACK & VERLOO, 1995). Some elements, such as Cd, for example, in natural sediments are strongly bound with organic substances (sulfides), but heavy metals with anthropogenic origin are bound very weakly (RULE & ALDEN, 1992). Nevertheless, the content of heavy metals in sediment depends on different factors such as pH (TACK *et al.*, 1996), type of sediment, ionic strength, conductivity, (ZHOU & KOTT, 1995), algal biofilms (WOODROOFF *et al.*, 1999), redoxpotential (VERLOO & COTTENIE, 1985), size fractions (KRALIK, 1999), etc.

Analyses of sediment in the River Vardar (Table 2) show that on sampling sites located in the upper part of the River Vardar (from source to Skopje) the anthropogenic pressure is low. Decreasing of values on sampling sites located after the communal channels of the towns Tetovo and Gostivar (T₂ and T₃), is a

Table 2
Average, maximal and minimal values of content of heavy metals in the River Vardar sediment during investigated period 1998-1999.

	T1	T2	T3	T4	T5	T6	T7	T8	T9	T10
Cd (µg/L)	0.92 (0.20-3.80)	0.90 (0.2-2.0)	1.08 (0.1-2.5)	2.32 (0.1-3.8)	2.09 (0.1-5.1)	2.23 (0.4-3.2)	133.98 (18.3-212.9)	35.40 (4.95-27.4)	8.79 (1.7-25.0)	4.85 (0.5-14.0)
Co (µg/L)	13.99 (10.5-16.6)	13.99 (9.4-20.4)	11.36 (7.3-17.5)	13.15 (9.7-17.1)	14.62 (12.5-17.8)	13.91 (10.3-17.0)	13.13 (10.3-18.4)	14.39 (8.6-22.1)	12.97 (9.1-15.5)	11.73 (6.7-17.1)
Cu (µg/L)	46.83 (24.8-66.4)	45.29 (23.5-65.0)	38.37 (24.3-54.4)	52.52 (23.5-104.5)	53.21 (27.9-104.5)	49.29 (22.6-84.3)	146.11 (44.8-260.0)	74.14 (31.1-112.2)	37.55 (27.2-49.9)	39.22 (14.1-129.3)
Fe (µg/L)	3234.88 (2509-4925)	3619.39 (3050-4913)	3404.92 (2645-4490)	2889.99 (2256-4106)	2975.52 (2687-4131)	3127.94 (2282-4627)	2851.80 (1975-4095)	3255.65 (2794-4898)	3004.00 (2041-4795)	3127.73 (2058-4361)
Mn (µg/L)	790.95 (600-1625)	1068.20 (543-1446)	858.09 (453-938)	638.78 (440.2-879)	929.74 (765-1184)	864.93 (561-1516)	872.60 (532.5-1141)	956.14 (590-1270)	775.60 (494-1492)	602.86 (413-1368)
Pb (µg/L)	84.81 (15.0-157)	122.93 (70.0-185.0)	116.08 (66.0-210.0)	163.03 (86.3-271.5)	136.13 (30.0-271.5)	117.43 (27-207.5)	630.41 (96.7-1257)	170.28 (101.0-242)	139.63 (68.0-201.5)	121.39 (75.5-214.5)
Zn (µg/L)	102.40 (83.8-143.3)	87.13 (53.8-106.5)	76.35 (47.5-120.0)	173.98 (85.3-257.3)	214.66 (96.3-454.4)	196.77 (96.8-369.5)	1905.31 (524-5573)	420.83 (286.8-676)	196.24 (121.8-263)	134.24 (72.5-177.1)

result of physico-chemical characteristics of sediment on those locations. Nevertheless, the sediment is composed mainly of inorganic silica particles which have very a weak ability to adsorb heavy metals (ZHOU & KOTT, 1995; BIJLSMA *et al.*, 1994). Starting from sampling site T4 (located in Skopje), the content of heavy metals starts to increase. The highest values for all investigated heavy metals are determined at T7 (Veles). When obtained results from this sampling site are compared with reference dates (GONÇALVES *et al.*, 1992) enormous difference can be noticed. The values for Cd are about 670 times greater than background concentration. This remarkable increase in values also could be noticed for Pb and Zn (VOGEL & CHOVANEC, 1992; DAUVALTER, 1992; FACETTI *et al.*, 1998). According to these, characteristics sediment at this sampling site should be classified and treated as extremely dangerous (KELDERMAN & DROSSAERT, 1999).

WHITTON *et al.*, (1981) suggested 10 plant species as appropriate for biomonitoring of heavy metals in river ecosystems, and four of them belong to the group of algae: *Cladophora glomerata*, *Enteromorpha sp.*, *Lemanea fluviatilis* and *Nitella flexilis*.

WHITTON (1970) had pointed out *Cladophora glomerata* as potential species for biomonitoring of heavy metals as a result of

wide distribution in water ecosystems. This feature of *Cladophora* was referred to by several authors (BLUM, 1956; ROSEMARIN, 1985; WHITTON *et al.* 1989; DODDS, 1991; DODDS & GUDER, 1992; SHEATH & COLE, 1992; WHITTON & KELLY, 1995). The Natural population of *Cladophora* in rivers (Table 3) shown great ability to accumulate of heavy metals (KEENEY *et al.* 1976; ADO-RADY, 1980; WHITTON *et al.*, 1981, 1989; KELLY & WHITTON, 1989, JACKSON *et al.*, 1990; OERTEL, 1991, 1993). Obtained high significant statistical correlation between concentration of heavy metals in water and *Cladophora* suggest that this species is a very good object for the monitoring of heavy metals (WHITTON *et al.*, 1989; LEVKOV, 2001). In addition, *Cladophora* could be used in the monitoring of eutrophication, e.g. long-term biomonitoring of concentration of nitrogen and phosphorus (WIESE *et al.*, 1986; FREEMAN, 1986).

KELLY & WHITTON (1989) had compared the accumulation ability of aquatic mosses (*Rhynostegium riparoides*, *Amblistegium riparium* and *Fontinalis antipyretica*) and macrophytic algae (*Cladophora glomerata*, *Lemanea fluviatilis* and *Stigeoclonium tenue*) and concluded that the background concentrations for algae are lower than mosses. Accumulation processes increase with the

Table 3
Average, maximal and minimal values for content of heavy metals in *Cladophora* on nine sampling sites during investigated period.

	T2	T3	T4	T5	T6	T7	T8	T9	T10
Cd (µg/L)	0.50 (0.01-1.30)	0.75 (0.01-3.12)	1.38 (0.1-3.97)	1.55 (0.1-7.39)	1.12 (0.1-2.79)	166.36 (12.99-237.9)	31.31 (0.560-67.37)	8.78 (0.1-31.95)	5.11 (0.1-15.83)
Co (µg/L)	1.54 (0.30-2.48)	3.91 (0.1-4.55)	4.27 (0.1-8.75)	5.31 (0.1-11.17)	4.45 (0.1-10.15)	6.19 (0.25-18.32)	8.33 (2.50-22.6)	3.83 (1.9-12.1)	2.78 (0.2-7.3)
Cu (µg/L)	18.50 (9.52-28.5)	20.83 (14.43-33.05)	46.58 (12.9-153.3)	29.48 (18.98-45.4)	29.74 (14.95-37.19)	68.42 (16.55-206.5)	52.22 (15.5-143.8)	26.08 (11.90-52.6)	31.69 (11.37-52.6)
Fe (µg/L)	7.78 (3.58-12.8)	7.67 (2.56-11.78)	7.46 (4.06-12.91)	6.96 (3.58-11.57)	6.31 (4.20-9.13)	10.65 (4.20-17.7)	12.77 (4.91-20.42)	9.27 (3.72-12.06)	9.76 (3.84-16.85)
Mn (µg/L)	699.2 (272-1258.8)	1035.5 (230.3-3244)	887.9 (226.4-1773)	866.4 (206.6-1936)	1141.0 (336.3-2112)	2094.2 (411.3-6214)	1332.2 (273-2241)	1079.5 (285-2773)	625.4 (191.25-1668)
Pb (µg/L)	31.70 (9.50)	33.81 (6.0-76.5)	53.53 (18.0-213.5)	60.49 (11.1-151.1)	41.67 (27.49-92.0)	259.97 (71.5-891.8)	195.28 (55.5-591.7)	57.16 (31.0-102.0)	46.60 (7.0-86.0)
Zn (µg/L)	36.47 (20.3-72.1)	62.60 (28.27-140.4)	138.18 (33.07-239.6)	194.36 (51.3-322.2)	115.43 (28.10-197.5)	1385.41 (265.3-5049)	372.23 (88.6-807.4)	163.75 (92.0-382.9)	145.18 (35.86-264.0)

increase of the concentration of particular heavy metals (WEHR & WHITTON 1983a, 1983b; WEHR *et al.*, 1987). But the intensity of heavy metals accumulation in algae is higher than in aquatic mosses. This feature makes algae a better and more sensitive object for the monitoring of heavy metals (LEVKOV 2001).

Many studies for detection of heavy metal uptake and accumulation in *Cladophora* are based on a single collection (or several field investigations) of material in the summer period when the population of alga is dominant in rivers. Data for seasonal changes in the content of heavy metals in *Cladophora* are scarce (RANG & STIKES, 1987). This situation is mainly the result of the decrease in *Cladophora's* population especially in winter months and it is not most the representative (dominant) alga in the river ecosystem. In this period, basal parts of the alga only can be found, but they are not appropriate for chemical analysis due to calcification (WHITTON, 1981). Additionally, large biomass of epiphytic and grazer species makes cleaning of the material more difficult and increase the possibility of contamination (LEVKOV, 2001). Nevertheless, enough material for chemical analysis can be found with a detailed search of the river bottom.

Intensity and kinetics of the bioaccumulation processes of heavy metals were also tested in laboratory conditions (SCHANZ & THOMAS, 1981; VYMAZAL 1987, 1990a, 1990b, 1995; SOBHAN & STENBERG 1999). The kinetics of accumulation of heavy metals consists of two phases; (i) the initial phase is very rapid, occurring immediately after exposure to heavy metal; usually lasting for less than 30 minutes (VYMAZAL, 1990; GENTER, 1996; WANG & DEI, 2001a). This phase is probably passive, involving physical sorption of heavy metals. According to PICKERING & PUA (1969), this phase can be divided in two separate phases: a) intensive accumulation in the first 20 minutes and b) linear increasing in a period of 90 minutes. (ii) second phase, named as metabolism-depend

uptake (GENTER, 1996), is extended and slow with duration of more than one month. The accumulation trend could be linear in laboratory conditions (SAKAGUCHI *et al.*, 1979) or hyperbolic in natural conditions (CONWAY & WILLIAMS 1979). Also, nutrients such as N, P and Si have a great influence on heavy metal uptake and kinetics (WANG, 1987, WANG & DEI, 2001b, 2001c).

Comparing the literature data for heavy metal content it could be noticed that for all investigated metals obtained values for the River Vardar are much higher (Table 4). Nevertheless, VYMAZAL (1987, 1990, 1995) investigated the processes under laboratory conditions with exposure of *Cladophora* to concentrations of 200 mg/L of a particular heavy metal over period of six hours. This concentration is much higher than values detected in the River Vardar (and by other researchers), and it is very possible that it results in inhibition of metabolic processes in *Cladophora* and accumulation is due to passive (osmotic) transport of heavy metals and existing high gradient of concentration. These data are very important for explaining the kinetics of heavy metal uptake in *Cladophora*. Beside, algal populations in the River Vardar have permanent heavy metal pressure throughout the year and accumulation processes depend both on the metabolic condition of algae and environmental

Table 4
Maximal detected values for heavy metal content in *Cladophora glomerata* ($\mu\text{g/g}$ dry weight).

	Cd	Cu	Pb	Zn
Vyzamal (1995)*	109.6	115.2	157	117.6
Vyzamal (1987)*	/	/	/	643.0
Vyzamal (1990)*	119.1	/	159.1	
Whitton <i>et al.</i> (1989)**	6.97	33	330	1130.0
Oertel (1991)**	6.09	28.2	128	270.0
Keeny <i>et al.</i> (1979)**	3.9	7.2	12.5	23.7
Abo-Rady (1980)**	0.94	9.1	5.2	62.0
Levkov (2001)**	799.5	206.5	891.5	5048

* Data from culture exposed to high concentrations of heavy metal under laboratory conditions

** Natural populations of *Cladophora*

conditions. This is especially expressed at sampling site T8 (CIV) where due to the low pH value of river water (Table 1) accumulation processes are decreased. The obtained values from the River Vardar, compared with WHITTON *et al.* (1989) and OERTEL (1991), show the real heavy metal load in the river. Nevertheless, of cadmium and zinc content is more than 110 and 20 times higher respectively, than most polluted sites in Great Britain and the river Danube.

Beside numerous advantages that *Cladophora sp.* has as a potential biomonitoring object, it also has several limitations and disadvantages. Namely, the growth of *Cladophora* is mainly limited to mesotrophic to eutrophic waters, and depends on concentration of nitrogen and phosphorus (NEIL & JACKSON 1982; MANTAI *et al.*, 1982; MILLNER *et al.*, 1982), temperature, (WHITTON, 1970), light (GRAHAM *et al.*, 1982), water velocity and composition of the bottom (DODDS, 1991). The growth is seasonal and representative material for chemical analysis can be found only in seven to eight months during the year.

Several macrophytic algal species, beside *Cladophora*, were used to determine the effect of high heavy metal concentrations and the possibility of using those species as biomonitoring objects. Due to several features, such as easy recognition and collection of the talus and good ability to accumulate heavy metals, *Lemanea fluviatilis* is one of the potential algal species for biomonitoring (HARDING & WHITTON, 1981; KELLY & WHITTON 1989a) which shows special resistance to zinc and lead (WHITTON & SAY, 1975).

Very similar accumulation characteristics to *Cladophora*, have *Ulotrix zonata* and *Stigeoclonium tenue*, but as a result of limited ecological valence and distribution, application of these species for biomonitoring of heavy metals in river ecosystems is very limited (JACKSON *et al.*, 1990; KELLY & WHITTON, 1989a, 1989b).

The limited ecological valence of *Cladophora glomerata* to pollution parameters and the seasonal pattern of the life cycle are the main factors that make the biomonitoring of heavy metals based on this species uncertain and difficult. This is characteristic mainly of the winter months when *Cladophora* populations decrease. To avoid these problems, an alternative solution is the method based on epilithic communities (biofilms). According to Round (1991) epilithic communities have several advantages as biomonitoring tool, as (i) there is significant difference in content of heavy metal in epilithon for natural (unpolluted) and polluted sites (Table 5); (ii) the ecological valence of epilithon is much wider than filamentous species such as *Cladophora* (iii) the biomass of epilithon, in general, is increases with the increase of the pollution level.

Effects of increased levels of heavy metals on epilithic algal communities is studied under laboratory and natural conditions. Generally in natural communities effects are investigated at chronic (long term) exposures, what is "more realistic, what algae will experience in nature" (GENTER, 1996). They react more completely than filamentous algae or macrophytes (IVORRA *et al.*, 1999). On the other hand, there are very few literature data concerning accumulation of heavy metals in natural diatom communities (LEVKOV, 2001). IVORRA (2000) shows that there is a large difference in the content of heavy metals in algal communities from unpolluted and polluted sites, and mainly shows linear correlation with concentration of adequate heavy metal in ambient water (ABSIL & van SCHEPPINGEN, 1996). This difference is due to the different species composition in diatom communities (ADMIRAAL *et al.* 1997; GENTER, 1987). Some diatom species have developed tolerance mechanisms against cytotoxic effects of heavy metals (TORRES *et al.*, 1995, 1997) to reduce heavy metal toxicity by producing intracellular and extracellular binding components (AHNER *et al.*, 1995; AHNER & MOREL, 1995).

Table 5

Average, maximal and minimal values for content of heavy metals in epilithon on 10 sampling sites during investigated period .

	T1	T2	T3	T4	T5	T6	T7	T8	T9	T10
Cd (µg/L)	0.35 (0.15-3.5)	0.24 (0.05-0.55)	0.17 (0.1-0.25)	10.05 (3.60-17.85)	5.00 (0.60-7.85)	3.14 (1.35-8.05)	62.93 (41.32-83.9)	21.05 (5.25-29.0)	2.98 (1.60-4.30)	1.96 (1.80-2.15)
Co (µg/L)	5.81 (4.88-6.65)	8.46 (7.60-9.75)	9.33 (8.55-10.30)	11.18 (8.85-12.95)	11.11 (9.05-15.80)	11.90 (10.7-17.6)	13.35 (10.7-17.6)	11.98 (9.80-13.95)	11.76 (10.95-13.8)	12.28 (11.7-12.95)
Cu (µg/L)	42.74 (26.20-72.3)	31.22 (25.65-41.8)	34.43 (30.5-40.1)	49.11 (32.20-61.3)	20.95 (17.30-25.6)	24.89 (21.2-32.15)	102.01 (94.8-119.5)	68.55 (44.5-115.7)	32.34 (28.0-41.0)	25.76 (24.05-27.5)
Fe (µg/L)	6.73 (5.37-8.95)	12.42 (11.9-13.96)	16.28 (11.8-21.40)	20.82 (17.99-25.4)	14.45 (11.55-19.3)	18.03 (10.9-25.41)	21.06 (19.85-25.5)	20.57 (19.25-21.7)	23.24 (20.5-26.27)	20.34 (19.26-21.1)
Mn (µg/L)	438.1 (338.8-66.3)	771.5 (659-914)	1667.9 (1145-2399)	1343.7 (943.8-2114)	1095.3 (981-1202)	1046.3 (965-1150)	1255.9 (1093-1337)	971.7 (601-1325)	1381.7 (1273-1496)	923.4 (636-1113)
Pb (µg/L)	7.26 (4.50-12.0)	15.29 (9.15-20.00)	13.13 (9.50-16.50)	20.94 (17.75-25.5)	19.94 (18.50-23.0)	24.30 (20.50-31.5)	247.63 (107.5-595)	50.50 (46.0-56.60)	28.31 (26.25-29)	22.74 (19.75-28.0)
Zn (µg/L)	113.10 (67.5-17.9)	65.74 (54.2-72.5)	64.33 (53.7-75.63)	160.48 (126.9-200)	107.19 (88.5-148.8)	128.94 (82.5-165.0)	1619.53 (320.5-2743)	299.18 (255-316.3)	204.64 (166.3-261)	129.13 (90.0-230.0)

Statistical Analysis

The heavy metal content in *Cladophora* was correlated with concentrations of heavy metals in water and sediment. WHITTON (1971) noticed that content of heavy metals in sediment does not have an effect on *Cladophora glomerata*, but the author noticed that populations of *Cladophora* could be found at sites with high levels (content) of zinc in sediment. WHITTON *et al.* (1989) show that the content of heavy metals in river water has great impact on the content of heavy metals in *Cladophora*. There are positive linear correlations with high statistical significance between the concentration of a particular ion in water and algae. Besides these types of the correlation, the authors show that other ions have an impact on heavy metal accumulation, too. Obtained results on the River Vardar show the great impact of Zn on the accumulation of other elements such as Cd, Fe, and Pb (Table 6).

Statistical analysis of the River Vardar data shows that much higher correlations are establish between data sets obtained for sediment and *Cladophora*, than to the data sets for water and *Cladophora*.

Conducted statistical analysis represents the final evidence for the validity of epilithic communities as biomonitoring objects. Comparing the obtained correlation coefficients

(r) for epilithon and *Cladophora* it could be noticed that both of them have very similar values and patterns of behavior at pollution stress caused by heavy metals. This fact could be noticed when data sets for *Cladophora* and epilithon were compared for each heavy metal (Figs 2,3).

Obtained higher values for correlation coefficients for epilithon suggest that epilithic communities are better biomonitoring objects for heavy metal pollution than algae such as *Cladophora glomerata*, *Lemanea fluviatilis* and *Fritschella tuberosa* or aquatic mosses *Fontinalis antipyretica* and *Rhynostegium riparoides*.

Conclusions

Bearing in mind all the problems that a proper monitoring system faces today in relation to the heavy metal pollution of river ecosystems, according to presented results, we would like to recommend epilithon communities as a monitoring tool due to following conclusions based on investigations conducted on the River Vardar:

- The River Vardar in FYR Macedonia represents an ecosystem of severe human impact, originating both from industrial and communal waste water systems. Con-

Table 6
Correlation coefficient (r) between metal concentrations in water and sediment and the corresponding metals in *Cladophora* and epilithon.

	Cd _{epi}	Co _{epi}	Cu _{epi}	Fe _{epi}	Mn _{epi}	Pb _{epi}	Zn _{epi}	Cd _{clad}	Co _{clad}	Cu _{clad}	Fe _{clad}	Mn _{clad}	Pb _{clad}	Zn _{clad}
Cd _{aq}	0.69	0.51	0.47	0.26	-0.26	0.73	0.41	0.55	0.43	0.38	0.43	0.17	0.32	0.25
Co _{aq}	-0.07	-0.27	0.03	-0.32	-0.20	-0.09	-0.13	-0.10	-0.13	0.03	-0.19	-0.11	-0.10	-0.10
Cu _{aq}	0.12	0.22	0.08	0.28	-0.22	0.04	0.12	0.13	0.34	0.27	0.06	0.22	0.17	0.19
Fe _{aq}	-0.02	0.05	0.43	0.09	-0.24	0.06	-0.06	0.00	0.20	0.25	0.11	-0.03	0.10	-0.06
Mn _{aq}	0.56	0.55	0.58	0.40	-0.10	0.86	0.48	0.30	0.44	0.52	0.28	0.29	0.35	0.31
Pb _{aq}	0.27	0.02	0.35	-0.09	-0.05	0.65	0.556	0.53	0.23	0.37	0.18	0.50	0.53	0.63
Zn _{aq}	0.47	0.17	0.54	0.14	-0.28	0.45	0.78	0.69	0.55	0.51	0.37	0.55	0.69	0.75
Cd _{sed}	0.80	0.45	0.88	0.31	-0.03	0.87	0.76	0.71	0.48	0.83	0.28	0.35	0.57	0.60
Co _{sed}	-0.03	-0.13	0.13	-0.17	0.14	-0.01	-0.09	-0.08	-0.05	0.06	-0.10	-0.15	-0.05	-0.13
Cu _{sed}	0.62	0.22	0.82	0.13	-0.17	0.69	0.78	0.74	0.55	0.80	0.24	0.67	0.77	0.78
Fe _{sed}	-0.24	-0.30	0.18	-0.20	-0.32	-0.11	-0.18	-0.15	-0.10	-0.01	-0.24	-0.17	0.01	-0.11
Mn _{sed}	0.11	-0.08	0.12	-0.34	-0.23	0.10	0.07	0.10	0.22	-0.04	-0.12	0.17	0.12	0.08
Pb _{sed}	0.65	0.21	0.71	0.18	0.03	0.69	0.95	0.88	0.50	0.76	0.20	0.75	0.82	0.98
Zn _{sed}	0.80	0.52	0.73	0.28	0.02	0.98	0.75	0.61	0.45	0.73	0.26	0.44	0.51	0.56




Fig. 2: Correlation coefficients (r) for content of heavy metals in epilithon and *Cladophora* versus river water.

centrations of heavy metals in water and especially in sediment are probably the highest recorded recently, thus putting the river biota under enormous survival stress or maximal accumulation rates. Due to a wide range of influences (pH, phosphates, sulfates, organic compounds, etc.) there are frequent records of complete extinction of river bottom life, which is another reason to monitor epilithon since it is the first to develop after devastating human influence;

- River sediment has a great ability to accumulation of heavy metals and could be used for the monitoring of heavy metal pollution;




Fig. 3: Correlation coefficients (r) for content of heavy metals in epilithon and *Cladophora* (dependent variable) versus sediment (independent variable).

- *Cladophora glomerata* is a precise biomonitoring tool for determination and quantification of heavy metal pollution in the River Vardar. Concentrations of particular heavy metals in *Cladophora* reflect the heavy metal load in the River Vardar;
- Statistical correlation matrices reflect significant correlation coefficient regarding heavy metal concentration both in *Cladophora* and epilithon, versus concentrations of heavy metals in waters or sediment (except for Mn which are not statistically relevant after all). These findings suggest that epilithon can be relevantly

chosen for a biomonitoring tool of heavy metal pollution in rivers;

- Concentrations of heavy metals in epilithon were usually very much related to those recorded for the sediment, obviously pointing to the concentration of the particular metal in the sampling site and thus representing a better monitoring tool for a prolonged period of time;
- Epilithon (predominantly diatoms) is present in all river systems, although sometimes difficult to locate in large rivers, but can easily be introduced onto natural stones, and does not depend on most factors that influence filamentous algae, bryophytes or higher plants. It is also present throughout the year, but left alone tends to predominate in winter, when all the other possible monitoring organisms disappear.

References

- ABO-RADY, M.D.K., 1980. Makrophytische Wasserpflanzen als Bioindikatoren für die Schwermetallbelastung der oberen Leine. *Archiv für Hydrobiologie*, 89, (3): 287-404.
- ABSIL, M.C.P. & VAN SCHEPPINGEN, Y. 1996. Concentration of Selected Heavy Metals in Benthic Diatoms and Sediment in the Westerschelde Estuary. *Bulletin of Environmental Contamination and Toxicology*, 56: 1008-1015.
- ADMIRAAL, W., IVORRA, N., JONKER, M., BREMER, S., BARRANGUET, C. & GUASCH, H., 1997. Distribution of diatom species in metal-polluted Belgian-Dutch river: An experimental analysis. p.240-244. In: Use of algae for monitoring rivers III, edited by Prygiel J, Whitton BA, Bukowska.
- AHNER, B.A., KONG, S. & MOREL, F.M.M., 1995. Phytochelatin production in marine algae. 1. An interspecies comparison. *Limnology and Oceanography*, 40(4): 649-657.
- AHNER, B.A. & MOREL, F.M.M. 1995. Phytochelatin production in marine algae. 2. Induction by various metals. *Limnology and Oceanography*, 40 (4): 658-665.
- BLUM, J.L., 1956. The ecology of river algae. *Botanical Review*, 22: 291-341.
- BIJLSMA, M., GALIONE, A.L.S., KELDERMAN, P., ALAERTS, G.J. & CLARISSE, I.A., 1994. Assessment of heavy metal pollution in inner city canal sediments. Technical report. 7 pp.
- BOTELHO, C.M.S., BOAVENTURA, R.A.R., GONÇALVES, M.L.S.S. & SIGG, L., 1994a. Interactions of Lead (II) with Natural River Water. Part II: *Particulate Matter. Science of the Total Environment*, 151: 101-112.
- BOTELHO, C.M.S., BOAVENTURA R.A.R. & GONÇALVES, M.L.S.S., 1994b. Interactions of Lead (II) with Natural River Water. Part I: Soluble Organics. *Science of the Total Environment*, 149: 69-81.
- CONWAY, H.L. & WILLIAMS, S.C., 1979. Sorption of cadmium and its effect on growth and the utilization of inorganic carbon and phosphorus of two freshwater diatoms. *Bulletin of the Fisheries Research Board of Canada*, 36: 579-586.
- COX, E.J., 1991. What is the basis for using diatoms as monitors of river quality. p.33-40. In: Use of Algae for Monitoring rivers, edited by B.A. Whitton, E. Rott and G. Friedrich.
- DAUVALTER, V., 1992. Concentration of Heavy Metals in Superficial Lake Sediments of Pechenga District, Murmansk region, Russia. *Vatten* 48: 141-145.
- DODDS, W.K., 1991. Factors associated with dominance of the filamentous green alga *Cladophora glomerata*. *Water Research*, 25 (11): 1325-1332.
- DODDS, W.K. & GUDDER D.A., 1992. The ecology of *Cladophora*. Review. *Journal of Phycology*, 28: 415-427.
- FACETTI, J., DEKOV, V.M. & VAN GRIEKEN R., 1998. Heavy metals in sediments from the Paraguay River: a preliminary study. *Science of the Total Environment*, 209: 79-86.
- FORSTER, P., 1982. Species associations and metal contents of algae from rivers polluted by heavy metals. *Freshwater Biology*, 12: 17-39.
- FREEMAN, M.C., 1986. The role of nitrogen and phosphorus in the development of *Cladophora glomerata* (L.) Kützing in the Manawatu River, New Zeland. *Hydrobiologia*, 131 (1): 23-30.
- GENTER, R.B., CHERRY, D.S., SMITH E.P. & CAIRNS J., 1987. Algal-periphyton population and community changes from zinc stress in stream mesocosms. *Hydrobiologia*, 153: 261-275.
- GENTER, R.B., 1996. Ecotoxicology of Inorganic Chemical Stress to Algae. p.403-468. In: Algal

- Ecology, edited by Stevenson R.J., Bothwell M.L. and Lowe R.L., Academic Press.
- GONÇALVES, E.P.R., BOAVENTURA, R.A.R. & MOUVET, C., 1992. Sediments and aquatic mosses as pollution indicators for heavy metals in the Ave River basin (Portugal). *Science of the Total Environment*, 114: 7-24.
- GRAHAM, J.M., AUER, M.T., CANALE, R.P. & HOFFMAN, J.P., 1982. Ecological studies and mathematical modeling of *Cladophora* in Lake Huron: 4. Photosynthesis and respiration as function of light and temperature. *Journal of Great Lakes Research*, 8: 100-112.
- GRIZO, A., 1995. Heavy metals in the River Vardar. *Ekologija i Zastita na Zivotnata Sredina*, 3, (1-2): 43-49. (in Macedonian).
- HARDING, J.P. & WHITTON, B.A., 1981. Accumulation of zinc, cadmium, and lead by field populations of *Lemanea*. *Water Research*. 15: 301-319.
- IVORRA, N., HETTELAR, J., TUBBING, G.M.J., KRAAK, M.H.S., SABATER, S. & ADMIRAAL, W., 1999. Translocation of Microbenthic Algal Assemblages Used for In Situ Analysis of Metal Pollution in Rivers. *Archive of Environmental Contamination and Toxicology*, 37: 19-28.
- IVORRA, N.C., 2000. Metal induced succession in benthic diatom consortia. Doctor Dissertation. Faculty of Sciences, University of Amsterdam, The Netherlands. 161p.
- JACKSON, M.B., VANDERMEER, E.M. & HEINTSCH, L.S., 1990. Attached filamentous algae of Northern Lake Superior: Field ecology and biomonitoring potential during 1983. *Journal of Great Lakes Research*, 16 (1): 158-168.
- KELLY, M.G. & WHITTON, B.A., 1989a. Interspecific differences in Zn, Cd and Pb accumulation by freshwater algae and bryophytes. *Hydrobiologia*, 175: 1-11.
- KELLY, M.G. & WHITTON, B.A., 1989b. Relationship between accumulation and toxicity of zinc in *Stigeoclonium* (Chaetophorales, Chlorophyta). *Phycologia*, 28: 512-517.
- KEENEY, W.L., BRECK, W.G., VAN LOON, G.W. & PAGE, J.A., 1976. The determination of trace metals in *Cladophora glomerata*. C. *glomerata* as potential biological monitor. *Water Research*, 10: 981-984.
- KELDERMAN, P. & DROSSAERT, W.M.E., 1999. Characterization of polluted sediments in the city canals of Delft, The Netherlands. Proceedings of CATS IV Conference of Characterization and treatment of sediments, Antwerpen, Belgium, September 1999.
- KRALIK, M., 1998. A rapid procedure for environmental sampling and evaluation of polluted sediments. *Applied Geochemistry*, 1-10.
- KRSTIC, S. & MELOVSKI, Lj., 1994. Preliminary results of the saprobiological investigations on the River Vardar. Proceedings of Workshop: Monitoring of the environment in Republic of Macedonia. The Society of Ecologists of the Republic of Macedonia, 2:151 - 159. (in Macedonian).
- KRSTIC S., MELOVSKI LJ., LEVKOV Z. & STOJANOVSKI P., 1994. Complex investigations of the River Vardar. II. The most polluted sites in the first three mounts. *Ekologia i Zastita na Zivotnata Sredina*, 2.(2): 13-29.
- KRSTIC, S., 1995. Saprobiological characteristics of microflora in reka Vardar as indicator of the intensity of antropogenic influence. Ph.D. Thesis Faculty of Natural Sciences - Skopje. 387p. (in Macedonian).
- KRSTIC, S., LEVKOV, Z. & STOJANOVSKI, P., 1996. Diatoms in monitoring of the River Vardar (Macedonia). *Ekologia*, 32 (2): 1-16.
- KRSTIC, S., LEVKOV, Z. & STOJANOVSKI, P., 1997. Saprobiological characteristics of diatom microflora in river ecosystems in Macedonia as a parameter for determination of the intensity of antropogenic influence. In: Use of Algae for Monitoring Rivers III, edited by Prygiel J, Whitton BA Bukowska J. 145-153.
- LEVKOV, Z. (2001): Distribution of heavy metals in the River Vardar and their influence on microflora and aquatic mosses. MSci thesis, "St. Cirilus and Methodius" University, Skopje, p 203p.
- MANTAI, K.E., GARWOOD, P.E. & PEGLOW-SKI L.E., 1982. Environmental factors controlling physiological changes in *Cladophora* in Lake Erie. *Journal of Great Lakes Research*, 8 (1): 61-65.
- MARKERT, B., OEHLMANN, J. & ROTH, M., 1997. General Aspects of Heavy Metal Monitoring by Plants and Animals. In: Environmental Monitoring - Exposure Assessment and Specimen Banking, edited by K.S. Subramanian and G.V. Iyengar, ACS Symposium Series 654, Washington DC, 18-29.
- MELOVSKI, LJ., LEVKOV, Z. & KRSTIC, S., 1997. Concentration of phosphorus and heavy metals in river sediment in the River Vardar. *Ekologia i Zastita na Zivotnata Sredina*, 5 (2): 127-132. (in Macedonian).

- MILLNER, G.C., SWEENEY, R.A. & FREDERICK, V.R., 1982. Biomass and distribution of *Cladophora glomerata* in relation to some physical-chemical variables at two sites in Lake Erie. *Journal of Great Lakes Research*, 8 (1): 35-41.
- MOORE, P. & CHAPMAN, S., 1986. Methods in plant Ecology. Blackwell Scientific Publications. Oxford, 590p.
- NEIL, J.H. & JACKSON, M.B., 1982. Monitoring *Cladophora* growth conditions and the effect of phosphorus additions at a shoreline site in the northeastern Lake Erie. *Journal of Great Lakes Research*, 8 (1): 30-34.
- OERTEL, N., 1991. Heavy-metal accumulation in *Cladophora glomerata* (L.) Kütz. in the River Danube. *AMBIO* 20 (6): 264-268.
- OERTEL, N., 1993.: The applicability of *Cladophora glomerata* (L.) Kütz. in an active bio-monitoring technique to monitor heavy metals in the river Danube. *Science of the Total Environment*, Nr. Suppl. Part 2, 1293-1304.
- PICKERING, D.C. & PUIA, I.L., 1969. Mechanisms for the uptake of zinc by *Fontinalis antipyretica*. *Physiologia Plantarum*, 22, 653-661.
- PELLETIER, G., 1996. Applying Metals Criteria to Water Quality- Based Discharge Limits. Empirical Models of the Dissolved Fraction of Cadmium, Copper, Lead, and Zinc. Washington State Department of Ecology. Publication No. 96-339, pp. 24.
- RANG, S.A. & STOKES, P.M., 1987. Seasonal variation and uptake and loss of cadmium, lead and mercury in *Cladophora* in Niagara River: In: Heavy metals in the environment, Vol.2, Proceedings of International Conference, edited by Lindberg S.E. Hutchinson T.C., New Orleans, September 1987, 259-261.
- RECZYNSKA-DUTKA, M., 1991. Evaluation of chemical factors controlling heavy metal concentrations in waters of three selected reservoirs. *Polskie Archive for Hydrobiology*, 38 (1): 1-10.
- ROSEMARIN, A.S., 1985. Reproductive strategy in the filamentous alga *Cladophora glomerata* (L.) Kütz. - an explanation for its widespread distribution. Proceedings of Congress of International Association of Theoretical and Applied Limnology, 22, 2872-2877.
- ROUND F.E., 1991. Methods for use of epilithic diatoms for detection and monitoring changes in river water quality. *Journal of Applied Phycology*, 3: 129-145.
- RULE, J.H. & ALDEN, R.W., 1992. Partitioning of Cd in Geochemical Fractions of Anaerobic Estuarine Sediments. *Estuarine, Coastal and Shelf Science*, 34: 487-499.
- RULE, J.H. & ALDEN, R.W., 1996. Interaction of Cd and Cu in anaerobic estuarine sediments. I. Partitioning in geochemical fractions of sediments. *Environmental Toxicology and Chemistry*, 15 (4):460-465.
- SAKAGUCHI, T., TSUJI, T., NAKAJIMA, A. & HORIKOSHI, T., 1979. Studies in the accumulation of heavy metal elements in biological systems. Accumulation of cadmium by green microalgae. *European Journal of Applied Microbiology and Biotechnology*, 8: 207-215.
- SCHANZ, F. & THOMAS, E.A., 1981. Cultures of *Cladophoraceae* in water pollution problems. Proceedings of Congress of International Association of Theoretical and Applied Limnology, 21: 57-64.
- SHEATH, R.G. & COLE, K.M. 1992. Biogeography of stream macroalgae in North America. *Journal of Phycology*, 28: 448-460.
- SOBHAN, R. & STERENBERG, S.P.K, 1999. Cadmium removal using *Cladophora*. *Journal of Environmental Science and Health*, 34 (1): 53-72.
- StatSoft, Inc., 1995. STATISTICA for Windows [Computer program manual]. Tulsa, OK, USA.
- TACK, F.M.G. & VERLOO, M.G., 1995. Chemical speciation and fractionation in soils and sediment heavy metal analysis: A review. *International Journal of Environmental Analytical Chemistry*, 59: 225-238.
- TACK, F.M., CALLEWAERT, O.W.J.J. & VERLOO, M.G., 1996. Metal solubility as a function of pH in a contaminated, dredged sediment affected by oxidation. *Environmental Pollution*, 91 (2): 199-208.
- TORRES, E., CID, A., FIDALGO, P., HERRERO, C. & ABALDE, J., 1995. Tolerance and detoxification mechanisms in marine diatom *Phaeodactylum tricorutum* exposed to cadmium. *Journal of Marine Biotechnology*, 3: 176-178.
- TORRES, E., CID, A., FIDALGO, P., HERRERO, C. & ABALDE, J., 1997. Long-chain class III methallothioneins as a mechanism of cadmium tolerance in the marine diatom *Phaeodactylum tricorutum* Bohlin. *Aquatic Toxicology*, 39: 231-246.
- VELJANOVSKI, A., 1982. Content of cadmium in the River Vardar water during period of 1978-1980. Proceedings of Meeting "Protection, management

- and improvement of freshwater quality". Ohrid, R. Macedonia. 3-4.06.1982. (in Macedonian).
- VELJANOVSKI, A., 1987. Content of lead, cadmium and zink in surface waters in periods 1976-1980 and 1981-1985. Report for IBID. pp.73-80.
- VERLOO, M.G. & COTTENIE, A., 1985. Influence of redoxpotential and pH on the transfer of heavy metals from solid to the liquid phase in river sediments. *Med Fac Landbouww Rijksuniv Gent* 50 (1): 47-53.
- VOGEL, W.R. & CHOVANEC, A. 1992. Sediment analysis as a method of monitoring industrial emissions. *Hydrobiologia*, 235/236: 723-730.
- VYMAZAL, J., 1987.: Zn uptake by *Cladophora glomerata*. *Hydrobiologia*, 148, 97-101.
- VYMAZAL, J., 1990a. Uptake of heavy metals by *Cladophora glomerata*. *Acta Hydrochimica et Hydrobiologica*, 18, 657-665.
- VYMAZAL, J., 1990b. Uptake of lead, chromium, cadmium and cobalt by *Cladophora glomerata*. *Bulletin of Environmental Contamination and Toxicology*, 44: 468-472.
- VYMAZAL, J., 1995. Influence of pH on heavy metals uptake by *Cladophora glomerata*. *Polskie Archive for Hydrobiology*, 43 (3): 231-237.
- WANG, W., 1987. Factors affecting metal toxicity to (and accumulation by) aquatic organism - overview. *Environment International*, 13. 437-457.
- WANG, W. & LEWIS, A.M., 1997. Metal accumulation by aquatic macrophytes. p.367-416. In: *Plants for Environmental Studies*, edited by W. Wang, J.W. Gorsuch, and J.S. Hughes. Lewis Publishers. New York.
- WANG, W.X. & DEI, R.C.H., 2001a. Metal uptake in a coastal diatom influenced by major nutrients (N,P, and Si). *Water Research*, 35: 315-321.
- WANG, W.X. & DEI, R.C.H., 2001b. Influence of phosphate and silicate on Cr (VI) and Se (IV) accumulation in marine phytoplankton. *Aquatic Toxicology*, 52: 39-47.
- WANG, W.X. & DEI, R.C.H., 2001c. Effects of major nutrient additions on metal uptake in phytoplankton. *Environmental Pollution*, 111: 233-240.
- WEHR, J.D. & WHITTON, B.A., 1983a. Accumulation of heavy metals by aquatic mosses. 2: *Rhinostegium riparoides*. *Hydrobiologia* 100, 261-284.
- WEHR, J.D. & WHITTON, B.A., 1983b. Accumulation of heavy metals by aquatic mosses. 3: Seasonal changes. *Hydrobiologia* 100, 285-291.
- WEHR, J.D., KELLY, M.G. & WHITTON B.A., 1987. Factors affecting accumulation and loss of zinc by the aquatic moss *Rhinostegium riparoides* (Hedw) C.Jens. *Aquatic Botany*, 19: 261-274.
- WHITTON, B.A., 1970. Biology of *Cladophora* in freshwaters. *Water Research*, 4: 457-476.
- WHITTON, B.A. & SAY P.J. 1975. Heavy metals. p.286-311. In: *River Ecology*, edited by B.A. Whitton, Blackwell, Oxford.
- WHITTON, B.A., SAY, P.J. & WEHR, J.D., 1981. Use of plants to monitor heavy metals in rivers. p.135-146. In: *Heavy metals in Northern England: Environmental and Biological Aspects*, edited by P.J.Say and B.A. Whitton, Department of Botany, University of Durham.
- WHITTON, B.A., BURROWS, I.G. & KELLY, M.G., 1989. Use of *Cladophora glomerata* to monitor heavy metals in rivers. *Journal of Applied Phycology*. 1: 293-299.
- WHITTON, B.A., HARDING, J.P.C., KELLY, M.K. & SAY, P.J., 1991. Use of Plants to Monitor Heavy Metals in Freshwaters. Standing Committee of Analysts 1990. HMSO, 43p.
- WHITTON, B.A. & KELLY, M.G., 1995. Use of algae and other plants for monitoring rivers. *Australian Journal of Ecology*, 20: 45-56.
- WIESE, G., BAHR, I., FUCHS, S. & SCHUER-MANN, L., 1986. Phosphate uptake and gas metabolism of *Myriophyllum spicatum* and *Cladophora glomerata*. *Acta Hydrochimica et Hydrobiologica*. 14 (5): 475-484.
- WOODROOFF, S.L., HOUSE, W.A., CALLOW, M.E. & LEADBEATER B.S.C. (1999): The effects of biofilms on chemical processes in surficial sediments. *Freshwater Biology*, 41: 73-89.
- ZHOU X.D. & KOT S.C. (1995): Heavy metal ion adsorption on sediments of the Weiho and Hanjiang rivers, China. *Journal of Environmental Hydrology*,. 3 (2): 1-12.