



Mediterranean Marine Science

Vol 2, No 1 (2001)



Mediterranean climate variability during the Holocene

J.S.L. CASFORD, R. ABU-ZIED, E.J. ROHLING, S. COOKE, K.P. BOESSENKOOL, H. BRINKHUIS, C. DE VRIES, G. WEFER, M. GERAGA, G. PAPATHEODOROU, I. CROUDACE, J. THOMSON, V. LYKOUSIS

doi: 10.12681/mms.275

To cite this article:

CASFORD, J., ABU-ZIED, R., ROHLING, E., COOKE, S., BOESSENKOOL, K., BRINKHUIS, H., DE VRIES, C., WEFER, G., GERAGA, M., PAPATHEODOROU, G., CROUDACE, I., THOMSON, J., & LYKOUSIS, V. (2001). Mediterranean climate variability during the Holocene. *Mediterranean Marine Science*, *2*(1), 45–56. https://doi.org/10.12681/mms.275 Vol. 2/1, 2001, 45-55

Mediterranean climate variability during the Holocene

J.S.L. CASFORD¹, R. ABU-ZIED¹, E.J. ROHLING¹, S. COOKE¹, K.P. BOESSENKOOL², H. BRINKHUIS², C. DE VRIES², G. WEFER³, M. GERAGA⁴, G. PAPATHEODOROU⁴, I. CROUDACE¹, J. THOMSON¹ and V. LYKOUSIS⁵

¹ Southampton Oceanography Centre, Southampton SO14 3ZH, UK e-mail: E.Rohling@soc.soton.ac.uk

- ² Laboratory for Paleobotany and Palynology, Utrecht University, 3584CD Utrecht, Netherlands
- ³ Fachbereich Geowissenschaften, Universitat Bremen, Denmark
- ⁴ Department of Geology, University of Patras, Greece

⁵ National Centre for Marine Research, Athens, Greece

Manuscript received: 13 July 1999; accepted in revised form: 28 December 2000

Abstract

We present a study on four high sedimentation-rate marine cores with suppressed bioturbation effects, recovered along the northern margin of the eastern Mediterranean. We demonstrate that this region, central to the development of modern civilisation, was substantially affected throughout the Holocene by a distinct cycle of cooling events in the order of 2° C. In the best-preserved cases the onset of these events appears particularly abrupt, within less than a century. The cooling events typically lasted several centuries, and there are compelling indications that they were associated with increased aridity in the Levantine/NE African sector (Rossignol-Strick, 1995; 1998; Alley et al., 1997; HASSAN, 1986; 1996; 1997a,b; MCKIM MALVILLE et al., 1998). Several of these episodes appear to have been coincident with cultural reorganisations, with indigenous developments (eg. cattle domestication, new technologies) and population migrations and fusion of peoples and ideas (HASSAN, 1986; 1996; 1997a,b; MCKIM MALVILLE, 1998). We infer that climatic events of a likely high-latitude origin (O'BRIEN et al., 1995; BOND et al., 1997; MAYEWSKI et al., 1997; ALLEY et al., 1997) caused cooling and aridity in and around the eastern Mediterranean via a direct atmospheric link, and therefore played an important role in the development of modern civilisation.

Keywords: Climatic variability, Palaeoceanography, Aegean, Holocene, Foraminifera, Stable isotopes.

Introduction

The Holocene is customarily regarded as et al., 1995; BOND et al., 1997; CAMPBELL et a period of climate stability, but recently a *al.*, 1998; BIANCHI & MCCAVE, 1999). This distinct millenium-scale repetition of Holo- climatic variability may have driven migracene cold events has been recognised (O'BRIEN tory patterns and settlement behaviour in

early societies (HASSAN, 1986; 1996; 1997a, b), but it remains to be established how high-latitude Holocene climate variability affected the near/middle East region. One measure of such climatic variability it is fluctuations in past sea surface temperature (SST) estimated from marine planktonic microfossil records (CITA, 1977; THUNELL, 1977; ROHLING et al., 1993; PFLAUMANN et al., 1996; KALLEL et al., 1997; ORTIZ & MIX, 1997; ROHLING et al., 1997; TARGARONA, 1997; HAYES et al., 1999). We assess the Late Glacial-Holocene SST changes from planktonic foraminiferal abundance variations in four cores (Table 1). IN68-9 from the S Adriatic Sea; C40 from the SW Aegean Sea; LC21 from the SE Aegean Sea; and LC31 from W of Cyprus (Fig. 1), using a simple a-priori grouping of warm versus cool species (expressed as % warm species/ (%warm + %cool species), Table 2; Fig. 2 and 3). Similar groupings are found from multivariate statistical analyses of Mediterranean records (CITA, 1977; THUNELL, 1977; ROHLING et al., 1993; HAYES et al., 1999). In subtropical waters, the species of the "warm" group predominate within the shallow, warm, and nutrient-depleted summer mixed layer. Many are spinose and contain photosynthetic symbionts. Species of the "cool" group are mostly present in winter, when SST is reduced and the mixed layer has considerably

Core characteristics.						
Core	Depth					
LC-21	350 39.71' N	1522 m				
	260 34.96' E					
LC-31	340 59.76' N	2298 m				
	31o 09.81' E					
IN68-9	410 47.5' N	1234 m				
	17o 54.5' E					
C-40	360 56.12' N	852 m				
	240.04.69' F					

Table 1

deepened (> 100 m) and become enriched in nutrients. Despite an obvious overprint of the degree of eutrophication related to the development of the seasonal thermocline, recent modern analogue studies show that such groupings provide useful first-order information on SST fluctuations (PFLAU-MANN et al., 1996; KALLEL et al., 1997; ORTIZ & MIX, 1997). In corroboration of the SST fluctuations based on planktonic foraminifera (zooplankton), we include a similar warm and cool water grouping for core IN68-9 of selected organic walled cyst species derived from auto-trophic dinoflagellates (phytoplankton) that generally preserve well (Table 2). Both qualitative records differ in detail due to the effects of productivity and preservation constraints, but the basic patterns in intervals with comparable sampling resolution are quite similar (Fig.3).



Fig. 1: Map of Eastern Mediterranean, showing locations of cores used in this study.







Fig. 3: Records of "warm" versus "cold" planktonic foraminiferal species as discussed in the text. LC-21 is plotted versus time, plots of IN68-9 for clarity show depth subtracting thicknesses of turbidites and ash-layers, and for LC31 and C40 versus true depth. Planktonic foraminiferal biozone boundaries after ORISSEN et al. (1993). Cooling intervals are numbered sequentially in bold. Main cooling events in the dinoflagellate-based warm-cold record (see text) for IN68-9 are numbered sequentially in bold italics, with the prefix "d". In LC21, the dinoflagellate-based temperature record is near 100% throughout the phies are based on G. bulloides for IN68-9 and G. ruber for LC21 and LC31, from the 250-350 mm size-window. Error-bars in LC31 represent range of variability observed in replicate analyses, for analyses added at the S.O.C. isotope facility to an originally less detailed record analysed at Bremen. The two Ferminations T1a and T1b. Within the basal sapropel unit, relatively small effects of temperature change on δ^{18} O are overwhelmed by those of the dra-Grey bands represent sapropelic intervals. The cross-hatched interval within the sapropel in LC31 represents a slump (Table 3). Oxygen isotope stratigraacilities were intercalibrated by exchange of laboratory standards and replicate sample analyses. Bold lines indicate the two main deglaciation steps: matic summer rainfall increase witnessed by a peak in deciduous forest pollen in both IN68-9 (TARGARONA, 1997) and LC21 (preliminary record; not shown). Warm-cold record for C40 excludes Turborotalita quinqueloba, a small-size species that is over represented in this > 120 mm study, relative to the Holocene, and is compromised by low cyst abundances outside the sapropelic intervals; levels where minor cooling events were found are marked "d--other three $>150 \ \mu m$ studies.

Planktonic foraminifera (zooplankton)				
Warm group	Cool group			
Globigerinoides ruber	Globorotalia scitula			
Globigerinoides sacculifer	Turborotalita quinqueloba			
Hastigerina pelagica	Globorotalia inflata			
Globoturborotalita rubescens	Neogloboquadrina pachyderma			
Orbulina universa				
Globigerinella digitata				
Globoturborotalita tenella				
Globigerinella siphonifera				
Selected autotrophic dinoflagellate cyst species (phytoplankton)				
Warm group	Cool group			
Impagidinium aculeatum	Bitectatodinium tepikiense			
Impagidinium strialatum	Spiniferites elongatus			
Impagidinium paradoxum	Impagidinium pallidum			
Spiniferites mirabilis				

 Table 2

 Composition of species-groups used in planktonic foraminiferal and dinoflagellate warm-cold plots.

Time-stratigraphic framework

The chronostratigraphy is determined by several AMS14C analyses in each of cores IN68-9, LC21, LC31 and C40, all performed on hand-picked clean planktonic foraminiferal tests (Table 3). A calendar age of 3.578 ka BP (KUNIHOLM et al., 1996) is used for the Santorini ash layer at 81-91 cm in LC21, and 33-36 cm in C40. We calibrate all radiocarbon convention ages using the marine mode of the Calib3.03 programme (STUIVER & REIMER, 1993). It is immediately obvious that some of the AMS14C datings tend towards older ages than expected from the well-established ages of the biozonal boundaries and the Santorini ash layer (Table 3, Fig. 3). This tendency is mostly minor (e.g. the ages around the Santorini ash in LC21), but in some cases a substantial offset is seen. For example, the age immediately below the 5 cm thick slump within the sapropel in LC31 appears about 1000 years too old. This is probably due to the immediate proximity of the mass-transport layer, which itself is about 3000 years older than surrounding sediments. Offsets towards older radiocar-

bon ages are potentially caused by addition of an uncertain component of older tests to the material analysed, which seems possible in view of the high accumulation rates of the cores and their proximity to the continental slopes. Calibrated ages for LC21 are then plotted versus depth subtracting the thicknesses of the interbedded Santorini ash layer and turbidites. As LC21 has the highest resolution of the four cores in the study and appears to be the most complete record, it is used as a master chronology in the following discussion (Fig. 2). All ages are reported as calibrated AMS¹⁴C years BP unless otherwise stated.

Bio-stratigraphic control also provides two well-dated correlation horizons, the biozone I-II and II-III boundaries, as defined on the basis of 11 central Mediterranean marine records with a total of 50 AMS¹⁴C ages (JORISSEN *et al.*, 1993). Subsequent studies emphasised Mediterranean-wide applicability and synchroneity of both boundaries (CAPOTONDI *et al.*, 1999; HAYES *et al.*, 1999). This includes the Aegean Sea

Table 3

Age control-points used in the present study, with true depths in the cores, and corrected depths (subtracting thicknesses of turbidites and ash-layers). I-II and II-III are planktonic foraminiferal biozone boundaries (JORISSEN *et al.*, 1993). Radiocarbon convention ages calibrated using marine mode of programme Calib3.03 (STUIVER & REIMER, 1993). Santorini age after KUNIHOLM *et al.* (KUNIHOLM *et al.*, 1996). Core LC-31 dates include a local ΔR correction of 149±39 years after FACORELLIS et al., (FACORELLIS *et al.*, 1998). Dates in LC-21 are after MERCONE *et al.*, (MERCONE *et al.*, 1999) and in C-40 after GERAGA *et al.*, (GERAGA *et al.*, 2000).

True depth (cm)	Corrected depth (cm)	AMS lab.code	Uncorrected AMS ¹⁴ C age from direct dating or dated horizon (kaBP)	cal.age (ka BP)	cal.age range (1σ) (kaBP)	
IN68-9						
11.5	7.5	UTC-500	3.16 ± 0.12	2.93	2.78-3.10	
54.5	38.5	UTC-1607	6.39 ± 0.06	6.85	6.77-6.90	
157.25	81.25	UTC-501	9.28 ± 0.18	9.93	9.80-10.08	
241.5	157.5	UTC-502	13.10 ± 0.20	14.95	14.61-15.30	
322.5	201.5	UTC-503	14.20 ± 0.30	16.55	16.15-16.92	
510.5	247.5	UTC-504	17.20 ± 0.30	19.82	19.40-20.30	
LC-21						
50	50	CAMS-41314	3.35 ± 0.06	3.20	3.11-3.27	
95.5	85.5	CAMS-41313	4.29 ± 0.06	4.40	4.32-4.45	
137.75	127.75	CAMS-41311	5.59 ± 0.06	5.95	5.90-6.03	
161.5	151.5	CAMS-41315	7.48 ± 0.06	7.90	7.94-7.84	
174.25	164.25	CAMS-41312	8.12 ± 0.06	8.52	8.59-8.44	
179.5	169.5	AA-30364	9.01 ± 0.07	9.54	9.22-9.86	
209	199	AA-30365	11.77 ± 0.08	13.14	12.77-13.51	
252.5	242.5	CAMS-41316	14.45 ± 0.06	16.85	16.96-16.74	
81-91	81	Santorini	3.578			
LC-31						
28.5	28.5	CAMS-45864	3.45 ± 0.05	3.33	3.26-3.36	
60.5	60.5	CAMS-45863	6.12 ± 0.05	6.54	6.47-6.62	
84.25	81 (slump)	AA-30367	10.90 ± 0.10	12.42	12.27-12.55	
87.5	82.5	CAMS-45861	8.74 ± 0.05	9.38	9.34-9.43	
96.5	91.5	CAMS-45862	8.50 ± 0.05	9.06	8.98-9.19	
131.5	126.5	CAMS-45860	12.04 ± 0.05	13.57	13.47-13.69	
247.5	242.5	CAMS-45859	32.96 ± 0.05	35.96		
				(simple addition	(simple 3kyr addition)	
C-40						
73.5	73.5	Beta-110420	6.83 ± 0.11	7.30	7.20-7.39	
82.5	82.5	Beta-110419	7.83 ± 0.14	8.26	8.10-8.38	
131	131	Beta-110418	12.35 ± 0.16	13.93	13.72-14.16	
33-36	33-36	Santorini	3.578			

(ZACHARIASSE et al., 1997), as is shown also by the results of core C40 presented here. The II/III biozonal boundary in the Adriatic however, does not appear to be entirely synchronous with the corresponding faunal break in our other cores. The chronostratigraphy of Adriatic core IN68-9 gives ages for the zone I-II and II-III boundaries of 10.5 cal ka BP and 14.5 cal ka BP, respectively (JORISSEN et al., 1993; ROHLING et al., 1993, 1997). We note that IN68-9 presents a date almost 1000 years younger for the II/III boundary than that observed in their other cores (JORISSEN et al., 1993) or in the cores presented here. As yet, there is no satisfactory explanation but it is possible that a small hiatus is present in IN68-9 over the II/III boundary.

Cooling events

We identify several cooling events, these are sequentially numbered (in bold), noting their positions relative to the biozonal boundaries, ash layers, and sapropelic units (Fig. 2 and 3). Events 1 and 6 are less accurately timed because of their occurrence near the limits of the investigated records where resolution is lowest. The 2a event (9.6-9.8 ka BP) is only identified positively from core IN68-9 with limited support from the other cores outside the Adriatic. The 1a event (13.4-13.6 ka BP) is only identified in the base of core LC-31, and needs to be considered with caution. The age-ranges determined for the NE Mediterranean cooling events are 1.5-2.0, 3.6-4.0, 5.8-6.2, 8.2-8.6, 9.6-9.8, 11.4-12.4, 13.4-13.6, and 17.0-18.5 ka BP, similar to ages reported for high-latitude N. Atlantic cold events in the Holocene (Table 4) (O'BRIEN et al., 1995; BOND et al., 1997).

The magnitude of the 8 ka BP event (our event 3) in IN68-9 has been estimated in the order of 2 C-degrees (ROHLING *et al.*, 1997). For a full glacial-interglacial temperature difference in the Aegean Sea of about

6 C-degrees (ROHLING & DE RIJK, 1999), comparison of the magnitudes of the cooling events with the full glacial-interglacial amplitude of change in the warm/cold plot of LC21 (Fig. 3) also suggests effects of the order of 2 C-degrees.

The rapidity of the onset of the cooling events is best illustrated using the 8 ka BP event (event 3), since it follows the deposition of the unbioturbated (abenthic) lower sapropelic unit. In LC21, this event starts from one sample to the next. Due to the high accumulation rate in this core we infer that the period of onset for this event was less than a century (Fig. 2). The cooling events typically last several centuries, and appear to be repeated at regular intervals.

Discussion

Together, the timing of the events and the apparent rapidity of changes in the eastern Mediterranean suggest a direct atmospheric link between high latitudes and the NE Mediterranean. Regionally - expressions of cold Atlantic Heinrich events in the glacial Gulf of Lions (NW Mediterranean) also suggest an efficient atmospheric teleconnection between that area and the high latitudes (ROHLING et al., 1998). Today, high latitude Arctic/N.Atlantic perturbations are rapidly transmitted to the northern edge of the Mediterranean via orographically channelled winter outbreaks of cold and dry air high latitudes over the from NW Mediterranean, the Adriatic and the Aegean Sea (LEAMAN & SCHOTT, 1991; MARIOPOULOS, 1961; POULOS et al., 1997). An increased duration and/or intensity of these atmospheric outbreaks would cause not only cooling, but also aridity. Palaeobotanical records around the Aegean Sea (ROSSIGNOL-STRICK, 1995), and the $\delta^{18}O$ and $\delta^{13}C$ profiles of Soreq Cave (Israel) (BAR MATTHEWS et al., 1999) both suggest notable cooling and aridity during the Younger Dryas and around 8 ka BP.

Table 4

Comparison of ages of abrupt climate events. This paper: planktonic foraminifera-based cooling events in the Adriatic and Aegean/NE Levantine Seas. BOND *et al.*, 1997: high latitude N. Atlantic cold events from marine records compared with GISP2 ice record. HASSAN *et al.*, 1997a,b and MCKIM MALVILLE *et al.* 1998: NE African arid/cold spells and impact on early societies. GASSE *et al.*, 1996: aridity in the Lake Bangong area (W. Tibet). O'BRIEN *et al.*, 1995: cold events with increased meridional atmospheric circulation affecting the Greenland Ice Sheet. Last column: mean ages based on boundaries of discrete events listed in the table, with standard error of the mean (cal. ka BP).

This paper	Bond et	Hassan	McKim. et al,	Gasse et al.,	O'Brien	Event
	al.,1997	1997a,b	1998	1996	et al., 1995	
				0.7	0.0-0.6	
1.5	1.2-1.7	1.5-2.0		1.3-2.1		6?
3.2-4.0	2.5-3.1				2.4-3.1	5
	4.1-4.8	4.0-4.4		<3.9		
5.8-6.2	5.6-6.3	5.5-6.7*	<5.8 arid	5.7-6.3	5.0-6.1	4
8.2-8.6	7.6-8.5	7.7-8.3	7.4-8.1***	7.7-8.2	7.8-8.8	3
9.6-9.8	9.4-9.7	9.2-9.5		9.6-9.9		2a‡
11.4-12.4	10.7-12.6**				>11.3	2
13.4-13.6	13.0-13.4	13.4-14.0				1a ^{‡‡}
	14.6-15.0					
17.0-18.5						1

^{*} Onset of major drought and cold conditions around 6.7 ka BP, culminating around 5.5 ka BP with shift of societies into active Nile channels (HASSAN, 1997b).

** Their event 8 and Y.D. combined (BOND et al., 1997).

^{***} Combination of two separately reported events (7.4-7.5 and 7.9-8.1) (MCKIM MALVILLE *et al.*, 1998).

[‡] Only weakly supported outside of Adriatic.

^{‡‡} Identified in LC-31 only, to be treated with caution.

Moreover, overviews of continental palaeoclimate and archaeological records from the Levant and NE Africa show arid spells of similar age to our Adriatic/Aegean cooling events (Table 4), coinciding with important reorganisations in settlement patterns and agricultural behaviour of early societies (HASSAN, 1997a,b; MCKIM MALVILLE et al., 1998). For example, "droughts ... between 8.3 and 7.7 ka BP encouraged movements of nomadic cattle keepers from the Egyptian Sahara and bands possessing sheep/goats and wheat/barley from Southwest Asia to the banks of the Nile Valley and Nile Delta. This event was crucial for the agricultural developments in the Nile Valley" (HASSAN, 1998) (our event 3). In addition, "It was during the

7th and 6th millennium bp episodes of severe droughts that dwellers of the desert drifted towards the Nile Valley, and elsewhere, in search of water, food, and fodder" (HASSAN, 1997), (our event 4, see Table 4). We infer that the development of modern civilisation was punctuated by abrupt cooling/aridity events due to an atmospheric teleconnection between the "cradle of modern civilisation" and the high latitudes. We propose that this connection operates through significant increases in winter northerly air flow over the eastern Mediterranean.

The observed temporal spacing of these cooling events appears (Table 4) similar to that of planet-induced sun-tide cycles (LAMB, 1972). Ice-core 14C residual series

indicate this periodicity to be around 1450 years, suggesting that fluctuations in solar forcing may indeed have affected glacial and Holocene climate on these time-scales (MAYEWSKI et al., 1997). However, any external forcing would be likely to drive internal oscillations within the earth's ocean/climate system, such as the recently invoked "oceanic oscillator", a fluctuation in the global thermohaline circulation, the nature of which remains to be determined (BIANCHI & MCCAVE, 1999). We tentatively propose that the Mediterranean may form part of the "oceanic oscillator". Although the Mediterranean appears to behave initially as a passive response-basin to high-latitude cold events, resultant increases in buoyancy loss would increase its saline outflow flux into the Atlantic. Mediterranean salt today contributes to the water-column destabilisation needed for North Atlantic Deep Water (NADW) formation (eg. REID, 1979; HILL & MITCHELSON-JACOB, 1993). If the Mediterranean salt flux into the Atlantic were sufficiently increased to trigger enhanced NADW formation (for assessment of current hypotheses, see BRYDEN & WEBB, 1998), then consequent increases in heat-advection to high latitudes could have contributed to termination of high-latitude cold events. This possibility merits consideration, particularly in view of the anthropogenic impact on Mediterranean deep water properties and the recently-observed and rapid major salt redistribution within the eastern Mediterranean (BETHOUX et al., 1990; LEAMAN & SCHOTT, 1991; ROHLING & BRYDEN, 1992; BETHOUX & GENTILLI, 1996; ROETHER, 1996).

Acknowledgements

This paper represents a joint effort of EC-MAST3 programmes "Climatic variability of the Mediterranean paleo-circulation" and "Sapropels and palaeoproductivity". Cores LC21 and LC31 were recovered

within EC-MAST2 programme PALAE-OFLUX and are held at the BOSCOR repository in Southampton. We thank M. Segl at the isotope laboratory in Bremen, F. Hassan, M. Rossignol-Strick, F.J. Jorissen, and J. Targarona for valuable discussions, the NERC laboratory for AMS radiocarbon analyses, the Egyptian Government for support to RA-Z, K.A.F. Zonneveld and J. Targarona for analysis and synthesis of the dinoflagellate and pollen records of IN68-9, and J. Kallmeyer for his contribution to study of LC31.

References

- ALLEY, R.B., MAYEWSKI, P.A., SOWERS, T., STUI-VER, M., TAYLOR, K.C. & CLARCK, P.U., 1997. Holoce ne climatic instability: a prominent, widespread event 8200 yrs ago, *Geology* 25, 483-486.
- BAR-MATTHEWS, M., AYALON, A., KAUFMAN, A. & WASSERBURG, G.J., 1999. The eastern Mediterranean paleoclimate as a reflection of regional events: Soreq cave, Israel, *Earth and Planetary Science letters* 166, 85-95.
- BETHOUX, J.P. & GENTILLI, B., 1996. The Mediterranean Sea, coastal and deep-sea signatures of climatic and environmental changes, *Journal of Marine Systems* 7, 383-394.
- BETHOUX, J.P., GENTILLI, B., RAUNET, J. & TAILLEZ, D., 1990. Warming trend in the western Mediterranean deep water, *Nature* 347, 660-662.
- BIANCHI, G.G. & MCCAVE, N., 1999. Holocene periodicity in North Atlantic climate and deepocean flow south of Iceland, *Nature* 397, 515-517.
- BOND, G., SHOWERS, W., CHESEBY, M., LOTTI, R., ALMASI, P., DEMENOCAL, P., PRIORE, P., CUL-LEN, H., HAJDAS, I. & BONANI, G., 1997. A pervasive millenial-scale cycle in North Atlantic Holocene and Glacial climates, *Science* 278, 1257-1266.
- BRYDEN, H.L. & WEBB, D.J., 1998. A perspective on the need to build a dam across the Straits of Gibraltar, *Newsletter European Geophysical Society* 69, 1-3.
- CAMPBELL, I.D., CAMPBELL, C., APPS, M.J., RUT-TER, N.W. & BUSH, A.B.G., 1998. Late Holocene ~1500 yr climatic periodicities and their impli-

cations, Geology 26, 471-473.

- CAPOTONDI, L., BORSETTI, A.M. & MORIGI, C., 1999. Foraminiferal ecozones, a high-resolution proxy for the late Quaternary biochronology in the central Mediterranean Sea, *Marine Geology* 153, 253-274.
- CITA, M.B., VERGNAUD-GRAZZINI, C., ROBERT, C., CHAMLEY, H., CIARANFI, N. & D'ONOFRIO, S., 1977. Paleoclimatic record of a long deep sea core from the eastern Mediterranean. Quaternary Research 8, 205-235.
- FACORELIS, Y., MANIATIS, Y. & KROMER, B., 1998. Apparent 14C ages of marine mollusk shells from a Greek island: calculation of the marine reservoir effect in the Aegean Sea. *Radiocarbon* 40, 963-973.
- GASSE, F., FONTES, J.CH., VAN CAMPO, E. & WEI, K., 1996. Holocene environmental changes in the Bangong Co basin (western Tibet). Part 4: Discussion and conclusions, *Palaeogeography*, *Palaeoclimatology*, *Palaeoecology* 120, 79-92.
- GERAGA M., TSAILA-MONOPOLIS, ST., IOAKIM, C., PAPATHEODOROU, G., FERENTINOS, G., 1999. Evaluation of palaeoenvironmental changes during the last 18,000 years in the Myrtoon basin, SW Aegean Sea. *Palaeogeography, Palaeoclimatology, Palaeoecology.*, 156, 1-17
- HASSAN, F.A., 1986. Desert environment and origins of agriculture in Egypt, *Norwegian Archae*ological Review 19, 63-76.
- HASSAN, F.A., 1996. Abrupt Holocene climatic events in Africa, in *Aspects of African Archaeology* (eds Pwiti, G. & Soper, R.), 83-89. (University of Zimbabwe, Harare)
- HASSAN, F.A., 1997. Holocene palaeoclimates of Africa, African Archaeological Review 14, 213-230.
- HASSAN, F.A., 1997. Nile Floods and political dis order in early Egypt, in third millenium BC cli-
- mate change and old world collapse (eds Nuzhet Dalfes, H., Kukla, G. & Weiss, H.), NATO ASI Series I 49, 1-23 (Springer-Verlag, Berlin).
- HASSAN, F.A., 1998. Holocene climatic change and riverine dynamics in the Nile Valley, in Before food production in North Africa, Union Internationale de Sciences Préhistoriques et protohistoriques, XIII world congress, Forli, 1996 (eds di Lernia, S. & Manzi, G.), 43-51 (A.B.A. C.O Edizioni, Rome.
- HAYES, A., ROHLING, E.J., DE RIJK, S. & ZACHARIA-SSE, W.J., 1999. Mediterranean planktonic foraminiferal faunas during the last glacial cycle.

Marine Geology 153, 239-252.

- HILL, A.E. & MITCHELSON-JACOB, E.G., 1993. Observations of a poleward-flowing saline core on the continental slope west of Scotland, *Deep-Sea Research* 40, 1521-1527.
- JORISSEN, F.J., ASIOLI, A., BORSETTI, A.M., CA-POTONDI, DE VISSER, J.P., HILGEN, F.J., ROH-LING, E.J., VAN DER BORG, K., VERGNAUD-GRAZ-ZINI, C. & ZACHARIASSE, W.J., 1993. Late Quaternary central Mediterranean biochronology. *Marine Micropaleontology* 21, 169-189.
- KALLEL, N., PATERNE, M., DUPLESSY, J.C., VERG-NAUD-GRAZZINI, C., PUJOL, C., LABEYRIE, L.D., ARNOLD, M., FONTUGNE, M. & PIERRE, C., 1997. Enhanced rainfall in the Mediterranean region during the last sapropel event, *Oceanologica Acta* 20, 697-712.
- KUNIHOLM, P.I., KROMER, B., MANNING, S.W., NEW-TON, M., LATINI, C.E. & BRUCE, M.J., 1996. Anatolian tree rings and the absolute chronology of the eastern Mediterranean, 2220-718 BP, *Nature* 381, 780-783.
- LAMB, H.H., 1972. *Climate; past, present and future* 1, 613 pp. (Methuen & Co, London).
- LEAMAN, K.D. & SCHOTT, F.A., 1991. Hydrographic structure of the convection regime in the Gulf of Lions: Winter 1987, *Journal of Physical Oceanography* 21, 575-598.
- MARIOLOPOULOS, E.G., 1961. An outline of the climate of Greece, *Publications of the Meteorological Institute of the University of Athens* 6, 51pp.
- MAYEWSKI, P.A., MEEKER, L.D., TWICKLER, M.S., WHITLOW, S., YANG, Q., LYONS, W.B. & PRENTICE, M., 1997. Major features and forcing of highlatitude northern hemisphere atmospheric circulation using a 110,000-year-long glaciochemical series, *Journal of Geophysical Research* 102, 26345-26366.
- MCKIM MALVILLE, J., WENDORF, F., MAZAR, A.A. & SCHILD, R., 1998. Megaliths and Neolithic astronomy in southern Egypt, *Nature* 392, 488-491.
- MERCONE, D., THOMSON, J., CROUDACE, I.W., SI-ANI, G., PATERNE, M. & TROELSTRA, S., 2000. Duration of S1, the most recent sapropel in the Mediterranean Sea, as indicated by accelerator mass spectroscopy radiocarbon and geochemical evidence. *Paleoceanography* 15, 336-347.
- ORTIZ, J.D. & MIX, A.C., 1997. Comparison of Imbrie-Kipp transfer function and modern analog temperature estimates using sediment trap and core top foraminiferal faunas, *Paleoceanogra*-

phy 12, 175-190.

- O'BRIEN, S.R., MAYEWSKI, P.A., MEEKER, L.D., MEESE, D.A., TWICKLER, M.S. & WHITLOW, S.I., 1995. Complexity of Holocene climate as reconstructed from a Greenland ice core. *Science* 270, 1962-1964.
- PFLAUMANN, U., DUPRAT, J., PUJOL, C. & LABEY-RIE, L.D., 1996. SIMMAX: a modern analog technique to deduce Atlantic sea surface temperatures from planktonic foraminifera in deepsea sediments, *Paleoceanography* 11, 15-35.
- POULOS, S.E., DRAKOPOULOS, P.G. & COLLINS, M.B., 1997. Seasonal variability in sea surface oceanographic conditions in the Aegean Sea (eastern Mediterranean): an overview, *Journal* of Marine Systems 13, 225-244.
- REID, J.L., 1979. On the contribution of the Mediterranean outflow to the Norwegian-Greenland Sea, Deep-Sea Research 26, 1199-1223.
- ROETHER, W. H., MANCA, B. B., KLEIN, B., BRE-GANT, D., GEORGOPOULOS, D., BEITZEL, V., KO-VACEVIC, V. & LUCHETTA, A., 1996. Recent changes in eastern Mediterranean deep waters. *Science* 271, 333-335.
- ROHLING, E.J. & BRYDEN, H.L., 1992. Maninduced salinity and temperature increases in Western Mediterranean Deep Water. *Journal* of *Geophysical Research* 97, 11191-11198.
- ROHLING, E.J., JORISSEN, F.J., VERGNAUD-GRAZZI-NI, C. & ZACHARIASSE, W.J., 1993. Northern Levantine and Adriatic Quaternary planktic foraminifera; Reconstruction of paleoenvironmental gradients. *Marine Micropaleontology* 21, 191-218.
- ROHLING, E.J., JORISSEN, F.J. & DE STIGTER, H.C., 1997. 200 Year interruption of Holocene sapropel formation in the Adriatic Sea. *Journal of Micropalaeontology* 16, 97-108.
- ROHLING, E.J., HAYES, A., KROON, D., DE RIJK, S. & ZACHARIASSE, W.J., 1998. Abrupt cold spells

in the NW Mediterranean. *Paleoceanography* 13, 316-322.

- ROHLING, E.J. & DE RIJK, S., 1999. Holocene climate optimum and last glacial maximum in the Mediterranean: the marine oxygen isotope record. *Marine Geology* 153, 57-75.
- ROSSIGNOL-STRICK, M., 1995. Sea-land correlation of pollen records in the eastern Mediterranean for the glacial-interglacial transition: biostratigraphy versus radiometric timescale, Quaternary Science Reviews 14, 893-915.
- ROSSIGNOL-STRICK, M.,1998. Paléoclimat de la Méditerranée orientale et de l'Asie du Sud-Ouest de 15 000 a 6 000 BP, Paléorient 23, 175-186.
- STUIVER, M. & REIMER, P.J., 1993. Extended 14C data base and revised CALIB 3.0 14C age calibration program, *Radiocarbon* 35, 215-230.
- TARGARONA, J., 1997. Climatic and Oceanographic evolution of the Mediterranean region over the last glacial-interglacial transition; a palynological approach, *LPP Contributions* 7, 155 pp.
- THUNELL, R.C., WILLIAMS, D.F. & KENNETT, J.P., 1977. Late Quaternary paleoclimatology, stratigraphy, and sapropel history in eastern mediterranean deep-sea sediments. *Marine Micropaleontology* 2, 371-388.
- ZACHARIASSE, W.J., JORISSEN, F.J., PERISSORATIS, C., ROHLING, E.J. & TSAPRALIS, V., 1997. Late Quaternary foraminiferal changes and the nature of sapropel S1 in Skopelos Basin. Proceedings of the 5th Hellenic symposium on Oceanography and Fisheries, Kavalla, Greece, 15-18 April 1997 1, 391-394.