

Mediterranean Marine Science

Vol 14, No 2 (2013)

Vol 14, No 2 (2013)



Records of *Polydora cornuta* and *Streblospio gynobranchiata* (Annelida, Spionidae) from the Black Sea

V.I. RADASHEVSKY, Z.P. SELIFONOVA

doi: [10.12681/mms.415](https://doi.org/10.12681/mms.415)

To cite this article:

RADASHEVSKY, V., & SELIFONOVA, Z. (2013). Records of *Polydora cornuta* and *Streblospio gynobranchiata* (Annelida, Spionidae) from the Black Sea. *Mediterranean Marine Science*, 14(2), 261–269.
<https://doi.org/10.12681/mms.415>

Records of *Polydora cornuta* and *Streblospio gynobranchiata* (Annelida, Spionidae) from the Black Sea

V. I. RADASHEVSKY^{1,2} and Z. P. SELIFONOVA³

¹ A.V. Zhirmunsky Institute of Marine Biology, Far Eastern Branch of the Russian Academy of Sciences, Vladivostok 690059, Russia

² Far Eastern Federal University, Vladivostok 690091, Russia

³ Admiral Ushakov Maritime State University, Novorossiysk 353918, Russia

Corresponding author: radashevsky@gmail.com

Handling Editor: Melih Ertan Çinar

Received: 18 July 2012; Accepted: 1 April 2013; Published on line: 26 April 2013

Abstract

Two spionid polychaetes, *Polydora cornuta* and *Streblospio gynobranchiata*, were identified in benthic samples collected in the northern Black Sea and adjacent waters. These species have earlier been classified as the worst invaders in soft bottom communities in the Mediterranean Sea. *Polydora cornuta* had been previously misidentified and widely reported from the Black Sea and the Sea of Azov as *P. ciliata*, *P. ciliata limicola* and *P. limicola*. *Streblospio gynobranchiata* is a new invader currently extending its distribution into the Mediterranean, Black and Caspian Seas. Morphology, diagnostic characters and biology of the species are discussed and the history of their records in the Mediterranean, Black and Caspian Seas and the Sea of Azov is reviewed.

Keywords: *Polydora cornuta*, *Streblospio gynobranchiata*, biological invasions, Russia, Ukraine.

Introduction

In the last few decades marine biological invasions caused by human activities and global climate changes have become grave matters of concern due to their ecological and economic ramifications, causing intensified research efforts in this field (Occhipinti-Ambrogi & Savini, 2003; Carlton, 2009; Katsanevakis *et al.*, 2013). A total of 986 alien or non-indigenous species have been recognized in the Mediterranean basin, of which 560 (57%) species have been reported as established (Zenetos *et al.*, 2012). In the Black Sea, about 160 alien species have been identified since the first scientifically documented invasions of the barnacles *Amphibalanus improvisus* (Darwin, 1854) and *A. eburneus* (Gould, 1841) in Sevastopol Bay in 1844 and 1892, respectively (Gomoiu & Skolka, 1996; Zaitsev & Öztürk, 2001; Leppäkoski *et al.*, 2009). A six fold increase in the number of recognized alien species in the Black Sea in the last decade has resulted from not only an increased number of studies but also from the rapid growth of human activities in the region (Alexandrov, 2004).

Lists of marine alien species always include polychaetous annelids, among which spionid polychaetes (Annelida: Spionidae) are common invaders (Çinar *et al.*, 2005a, 2011; Şahin & Çinar, 2012). For example, intensive studies of polychaetes along the Turkish coast

of the Aegean Sea revealed a total of 35 spionid species of which nine were recognized as alien (Dağlı *et al.*, 2007, 2011). The massive development of *Polydora cornuta* Bosc, 1802 in Odessa Harbour in the early 1960s (Losovskaya & Nesterova, 1964, as *P. ciliata limicola* Annenkova, 1934) was probably one of the most conspicuous marine invasions in the Black Sea. Since then, *P. cornuta* has been reported from harbours in various countries surrounding the Black Sea and is now considered to be one of the worst invasive species in soft bottom communities in the Mediterranean Sea (Zenetos *et al.*, 2005). Another spionid on the list of worst invaders, *Streblospio gynobranchiata* Rice & Levin, 1998, was recently introduced and is currently spreading in the Mediterranean, Black and Caspian Seas.

Two important Russian international seaports, Novorossiysk and Tuapse, are located in the northeast Black Sea (Fig. 1). Their annual traffic capacity is about 5000 and 1000 ships per year, respectively, and their carrying capacity is about 100 and 20 million tons of cargo, respectively. The whole region is therefore subjected to high levels of pollution, ballast water discharge (more than 50 million tons per year in Novorossiysk Bay) and species movement via ship fouling, all factors that favour the introduction and establishment of non-indigenous species. As a consequence, 23 alien and 44 cryptogenic species have been reported from the port areas (Selifon-



Fig. 1: Maps showing previous (A) and present (B) records of spionid polychaetes in the Mediterranean, Black and Caspian Seas and the Seas of Azov and Marmara: *Polydora cornuta* (yellow circles): 1 – Tena *et al.* (1991); 2 – Radashevsky (2005), Surugiu (2005, 2006, 2012), Surugiu & Novac (2007); 3 – Çinar *et al.* (2005b, 2012a); 4 – Dağlı & Ergen (2008), Karhan *et al.* (2008), Çinar *et al.* (2009); 5 – Boltacheva & Lisitskaya (2007); 6 – Simboursa *et al.* (2008); 7 – Bondarenko (2011); 8 – Losovskaya (2011); 9 – Selifonova (2011); 10 – Çinar *et al.* (2012b). *Streblospio gynobranchiata* (red squares): 11 – Çinar *et al.* (2005 a,b, 2012a), Dağlı *et al.* (2011); 12 – Boltacheva (2008); 13 – Taheri *et al.* (2008), Taheri & Foshtomi (2011); 14 – Çinar *et al.* (2009).

ova, 2012). Sevastopol and Odessa are among the largest seaports in Ukraine in the north and north-western parts of the Black Sea, respectively. Recent sampling in Novorossiysk Bay and the port of Tuapse, and samples previously collected in Sevastopol, Odessa and adjacent areas, provided good material for taxonomic revisions and better insight into the regional distribution of invasive species of Spionidae.

The purpose of this study is to report new records of spionid polychaetes in the northern Black Sea and adjacent waters, re-examine old material and present an overview of the expansion of invasive spionids in the Mediterranean, Black and Caspian seas and the Sea of Azov.

Materials and Methods

Field collections were made in Snake Lake Liman (Big Utrish) (Zmeinoe Ozero Liman, Bolshoi Utrish), Novorossiysk Harbour (Novorossiysk Bay), Tuapse Harbour and Sochi, all in the north-eastern part of the Black Sea, Russia, by Zhanna P. Selifonova in September 2010, August 2011, November 2011, and July 2012. Sediment

samples were collected using a Petersen grab (0.04 m²) and screened in the field with a 500 µm mesh sieve. The residue was fixed in 10% formaldehyde solution, rinsed in fresh water, and then transferred to 70% ethanol. Polychaetes were removed from the residue under a stereomicroscope at the Admiral Ushakov Maritime State University, Novorossiysk. Additional material from the Sea of Azov and the Black Sea (Sevastopol Bay, Odessa and adjacent regions) was provided by Marta I. Kiseleva, Ludmila V. Vorobyova, Ivan V. Sinigub, Elena V. Lisitskaya and Natalia A. Boltacheva. Photographs were taken using a Zeiss Axio Imager 7.2 microscope equipped with a digital camera. Specimens were stained with a solution of methylene green in alcohol prior to photography.

Supplementary material on *P. cornuta* from the Aegean Sea, Greece, was provided by Nomiki Simboursa. All the material was deposited at the Museum of the Institute of Marine Biology (MIMB), Vladivostok, Russia, and the Natural History Museum of Los Angeles County (LACM-AHF), Los Angeles, U.S.A. Information about the samples is given below along with descriptions of the specimens. The number of specimens in each sample is

given in parentheses after the museum abbreviation and registration number.

Taxonomic Account

Spionidae Grube, 1850

Polydora cornuta Bosc, 1802

Polydora cornuta Bosc, 1802: 150–153, pl. 12, figs 7–8. Blake & Maciolek, 1987: 12–14, fig. 1. Tena *et al.*, 1991: 32–35, fig. 3. Çinar *et al.*, 2005b: 824–826, figs 3, 4; 2006 (ecol.): 231–241; 2008 (ecol.): 5–15; 2009 (ecol.): 13–16; 2012a (ecol.): 1462; 2012b (ecol.): 960. Radashevsky, 2005: 3–19, figs 1–4 (adult and larval morphology). Surugiu, 2005a: 66–67; 2012 (Part.): 47–50, fig. 2. Boltacheva & Lisitskaya, 2007: 33–35, fig. 1. Dağlı & Ergen, 2008: 231–233, fig. 3. Simboursa *et al.*, 2008: 123–124, fig. 2. Bondarenko, 2011 (ecol.): 12–16. Losovskaya, 2011 (ecol.): 50–54. Selifonova, 2011: 48, fig. 1.

Polydora ligni Webster, 1879: 119. *Fide* Blake & Maciolek, 1987: 12, 14.

Polydora ciliata: Kiseleva, 2004 (Part.): 264–266, fig. 107. Not Johnston, 1838.

Polydora ciliata limicola: Losovskaya & Nesterova, 1964 (ecol.): 1559–1560. Losovskaya, 1969 (ecol.): 58. Moroz, 1977 (ecol.): 1104–1105. Kiseleva, 1987 (ecol.): 42; 1988 (ecol.): 27. Sinegub, 2001 (ecol.): 301–315; 2002a (ecol.): 341–342; 2002b (ecol.): 31–32. Vorobyova *et al.*, 2008 (ecol.): 46–53. Not Annenkova, 1934.

Polydora limicola: Losovskaya, 1976 (ecol.): 102–103; 1977 (ecol.): 44–45; 2008 (ecol.): 358–360. Moroz, 1985 (ecol.): 104–107; 1992 (ecol.): 150–151. Losovskaya & Sinegub, 2002 (ecol.): 5–9. Losovskaya & Zolotarev, 2003 (ecol.): 248–249. Losovskaya *et al.*, 2004 (ecol.): 51–57; 2005 (ecol.): 618–621. Bondarenko, 2009 (ecol.): 22–26. Vorobyova & Bondarenko, 2009 (ecol.): 113–119. Not Annenkova, 1934.

Material

SEA OF AZOV, R/V *Gidrolog*, coll. V.P. Zakutsky: sta. 40, 46°12'N, 37°18'E, 25 Jun 1983, MIMB 27148 (3); sta. 54, 46°12'N, 37°48'E, 26 Jun 1983, MIMB 27149 (11); sta. 55, 45°43'N, 36°43'E, 26 Jun 1983, MIMB 27150 (1); sta. 69, 45°33'N, 35°48'E, 29 Jun 1983, MIMB 27151 (3); sta. 74, 45°43'N, 35°12'E, 29 Jun 1983, MIMB 27152 (6).

RUSSIA, north-eastern Black Sea, **Bolshoi Utrish**, Snake Lake Liman, 44°45.92'N, 37°23.24'E, 3 m, mud, coll. Zh.P. Selifonova, 28 Aug 2011, MIMB 27127 (1).

UKRAINE, northern Black Sea, Crimea: **Karkinitzky Gulf**, Swan Islands, 45°51.4'N, 33°29.8'E, intertidal, fouling on a concrete pipe, coll. V.A. Grintsov, 12 Aug 2005, MIMB 14114 (6); 0.3 m, muddy sand, coll. E.V. Lisitskaya, 14 Aug 2007, MIMB 27202 (3). **Sevastopol Bay**: Martyn Bight, 44°37'N, 33°30.5'E,

coll. V.I. Radashevsky, 2 Dec 2005, MIMB 14115 (1); 44°37.09'N, 33°31.41'E, reared in laboratory from larvae collected from plankton in Sebastopol Bay, M.I. Kiseleva, 10 Nov 1988, MIMB 27153 (20). Reared in laboratory from larva collected from plankton in **Karantinnaya Bay**, 44°37.02'N, 33°30.25'E, 10 Aug 2006, E.V. Lisitskaya, MIMB 17927 (1). Reared in laboratory from larvae collected from plankton in **Balaklava Bay**, 44°30.12'N, 33°35.86'E, on 2 May 2006, fixed on 21 Aug 2006, E.V. Lisitskaya, MIMB 17926 (10); collected from plankton on 21 Aug 2006, fixed on 13 Oct 2006, E.V. Lisitskaya, MIMB 17925 (7).

UKRAINE, north-western Black Sea, **Odessa area**: off Odessa, 46°30'N, 30°50'E, 7–26 m, muddy sand, coll. I.A. Sinegub, 23 Apr 2005, MIMB 10517 (85); Sukhoi Liman, 46°19.9'N, 30°39.7'E, 0.3 m, sand, shell fragments, coll. I.A. Sinegub, 26 Jun 2001, MIMB 14119 (40); 0.3–13 m, mud, shell fragments, coll. I.A. Sinegub, 04 Sep 2002, MIMB 14120 (25). **Dnieper-Danube area**: 45°52'N, 30°20'E, 14 m, muddy sand and shell fragments, coll. I.A. Sinegub, Aug 1992, MIMB 14116 (300+); 45°19'N, 29°49'E, 23 m, mud, coll. I.A. Sinegub, 25 Nov 2005, MIMB 14118 (4).

Supplementary material

GREECE, Aegean Sea, **Saronikos Gulf**, near Psytalia Is., 37°55.42'N, 23°35.45'E, 65 m, sandy mud, coll. N. Simboursa, 9 Feb 2012, MIMB 27201 (2).

TURKEY, **Bosphorus Strait**, Kuzguncuk, 41°2.25'N, 29°1.667'E, 0.5–1 m, *Mytilus galloprovincialis* community on hard bottom, 21 May 2004, coll. E. Kalkan, MIMB 17924 (6). Sea of Marmara, **Izmit Bay**, sts 1, 2, 4–7 (see coordinates in Dağlı & Ergen, 2008: Annex 1), 3–10 m, mud, coll. E. Dağlı, 18 Aug 2006, MIMB 27142–27147 (18).

Adult morphology

Up to 13 mm long and 1.1 mm wide for 60 chaetigers. Prostomium anteriorly incised to bifid, posteriorly extending to end of chaetiger 3 as a low caruncle. Two pairs of small black eyes and a prominent occipital antenna present on prostomium. Palps with a fine continuous black line along each side of frontal groove. Chaetiger 1 with short capillaries in neuropodia and small postchaetal lamellae in both rami; notochaetae absent. Chaetiger 5 with heavy falcate spines and companion chaetae arranged in a slightly curved oblique row; falcate spines each with a lateral tooth and a narrow thin subdistal longitudinal flange; companion chaetae each with a broom-like distal tip closely adhering to convex side of falcate spine; dorsal superior and ventral capillaries absent. Bidentate hooded hooks in neuropodia from chaetiger 7, not accompanied by capillaries. Branchiae

from chaetiger 7 through most of body. Pygidium disc-like with dorsal gap.

Remarks

Polydora cornuta was originally described from Charleston Harbor, South Carolina, Atlantic USA by Bosc (1802) and later reported from estuaries and sea ports (mainly located in estuaries) all over the world (see Radashevsky, 2005). The first report of the species from the Black Sea was a misidentification. Pereyaslawzewa (1891) identified worms boring into gastropod shells occupied by hermit crabs off Sevastopol, Crimea, as *P. cornuta*, despite the fact that genuine *P. cornuta* is a tube-dwelling species unable to bore into calcareous substrata. The species was misidentified again when Losovskaya & Nesterova (1964) and Losovskaya (1969) reported large numbers of *Polydora ciliata limicola* Annenkova, 1934 in Sukhoi Liman Lagoon in the Odessa area. Samples collected at that time no longer exist but discussion and joint examination of recent material by Galina V. Losovskaya and Vasily I. Radashevsky during VIR's visit to Odessa in December 2005 helped to identify that material unambiguously as *P. cornuta* (see also Losovskaya, 2008: 360, 2011: 53). Adult worms were first recorded from the Odessa area in 1962 and, in 1963 population density reached 1155 ind.m⁻². In 2001, the population density of the species in Odessa Harbour reached 6600 ind.m⁻² in the benthic community and 9600 ind.m⁻² in the pier fouling communities (Losovskaya *et al.*, 2004). Remarkably, no tube-dwelling (non-boring) *Polydora* worms were reported from the Black Sea before the 1960s, despite several studies on polychaetes in this region. It is likely therefore that Losovskaya & Nesterova (1964) reported the first arrival of *P. cornuta* in the Black Sea, at least in its northern part, which might actually have happened in the late 1950s.

As *P. ciliata*, *P. ciliata limicola*, *P. limicola*, and ultimately as *P. cornuta*, the species was reported from the Odessa area, Dnieper-Danube and adjacent regions in the north-western Black Sea (Losovskaya, 1976, 1977, 2008, 2011; Moroz, 1977, 1985, 1992; Sinegub, 2001, 2002a,b; Losovskaya & Sinegub, 2002; Losovskaya & Zolotarev, 2003; Losovskaya *et al.*, 2004, 2005; Vorobyova *et al.*, 2008; Bondarenko, 2009, 2011; Vorobyova & Bondarenko, 2009), Sevastopol in Crimea (Moroz, 1985; Kiseleva, 1988, 2004; Boltacheva & Lisitskaya, 2007), the Romanian coast (Surugiu & Manoleli, 1999; Surugiu, 2005a,b, 2006, 2012; Surugiu & Novac, 2007), and Tuapse Harbour, Russia (Selifonova, 2011). Kiseleva (1987) reported *P. ciliata limicola* for the first time from the Sea of Azov (Fig. 1A). Descriptions provided by these authors and our examination of the material reported in most of these ecological surveys (see *Material* section above) undoubtedly confirmed the presence of *P. cornuta*, as re-

described by Blake & Maciolek (1987) and Radashevsky (2005). It is possible that earlier reports of *P. limicola* from the Romanian coast were also misidentifications of *P. cornuta* but the lack of the original material precludes verification (see Surugiu, 2005a).

Re-examination of old material from Izmir Bay on the Turkish coast of the Aegean Sea by Çinar *et al.* (2005b) revealed that *P. cornuta* was present in the region in 1986; sampling in this area in 2003 found a dense population of the species, up to 3170 ind.m⁻² (Çinar *et al.*, 2005b). Populations of *P. cornuta* up to 170 ind.m⁻² were reported from Izmit Bay (Dağlı & Ergen, 2008) and 3390 ind.m⁻² from Kalamiş Bay (Karhan *et al.*, 2008), both in the Sea of Marmara; a dense population of the species up to 4340 ind.m⁻² was reported from the Golden Horn Estuary in the Bosphorus Strait (Çinar *et al.*, 2009). In the Dniester estuary, the population density of the species reached 54904 ind.m⁻² (Vorobyova & Bondarenko, 2009), and in severely polluted areas on the Romanian coast it reached 95000 ind.m⁻² while the biomass attained 157.4 g.m⁻² (Surugiu, 2005b). In the Mediterranean Sea, *P. cornuta* was reported from Valencia Harbour in Spain (Tena *et al.*, 1991), and Elefsis Bay on the Greek coast of the Aegean Sea (Simboura *et al.*, 2008) (Fig. 1A).

Polydora cornuta is an opportunistic species tolerant of wide ranges of salinity and temperature, and characterized by early maturation, iterative breeding in water temperatures above 13°C, high larval production, ability to colonize disturbed and polluted substrata and to establish high density populations in a short time (Dauer *et al.*, 1981; Zajac, 1991a,b; Radashevsky, 2005). It is being widely distributed by shipping and aquaculture. Planktotrophic larvae of the species survive transportation in ballast water, and adults may form dense settlements through fouling of ship hulls (Radashevsky, in preparation). The routes of the species in the Mediterranean, Black Sea and the Sea of Azov remain uncertain. The limited number of recent reports from the Mediterranean Sea (in contrast to many reports from the Black Sea) may be due to the limited number of taxonomic studies rather than a limited distribution of the species in the region. It is possible that *P. cornuta* was either first introduced by ocean-going vessels to the north-western part of the Black Sea and then transported to the eastern Mediterranean or that there were several independent introductions into the Mediterranean Sea and Black Sea. The native distribution of the species is unknown and may no longer be possible to determine because of numerous transportations by human activities.

Distribution

Widespread, mainly in estuaries and sea ports.




Fig. 2: *Streblospio gynobranchiata* female morphology (formaldehyde-fixed specimens stained with a solution of methylene green in alcohol). A – anterior end, left lateral view. B – same, dorsal view. C – anterior end, left lateral view; palps missing. D – fertile chaetigers 11–13 without extensions of lateral body wall, left lateral view. E – fertile chaetigers 19–21 with extensions of lateral body wall, dorsal view. F – extension of lateral body wall with developed oocyte inside. G – fertile chaetigers 19–20 with extensions of lateral body wall, left lateral view. *br* – branchia; *ex* – extension of lateral body wall; *la* – lateral eye; *me* – median eye; *oo* – oocyte; *pa* – palp; *pe* – peristomium; *po* – dorsal transverse pouch on chaetiger 2; *pr* – prostomium; *sa* – sabre chaeta; *ve* – germinal vesicle. Scale bars: A – 200 μ m; B–E – 100 μ m; F, G – 50 μ m. A, B – Novorossiysk Bay, MIMB 27182; C, D – Snake Lake Liman, MIMB 27183; E–G – Snake Lake Liman, MIMB 27184.

Streblospio gynobranchiata Rice & Levin, 1998

Figs 2, 3

Streblospio gynobranchiata Rice & Levin, 1998: 694–707, figs. 1–13. Çinar *et al.*, 2005a (ecol.): 120–135; 2005b: 822–824, fig. 2; 2006 (ecol.): 231–241; 2008 (ecol.): 5–15; 2009 (ecol.): 13–16; 2012a (ecol.): 1462. Boltacheva, 2008: 12, fig. 1. Taheri *et al.*, 2008 (ecol.): 2–4. Dağlı *et al.*, 2011 (ecol.): 51–62. Taheri & Foshtomi, 2011 (ecol.): 609–612.

Streblospio sp.: Murina *et al.*, 2008: 46, fig. 1. Selifonova, 2009: 245, fig. 3.

Material

RUSSIA, north-eastern Black Sea: **Novorossiysk** Harbour, Novorossiysk Bay, 44°43.55'N, 37°47.1'E,




Fig. 3: *Streblospio gynobranchiata* female morphology (formaldehyde-fixed specimens stained with a solution of methylene green in alcohol). A, B – chaetigers 11–14, ventral view, showing median unpaired seminal receptacles; intersegmental lines marked by double-arrows. C – middle fertile chaetigers with extensions of lateral body wall, left lateral view. D – posterior end, left lateral view, showing anterior fertile chaetigers with extensions of lateral body wall and posterior fertile chaetigers without extensions of lateral body wall. *ch11–ch14* – chaetigers 11–14; *ex* – extension of lateral body wall; *oo* – oocyte; *re* – seminal receptacle. Scale bars: A–C – 100 μ m; D – 200 μ m. A, B – Snake Lake Liman, MIMB 27184; C – Novorossiysk Bay, MIMB 27182; D – Snake Lake Liman, MIMB 27183.

10 m, mud, 1 Aug 2007, MIMB 27180 (3); 31 Jul 2008, MIMB 27181 (70); 3 Sep 2010, MIMB 27182 (75). **Bolshoi Utrish**, Snake Lake Liman, 44°45.92'N, 37°23.24'E, 3 m, mud, 28 Aug 2011, MIMB 27183 (7); 15 Nov 2011, MIMB 27184 (18).

UKRAINE, north-western Black Sea, **Odessa area**, Sukhoi Liman, 46°19.9'N, 30°39.7'E, 0.3–13 m, mud, shell fragments, coll. I.A. Sinegub, 04 Sep 2002, MIMB 14124 (10).

Adult morphology

Up to 12 mm long and 0.5 mm wide for 55 chaetigers. Prostomium anteriorly rounded, posteriorly extending to chaetiger 1. Peristomium forming a prominent hood enveloping prostomium laterally and ventrally (Fig. 2A–C). Two pairs of small red eyes arranged trapezoidally. Occipital antenna absent on prostomium. Palps as long as 5–10 chaetigers, with up to 20 small transverse bands/spots of dark pigment. Short median papilla present on dorsal side of chaetiger 1. Chaetiger 1 with short capillaries in both rami and small postchaetal lamellae in neuropodia; notopodial lamellae absent. Notopodial postchaetal lamellae of chaetiger 2 interconnected by a prominent transverse membrane or collar, forming a characteristic pouch open anteriorly; median part of collar straight to slightly concave (Fig. 2C). Sabre chaetae

in neuropodia from chaetiger 7. Hooks in neuropodia from chaetiger 7, up to 9 in a series, accompanied by 1–3 alternating capillaries and 1–2 inferior sabre chaetae. Hooks multidentate, with 3–5 pairs of small upper teeth arranged in two vertical rows above main fang. Single pair of branchiae on chaetiger 1 situated just posterior to pair of palps; branchiae apinnate, flattened, with undulating lateral edges, almost equal in length to palps and appearing similar to them (Fig. 2A–C). Finger-like extensions of lateral body wall up to 350 μm long and 50 μm in diameter in mature females from chaetigers 16–19 to chaetigers 22–33; extensions oriented dorsally and posteriorly and containing mature oocytes (Figs 2E–G, 3C, D). Pygidium small, rounded, with one pair of very short ventral knobs (Fig. 3D). In males, sperm from chaetiger 7 to chaetigers 20–36. In females, oocytes from chaetiger 8 to chaetigers 25–33. Oocytes up to 85 μm in diameter, with germinal vesicle 30 μm , and single nucleolus 10 μm in diameter; up to about 10 large vitellogenic oocytes present in each ovary and 20 oocytes in each fertile segment (Figs 2D–F, 3C, D). Unpaired intraepithelial seminal receptacles present in females on ventral side of 2–3 chaetigers, from chaetigers 13–14 to chaetigers 14–15; receptacles oval to fusiform, up to 110 μm long and 30 μm wide, oriented longitudinally between ventral nerve cords (Fig. 3A, B).

Habitat

In Novorossiysk Harbour, adults of *S. gynobranchiata* usually occur in the estuary of River Tsemess, with water salinity at 12.9 ‰. They inhabit the upper layer of muddy sediments containing hydrogen sulphide. The population density of the species reached 980 ind.m⁻² in 2001, 9000 ind.m⁻² in 2007 and 10000 ind.m⁻² in 2008.

Remarks

Streblospio gynobranchiata was originally described from Tampa Bay, Florida, USA by Rice & Levin (1998) and later reported from the Aegean Sea, Turkey (Çinar *et al.*, 2005 a,b, 2008, 2012a; Dağlı *et al.*, 2011), Crimea in the northern Black Sea (Boltacheva, 2008), the southern part of the Caspian Sea (Taheri *et al.*, 2008; Taheri & Foshtomi, 2011), and the Golden Horn Estuary, Bosphorus Strait (Çinar *et al.*, 2009) (Fig. 1A). In all localities, worms occurred on muddy substrata near to and within the port areas, reaching maximum density of 60480 ind.m⁻² in Alsancak Harbour (Izmir Bay of the Aegean Sea) (Çinar *et al.*, 2009). Because the species was not found in earlier biological surveys conducted in these areas, it is classified as an alien, most likely introduced with ballast waters.

Worms collected within the framework of this study fit all the features of *S. gynobranchiata* (see Rice & Levin,

1998) and are characterized by the presence of a single pair of branchiae situated close to and appearing similar to palps, a dorsal pouch on chaetiger 2, lateral extensions of the body wall, and hooks in neuropodia consistently beginning from chaetiger 7. Remarkably, the body wall extensions are present only in mature females but absent in immature individuals and males.

Of the three currently recognized *Streblospio* species (see Mahon *et al.*, 2009), *S. benedicti* Webster, 1879 and *S. gynobranchiata* have been considered as native to the Americas, while *S. shrubsolei* (Buchanan, 1890) seems to be native to European waters. Adults of *S. gynobranchiata* are similar to *S. benedicti* in having neuropodial hooks starting from chaetiger 7, and both differ from *S. shrubsolei*, which have hooks starting from chaetigers 8–10. Males of *S. gynobranchiata* and *S. benedicti* appear quite similar to each other, but females differ in having lateral body wall extensions in *S. gynobranchiata* and dorsal brooding pouches in *S. benedicti*. As one of the most common polychaete species on both coasts of North America, and also due to poecilogonous ability to produce planktotrophic and lecithotrophic larvae, *S. benedicti* became a subject of numerous studies on ecology and reproductive biology (e.g. Levin & Huggett, 1991; McCain, 2008; Gibson *et al.*, 2010). The other two species have been less thoroughly studied and morphological differences between them have remained uncertain.

Distribution

Atlantic coast of North and South America from Florida, U.S.A. to Paraná, Brazil; Mediterranean, Black and Caspian Seas.

Conclusion

Alterations (addition - invasion, deletion - extinction, and altered population dynamics) to the biodiversity of terrestrial, freshwater and marine communities are the expected and inexorable consequences of human activity currently seen throughout the world (Carlton, 2009). However, recognizing these alterations is not an easy task except where large, well-known species are of concern. Although alien, non-indigenous species have become subjects of study in many countries in recent years, these studies are still mainly based on morphological examination and comparison of fixed specimens. When the biogeographic origin of a species and its morphological variability are unknown, identification of specimens from remote places often remains ambiguous and uncertain.

Spionid polychaetes demonstrate an amazing diversity of morphological, ecological and reproductive characteristics and show significant variability (both ontogenetic and individual) in important taxonomic characters (Radashevsky, 2012). The geographic morphological

variability of spionids is poorly studied but should be considered when making identifications. Molecular techniques, currently under rapid development, are certainly needed not only for the identification of species but also for recognition of their biogeographic origins and possible routes of distribution. Molecular comparisons of spionids from distant places will provide closer insight into their taxonomy and distribution.

Acknowledgements

Marta I. Kiseleva, Lyudmyla V. Vorobyova, Ivan V. Sinegub, Elena V. Lisitskaya, Natalia A. Boltacheva, Ertan Dağlı and Nomiki Simbura provided important material for this study. Galina V. Losovskaya shared with VIR her notes, remarks and memories about *Polydora* worms examined in the early 1960's; this helped to identify the worms in question as *P. cornuta*. Mark A. Popov organized buddy diving with VIR in Sevastopol Bay. Tim Worsfold and Melih E. Çinar provided valuable comments and edits to the manuscript. To all these persons we express our sincere gratitude. Financial support was provided by the Russian Foundation for Basic Research (Project 12-04-00609) through the A.V. Zhirmunsky Institute of Marine Biology; the Government of the Russian Federation (Project 11.G34.31.0010) through the Far Eastern Federal University; and the research program "Estimation of Ecological Risks of Invasion of Alien Species in Seas of Russia with Water Transport" of the Admiral Ushakov Maritime State University for 2009–2011.

References

- Alexandrov, B.G., 2004. Problem of aquatic organisms transportation by ships and some approaches for risk assessment of the new invasions. *Marine Ecological Journal*, 3 (1), 5-17. (In Russian).
- Annenkova, N., 1934. Kurze Übersicht der Polychaeten der Litoralzone der Bering-Insel (Kommandor-Inseln), nebst Beschreibung neuer Arten. *Zoologischer Anzeiger*, 106 (12), 322-331.
- Blake, J.A., Maciolek, N.J., 1987. A redescription of *Polydora cornuta* Bosc (Polychaeta: Spionidae) and designation of a neotype. *Bulletin of the Biological Society of Washington*, 7 (1), 11-15.
- Boltacheva, N.A., 2008. Finding of new alien species *Streblospio gynobranchiata* Rice et Levin, 1998 (Polychaeta: Spionidae) in the Black Sea. *Marine Ecological Journal*, 7 (4), 12. (In Russian).
- Boltacheva, N.A., Lisitskaya, E.V., 2007. About species of *Polydora* (Polychaeta: Spionidae) from the Balaklava bay (the Black Sea). *Marine Ecological Journal*, 6 (3), 33-35. (In Russian).
- Bondarenko, O.S., 2009. Species composition and features of polychaete distribution in the western Black Sea. *Ecology of the sea*, 78, 22-27. (In Russian).
- Bondarenko, O.S., 2011. Polychaete taxocene in the seaside estuary of Danube River. *Marine Ecological Journal*, 2, 11-16. (In Russian with English summary).
- Bosc, L.A.G., 1802. Histoire naturelle des vers, contenant leur description et leurs mœurs, avec figures dessinées d'après nature. Tome 1. Deterville, Paris, 324 pp.
- Carlton, J.T., 2009. Deep invasion ecology and the assembly of communities in historical time. p. 13-56. In: *Biological Invasions in Marine Ecosystems. Ecological Studies* 204. Rilov, G. & Crooks, J.A. (Eds). Springer, Berlin Heidelberg.
- Çinar, M.E., Balkis, H., Albayrak, S., Dağlı, E., Karhan, S.Ü., 2009. Distribution of polychaete species (Annelida: Polychaeta) on the polluted soft substrate of the Golden Horn Estuary (Sea of Marmara), with special emphasis on alien species. *Cahiers de Biologie Marine*, 50 (1), 11-17.
- Çinar, M.E., Bilecenoglu, M., Öztürk, B., Katagan, T., Aysel, V., 2005a. Alien species on the coast of Turkey. *Mediterranean Marine Science*, 6 (2), 119-146.
- Çinar, M.E., Dağlı, E., Açık, S., 2011. Annelids (Polychaeta and Oligochaeta) from the Sea of Marmara, with descriptions of five new species. *Journal of Natural History*, 45 (33-34), 2105-2143.
- Çinar, M.E., Ergen, Z., Dağlı, E., Petersen, M.E., 2005b. Alien species of spionid polychaetes (*Streblospio gynobranchiata* and *Polydora cornuta*) in Izmir Bay, eastern Mediterranean. *Journal of the Marine Biological Association of the United Kingdom*, 85 (4), 821-827.
- Çinar, M.E., Katağan, T., Koçak, F., Öztürk, B., Ergen, Z. et al., 2008. Faunal assemblages of the mussel *Mytilus galloprovincialis* in and around Alsancak Harbour (Izmir Bay, eastern Mediterranean) with special emphasis on alien species. *Journal of Marine Systems*, 71 (1), 1-17.
- Çinar, M.E., Katagan, T., Öztürk, B., Dağlı, E., Açık, S. et al., (2012a). Spatio-temporal distributions of zoobenthos in soft substratum of Izmir Bay (Aegean Sea, eastern Mediterranean), with special emphasis on alien species and ecological quality status. *Journal of the Marine Biological Association of the United Kingdom*, 92 (7), 1457-1477.
- Çinar, M.E., Katagan, T., Öztürk, B., Dağlı, E., Açık, S. et al., (2012b). Spatio-temporal distributions of zoobenthos in Mersin Bay (Levantine Sea, eastern Mediterranean) and the importance of alien species in benthic communities. *Marine Biology Research*, 8, 954-968.
- Çinar, M.E., Katağan, T., Öztürk, B., Egemen, Ö., Ergen, Z. et al., 2006. Temporal changes of soft-bottom zoobenthic communities in and around Alsancak Harbor (Izmir Bay, Aegean Sea), with special attention to the autecology of exotic species. *Marine Ecology*, 27 (3), 229-246.
- Dağlı, E., Çinar, M.E., Ergen, Z., 2011. Spionidae (Annelida: Polychaeta) from the Aegean Sea (eastern Mediterranean). *Italian Journal of Zoology*, 78 (S1), 49-64.
- Dağlı, E., Ergen, Z., 2008. First record of *Polydora cornuta* Bosc, 1802 (Polychaeta: Spionidae) from the Sea of Marmara, Turkey basin. *Aquatic Invasions*, 3 (2), 225-227.
- Dağlı, E., Ergen, Z., Çinar, M.E., 2007. The significance of Spionidae (Annelida, Polychaeta) species living in the benthic environment the coasts of Turkey. *Türk Sucul Yasam Dergisi*, 3-5 (5-8), 152-158. (In Turkish).
- Dauer, D.M., Maybury, C.A., Ewing, R.M., 1981. Feeding behavior and general ecology of several spionid polychaetes from the Chesapeake Bay. *Journal of Experimental Marine Biology and Ecology*, 54 (1), 21-38.
- Gibson, G., MacDonald, K., Dufton, M., 2010. Morphogenesis and phenotypic divergence in two developmental morphs of *Streblospio benedicti* (Annelida, Spionidae). *Invertebrate Biology*, 129 (4), 328-343.

- Gomoiu, M.-T., Skolka, M., 1996. Changements recents dans la biodiversité de la Mer Noire dûs aux immigrants. *GEO-ECOMARINA, RCGGM, "Danube Delta - Black Sea system under Global Changes Impact"*, 1, 34-47.
- Johnston, G., 1838. Miscellanea Zoologica. III.—The British Ariadæ. *Magazine of Zoology and Botany, Edinburgh*, 2, 63-73.
- Karhan, S.Ü., Kalkan, E., Simboura, N., Mutlu, E., Bekbölet, M., 2008. On the occurrence and established populations of the alien polychaete *Polydora cornuta* Bosc, 1802 (Polychaeta: Spionidae) in the Sea of Marmara and the Bosphorus Strait (Turkey). *Mediterranean Marine Science*, 9 (1), 5-19.
- Katsanevakis, S., Zenetos, A., Belchior, C., Cardoso, A.C., 2013. Invading European Seas: Assessing pathways of introduction of marine aliens. *Ocean & Coastal Management*, 76, 64-74.
- Kiseleva, M.I., 1987. Changes of composition and distribution of polychaetous worms from the Sea of Azov. *Hydrobiological Journal*, 23 (2), 40-45. (In Russian).
- Kiseleva, M.I., 1988. A characteristic of benthos changes of many years in the littoral zone of the area of Sevastopol. *Ecology of the sea*, 28, 26-32. (In Russian).
- Kiseleva, M.I., 2004. *Polychaetes (Polychaeta) of the Azov and Black Seas*. Apatity, KSC RAS, 409 pp. (In Russian).
- Leppäkoski, E., Shiganova, T., Alexandrov, B., 2009. European enclosed and semi-enclosed seas. p. 529-547. In: *Biological Invasions in Marine Ecosystems*. Rilov, G. & Crooks, J.A. (Eds). *Ecological Studies*, 204. Springer-Verlag, Berlin.
- Levin, L.A., Huggett, D.V., 1991. Implications of larval development mode for polychaete population dynamics: lessons from a poecilogonous species, *Streblospio benedicti*. *Bulletin of Marine Science*, 48 (2), 590-591.
- Losovskaya, G.V., 1969. On the changes in the bottom fauna of the Sukhoi liman after connecting it with the sea. p. 56-59. In: *Biological problems of the oceanography of the southern seas*. Vinogradov, K.A. (Ed.). Naukova Dumka, Kiev. (In Russian).
- Losovskaya, G.V., 1976. On the expansion of the area of distribution of *Polydora limicola* Annenkova - a polychaete species new for the Black Sea. *Hydrobiological Journal*, 12 (1), 102-103. (In Russian).
- Losovskaya, G.V., 1977. *Ecology of Polychaeta of the Black Sea*. Naukova Dumka, Kiev, 92 pp. (In Russian).
- Losovskaya, G.V., 2008. Spatial distribution of the polychaetes *Neanthes succinea* and *Polydora limicola* in areas with different level of anthropogenic impact. *Ecological security of coastal and shelf zones and complex exploitation of shelf resources*, 17, 357-361. (In Russian).
- Losovskaya, G.V., 2011. On using some polychaete taxocene parameters for monitoring environmental quality in the north-western Black Sea. *Marine Ecological Journal*, 10 (1), 50-55. (In Russian with English summary).
- Losovskaya, G.V., Nesterova, D.A., 1964. On the mass development of a form of Polychaeta, *Polydora ciliata* ssp. *limicola* Annenkova, new for the Black Sea, in the Sukhoi Liman (north-western part of the Black Sea). *Zoological Journal*, 43 (10), 1559-1560. (In Russian).
- Losovskaya, G.V., Sinegub, I.A., 2002. Detritivorous polychaetes in the ecosystem from the Odessa region of the Black Sea. *Ecology of the sea*, 62, 5-9. (In Russian).
- Losovskaya, G.V., Sinegub, I.A., Bondarenko, A.S., 2005. Diversity of polychaete fauna in fouling of hydrotechnical constructions and benthos of Odessa port. *Ecological security of coastal and shelf zones and complex exploitation of shelf resources*, 12, 671-621. (In Russian).
- Losovskaya, G.V., Sinegub, I.A., Rybalko, A.A., 2004. Comparison of species composition and quantitative development of polychaetes in fouling and benthos of Odessa port. *Marine Ecology of the sea*, 3 (1), 51-58. (In Russian).
- Losovskaya, G.V., Zolotarev, V.N., 2003. The polychaete *Polydora limicola* in benthic communities of the Black Sea. *Russian Journal of Marine Biology*, 29 (4), 248-250. (In Russian).
- Mahon, A.R., Mahon, H.K., Dauer, D.M., Halanynch, K.M., 2009. Discrete genetic boundaries of three *Streblospio* (Spionidae, Annelida) species and the status of *S. shrubsolei*. *Marine Biology Research*, 5 (2), 172-178.
- Mccain, E.R., 2008. Poecilogony as a tool for understanding speciation: Early development of *Streblospio benedicti* and *Streblospio gynobranchiata* (Polychaeta: Spionidae). *Invertebrate Reproduction and Development*, 51 (2), 91-101.
- Moroz, T.G., 1977. Polychaeta of river mouth regions in the north-west Black Sea coast. *Zoological Journal*, 56 (7), 1103-1107. (In Russian with English summary).
- Moroz, T.G., 1985. Polychaeta of estuaries on the north-west coast of Black Sea. p. 104-107. In: *Polychaeta: Morphology, Systematics, Ecology (Explor. Fauna Seas, 34(42))*. Buzhinskaja, G.N. & Khlebovich, V.V. (Eds). Zoological Institute of the USSR Academy of Sciences, Leningrad. (In Russian).
- Moroz, T.G., 1992. Ecological peculiarities of mass species of polychaetes from the river mouth regions of north-west coast of the Black Sea. p. 145-152. In: *Polychaeta and their ecological significance [Explor. Fauna Seas, 43 (51)]*. Buzhinskaja, G.N. (Ed.). Zoological Institute, Russian Academy of Sciences, St. Petersburg. (In Russian).
- Murina, V.V., Selifonova, Zh.P., Melnik, V.F., 2008. Finding of polychaete *Streblospio* sp. (Polychaeta: Spionidae) in the Novorossiysk port of the Black Sea. *Marine Ecological Journal*, 7 (1), 46. (In Russian).
- Occhipinti-Ambrogi, A., Savini, D., 2003. Biological invasions as a component of global change in stressed marine ecosystems. *Marine Pollution Bulletin*, 46, 542-551.
- Pereyaslawzewa, S., 1891. Addendum to the Fauna of the Black Sea. *Proceedings of the Kharkov University Society of Naturalists*, 25, 235-274. (In Russian).
- Radashevsky, V.I., 2005. On adult and larval morphology of *Polydora cornuta* Bosc, 1802 (Annelida: Spionidae). *Zootaxa*, 1064, 1-24.
- Radashevsky, V.I., 2012. Spionidae (Annelida) from shallow waters around the British Islands: an identification guide for the NMBAQC Scheme with an overview of spionid morphology and biology. *Zootaxa*, 3152, 1-35.
- Rice, S.A., Levin, L.A., 1998. *Streblospio gynobranchiata*, a new spionid polychaete species (Annelida: Polychaeta) from Florida and the Gulf of Mexico with an analysis of phylogenetic relationships within the genus *Streblospio*. *Proceedings of the Biological Society of Washington*, 111 (3), 694-707.
- Selifonova, Zh.P., 2009. Marine biological invasions in waters of the port of Novorossiysk in the Black Sea. *Russian Journal of Marine Biology*, 35 (3), 242-249. (In Russian).
- Selifonova, Zh.P., 2011. Record of *Polydora cornuta* (Polychaeta: Spionidae) from the Tuapse port (the Black Sea). *Marine Ecological Journal*, 10 (2), 48. (In Russian).
- Selifonova, Zh.P., 2012. *Marine ecosystems of the ports of Novorossiysk and Tuapse of the Black Sea*. Nauka, St. Petersburg, 228 pp. (In Russian).
- Simboura, N., Sigala, K., Voutsinas, E., Kalkan, E., 2008. First occurrence of the invasive alien species *Polydora cornuta* Bosc, 1802 (Polychaeta: Spionidae) on the coast of Greece

- (Elefsis Bay; Aegean Sea). *Mediterranean Marine Science*, 9 (2), 121-126.
- Sinegub, I.A., 2001. Macrozoobenthos of the coastal waters of Zmeinny Island (Black Sea). *Ecological security of coastal and shelf zones and complex exploitation of shelf resources*, 2, 301-315. (In Russian).
- Sinegub, I.A., 2002a. Macrozoobenthos of the Sukhoi liman and adjacent part of the Black Sea. *Ecological security of coastal and shelf zones and complex exploitation of shelf resources*, 1 (6), 338-345. (In Russian).
- Sinegub, I.A., 2002b. Macrozoobenthos of a coastal ecotone Al-ibei liman (northern Black Sea). *Ecology of the sea*, 62, 30-33. (In Russian).
- Surugiu, V., 2005a. Inventory of inshore polychaetes from the Romanian coast. *Mediterranean Marine Science*, 6 (1), 51-73.
- Surugiu, V., 2005b. The use of polychaetes as indicators of eutrophication and organic enrichment of coastal waters: a study case-Romanian Black Sea Coast. *Analele Științifice ale Universității "Al.I. Cuza" Iași, s. Biologie animal*, 51, 55-62.
- Surugiu, V., 2006. New data on the polychaete fauna from the Romanian coast of the Black Sea. *Analele științifice ale Universității "Al. I. Cuza" Iași, serie Biologie animala*, Supplement: 47-56.
- Surugiu, V., 2012. Systematics and ecology of species of the *Polydora*-complex (Polychaeta: Spionidae) of the Black Sea. *Zootaxa*, 3518, 45-65.
- Surugiu, V., Manoleli, D., 1999. Nouvelle contribution à l'étude des Annélides Polychètes de la région d'Agiea (littoral Roumain). *Analele științifice ale Universității "Al. I. Cuza" Iași, serie Biologie animala*, 44-45, 21-25.
- Surugiu, V., Novac, A., 2007. Biodiversity and vertical distribution of polychaetes associated with mussel beds on artificial hard substrates (Romanian coast, Black Sea). *Rapport de la Commission Internationale pour l'Exploration Scientifique de la Mer Méditerranée*, 38, 609.
- Şahin, G.K., Çinar, M.E., 2012. A check-list of polychaete species (Annelida: Polychaeta) from the Black Sea. *Journal of the Black Sea/Mediterranean Environment*, 18 (1), 10-48.
- Taheri, M., Foshtomi, M.Y., 2011. Community structure and biodiversity of shallow water macrobenthic fauna at Noor coast, South Caspian Sea, Iran. *Journal of the Marine Biological Association of the United Kingdom*, 91 (3), 607-613.
- Taheri, M., Seyfabadi, J., Abtahi, B., Foshtomi, M.Y., 2008. Population changes and reproduction of an alien spionid polychaete, *Streblospio gynobranchiata*, in shallow waters of the south Caspian Sea. *JMBA2 - Biodiversity Records*, Published on-line: 1-5.
- Tena, J., Capaccioni-Azzati, R., Porras, R., Torres-Gavilá, F.J., 1991. Cuatro especies de poliquetos nuevas para las costas mediterráneas españolas en los sedimentos del antepuerto de Valencia. *Miscellánia Zoológica, Barcelona*, 15, 29-41.
- Vorobyova, L.V., Bondarenko, O.S., 2009. Meiobenthic bristle worms (Polychaeta) of the western Black Sea shelf. *Journal of the Black Sea/Mediterranean Environment*, 15, 109-121.
- Vorobyova, L., Bondarenko, O., Izaak, O., 2008. Meiobenthic polychaetes in the northwestern Black Sea. *Oceanological and Hydrobiological Studies*, 37 (1), 43-55.
- Webster, H.E., 1880. The Annelida Chaetopoda of New Jersey. *Annual Reports of the New York State Museum of Natural History*, 32, 101-128.
- Zaitsev, Y., Öztürk, B. (Eds), 2001. *Exotic species in the Aegean, Marmara, Black, Azov and Caspian Seas*. Turkish Marine Research Foundation, Istanbul, 267 pp.
- Zajac, R.N., 1991a. Population ecology of *Polydora ligni* (Polychaeta: Spionidae). I. Seasonal variation in population characteristics and reproductive activity. *Marine Ecology - Progress Series*, 77 (2-3), 197-206.
- Zajac, R.N., 1991b. Population ecology of *Polydora ligni* (Polychaeta: Spionidae). II. Seasonal demographic variation and its potential impact on life history evolution. *Marine Ecology - Progress Series*, 77 (2-3), 207-220.
- Zenetos, A., Çinar, M.E., Pancucci-Papadopoulou, M.A., Harmelin, J.G., Furnari, G. et al., 2005. Annotated list of marine alien species in the Mediterranean with records of the worst invasive species. *Mediterranean Marine Science*, 6 (2), 63-118.
- Zenetos, A., Gofas, S., Morri, C., Rosso, A., Violanti, D. et al., 2012. Alien species in the Mediterranean Sea by 2012. A contribution to the application of European Union's Marine Strategy Framework Directive (MSFD). Part 2. Introduction trends and pathways. *Mediterranean Marine Science*, 13 (2), 328-352.