

Mediterranean Marine Science

Vol 15, No 2 (2014)



Catch and discardfish species of trawl fisheries in the Iskenderun Bay (Northeastern Mediterranean) with emphasis on lessepsian and chondrichthyan species

E. YEMISKEN, C. DALYAN, L. ERYILMAZ

doi: [10.12681/mms.538](https://doi.org/10.12681/mms.538)

To cite this article:

YEMISKEN, E., DALYAN, C., & ERYILMAZ, L. (2014). Catch and discardfish species of trawl fisheries in the Iskenderun Bay (Northeastern Mediterranean) with emphasis on lessepsian and chondrichthyan species. *Mediterranean Marine Science*, 15(2), 380–389. <https://doi.org/10.12681/mms.538>

Catch and discard fish species of trawl fisheries in the Iskenderun Bay (North-eastern Mediterranean) with emphasis on lessepsian and chondrichthyan species

E. YEMISKEN, C. DALYAN and L. ERYILMAZ

Science Faculty, Department of Biology, Istanbul University, Turkey

Corresponding author: emreyemisken@yahoo.com.tr

Handling Editor: Costas Papaconstantinou

Received: 16 June 2013; Accepted: 3 April 2014; Published on line: 24 June 2014

Abstract

Fish species in catch and discard of trawl fisheries in and around Iskenderun Bay were examined within the fishing closure period and fishing period. The sampling was performed from May 2010 to January 2011 by a commercial trawl vessel. Chondrichthyan species accounted for 51 % of discard catch biomass while *Gymnura altavela* and *Dasyatis pastinaca* were dominant in hauls. 27 lessepsian fish species were captured during the study, nine of them being target species for trawl fisheries. In total, 14 of the lessepsian fish species were determined as discard species. In both sampling periods, *Equulites klunzingeri* and *Citharus linguatula* contributed to discard fish catch dissimilarity among depth ranges (deeper and shallower than 60 m). *E. klunzingeri* showed high abundance in discard catch. There were no significant differences in the distribution of the discard fish biomass between the sampling periods (ANOVA test, $p > 0.05$). However, depth range highlighted significant differences in discard fish catch composition ($p < 0.05$). Among major commercial fish species of trawl fisheries, *Mullus surmuletus* and *Sparus aurata* were not separated as discard in any haul by fishermen. Any size of these two species was included in the commercial catch (Total length ranged from 61 to 721 mm).

Keywords: Discard Fish, Trawl Fisheries, North-eastern Mediterranean Sea, Iskenderun Bay.

Introduction

World fisheries have not considered the sustainability of wild food resources until a few decades ago. Destruction of fish stocks was discovered by researchers in 1980s. Thereafter, the discard problem in marine fisheries was described (Alverson *et al.*, 1994; Hall, 1996). Non-target or non-commercial catches are defined as discard, generally brought onto the deck of fishing vessels and then returned to the sea. Although some species are included in the commercial part, they might be determined as discard because of unmarketable consideration (Hall *et al.*, 2000). Unmarketable sizes may change with fishing time and depth depending on fishermen (Kelleher, 2005).

Discard rates in multispecies fisheries are higher than for other fishing methods and the Mediterranean Sea fishery is characterized by multispecies fisheries (Tudela, 2004). Hence, discard rates geographically estimated ranged from 23 to 67 %, and belong to different Mediterranean Sea habitats (Machias *et al.*, 2001; D'Onghia *et al.*, 2003; Sanchez *et al.*, 2004; Tudela, 2004; Alsayes & Fatthouh 2009, Damalas *et al.*, 2010, Edelist *et al.*, 2011). Also, in the Turkish Seas, research has been focused on studies concerning discard catch in the last decade, and most of these studies were carried out using beam trawls (Kınacıgil, 1999a, 1999b; Demirci, 2003; Yazıcı *et al.*,

2006; Bayhan *et al.*, 2006; Soykan *et al.*, 2006; Gökçe & Metin, 2006). Gurbet *et al.* (2013) mentioned that the total discard biomass ratio was 30.5 % in the trawl fisheries of Izmir Bay (Aegean Sea). Gucu (2012) pointed out the status of bottom trawl fisheries from 1980s onwards in Levantine Sea, and compared data considering temporal and depth differences.

The objective of this study is to detect the fish composition of trawl fisheries in Iskenderun Bay and to analyze the commercial and discard fish species with their rates in different periods with emphasis on Lessepsian and Chondrichthyan fish species.

Material and Methods

Study Area

The study was carried out in and around Iskenderun Bay, which is located in the Levantine Sea (North-eastern Mediterranean) (Fig. 1), an important trawling area in the Eastern Mediterranean Sea. The bay shows high productivity caused by local wind effects and movements such as upwelling and rich terrestrial nutrient inputs. Surface water temperature ranges from 16°C to 33°C, and salinity between 37 and 40 psu (Polat & Piner, 2002; Polat, 2010). Regional fish communities are exposed to fishing pressure from almost one hundred trawlers vessel every




Fig. 1: Study area in and around Iskenderun Bay (Levantine Sea).

year. Marine fisheries along Turkish south coast (North of the Levantine Sea) include 6.93 % of marine fisheries productivity in Turkish seas (Anonymous, 2012a). The trawling ban period is between April 15 and September 15 in Turkish seas, in accordance with the decision of the Ministry of Agriculture and Rural Affairs, General Directorate of Protection and Control Vision of the Republic of Turkey. Trawling is permitted over two miles in the bay and one mile off Samandag coast (Anonymous, 2012b). Thus, in general, trawl fishery activities in the bay take place on bottoms deeper than 60 m.

Sampling Data

Field work was performed from May 2010 to January 2011 in Iskenderun Bay. The May and August surveys were included in the trawl fishing closure period, while the October and January surveys took place during the fishing period. Data was collected using a commercial trawl vessel (23 m in length) with engine power up to 400 HP, and the stretched codend mesh size of the trawl net was 44 mm. A total of 32 hauls were performed at depths between 31 and 110 m determined by the fishermen. Ten hauls were carried out in May, seven hauls in August, seven hauls in October and eight hauls in January. The mean haul duration was 1.5 hours at an average speed of 2.5 knots. The commercial catch in each haul was selected by the crew on board, and the remaining catch was considered as discard. Biomass and abundance data of the fish species was recorded as discard and commercial catch by researchers. We randomly applied subsampling for the most abundant species such as *E. klunzingeri*. The commercial and discard catches were stored at -20°C at the laboratory. After identification of the fish species, total length (mm) and body weight (g) were measured.

Data Analysis

The catch data of each haul was standardized with the catch per unit effort (CPUE) (D'Onghia *et al.*, 2003). It was performed as fish weight divided by catch time, where C_w is the weight of the fish catch and C_t is the towing time.

$$\text{CPUE} = C_w / C_t$$

The number of species (S) was counted for each haul. The diversity indices were computed using the number of specimens to standardize with catch hour per haul. The estimation was made using the Simpson diversity index, Species Richness, Shannon-Wiener diversity index H' (\log_2) and Pielou evenness index;

$$\text{Simpson Diversity Index} = 1 - \lambda \rightarrow \lambda =$$

$$\frac{\sum n_i(n_i - 1)}{N(N - 1)} \rightarrow 1 - \frac{\sum n_i(n_i - 1)}{N(N - 1)}$$

" λ " is the Simpson Index and Simpson diversity index is found $1 - \lambda$, where " n_i " is the total number of specimens of each species and " N " is the total number of specimens of all species.

Margalef Species Richness index (d) = $(S - 1) / \ln N$, where " S " is the number of species in the samples (Beisel and Moreteau, 1997).

Shannon Wiener Diversity Index (H') = $-\sum p_i \ln(p_i)$ where " p_i " is the proportion of total samples belonging to the species,

$$\text{Evenness Index } (J') = H' / \log S$$

Diversity indices were compared with sampling months. Eight commercial fish species: *Saurida undosquamis*, *Nemipterus randalli*, *Mullus barbatus*, *M. surmuletus*, *Pagellus erythrinus*, *Upeneus moluccensis*,




Fig. 2: MDS plot of discard fish catch composition. Each haul was assigned as ▲ ; deeper than 60 m and ▼ ; shallower than 60 m.

U. pori, *Sparus aurata* were analyzed to total length measurement in all hauls.

Haul depths were divided in two: hauls with depths shallower than 60 m and deeper than 60 m. Homogeneity of variance was tested with Levene's test. Variances were transformed using the $\log(x+1)$ formula. After data transformation, the factor on fish discard biomass was analyzed with the one-way ANOVA statistical test. The analysis was carried out using the statistical programme SPSS, version 17 (Levesque, 2007).

Abundance of fish discard data was transformed $[\log(x+1)]$ and then rank-ordered using the Bray Curtis similarity matrix. The Non-parametric Multidimensional Scaling test was performed using the Primer 6 statistical package programme (Clarke & Warwick, 2001). The SIMPER test was performed to determine the level of similarity and dissimilarity according to trawl fishing sampling depth.

Results

77994 individuals were obtained in 32 hauls, belonging to 97 fish species (Table 1). 69 (n=54119) of them were determined as discard. 43 (n=52655) species were thrown back into sea regardless of their size, while 26 species (n=1464) have been considered as unmarketable size of commercial species. 993.4 kg of the total 2491.3 kg of fish catch was discard (Table 1).

Total fish biomass caught per haul was between 5.7 and 119.5 kg/h during the fishing period and between 1.1 and 40.9 kg/h of that amount was discard catch. However, total fish biomass caught per haul ranged from 3.4 to 211.2 kg/h during the fishing closure time, and 0.6 to 151.2 kg/h of that amount was discard catch. The average discard catch rate was found to be 32.2 ± 16.5 % during the study. The estimates of fish biomass have been separately studied in terms of depth ranges (Table 2).

Ecological Index

The ecological indices of total and discard fish catch per month are shown in Table 3. The evenness index (J') ranged from 0.40 to 0.63 in total catch. The Shannon - Weiner diversity index per haul was estimated to be 2.46 in May 2010, 1.91 in August 2010, 2.55 in October 2010 and 2.68 in January 2011.

Abundance of discard catches showed differences according to depth (ANOSIM. $R=0.531$, $p<0.001$). The SIMPER analysis indicated that *C. linguatula* displayed highest average similarity among discard fish. In waters shallower than 60 m, *E. klunzingeri* showed highest average abundance and nine Lessepsian species contributed mainly to average similarity. Also, these two species contributed to dissimilarity in the sampling depth groups (Table 4).

Comparison of Discard Fish Biomass

The computation of the discard fish rates indicated that 44.5 % of the catch was in the depth range of 30-60 m and 28.1% was in the depth range of 60-110 m. The average of discard rates among sampling months showed close values except of January. The highest discard CPUE was determined in August (average: 44.86 kg/h), while the lowest was in October (average: 9.04 kg/h). Although a significant relationship was found between depth and discard fish biomass ($p < 0.05$, $p = 0.0001$), there is no differences between sampling periods and discard biomass ($p > 0.05$, $p = 0.106$).

Chondrichthyan species

Chondrichthyan species abundance represented 0.9 % in total discard catch. (n=465). Four species were obtained in the commercial catch: *Mustelus mustelus* (n=1). *Carcharhinus plumbeus* (n=2) and *Rhinobatus cemiculus* (n=1), Chondrichthyan species are included in discard catch biomass at a rate of 51 %. *G. altavela* (n=203) and *D. pastinaca* (n=125) were the main species among the chondrichthyans in the study area (Table 1). Discard rates ranged from 27 % to 77 % depending on the sampling periods. Discard of chondrichthyan species was estimated to be 12.5 (May), 19.1 (August), 2.3 (October) and 5.08 (January) (Fig. 3.).

Lessepsian Species

In total, 27 Red Sea originated fish species were found in the study. 13 of them were commercial and 14 species were discard. Lessepsian fish constituted 71 % of total catch abundance. They constituted 55 % of total catch biomass and dominated the commercial catch biomass at the rate of 69.5 %. In January, commercial biomass of Lessepsian fish reached 93.5% in a haul. Furthermore, they represented between 0.7 % and 92.8 % (on average 38.5 %) of the discard catch.

Thirteen (13) specimens of the pelagic pony fish, *E. klunzingeri*, were found, representing with 0.4 % in total catch biomass deeper than 60 m depth, and consti-

Table 1. Commercial (C) or Discard (D); N, number of specimens; F %, frequency of species occurrence in total hauls; Discard Rate % (per each discard species), percentage of discard abundance in total.

| Species | N | F % | Total Weight (kg) | Discard Weight (kg) | Discard Rate % | Discard Size (mm) | Commercial Size (mm) | Legal Minimum Landing Size | Value |
|-----------------------------------|-------|-----|-------------------|---------------------|----------------|-------------------|----------------------|----------------------------|-------|
| Lessepsian Fishes | 67324 | | 1469.5 | 399.7 | 27.2 | | | | |
| <i>Apogon fasciatus</i> | 69 | 40 | 0.822 | 0.822 | 100 | 53-106 | - | - | D |
| <i>Apogon queketti</i> | 77 | 53 | 1.262 | 1.262 | 100 | 63-128 | - | - | D |
| <i>Apogon smithii</i> | 138 | 39 | 3.244 | 3.244 | 100 | 61-146 | - | - | D |
| <i>Callionymus filamentosus</i> | 140 | 30 | 2.753 | 2.753 | 100 | 76-148* | - | - | D |
| <i>Champsodon sp.</i> | 796 | 51 | 9.022 | 9.022 | 100 | 46-149 | - | - | D |
| <i>Cynoglossus sinuababici</i> | 1 | 4 | 0.003 | 0.003 | 100 | 82 | - | - | D |
| <i>Decapturnus russeli</i> | 2 | 3 | 0.154 | 0.075 | 100 | 90-92 | - | - | C |
| <i>Dussumieria elopoides</i> | 4 | 7 | 0.1 | 0 | 0 | - | 146-161 | - | C |
| <i>Equulites klunzingeri</i> | 46841 | 62 | 354.895 | 354.895 | 100 | 46-102 | - | - | D |
| <i>Etmeneus teres</i> | 5 | 12 | 0.293 | 0 | 0 | - | 158-240 | - | C |
| <i>Fistularia commersoni</i> | 6 | 7 | 0.792 | 0.057 | 100 | 258-338 | - | - | D |
| <i>Lagocephalus scleratus</i> | 3 | 6 | 0.655 | 0.655 | 100 | 71-385 | - | - | D |
| <i>Lagocephalus spadiceus</i> | 103 | 41 | 7.541 | 7.541 | 100 | 60-291 | - | - | D |
| <i>Lagocephalus suezensis</i> | 136 | 39 | 5.548 | 5.548 | 100 | 106-177 | - | - | D |
| <i>Nemipterus randalli</i> | 3372 | 100 | 166.046 | 3.61 | 6 | 52-130* | 85-194 | - | C |
| <i>Oxyrichthys papuensis</i> | 53 | 32 | 0.763 | 0.763 | 100 | 113-170 | - | - | D |
| <i>Pomadasys stradiens</i> | 19 | 3 | 0.996 | 0 | 0 | - | 132-186 | - | C |
| <i>Saurida undosquamis</i> | 7177 | 100 | 653.681 | 1.747 | 1.8 | 61-172 | 131-362 | - | C |
| <i>Siganus rivulatus</i> | 3 | 7 | 0.052 | 0.052 | 100 | 106-124 | - | - | C |
| <i>Sillago sihama</i> | 4 | 7 | 0.164 | 0 | 0 | - | 170-181 | - | C |
| <i>Sphyræna chrysotenia</i> | 22 | 8 | 1.949 | 0 | 0 | - | 197-275 | - | C |
| <i>Sphyræna flavicauda</i> | 2 | 4 | 0.193 | 0 | 0 | - | 250-278 | - | C |
| <i>Stephanolepis diaspros</i> | 158 | 58 | 4.534 | 4.534 | 100 | 80-202 | - | - | D |
| <i>Torquigener flavimaculosus</i> | 9 | 14 | 0.104 | 0.104 | 100 | 72-109 | - | - | D |
| <i>Trachurus indicus</i> | 1 | 3 | 0.048 | 0 | 0 | 171 | - | - | C |
| <i>Upeneus moluccensis</i> | 613 | 68 | 10.147 | 1.203 | 28 | 66-115 | 70-218 | 100 | C |
| <i>Upeneus pori</i> | 7570 | 54 | 243.776 | 1.859 | 2 | 80-154 | 106-167 | - | C |
| Native Fishes | | | 1021.8 | 593.7 | | | | | |
| Chondrichthyes | 474 | | 519.8 | 503.0 | 96.8 | | | | |
| <i>Carcharhinus plumbeus</i> | 2 | 4 | 4.9 | 0 | 0 | - | 708-721 | - | C |
| <i>Dasyatis marmorata</i> | 32 | 28 | 20.314 | 20.314 | 100 | 458-668 | - | - | D |
| <i>Dasyatis pastinaca</i> | 125 | 52 | 162.685 | 162.685 | 100 | 450-1100 | - | - | D |
| <i>Mustelus mustelus</i> | 1 | 3 | 4.358 | 0 | 100 | 1030 | - | - | C |
| <i>Gymnura altavela</i> | 203 | 29 | 246.133 | 246.133 | 100 | 265-1040 | - | - | D |
| <i>Pteromyia laeus bovinus</i> | 5 | 14 | 12.08 | 12.08 | 100 | 535-1490 | - | - | D |
| <i>Raja miraletus</i> | 16 | 23 | 1.375 | 1.375 | 100 | 80-394 | - | - | D |
| <i>Raja radula</i> | 31 | 23 | 19.061 | 19.061 | 100 | 208-505 | - | - | D |
| <i>Raja sp.</i> | 5 | 3 | 0.02 | 0.02 | 100 | 50-85 | - | - | D |
| <i>Rhinobatos cemiculus</i> | 51 | 26 | 48.347 | 40.847 | 98 | 185-764 | 1300 | - | C |
| <i>Torpedo marmorata</i> | 2 | 7 | 0.196 | 0.196 | 100 | 162 | - | - | D |
| <i>Torpedo torpedo</i> | 1 | 4 | 0.327 | 0.327 | 100 | 295 | - | - | D |
| Teleosts | 10196 | | 502.6 | 90.7 | 18.1 | | | | |
| <i>Argyrosomus regius</i> | 6 | 7 | 1.297 | 0 | 0 | - | 225-337 | 250 | C |
| <i>Arnoglossus kessleri</i> | 4 | 7 | 0.041 | 0.041 | 100 | 82-100 | - | - | D |
| <i>Arnoglossus laterna</i> | 6 | 3 | 0.078 | 0.078 | 100 | 94-130 | - | - | D |
| <i>Arnoglossus sp.</i> | 9 | 13 | 0.047 | 0.047 | 100 | 64-102 | - | - | D |
| <i>Arnoglossus thori</i> | 36 | 34 | 0.661 | 0.661 | 100 | 80-117 | - | - | D |
| <i>Balistes capricornus</i> | 1 | 3 | 0.231 | 0 | 100 | 183 | - | - | D |
| <i>Blennius ocellaris</i> | 2 | 6 | 0.055 | 0.055 | 100 | 128 | - | - | D |
| <i>Boops boops</i> | 15 | 17 | 0.726 | 0 | 0 | - | 136-188 | - | C |
| <i>Bothus podas</i> | 73 | 22 | 1.981 | 1.981 | 100 | 78-179 | - | - | D |

(continued)

Table 1 (continued)

| Species | N | F % | Total Weight (kg) | Discard Weight (kg) | Discard Rate % | Discard Size (mm) | Commercial Size (mm) | Legal Minimum Landing Size | Value |
|--------------------------------------|------|-----|-------------------|---------------------|----------------|-------------------|----------------------|----------------------------|-------|
| <i>Caranx rhonchus</i> | 9 | 7 | 0.994 | 0 | 0 | - | 214 | - | C |
| <i>Chelidonichthys lucernus</i> | 376 | 82 | 33.765 | 0.342 | 4 | 107-148 | 150-286 | 180 | C |
| <i>Citharus linguatula</i> | 3219 | 78 | 58.643 | 58.643 | 100 | 75-206 | - | - | D |
| <i>Conger conger</i> | 1 | 3 | 0.404 | 0.404 | 100 | 661 | - | - | D |
| <i>Dactylopterus volitans</i> | 1 | 3 | 0.13 | 0.13 | 100 | 233 | - | - | D |
| <i>Deltentosteus quadrimaculatus</i> | 46 | 20 | 0.215 | 0.215 | 100 | 48-95 | - | - | D |
| <i>Dentex dentex</i> | 1 | 4 | 0.138 | 0 | 0 | - | 217 | 350 | C |
| <i>Diplodus annularis</i> | 41 | 19 | 2.182 | 0.171 | 12 | - | 114-182 | - | C |
| <i>Diplodus sargus</i> | 4 | 3 | 0.228 | 0 | 0 | - | 144-165 | 210 | C |
| <i>Diplodus vulgaris</i> | 6 | 5 | 0.353 | 0 | 0 | - | 149-164 | 180 | C |
| <i>Echeneis naucrates</i> | 2 | 6 | 0.374 | 0.374 | 100 | 360-448 | - | - | D |
| <i>Engraulis encrasicolus</i> | 26 | 12 | 0.276 | 0.037 | 100 | 62-133 | - | 90 | C |
| <i>Epinephelus aeneus</i> | 86 | 33 | 37.828 | 0.054 | 49 | 120-170 | 196-920 | 450 | C |
| <i>Epinephelus haifensis</i> | 2 | 3 | 0.074 | 0 | 0 | - | 125-133 | - | C |
| <i>Epinephelus marginatus</i> | 3 | 6 | 1.065 | 0 | 0 | - | 179-390 | - | C |
| <i>Gobius niger</i> | 4 | 14 | 0.069 | 0.069 | 100 | 82-137 | - | - | D |
| <i>Lepidotrigla cavillone</i> | 67 | 46 | 0.773 | 0.773 | 100 | 62-111 | - | - | D |
| <i>Lithognathus mormyrus</i> | 110 | 10 | 11 | 0 | 0 | - | 145-212 | - | C |
| <i>Merluccius merluccius</i> | 1 | 4 | 0.04 | 0 | 0 | - | 185 | 250 | C |
| <i>Microchirus ocellatus</i> | 61 | 7 | 1.348 | 1.348 | 100 | 106-131 | - | - | D |
| <i>Mullus barbatus</i> | 253 | 76 | 16.865 | 0.253 | 9.37 | 71-116 | 114-257 | 130 | C |
| <i>Mullus surmuletus</i> | 63 | 28 | 4.303 | 0 | 0 | - | 103-235 | 110 | C |
| <i>Pagellus acarne</i> | 281 | 26 | 4.876 | 2.413 | 62 | 73 -123 | 122 - 162 | - | C |
| <i>Pagellus erythrinus</i> | 3937 | 86 | 219.346 | 2.85 | 0.38 | 71-135 | 105-245 | - | C |
| <i>Pagrus caeruleostictus</i> | 16 | 9 | 2.629 | 0.036 | 15 | 98-110 | 190-288 | - | C |
| <i>Phycis phycis</i> | 1 | 3 | 0.006 | 0.006 | 100 | 92 | - | - | C |
| <i>Pomadasys incisus</i> | 27 | 6 | 2.144 | 0 | 0 | - | 69-183 | - | C |
| <i>Sardinella aurita</i> | 1 | 3 | 0.001 | 0.001 | 100 | 130 | - | 110 | C |
| <i>Sarpa salpa</i> | 1 | 3 | 0.05 | 0 | 100 | 151 | - | - | C |
| <i>Scomber japonicus</i> | 5 | 9 | 0.314 | 0 | 0 | - | 139-191 | 180 | C |
| <i>Scorpaena notata</i> | 2 | 6 | 0.246 | 0.061 | 50 | 141 | 228 | - | C |
| <i>Scorpaena scrofa</i> | 2 | 3 | 0.114 | 0.114 | 100 | 152 | - | 150 | C |
| <i>Seriola dumerili</i> | 1 | 4 | 0.186 | 0 | 0 | - | 245-336 | 300 | C |
| <i>Serranus cabrilla</i> | 18 | 16 | 0.302 | 0.269 | 100 | 85-161 | - | - | D |
| <i>Serranus hepatus</i> | 88 | 49 | 0.677 | 0.677 | 100 | 58-101 | - | - | D |
| <i>Solea kleini</i> | 4 | 7 | 0.16 | 0.03 | 25 | 152 | 158-176 | - | C |
| <i>Solea lascaris</i> | 1 | 3 | 0.047 | 0.047 | 100 | 172 | - | - | D |
| <i>Solea solea</i> | 173 | 79 | 13.397 | 0.82 | 13 | 139-217 | 141-323 | 200 | C |
| <i>Sparus aurata</i> | 544 | 94 | 50.855 | 0 | 0 | - | 165-209 | 200 | C |
| <i>Sphyræna sphyræna</i> | 54 | 18 | 6.73 | 0 | 0 | - | 280-409 | - | C |
| <i>Spicara flexuosa</i> | 14 | 15 | 0.482 | 0.185 | 0 | - | 129-170 | - | C |
| <i>Spicara maena</i> | 2 | 3 | 0.127 | 0 | 0 | - | 157-180 | - | C |
| <i>Spicara smarís</i> | 1 | 4 | 0.005 | 0 | 100 | 80 | - | - | D |
| <i>Spondylíosa cantharus</i> | 8 | 3 | 0.393 | 0 | 0 | - | 121-170 | - | C |
| <i>Synodus saurus</i> | 14 | 10 | 0.518 | 0.518 | 100 | 126-226 | - | - | C |
| <i>Trachinus draco</i> | 30 | 32 | 2.429 | 2.429 | 100 | 171-296 | - | - | D |
| <i>Trachurus trachurus</i> | 43 | 36 | 1.718 | 0.068 | 9 | 91-223 | 91-236 | 130 | C |
| <i>Trichiurus lepturus</i> | 125 | 42 | 9.172 | 9.172 | 100 | 409-556 | - | - | D |
| <i>Trigloporus lastoviza</i> | 81 | 28 | 3.914 | 0.657 | 53 | 74-159 | 152-228 | - | C |
| <i>Uranoscopus scaber</i> | 65 | 39 | 2.072 | 2.072 | 100 | 83-115 | - | - | D |
| <i>Zeus faber</i> | 123 | 41 | 3.473 | 2.64 | 95 | 118-164 | 188-226 | - | C |

Table 2. Average total and discard catch biomass (kg/h) and discard rates according to depth ranges and sampling months.

| | May | | August | | October | | January | | Total | |
|---------------------|-------|--------|--------|--------|---------|--------|---------|--------|---------------|---------------|
| Depths (m) | 30-60 | 60-110 | 30-60 | 60-110 | 30-60 | 60-110 | 30-60 | 60-110 | 30-60 | 60-110 |
| Sampling Hauls | 4 | 6 | 4 | 3 | 1 | 6 | 3 | 5 | 12 | 20 |
| Total Catch Biomass | 70.4 | 13.8 | 135.8 | 71 | 21.6 | 21.9 | 68 | 21.7 | 87.6 ± 61.01 | 26.78 ± 22.86 |
| Discard Biomass | 31.4 | 5.1 | 71.7 | 9.1 | 10 | 8.9 | 17.7 | 7.8 | 39.61 ± 39.23 | 7.5 ± 8.06 |
| Discard Rate (%) | 44.5 | 37.2 | 52.8 | 12.8 | 46.3 | 40.6 | 26 | 35.9 | 39.83 ± 16.13 | 27.6 ± 14 |

Table 3. Ecological parameter means for the sampling periods in Iskenderun Bay. S: Species number (S), N (Total specimens), d (Species Richness), J' (Evenness Diversity index), H' (Shannon-Wiener Diversity Index), 1-λ (Simpson Diversity Index), Mean, SD (Standard Deviation).

| | May | | August | | October | | January | | Total | | | |
|-----|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|-------------|
| | Catch | Discard | Catch | Discard | Catch | Discard | Catch | Discard | Catch | | Discard | |
| | | | | | | | | | 30-60 | 60-110 | 30-60 | 60-110 |
| N | 10704 | 4651 | 41746 | 30826 | 5479 | 4271 | 20065 | 14357 | 64772 | 13222 | 48507 | 5598 |
| S | 21± 6 | 13±4 | 28 ± 4 | 20 ± 4 | 22 ± 5 | 16 ± 5 | 21 ± 3 | 12 ± 2 | 26 ± 4 | 22 ± 6 | 17 ± 3 | 14 ± 6 |
| H' | 2.46 ± 0.52 | 1.49 ± 0.72 | 1.91 ± 0.70 | 0.68 ± 0.56 | 2.55 ± 0.59 | 1.23 ± 0.37 | 2.68 ± 0.65 | 1.5 ± 0.64 | 1.85 ± 0.72 | 2.75 ± 0.31 | 0.92 ± 0.95 | 1.46 ± 0.3 |
| J | 0.57 ± 0.13 | 0.62 ± 0.27 | 0.40 ± 0.14 | 0.12 ± 0.08 | 0.59 ± 0.15 | 0.26 ± 0.14 | 0.63 ± 0.16 | 0.38 ± 0.14 | 0.39 ± 0.15 | 0.6 ± 0.09 | 0.33 ± 0.35 | 0.59 ± 0.15 |
| d | 3.51 ± 0.55 | 2.56 ± 0.9 | 3.47 ± 0.70 | 2.8 ± 0.92 | 3.40 ± 0.53 | 2.6 ± 0.7 | 3.08 ± 0.37 | 2.28 ± 0.83 | 3.33 ± 0.55 | 3.39 ± 0.57 | 2.52 ± 0.8 | 2.57 ± 0.91 |
| 1-λ | 0.71 ± 0.11 | 0.6 ± 0.27 | 0.55 ± 0.18 | 0.25 ± 0.22 | 0.69 ± 0.18 | 0.5 ± 0.18 | 0.76 ± 0.16 | 0.63 ± 0.25 | 0.55 ± 0.2 | 0.8 ± 0.09 | 0.33 ± 0.36 | 0.62 ± 0.13 |

Table 4: Results of SIMPER analysis of discard catch. Av. Ab.; Average of abundance, Av. Sim.; Average of Similarity, Sim./SD; Similarity of Standard Deviation, Contrib.; Contribution Percentage, Cum.; Cumulative Percentage. Av. Diss.; Average of Dissimilarity.

| Species | Av.Ab. | Av.Sim | Sim/SD | Contrib% | Cum.% | |
|--|----------|----------|---------|----------|------------|--------|
| Deeper than 60 m depth (60-110 m) average similarity : 38.78 | | | | | | |
| <i>Citharus linguatula</i> | 3.88 | 13.7 | 2.22 | 35.34 | 35.34 | |
| <i>Saurida undosquamis</i> | 1.32 | 4.22 | 1.49 | 10.89 | 46.23 | |
| <i>Champsodon sp.</i> | 2.04 | 4.14 | 0.61 | 10.68 | 56.91 | |
| <i>Nemipterus randalli</i> | 1.46 | 3.6 | 1.22 | 9.29 | 66.2 | |
| <i>Lepidotrigla cavillione</i> | 0.87 | 2.05 | 0.72 | 5.28 | 71.48 | |
| <i>Serranus hepatus</i> | 1.01 | 1.65 | 0.71 | 4.27 | 75.74 | |
| <i>Zeus faber</i> | 0.86 | 1.65 | 0.45 | 4.25 | 79.99 | |
| <i>Uranoscopus scaber</i> | 0.74 | 1.08 | 0.54 | 2.79 | 82.77 | |
| <i>Upeneus moluccensis</i> | 0.88 | 0.89 | 0.4 | 2.28 | 85.06 | |
| <i>Apogon quaketti</i> | 0.64 | 0.88 | 0.48 | 2.28 | 87.34 | |
| Shallower than 60 m depth (30-60) average similarity : 38.29 | | | | | | |
| <i>Equulites klunzingeri</i> | 5.43 | 11.36 | 1.01 | 29.68 | 29.68 | |
| <i>Apogon smithii</i> | 1.64 | 3.47 | 0.89 | 9.07 | 38.75 | |
| <i>Upeneus pori</i> | 1.44 | 2.85 | 1.01 | 7.45 | 46.2 | |
| <i>Dasyatis pastinaca</i> | 1.41 | 2.83 | 1.04 | 7.4 | 53.6 | |
| <i>Nemipterus randalli</i> | 0.99 | 2.09 | 0.91 | 5.47 | 59.07 | |
| <i>Stephanolepis diaspros</i> | 0.94 | 1.95 | 1.48 | 5.08 | 64.15 | |
| <i>Saurida undosquamis</i> | 0.83 | 1.4 | 0.79 | 3.64 | 67.8 | |
| <i>Apogon fasciatus</i> | 0.85 | 1.39 | 0.56 | 3.63 | 71.43 | |
| <i>Lagocephalus spadiceus</i> | 1.1 | 1.16 | 0.55 | 3.03 | 74.46 | |
| <i>Citharus linguatula</i> | 1.07 | 1.13 | 0.45 | 2.95 | 77.41 | |
| Shallower & Deeper than 60 m average dissimilarity: 78.08 | | | | | | |
| | > 60 m | 60 m > | | | | |
| | Av.Abund | Av.Abund | Av.Diss | Diss./SD | Contrib. % | Cum. % |
| <i>Equulites klunzingeri</i> | 0.64 | 5.43 | 9.99 | 1.48 | 12.79 | 12.79 |
| <i>Citharus linguatula</i> | 3.88 | 1.07 | 6 | 1.95 | 7.69 | 20.48 |
| <i>Champsodon sp.</i> | 2.04 | 0.39 | 4.15 | 1.05 | 5.32 | 25.8 |
| <i>Apogon smithii</i> | 0.11 | 1.64 | 3.37 | 1.26 | 4.31 | 30.11 |
| <i>Upeneus pori</i> | 0.29 | 1.44 | 2.67 | 1.38 | 3.41 | 33.53 |
| <i>Dasyatis pastinaca</i> | 0.38 | 1.41 | 2.66 | 1.27 | 3.41 | 36.94 |
| <i>Nemipterus randalli</i> | 1.46 | 0.99 | 2.16 | 1.33 | 2.76 | 39.7 |
| <i>Upeneus moluccensis</i> | 0.88 | 0.6 | 2.12 | 0.91 | 2.72 | 42.42 |
| <i>Gymnura altavela</i> | 0.25 | 0.99 | 2.11 | 0.88 | 2.7 | 45.12 |
| <i>Stephanolepis diaspros</i> | 0.68 | 0.94 | 2.07 | 1.46 | 2.65 | 47.77 |




Fig. 3: Chondrichthyan species in fish discard.

tuting 20 % of the total fishing rate in terms of abundance values. When its CPUE is compared with the its depth range, the pony fish had 17.4 kg/h at depths between 30 to 60 m and 0.02 kg/h in waters deeper than 60 m.

Some of the other Lessepsian species such as *N. randalli*, *U. moluccensis*, *U. pori* and *S. undosquamis* were mainly commercial species for regional fisheries. Biomass values of these species in total catch were 3.4 kg/h,

0.2 kg/h, 5 kg/h and 13.6 kg/h, respectively. While *U. moluccensis* was found in waters deeper than 60 m, *U. pori* was detected at depths between 30 and 60 m.

Length Ranges of the Commercial Species

S. undosquamis (654 kg), *U. pori* (244 kg), *P. erythrinus* (219 kg), *N. randalli* (166 kg) and *S. aurata* (51 kg) constitute the major target species in commercial trawl fisheries of Iskenderun Bay. These five species consisted of 54 % of the commercial catch these species. Besides, these fishes have low discard weight rates (0 to 2.2%). *S. undosquamis*, the highest biomass commercial species, attained a length range of 61 to 172 mm (mean = 129 mm, SD= 45.2) in discard. The length of *P. erythrinus* ranged between 71 mm and 135 mm (mean = 113 mm, SD = 21.8) in discard. However, individuals that were bigger than 105 mm were evaluated to have economic value. Discard length of *N. randalli* was between 52 mm and 130 mm (mean = 97.2 mm, SD= 14.3), while *Upeneus moluccensis* attained a discard length of 66 to 115 mm (mean = 89 mm, SD= 12.2) (Fig. 5).




Fig. 4: Commercial and discard catch length ranges of eight commercial fish species in trawl fisheries; mean, standard deviation (SD), number of specimens (N).

S. aurata and *M. surmuletus* were not in the discard catch. Body size of the commercial fish ranged from 69 mm to 721 mm (Fig. 5).

Discussion

The variations of abundance and biomass data demonstrate the direct effect of fisheries on fish fauna. The percentages of discard fish species showed different results due to the changeable factors such as depth and time in Mediterranean Sea trawl fisheries (Machias *et al.*, 2001; D'Onghia *et al.*, 2003; Sanchez *et al.*, 2004; Tudela, 2004; Alsayes & Fathouh, 2009; Damalas *et al.*, 2010; Edelist *et al.*, 2011).

The study area is highly exploited during the fishing season (September 15 - April 15). High rates of discard were accompanied by low homogeneity in August (Table 2). The fish catch shown a more homogenous structure in fishing season. Also, the number of discard species declined toward the end of fishing season (Table 3).

Sanchez *et al.* (2007) showed that *M. barbatus* had low discard rates in the Adriatic and Catalan sea. Gucu (2012) verified that discard rates increased toward shallower depths in Mersin Bay. *Bothus podas* and the unmarketable size of *M. barbatus* were included in discard composition shallower than 100 m in Mersin Bay (North-Western Levantine Sea) (Gucu, 2012). In our study, *M. barbatus* also had a low ratio in discard catch. Besides, *S. aurata* and *M. surmuletus* was not included in discard catch. All specimens were considered as marketable, but some were not suitable for the minimum landing size regulation (Fig. 4).

Discard biomass ratios were calculated to be 44 % for the shallower than 60 m depth, and its value was 75 % of all the discard catch. 28 % of the total catches consisted of biomass of trawl fishing area (deeper than 60 m). These ratios indicate that shallow water has higher fish productivity than the trawl fishing area. We can support this opinion with biomass of *S. undosquamis*. It was found to be 9.94 kg/h and 19.57 kg/h in shallower water and deeper depths, respectively. It could be followed by long-term researches on commercial species biomass such as *S. undosquamis* in order to get more information in the fishing closed area of Iskenderun bay.

Chondrichthyan Fishes

The biomass percentages of cartilaginous species constituted the main discard group within the discard catch (Fig.3). *Carcharhinus plumbeus*, *Gymnura altavela* and *Mustelus mustelus* are classified as "vulnerable", while *Rhinobatos cemiculus* is "endangered" on the IUCN list (IUCN, 2013). There are some studies on *Carcharhinus plumbeus* in the Turkish Seas. One of these studies was carried out in Boncuk Bay (South-eastern Aegean Sea), which is thought to be a spawning and hatching area for the species (Bilecenoglu, 2008; Akca, 2010). Although the marketing of this specie is banned by law in

Turkey (Anonymous, 2012b). *C. plumbeus* is still caught and sold on the markets due to the poor control mechanism. Damalas & Vassilopoulou (2009) pointed out that *Dasyatis pastinaca* and *Mustelus sp.* are mostly marketed in the central Aegean Sea. However, Gurbet *et al.* (2013) mentioned that cartilaginous species were not included in the commercial catch composition in Izmir Bay (Aegean Sea). *Mustelus* species are rarely found, and it is evaluated as an incidental commercial catch in Iskenderun Bay. *Dasyatis* is not considered as a marketable species in the area.

Lessepsian fishes

The structure of the coastal fish community shows variability in the area because of new immigrant species entering from the Red Sea. In addition, the majority of the total catches consisted of these species in the regional fisheries. For example, *N. randalli* was recorded in 2008 (Bilecenoglu & Russell, 2008), and is now considered as a commercial species by fishermen. The biomass of *N. randalli* was found to be 2.42 kg/h in the study. Gucu *et al.* (2010) found that the percentages of Lessepsian fish species decreased in total catch when compared to the 1980s and 2000s, but the number of Lessepsian fish species increased. Gucu *et al.* (1994) reported 20 Lessepsian fish species in the 1980s. Recently, we captured 27 Lessepsian fish species in Iskenderun Bay. Until now, over 50 Lessepsian fish species are known in Turkish Seas (Bilecenoglu *et al.*, 2002; Erguden *et al.*, 2012; Dalyan *et al.*, 2012; Bodilis *et al.*, 2014). In the 1980s, *S. undosquamis* and *E. klunzingeri* formed the highest biomass at the rate of 23 % and 14 % of total catch in the Levantine Sea (Gucu *et al.*, 2010). We obtained practically the same results for these species (*S. undosquamis* 26 % and *E. klunzingeri* 13 % of total fish catch). *Equulites klunzingeri* was a major species within the discard catch. On the continental shelf of the North Eastern Levantine Sea, *E. klunzingeri* was found at the rates of 9.4 %, 2.3 % and 20.9 % of the fish fauna in fall 1983, spring 1984 and fall 1984, respectively (Gucu *et al.*, 2010). Çiçek (2006) mentioned that *E. klunzingeri* was 0.81 kg/h in Tasucu Bay (The west of Iskenderun Bay). The distribution of Pony fish probably decreases towards the west of Iskenderun Bay. In the sampling area, its distribution depends on the depth range (Table 3). In our study, it was observed that *Citharus linguatula* and *Champsodon sp.* replaced *E. klunzingeri* within the discard catch of deeper than 60 m. It was also observed that *Champsodon sp.* and *C. linguatula* constituted 11.26 and 9.13 % of total catch, respectively. In the last five years, three *Champsodon* species were recorded from the Mediterranean Sea (Cicek & Bilecenoglu 2009; Dalyan *et al.*, 2012; Gokoglu & Ozvarol, 2013). Our laboratory studies showed that the obtained *Champsodon* species belonged to *C. capensis* and *C. nudivittis*. However, samples were classified as

Champsodon sp. because it was not possible to identify the differences between species on board. In this way, incorrect values were not employed while calculating abundance and biomass values. They were found in increasing rates during sampling.

The minimum landing sizes of the commercial fishes could be considered inappropriate according to legislative regulations of fisheries management in Turkey (Table 1). The reason is that weak control mechanism of trawl fisheries in the studying area. In the last declaration of minimum landing size of commercial fish species, one Lessepsian fish species, namely *U. moluccensis*, was on the list. But almost 60 % of the commercial fish species biomass consisted of Lessepsians in Iskenderun Bay. Biological research on commercial species should increase not only for the conservation of commercial species but also for taking management measures for the fisheries in the area.

Studies on discard catches are important for the management of fisheries activities, especially for multispecies trawl fisheries. Additional comprehensive studies on fisheries and their problems in the area, which is characterised by a dynamic structure due to the entry of new Red Sea originating fish species.

Acknowledgements

We thank the fishermen of AZAKLAR for their collaboration in this study and Prof. Dr. A. Tursi for giving us a chance to evaluate our data at his laboratory; and Dr. R. Carlucci for his valuable comments. And we also thank Suna TUZUN for editing the English. This work was supported by Istanbul University (Project no: 6685).

References

- Akca, N., 2010. *Underwater observations on the bioecology of Carcharhinus plumbeus (Nardo, 1827) inhabiting Boncuk Bay (Gökova Gulf)*. Master Thesis. Adnan Menderes University, Aydın, Turkey, 63 pp.
- Alsayes, A., Fattouh, S.A.E., 2009. By-Catch and discarding of trawl fisheries at the Mediterranean coast of Egypt. *World Journal of Fish and Marine Sciences*, 1 (3), 199 - 205.
- Alverson, D.L., Freeberg, M.H., Murawski, S.A., Pope, J.G., 1994. *A fisheries technical paper global assessment of fisheries bycatch and discard*. Food and Agriculture Organization of the United Nations, Rome, no 339, 233 pp.
- Anonymous, 2012a. *Fisheries Statistics*. <http://www.tuik.org.tr>, (Accessed 29 May 2013).
- Anonymous, 2012b. *Fisheries regulation laws for Turkish seas Communiqué No: 3/1*. www.tarim.gov.tr (Accessed 29 May 2013).
- Bayhan, K.,Y., Çiçek, E., Unluer, T., Akkaya, M., 2006. Guneydogu Marmara'da Algarna ile Karides Avcılığında Av Kompozisyonu ve Hedef Dışı Av. *Ege Universitesi Su Urunleri Fakultesi Yayınları*, 23 (3-4), 277-283.
- Beisel, N.J., Moreteau, C.J., 1997. A Simple Formula for Calculating the Lower Limit of Shannon's Diversity Index. *Ecological Modelling*, 99, 289-292.
- Bilecenoglu, M., Taşkavak, E., Mater, S., Kaya, M., 2002. *Checklist of the marine fishes of Turkey*, Zootaxa, 113, 1-194.
- Bilecenoglu, M., 2008. *Conservation and monitoring project of sandbar sharks (Carcharhinus plumbeus) in Boncuk Bay, Gökova special environmental protection area*. environmental protection agency for special areas, 32 p.
- Bilecenoglu, M., Russell, B.C., 2008. Record of *Nemipterus randalli* Russell, 1986 (Nemipteridae) from Iskenderun Bay, Turkey. *Cybiurn*, 32 (3), 279-280.
- Bodilis, P., Louisy, P., Draman, M., Arceo, H.O., Francour, P., 2014. Can Citizen Science Survey Non-indigenous Fish Species in the Eastern Mediterranean Sea? *Environmental management*, 53 (1), 172-180.
- Clarke, K.R., Warwick R.M., 2001. *Change in Marine Communities, an Approach to Statistical Analysis and Interpretation. PRIMER-E, II Edition*. Plymouth, UK.
- Çiçek, E., 2006. *Study on the potancially economical important species trawled from Karataş (Adana) Coasts*, PhD. Thesis. Çukurova University, Adana, Turkey, 146 pp.
- Çiçek, E., Bilecenoglu, M., 2009. A new alien fish in the Mediterranean Sea: *Champsodon nudivittis* (Ogilby, 1895) (Champsodontidae). *Acta Ichthyologica et Piscatoria*, 39 (1), 67-69.
- Dalyan, C., Yemiskien, E., Eryilmaz, L., 2012. A new record of gaper (*Champsodon capensis* Regan, 1908) in the Mediterranean Sea. *Journal of Applied Ichthyology*, 28, 5, 834-835.
- Damalas, D., Vassilopoulou, V., 2009. Chondrichthyan by-catch and discards in the bottom trawl fishery of the central Aegean Sea (Eastern Mediterranean) pp. 935-940. In: *9th Symposium on Oceanography and Fisheries*, 2009, Volume II, HCMR, Athens.
- Damalas, D., Maravelias, D.C., Katsanevakis, S., Karageorgis, P. A., Papaconstantinou, C., 2010. Seasonal abundance of non-commercial demersal fish in the Eastern Mediterranean Sea in relation to hydrographic and sediment characteristics, *Estuarine, Coastal and Shelf Science*, 89, 107-118.
- Demirci, A., 2003. *Non target demersal species inhabiting Iskenderun Bay and their biomass estimation*, Msc. Thesis, Mustafa Kemal Universitesi, Antakya, Turkey, 40 pp.
- D'Onghia, G., Carlucci, R., Maiorano, P., Panza, M., 2003. Discards from deep-water bottom trawling in the Eastern-Central Mediterranean Sea and effects of mesh size changes. *Journal of Northwestern Atlantic Fisheries Science*, 31, 245-261.
- Edelist, D., Sonin, O., Golani, D., Rilov, G., Spanier, E., 2011. Spatiotemporal patterns of catch and discards of the Israeli Mediterranean trawl fishery in the early 1990's: Ecological and conservation perspectives. *Scientia Marina*, 75 (4), 641-652.
- Erguden, D., Bayhan, K.,Y., Turan, C., 2012. First record of spotbase burrfish, *Cylichthys spilostylus* (Actinopterygii: Tetraodontiformes: Diodontidae), from the marine waters of Turkey. *Acta Ichthyologica et Piscatoria*, 42 (2), 137-140.
- Gökçe, G., Metin, C., 2006. Balıkçılıkta hedef dışı Av sorunu üzerine bir inceleme. *Ege Universitesi Su Urunleri Dergisi*, 23 (3-4), 457-462.
- Gokoglu, M., Ozvarol, Y., 2013. Additional Records of *Champsodon vorax* and *Champsodon capensis* (Actinopterygii: Perciformes: Champsodontidae) from the Eastern Mediterranean Sea. *Acta Ichthyologica Et Piscatoria*, 43 (1), 79-82.

- Gurbet, R., Akyol, O., Yalçın, E., Özaydın, O., 2013. Discards in bottom trawl fishery in the Aegean Sea (Izmir Bay, Turkey). *Journal of Applied Ichthyology*, 29, 6, 1269-1274.
- Gucu, A.C., Bingel, F., Avsar, D., Uysal, N., 1994. Distribution and Occurrence of Red Sea Fish At The Turkish Mediterranean Coast-Northern Cilician Basin. *Acta Adriatica* 34 (1/2), 103-113.
- Gucu, A.C., Ok, M., Sakinan, 2010, *Sub Regional Technical Meeting On The Leseptian Migration And Its Impact On Eastern Mediterranean Fishery, Past And Present Of Fish Fauna In The NE Levant Sea And Factor Facilitating The Colonization By Leseptian Fishes*. FAO Eastmed Working Document, 7-9 December 2010 Nicosia 88-108.
- Gucu, A. C., 2012, Impact of depth and season on the demersal trawl Discard. *Turkish Journal of Fisheries and Aquatic Sciences*, 12, 817-830.
- Hall, M.A., 1996. *Reviews in fish biology and fisheries*. 6, 319-352.
- Hall, A.M., Alverson, LD., Metuzals, I.K., 2000. By-Catch: problems and solutions. *Marine Pollution Bulletin*, 41, 1, 6, 204-219.
- IUCN, 2013. *The IUCN Red List of Threatened Species*. Version 2013.2. <http://www.iucnredlist.org> Downloaded on 21 November 2013.
- Kelleher, K., 2005, *FAO Fisheries Technical Paper: Discards In The World's Marine Fisheries*, Food and Agriculture Organization Of The United Nations, Rome No 470, 131 pp.
- Kinacıgil, T.H., Çıra, E., İlkayaz, T.A., 1999a. Balıkçılıkta Hedeflenmeden Avlanan Turler Sorunu. *Ege Üniversitesi Su Ürünleri Dergisi*, 16 (3-4), 437-444.
- Kinacıgil, H.T., Çıra, E., İlkayaz, A.T., 1999b. A Preliminary Study on the Shrimp Trawling Bycatch In Tasucu Bay (Northeastern Mediterranean). *Journal of Fisheries and Aquatic Science*, 16 (1-2), 99-105.
- Levesque, R., 2007, *SPSS Programming and Data Management: A Guide for SPSS and SAS Users*, Fourth Edition, SPSS Inc., Chicago III, USA.
- Machias, A., Vassilopoulou, V., Vatsos, D., Bekas, P., Kallianiotis, A., Papaconstantinou, C., Tsimenides, N., 2001, Bottom Trawl Discards in the Northeastern Mediterranean Sea. *Fisheries Research*, 53 (2), 181-195.
- Polat, S., 2010, Abundance and distribution patterns of phytoplankton in the coastal waters of Northeastern Mediterranean, p.27-32, In: *Phytoplankton responses to Mediterranean environmental changes CIESM Workshop*, 7-10 October 2009, Tunis.
- Polat, S., Piner, P., M., 2002. Seasonal variations in biomass, abundance and species diversity of phytoplankton in the Iskenderun Bay (Northeastern Mediterranean). *Pakistan Journal of Botanic*, 34 (2), 101-112.
- Sanchez, P., Demestre, M., Martin, P., 2004. Characterisation of the Discards Generated By Bottom Trawling in the North-western Mediterranean. *Fisheries Research*, 67, 71-80.
- Sanchez, P., Sartor, P., Recasens, L., Ligas, A., Martin, R., Ranieri, D. S., Demestre, M., 2007. Trawl catch composition during different fishing intensity periods in two Mediterranean demersal fishing grounds. *Scientia Marina*. 71 (4), 765-773.
- Soykan, O., Kinacıgil, T.H., Tosunoğlu, Z., 2006. Taşucu Körfezi (Doğu Akdeniz) Karides Trollerinde Hedefdışı Av. *Ege Üniversitesi Su Ürünleri Fakültesi Yayınları*, 23 (1-2), 67-70.
- Tudela, S., 2004, Ecosystem effects of fishing in the Mediterranean: An analysis of the major threats of fishing gear and practices to biodiversity and the marine habitat. *Food and Agriculture Organization of the United Nations*, Rome no 74, 44 pp.
- Yazıcı, F., M., İşmen, A., Altınağaç, U., Ayaz, A., 2006. Marmara Deniz'inde Karides Algarnasının Av Kompozisyonu ve Hedeflenmeyen Av Uzerine Bir Çalışma. *Ege Üniversitesi Su Ürünleri Fakültesi Yayınları*, 23, (3-4), 269-275.