

## Mediterranean Marine Science

Vol 14, No 2 (2013)

Vol 14, No 2 (2013)



### Alien marine species of Libya: first inventory and new records in El-Kouf National Park (Cyrenaica) and the neighbouring areas

H. BAZAIRI, Y.R. SGHAIER, I. BENAMER, H. LANGAR, G. PERGENT, E. BOURAS, M. VERLAQUE, J.B. SOUSSI, A. ZENETOS

doi: [10.12681/mms.555](https://doi.org/10.12681/mms.555)

#### To cite this article:

BAZAIRI, H., SGHAIER, Y., BENAMER, I., LANGAR, H., PERGENT, G., BOURAS, E., VERLAQUE, M., SOUSSI, J., & ZENETOS, A. (2013). Alien marine species of Libya: first inventory and new records in El-Kouf National Park (Cyrenaica) and the neighbouring areas. *Mediterranean Marine Science*, 14(2), 451–462. <https://doi.org/10.12681/mms.555>

## Alien marine species of Libya: first inventory and new records in El-Kouf National Park (Cyrenaica) and the neighbouring areas

H. BAZAIRI<sup>1</sup>, Y. R. SGHAIER<sup>2</sup>, I. BENAMER<sup>3</sup>, H. LANGAR<sup>4</sup>, G. PERGENT<sup>5</sup>, E.M. BOURASS<sup>6</sup>,  
M. VERLAQUE<sup>7</sup>, J. BEN SOUSSI<sup>8</sup> and A. ZENETOS<sup>9</sup>

<sup>1</sup> Faculty of Sciences, University Mohammed V-Agdal, Rabat, Morocco

<sup>2</sup> Regional Activity Centre for Specially Protected Areas (RAC/SPA), Boulevard du Leader Yasser Arafat, BP 337, 1080 Tunis cedex, Tunisia

<sup>3</sup> Faculty of Natural Resources and Environmental Science, University of Omar Mukhtar (OMU), Libya

<sup>4</sup> Faculty of Sciences, University of Tunis El Manar 2092, Tunisia

<sup>5</sup> FRES 3041, EqEL, Faculty of Science, University of Corsica, BP 52, 20250 Corte, France

<sup>6</sup> Environment General Authority Box 13793 Tripoli, Libya

<sup>7</sup> Aix-Marseille University, CNRS, Mediterranean Institute of Oceanography (MIO), Marseille cedex 9, France

<sup>8</sup> Institut National Agronomique de Tunisie, 43 Avenue Charles Nicolle, Tunis, Tunisia

<sup>9</sup> Institute of Marine Biological Resources, Hellenic Centre for Marine Research (HCMR), Anavissos, Attica, Greece

Corresponding author: [hoceinbazairi@yahoo.fr](mailto:hoceinbazairi@yahoo.fr)

Received: 24 July 2013; Accepted: 12 October 2013; Published on line: 31 October 2013

### Abstract

The presence of marine alien species in El-Kouf National Park and the neighbouring areas was assessed using a compilation of available information and observations, a field survey conducted in October 2010 within the framework of the MedMPAnet project and results of further monitoring during June and September 2012. A total of 9 alien species were reported: namely Rhodophyta *Asparagopsis taxiformis* (Delile) Trevisan de Saint-Léon, Chlorophyta *Caulerpa racemosa* var. *cylindracea* (Sonder) Verlaque, Huisman & Boudouresque, the crab *Percnon gibbesi* (H. Milne-Edwards, 1853) and the fishes *Fistularia commersonii* Rüppell, 1838, *Siganus luridus* (Rüppell, 1829), *Siganus rivulatus* Forsskål, 1775, *Pempheris vanicolensis* Cuvier, 1831, *Lagocephalus sceleratus* (Gmelin, 1789) and *Sphyræna flavicauda* Rüppell, 1838. Several of these species were until now unknown for the National Park. The list of alien marine species of Libya is updated and discussed. Until now 63 marine alien species have been recorded along the Libyan coastline. These include 3 Foraminifera, 3 Ochrophyta, 5 Rhodophyta, 5 Chlorophyta, 1 Magnoliophyta, 11 Arthropoda, 13 Mollusca, 1 Echinodermata and 21 Chordata. Among these Non Indigenous Species, 43 are known as established along the Libyan coast including 8 invasive, 11 casual, 5 questionable, 3 cryptogenic and 1 unknown. An in-depth study of marine organisms would substantially increase the number of alien species occurring in Libya. Monitoring of marine assemblages of MPAs provides a valuable opportunity for delving further into the knowledge of native and introduced species.

**Keywords:** Marine aliens, marine protected area, Libya, Mediterranean Sea.

### Introduction

Marine species have been accidentally or intentionally transferred among the world's seas for as long as humans have crossed the oceans for exploration, colonization, commerce and war. However, biological invasions are being nowadays recognized as a major agent of global change following the spectacular increase of invasions by non-native marine and estuarine species in various regions of the world (Occhipinti-Ambrogi, 2007). Furthermore, biological invasions can interact with climate change and other components of global change such as increasing deposition of nitrogen and pollutants, and habitat disturbance by human activities (Vitousek *et al.*, 1997; Dukes & Mooney, 1999; Boudouresque & Verlaque, 2010). Many of the marine alien species exhibit aggressive invasive behaviour and represent significant

ecological pressures on marine and estuarine communities (Ruiz *et al.*, 1999; Streftaris & Zenetos, 2006; Sala *et al.*, 2011). In addition to alterations in ecosystem functioning and biodiversity loss, alien species can also cause serious economic (fisheries, tourism and aquaculture) and health impacts (Pimentel *et al.*, 2000).

The Mediterranean Sea is highly concerned by species introductions. Based on updated checklists (Zenetos *et al.*, 2010, 2011, 2012), a total of 986 alien species were known in the Mediterranean by December 2012: 775 in the Eastern basin, 308 in the Western basin, 249 in the Central basin and 190 in the Adriatic Sea. According to the Mediterranean Action Plan for Invasive Species (UNEP-MAP-RAC/SPA, 2005), the main known pathway/vector of species introduction into the Mediterranean Sea is the Suez Canal followed by shipping (ballast water and sediments, anchoring and fouling), aquaculture

(both marine and brackish species) and trade in live marine species (aquarium activities, fishing baits, seafood). This is in agreement with recent evaluation of pathways at Pan European level (Katsanevakis *et al.*, 2013a). Mediterranean biota are driven towards a ‘tropicalization’ as most of the alien species introduced there are of tropical affinity and origin (Bianchi & Morri, 2003). An up-to-date inventory of the alien species in the Mediterranean Sea, apart from its scientific merits, can fulfil the needs of regulatory requirements such as the EU Biodiversity Strategy to 2020 (EU, 2011) and environmental management options (Katsanevakis *et al.*, 2013b).

Libya is located in the Central Mediterranean Basin as adopted and described under the Marine Strategy Framework Directive (MSFD) (EU, 2008). Data regarding alien species are scarce and fragmented due to the lack of long term monitoring survey undertaken in a framework of national initiatives on alien species. It is worth mentioning the inventory of exotic fish reporting 16 introduced species (Ben Abdallah *et al.*, 2005) updated by Shakman & Kinzelbach (2007c).

This work reports on the alien marine species observed in El-Kouf National Park and the neighbouring areas. An updated list of alien marine species recorded from Libya is presented and discussed.

## Material and Methods

### El-Kouf National Park

El-Kouf National Park (herein: El-Kouf NP) is part of the Libyan coast of Cyrenaica. Located west off the city of El-Beida (N 32.7586, E 21.7484), it occupies an area of 32 000 ha, from the middle of Cyrenaica to Jar-Jar Ummah (Fig. 1). El-Kouf NP was established in 1970 under the supervision of the Arab Centre for the Studies of


Arid Zones and Dry Lands (ACSAD) and various surveys regarding the geology, wildlife and marine environment were conducted until the mid 1970s (Armsby, 1980). The coastline of El-Kouf NP covers ca. 20 km between Alhanya Village in the east (N 32.8412, E 21.5206) and Alauglah Beach in the west (N 32.7712, E 21.3303). In El-Kouf NP, the marine environment differs from west to east. The western part is formed of open and long sandy beaches with sandy bottoms, rocky patches and seagrass meadows of *Posidonia oceanica* (L) Delile. The eastern part is more rugged with few sandy lagoons and beaches and a dominance of cliffs and rocky substrata.

### Data collection

The presence of marine alien species in El-Kouf NP and the neighbouring areas was assessed through field surveys conducted in October 2010 and two additional surveys conducted in June and September 2012 as part of the local monitoring program.

Marine surveys were carried out in the National Park and 9 Km towards the east and 6 Km towards the West. A total of 86 stations were surveyed ranging from 0 to 46 m depth. Investigations were conducted directly from the coast (direct observation mainly in the intertidal area), using a bathyscope, by snorkelling and by scuba diving (Fig. 1). The presence of alien species was recorded. Supplementary data were obtained from professional fishermen. Alien species observed were identified and photographed.

The nomenclature adopted in this paper follows the World Register of Marine Species (WoRMS Editorial Board, 2013) and contributing databases (AlgaeBase, DAISIE, FishBase). The acclimatization (establishment) status of alien species (Established, Casual, Questionable, Cryptogenic or Invasive) in the Central Mediterranean is according to Zenetos *et al.* (2010). The date of



**Fig. 1:** Map showing the localization of the coastline of El-Kouf National Park and the surveyed stations.

first sighting in Libya (or of publication when no date of observation/collection was indicated) is also presented as well as the origin of the species.

Finally, scientific literature, websites and unpublished data were analysed to establish an updated list of the alien marine species reported from Libya. Each record has been carefully analyzed, with recourse to the published literature and, where necessary and possible, interviews with the authors.


## Results and Discussion

### El-Kouf National Park region

Nine alien marine species have been identified in El-Kouf NP and the neighbouring areas; namely, Rhodophy-

ta *Asparagopsis taxiformis* (Delile) Trevisan de Saint-Léon, Chlorophyta *Caulerpa racemosa* var. *cylindracea* (Sonder) Verlaque, Huisman & Boudouresque, the crab *Percnon gibbesi* (H. Milne-Edwards, 1853) and the fishes *Fistularia commersonii* Rüppell, 1838, *Lagocephalus sceleratus* (Gmelin, 1789), *Pempheris vanicolensis* Cuvier, 1831, *Siganus luridus* (Rüppell, 1829), *Siganus rivulatus* Forsskål, 1775, and *Sphyraena flavicauda* Rüppell, 1838 (Table 1; Fig. 2).

*Asparagopsis taxiformis* is an alien red alga exhibiting an invasive behaviour on several coasts of the Mediterranean basin (Flagella *et al.*, 2003). First reported from Libya in 1895 (De Toni, 1895), it was largely observed in El-Kouf NP and the neighbouring areas. Two cryptic taxa coexist under the name of *A. taxiformis* in the Mediterranean Sea:



**Fig. 2:** Alien marine species recorded in El-Kouf National Park and neighbouring areas. A: *Asparagopsis taxiformis*; B: *Caulerpa racemosa* var. *cylindracea*; C: *Percnon gibbesi*; D: *Fistularia commersonii*; E: *Pempheris vanicolensis*; F: *Lagocephalus sceleratus*; G: *Siganus luridus*; H: *Siganus rivulatus*; I: *Sphyraena flavicauda*.

**Table 1.** Alien marine species reported from the El-Kouf National Park (\*) and Libyan waters with the date of first sighting in Libya; world distribution and establishment success in the Central Mediterranean according to Zenetos *et al.* (2010) amended. Abbreviations: Est = established, Cas = Casual, Cry = Cryptogenic, Ques = Questionable, and Unk = Unknown. 1<sup>st</sup> sightings in bold indicate species first reported in the CMED from Libya.

| Species  | Authority                                      | Distribution         | 1 <sup>st</sup> Libyan sighting <sup>1</sup> | Success | Reference  |
|--|--|----------------------|--|---------|--|
| <b>PROTOZOA</b>                                    |  |                      |  |         |  |
| <b>Foraminifera</b>                                |  |                      |  |         |  |
| <i>Amphisorus hemprichii</i>                       | Ehrenberg, 1839                                | Circumtropical       | <b>2008</b>                                  | Unk     | Langer (2008)  |
| <i>Amphistegina lobifera</i>                       | Larsen, 1976                                   | Circumtropical       | <b>1979</b>                                  | Est     | Blanc-Vernet <i>et al.</i> (1979)  |
| <i>Coscinospira hemprichii</i>                     | Ehrenberg, 1839                                | Indo-Pacific         | <b>2008</b>                                  | Cry     | Langer (2008)  |
| <b>MACROPHYTES</b>                                 |  |                      |  |         |  |
| <b>Ochrophyta - Phaeophyceae</b>                   |  |                      |  |         |  |
| <i>Padina boergesenii</i>                          | Allender & Kraft                               | Indo-Pacific         | 1974   | Est     | Nizamuddin (1981, as <i>P. gymnospora</i> );   |
| <i>Padina boryana</i>                              | Thivy  | Indo-Pacific         | <b>1974</b>                                  | Est     | Nizamuddin (1981, as <i>P. tenuis</i> )  |
| <i>Sypododium schimperii</i>                       | (Buchinger ex Kützing) Verlaque & Boudouresque | Indo West Pacific    | <b>1977</b>                                  | Est     | Nizamuddin (1981, as <i>Spatoglossum solieri</i> ); Nizamuddin & Godeh (1989, as <i>Sypododium tubruquense</i> ) |
| <b>Rhodophyta</b>                                  |  |                      |  |         |  |
| <i>Acanthophora nayadiformis</i>                   | (Delile) Papenfuss                             | Indo-Pacific         | <b>1888</b>                                  | Est     | De Toni & Levi (1888 as <i>A. delilei</i> ); Muschler (1910); Nizamuddin <i>et al.</i> (1979)                    |
| * <i>Asparagopsis taxiformis</i>                   | (Delile) Trevisan de Saint-Léon                | Indo-Pacific         | <b>1895</b>                                  | Est     | De Toni (1895); Nizamuddin <i>et al.</i> (1979)  |
| <i>Chondria coeruleascens</i> <sup>2</sup>         | (J. Agardh) Falkenberg                         | East Atlantic        | 1989-1990                                    | Est     | Godeh <i>et al.</i> (1992)   |
| <i>Lophocladia lallemandii</i>                     | (Montagne) F. Schmitz                          | Indo-Pacific         | 1918   | Est     | Petersen (1918); Nizamuddin <i>et al.</i> (1979)   |
| <i>Polysiphonia atlantica</i>                      | Kapraun & J.N. Norris                          | N Atlantic/N Pacific | 1989-1990                                    | CyQues  | Godeh <i>et al.</i> (1992, as <i>P. macrocarpa</i> )   |
| <b>Chlorophyta</b>                                 |  |                      |  |         |  |
| * <i>Caulerpa racemosa</i> var. <i>cylindracea</i> | (Sonder) Verlaque, Huisman & Boudouresque,     | Indo-Pacific         | <b>1990</b>                                  | Inv     | Nizamuddin (1991)  |
| <i>Cladophora herpestica</i>                       | (Montagne) Kützing                             | Red Sea              | <b>1986</b>                                  | Ques    | Nizamuddin (1988, 1991 as <i>Cladophoropsis gerloffii</i> <sup>3</sup> )   |
| <i>Codium fragile</i> subsp. <i>fragile</i>        | (Suringar) Hariot                              | NW Pacific           | 1984   | Est     | Nizamuddin (1991)  |
| <i>Codium Taylorii</i>                             | P.C. Silva                                     | Indo-Pacific         | <b>1977</b>                                  | Est     | Nizamuddin (1991, as <i>C. coralloides</i> )   |
| <i>Ulva fasciata</i>                               | Delile   | Warm-temperate Seas  | 1979   | Est     | Nizamuddin <i>et al.</i> (1979); Godeh <i>et al.</i> (1992)  |
| <b>Magnoliophyta</b>                               |  |                      |  |         |  |
| <i>Halophila stipulacea</i>                        | (Forsskål) Ascherson                           | Red Sea              | <2009  | Est     | UNEP/MAP-RAC/SPA (2009); Sghaier <i>et al.</i> (2011)  |
| <b>CRUSTACEA</b>                                   |  |                      |  |         |  |
| <b>Amphipoda</b>                                   |  |                      |  |         |  |
| <i>Cymadusa filosa</i>                             | Savigny, 1816                                  | Indo-Pacific         | 1939   | Cry     | Maccagno (1939)  |
| <i>Hamimaera hamigera</i>                          | (Haswell, 1879)                                | Indo-Pacific         | <b>1972</b>                                  | Cas     | Ortiz & Petrescu (2007)  |
| <b>Cirripedia</b>                                  |  |                      |  |         |  |
| <i>Tetraclita rufotincta</i>                       | Pilsbry, 1916                                  | Indo-Pacific         | <b>2007</b>                                  | Cas     | Zaouali <i>et al.</i> (2007a)  |
| <b>Copepoda</b>                                    |  |                      |  |         |  |
| <i>Paraeuchaeta concinna</i>                       | (Dana, 1849)                                   | Indo-Pacific         | <b>1990</b>                                  | Cas     | Halim, 1990  |

(continued)

(continued) Table 1.

| Species                            | Authority                   | Distribution   | 1 <sup>st</sup> Libyan sighting' | Success | Reference   |
|------------------------------------|-----------------------------|----------------|----------------------------------|---------|---|
| <b>Decapoda</b>                    |                             |                |                                  |         |   |
| <i>Grapsus granulosus</i>          | H. Milne-Edwards, 1853      | Red Sea        | 2006                             | Cas     | Zaouali et al. (2007b)                                |
| * <i>Percnon gibbesi</i>           | (H. Milne-Edwards, 1853)    | W Atlantic     | 2004                             | Inv     | Elkrwe et al. (2008)                                  |
| <i>Plagusia squamosa</i>           | (Herbst, 1790)              | Indo-Pacific   | 2006                             | Est     | Zaouali et al. (2007b)                                |
| <i>Eucrate crenata</i>             | (De Haan, 1835)             | Indo-Pacific   | 1999                             | Est     | Zgozi et al., 2002                                    |
| <b>Isopoda</b>                     |                             |                |                                  |         |   |
| <i>Apanthura sandalensis</i>       | Stebbing, 1900              | Indo-Pacific   | 1976                             | Est     | Negoescu (1981)                                       |
| <i>Paradella diana</i>             | (Menzies, 1962)             | Indo-Pacific   | 2001                             | Cas     | Zgozi et al. (2002)                                   |
| <b>Stomatopoda</b>                 |                             |                |                                  |         |   |
| <i>Erygosquilla massavensis</i>    | (Kossmann, 1880)            | Red Sea-Indian | 2002                             | Est     | Zgozi et al. (2002)                                   |
| <b>MOLLUSCA</b>                    |                             |                |                                  |         |   |
| <b>Bivalvia</b>                    |                             |                |                                  |         |   |
| <i>Malleus (Mahufundus) regula</i> | (Forsskål in Niebuhr, 1775) | Indo-Pacific   | 2001                             | Est     | Giannuzzi-Savelli et al. (2001)                       |
| <i>Pinctada imbricata radiata</i>  | (Leach, 1814)               | Indo-Pacific   | 1913                             | Inv     | Monterosato (1917)                                    |
| <i>Fuvia fragilis</i>              | (Forsskål in Niebuhr, 1775) | Indo-Pacific   | 1997                             | Inv     | Zgozi et al. (2002)                                   |
| <b>Gastropoda</b>                  |                             |                |                                  |         |   |
| <i>Bursatella leachi</i>           | Blainville, 1817            | Red Sea-Indian | 2000                             | Inv     | Zgozi et al. (2002)                                   |
| <i>Cellana rota</i>                | (Gmelin, 1791)              | Red Sea-Indian | 2007                             | Ques    | Zaouali et al. (2007b)                                |
| <i>Clypeomorus bifasciata</i>      | (G.B. Sowerby II, 1855)     | Indo-Pacific   | 1994                             | Cas     | Giannuzzi-Savelli et al. (1997)                       |
| <i>Conomurex persicus</i>          | (Swainson, 1821)            | Indian Ocean   | 2006                             | Est     | Ben Souissi et al. (2007as <i>Strombus persicus</i> ) |
| <i>Conus fumigatus</i>             | Hwass in Bruguière, 1792    | Red Sea        | 1986                             | Cas     | Röckel (1986)   |
| <i>Erosaria turdus</i>             | (Lamarck, 1810)             | Red Sea-Indian | 2007                             | Est     | Ben Souissi et al. (2007)                             |
| <i>Haliotis pustulata cruenta</i>  | Reeve, 1846                 | Red Sea-Indian | 1994                             | Cas     | Giannuzzi-Savelli et al. 1994                         |
| <i>Monetaria annulus</i>           | (Linnaeus, 1758)            | Indo-Pacific   | 2006                             | Ques    | Ben Souissi et al. (2007, as <i>Cypraea annulus</i> ) |
| <i>Nerita sanguinolenta</i>        | Menke, 1829                 | Red Sea        | 1994                             | Cas     | Giannuzzi-Savelli et al. (1994)                       |
| <b>Polyplacophora</b>              |                             |                |                                  |         |   |
| <i>Acanthopleura gemmata</i>       | (Blainville, 1825)          | Indo-Pacific   | 2006                             | Ques    | Zaouali et al. (2007b)                                |
| <b>ECHINODERMATA</b>               |                             |                |                                  |         |   |
| <i>Ophiocoma scolopendrina</i>     | (Lamarck, 1816)             | Red Sea-Indian | 2007                             | Ques    | Zaouali et al. (2007b)                                |
| <b>FISH</b>                        |                             |                |                                  |         |   |
| <b>Actinopterygii</b>              |                             |                |                                  |         |   |
| <i>Alepes djedaba</i>              | (Forsskål, 1775)            | Indo-Pacific   | 1990                             | Est     | Ben Abdallah et al. (2005)                            |
| <i>Atherinomorus forskalii</i>     | (Rüppell, 1838)             | Red Sea        | 1929                             | Est     | Norman (1929) as <i>Prasenus pinguis</i>              |
| <i>Crenidens crenidens</i>         | (Forsskål, 1775)            | Red Sea-Indian | 1999                             | Est     | Al-Hassan & El-Silimi (1999)                          |
| * <i>Fistularia commersonii</i>    | Rüppell, 1838               | Indo-Pacific   | 2004                             | Inv     | Ben Abdallah et al. (2005)                            |
| <i>Hemiramphus far</i>             | (Forsskål, 1775)            | Indo-Pacific   | 2006                             | Est     | Shakman & Kinzelbach (2006, 2007b,c), Shakman (2008)  |
| <i>Herklotsichthys punctatus</i>   | (Rüppell, 1837)             | Red Sea        | 2005                             | Est     | Shakman & Kinzelbach (2007b,c)                        |
| * <i>Lagocephalus sceleratus</i>   | (Gmelin, 1789)              | Indo-Pacific   | 2006                             | Cas     | Kacem-Snoussi et al. (2009)                           |
| <i>Lagocephalus suezensis</i>      | Clark & Gohar, 1953         | Red Sea-Indian | 2009                             | Cas     | Kacem-Snoussi et al. (2009)                           |

(continued)

(continued) Table 1.

| Species                         | Authority                | Distribution   | 1 <sup>st</sup> Libyan sighting <sup>1</sup> | Success | Reference                                     |
|---------------------------------|--------------------------|----------------|--|---------|---|
| <i>Liza carinata</i>            | (Valenciennes, 1836)     | Red Sea-Indian | 2005   | Est     | Shakman & Kinzelbach (2007b,c)                |
| <i>Parexocoetus mento</i>       | (Valenciennes, 1847)     | Indo-Pacific   | 1966   | Est     | Ben Tuvia (1966)                              |
| * <i>Pempheris vanicolensis</i> | Cuvier, 1831             | Indo-Pacific   | 2004   | Est     | Ben Abdallah <i>et al.</i> (2004)             |
| <i>Sargocentron rubrum</i>      | (Forsskål, 1775)         | Indo-Pacific   | 1968   | Cas     | Štirn (1970)                                  |
| <i>Saurida undosquamis</i>      | (Richardson, 1848)       | Indo-Pacific   | 1982   | Est     | Zupanovic & El-Buni (1982)                    |
| <i>Scomberomorus commerson</i>  | (Lacepède, 1800)         | Indo-Pacific   | 1974   | Est     | Ben Abdallah <i>et al.</i> (2003)             |
| * <i>Siganus luridus</i>        | (Rüppell, 1829)          | Red Sea-Indian | 1968   | Est     | Štirn (1970)                                  |
| * <i>Siganus rivulatus</i>      | (Forsskål, 1775)         | Red Sea        | 1968   | Est     | Štirn (1970)                                  |
| * <i>Sphyræna flavicauda</i>    | Rüppell, 1838            | Indo-Pacific   | 1998   | Est     | Ben Abdallah <i>et al.</i> (2003)             |
| <i>Sphyræna chrysoaenia</i>     | Klunzinger, 1884         | Indo-Pacific   | 1968   | Est     | Štirn (1970); Shakman & Kinzelbach (2007b, c) |
| <i>Stephanolepis diaspros</i>   | Fraser-Brunner, 1940     | Red Sea        | 1965   | Est     | Zupanovic & El-Buni (1982)                    |
| <i>Upeneus moluccensis</i>      | (Bleeker, 1855)          | Indo-Pacific   | 1968   | Est     | Štirn (1970)                                  |
| <i>Upeneus pori</i>             | Ben-Tuvia & Golani, 1989 | Red Sea-Indian | 1994   | Est     | Ben Abdallah <i>et al.</i> (2005)             |

1. When no date of record is mentioned, we report the date of publication.

2. Native and introduced populations probably occur in the Mediterranean Sea.

3. Described from Libya, *C. gerloffii* is very close to *Cladophora herpestica* and might be merely a junior synonym of this species. However, according to Leliaert and Coppejans (2006), *C. gerloffii* would be a junior synonym of *Cladophoropsis membranacea*.

4. *Sphyræna flavicauda* Rüppell, 1838 is considered a synonym of *S. obtusata* by Doiuchi and Nakabo (2005, 2007).

5. *Sphyræna chrysoaenia* is considered a synonym of *S. pinguis* by Doiuchi and Nakabo (2005, 2007).

the alga described in 1813 from Alexandria and confined to the eastern Mediterranean Basin (Egypt, Lebanon, Syria and likely Libya) and a more recently introduced taxon exhibiting an invasive behaviour in the western basin and the Adriatic Sea. Molecular studies showed that the first might be of Atlantic provenance, via the Strait of Gibraltar, whereas the second colonized the Mediterranean from the Indo-Pacific Ocean, probably via the Suez Canal (Lessepsian species) (Ni Chualáin *et al.*, 2004; Andreakis *et al.*, 2007). Identification of *A. taxiformis* spp occurring in El-Kouf NP requires further investigation.


*Caulerpa racemosa* var. *cylindracea* was first observed in the Mediterranean Sea in Libya (Tajora, Tripoli) in 1990 on sandy rocky platforms in association with *Cystoseira* species and *Flabellia petiolata* (Turra) Nizamuddin (Nizamuddin, 1991). Shipping (ballast water) and aquaria trade are considered as possible vectors of introduction from the Indo-Pacific Ocean. Twenty three years after its discovery in Libya, *C. racemosa* var. *cylindracea* is present all around the Mediterranean Basin and in the close Atlantic Ocean (Canary Islands) (Klein & Verlaque, 2008; Marc Verlaque unpublished data). It is probably one of the major invaders ever observed in the Mediterranean Sea. In the El-Kouf NP, *C. racemosa* var. *cylindracea* was observed during a diving session along the coast; it was found in small patches in two areas at a depth of less than three meters and mixed with other native algae species such as *Padina* sp. and *Cystoseira* sp. The two areas were sheltered by the small rocky islands close to the coastline.

*Percnon gibbesi* is one of the most widely distributed Percnidae (Crustacea: Decapoda) in the world, with a range extending from California to Chile, from Florida to Brazil, and from Madeira to the Gulf of Guinea (Manning & Holthuis, 1981). It was first reported in the Mediterranean Sea in 1999 at Linosa Island (Relini *et al.*, 2000). Its presence along the Libyan coast was first recorded at Alhanya in 2004 (Elkrwe *et al.*, 2008) before to be mentioned in many other localities in the Cyrenaica (Katsanevakis *et al.*, 2011).

Since then, its abundance has increased rapidly, and it has been observed at a few other sites. During our survey, a total of 107 individuals of various size classes were recorded in El-Kouf NP. Currently *P. gibbesi* seems well established up to Ras Lateen near the Gulf of Bombah (N 32.6250, E 23.1140) while its western distribution beyond the Cyrenaica remains to be investigated.

The bluespotted cornetfish *Fistularia commersonii* is an Indo-Pacific species that invaded the Mediterranean Sea through the Suez Canal. It was first observed in Israel in 2000 (Golani *et al.*, 2002). Since then, it has been spreading throughout the Mediterranean Sea (Azzurro *et al.*, 2012). Reported in Libya since 2004 (Ben Abdallah *et al.*, 2005), *F. commersonii* was not yet abundant in the El-Kouf NP region but it started to appear in local fishing markets.

Originating in the tropical Indo-Pacific Ocean,



**Fig. 3:** Marine alien species in Libyan waters by taxonomic group.

*Lagocephalus sceleratus* has been listed as one of the 100 “worst invasive species” of the Mediterranean Sea (Streftaris & Zenetos, 2006). Since its first report along the coast of Turkey in 2003 (Akyol *et al.*, 2005), new findings have revealed that this Lessepsian species is now very common along the Levantine coasts of Turkey (Bilecenoglu *et al.*, 2006) and the south Aegean (Corsini *et al.*, 2006; Petrou *et al.*, 2011). In Libya, *L. sceleratus* was first collected in 2006 (Kacem-Snoussi *et al.*, 2009). Fishermen caught the fish in 2010 inside and outside the Ain Al Ghazala Lagoon and off Ain Al Ghazala (Milazzo *et al.*, 2012). *L. sceleratus* was not observed *in situ* in El-Kouf NP but specimens were caught by fishermen working in the area in October 2010.

The sweeper fish *Pempheris vanicolensis* is an Indo-Pacific species that invaded the Mediterranean Sea through the Suez Canal. It was first recorded in Lebanon in 1979 (Mouneimne, 1979). In Libya where it was first reported in 2004 (Ben Abdallah *et al.*, 2004), *P. vanicolensis* is frequently observed in relatively high abundance (Shakman & Kinzelbach, 2007a-c). In the surveyed area, the species formed small schools in cracks and underwater caves. Locally, the species has no economic value.

The dusky spinefoot *S. luridus* and the marbled spinefoot *S. rivulatus* are widely distributed in the Indian Ocean and the Red Sea, respectively. First observed in the Mediterranean Sea in 1931 and 1924, respectively (Ben-Tuvia, 1964), these two Lessepsian species are considered as invasive in the Eastern and Central Mediterranean basins where they have become common commercial fishes (Golani *et al.*, 2002; Ben Abdallah *et al.*, 2003, Streftaris & Zenetos, 2006; Shakman *et al.*, 2008). *Siganus luridus* and *S. rivulatus* were first reported from Libya in 1968 (Štirn, 1970). According to Ben Abdallah *et al.* (2003), *S. luridus* appeared in 1973 in commercial catches and became one of the most abundant species in captures by trammel and gill nets, though its abundance decreases towards the west. Since then *Siganus* spp.

have become very common in Libyan waters probably outcompeting the two native herbivorous species *Sarpa salpa* (Linnaeus, 1758) and *Sparisoma cretense* (Linnaeus, 1758) (Štirn, 1970; Shakman & Kinzelbach 2007b). *Siganus rivulatus* is mostly recorded from the eastern littoral of Libya where it is found in small groups in inshore waters close to sandy bottoms covered by seagrasses and caught in small quantities mixed with *S. luridus* (Ben Abdallah *et al.*, 2003), while *S. luridus* seems to be more present in central and western parts of the littoral (Shakman & Kinzelbach, 2007c). Both species were abundant in El-Kouf NP, especially *S. luridus* - numerous schools were observed in different parts of the surveyed area.

Due to their similar morphology, alien barracuda species often cause confusion. According to the distribution of the two species in the CIESM Atlas of exotic species (Golani *et al.*, 2002), the common species in Libya should be *S. chrysotaenia* collected since 1968 (Štirn 1970); *S. flavicauda* was collected three decades later (Ben Abdallah *et al.*, 2003). To-date, both species are common (in terms of abundance) and largely distributed in Libyan waters (Shakman & Kinzelbach, 2007b, c). The yellowtail barracuda was not observed *in situ* in El-Kouf NP but specimens were caught by fishermen working in the area in June 2012.

### ***Inventory of the alien marine species of Libya***

A comprehensive inventory of marine alien species for Libyan waters, along with their acclimatization status, origin and date of first sighting (or publication when no date of record is mentioned) was compiled.

Until now 63 marine aliens species have been recorded along the Libyan coastline. These include 3 Foraminifera, 14 macrophytes, 11 Crustacea, 13 Mollusca, 1 Echinodermata and 21 fishes (Actinopterygii) (Table 1, Fig. 3). Among these Non Indigenous Species, 43 are known as established including 8 invasive, 11 casual, 5 questionable, 3 cryptogenic and 1 unknown (Table 1).



Fishes rank first probably because they are the most studied taxon among alien marine organisms, and they highly contribute to local fishery activities (Shakman & Kinzelbach, 2007b). Besides the 16 species recorded by Ben Abdallah *et al.* (2005), 4 more species were added to the list at a later date: *Hemiraphus far* (Shakman & Kinzelbach, 2006); *Liza carinata* (Shakmam & Kinzelbach, 2007b, c), *L. sceleratus* and *L. suezensis* (Kacem-Snoussi *et al.*, 2009; Ben Abdallah *et al.*, 2011). To these we should add *Atherinomorus forskalii* reported by Norman (1929) as *Prasenus pinguis* (Lacépède, 1803). Of the 21 alien fishes, that all are Actinopterygii, 18 have established populations while 3 species (*Sargocentron rubrum*, *Lagocephalus sceleratus* and *L. suezensis*) are only known from one or two records. However this could be the result of the limited number of studies. For example, *L. sceleratus* is established in the Central Mediterranean and is expanding its geographic range to the Ionian Sea (Zenetos *et al.*, 2013). Other species such as *Cephalopholis taeniops* (Valenciennes, 1828), *Seriola rivoliana* Valenciennes, 1833, *Sphoeroides pachygaster* (Müller & Troschel, 1848) and *Synaptura lusitanica* (Capello, 1868) were not considered in this study as they are of Atlantic origin and it has been argued that their presence in the Mediterranean is due to range expansion. The list of newly introduced fish species in Libya is non-exhaustive. The latest record is that of the Bermuda sea chub *Kyphosus sectatrix* (Linnaeus, 1758) (Elbarassi *et al.*, 2013). Zupanovic & El-Bini (1982) have reported *Pomadasy olivaceum* Day, 1975 based on a single specimen collected off Zwara, Tripolitanian coast. This record has been listed in the DAISIE inventory as *Pomadasy stridens* (Forsskål, 1775). In the absence of voucher specimens and based on the opinion of local experts who report only its congeneric *Pomadasy Incicus*, a native Mediterranean species, the record of *P. stridens* is herein excluded.

Macrophytes rank second on the list. The analysis of literature (Nizamuddin *et al.*, 1979, 1994; Nizamuddin, 1981, 1985, 1991, 1995; Shameel, 1983; Zarmouth & Nizamuddin, 1991; Godeh *et al.*, 1992, 2009; Ribera *et al.*, 1992; Gallardo *et al.*, 1993; Nizamuddin & El Menifi Fathalla, 1993; Babbini & Bressan, 1997; Gómez Garreta *et al.*, 2001; Taskin *et al.*, 2012) gave a list of 240 native marine taxa of macrophytes. Four species not corresponding to valid taxa have been excluded, *i.e.* *Chaetomorpha urbica* (Zanardini) Kützing, *Cladophora catenata* (J. Agardh) Hauck, *Cladophoropsis puillis* (Kütz.) Boergesen and *Gracilaria verrucosa* var. *ramulosa* (C. Agardh) M.P. Reis, as well as the nine following alien taxa reported by Nizamuddin (1991) and Godeh *et al.* (1992, 2009), which we considered as misidentifications [*Bryopsis caespitosa* Suhr ex Kützing (as *Bryopsis caespitosa* Suhr in Kützing), *Bryopsis pennata* var. *leprieurii* (Kützing) Collins & Hervey (as *Bryopsis leprieurii* Kützing), *Chaetomorpha indica* (Kützing) Kützing, *Chaetomorpha melagonium* (F.Weber & Mohr) Kützing, *Chon-*

*dria oppositoclada* E.Y.Dawson, *Dasya villosa* Harvey, *Gastroclonium ovatum* (Hudson) Papenfuss, *Laurencia laxa* (R.Brown ex Turner) Gaillon (as *Laurencia laxa* Kützing) and *Ulva taeniata* (Setchell) Setchell & N.L. Gardner]. In this study, we have listed a total of 14 alien macrophytes, which represent 5.5 % of the Libyan marine macroflora. All these alien species are established except for *Cladophora herpestica* and *Polysiphonia atlantica* that are questionable and cryptogenic/questionable, respectively. This group includes one of the most invasive alien species in the Mediterranean, namely *Caulerpa racemosa* var. *cylindracea*.

Molluscs are known to be one of the dominant alien groups in the Mediterranean (Zenetos *et al.* 2012). Here, they rank third on the list, with only six established species, including two invasive bivalves (*Pinctada imbricata radiata* and *Fulvia fragilis*) and one invasive gastropod (*Bursatella leachii*), the remaining ones being casual or questionable. The presence of the limpet *Cellana rota* and the cowry *Monetaria annulus* reported by Ben Souissi *et al.* (2007) is questionable because *C. rota* can be easily confused with the native limpets, and the majority of old records of *M. annulus* were based on transported shells (the species was one of the molluscan shells most often carried around, as currency or ornaments) (Gofas & Zenetos, 2003).

Crustaceans with 9 alien species rank fourth. Four species are established and the other ones are casual or questionable. This group includes one of the most invasive alien species in the Mediterranean, namely the decapod *Percnon gibbesi*.

Finally, Foraminifera and Echinodermata included 3 and 1 alien species and occupy the fifth and the sixth places, respectively. The absence of taxonomic groups such as the Annelids, one of the major phyla ranking third in terms of contributors to Mediterranean alien biota [with 132 species], is indicative of the lack of taxonomic experts in the area. It is also worth noting the absence or rarity of zooplanktonic species such as Cnidaria and Copepoda.

Up to December 2012, a total of 249 alien species were known from the Central Mediterranean (Zenetos *et al.*, 2012). The majority of alien species in the Libyan waters are of Indo-Pacific origin (53%), followed by Red Sea – Indian Ocean species (16%), Red Sea species (14%), Circumtropical (3%). Less than 5% of the alien introduced species in Libya originate in the Atlantic as opposed to the Atlantic species expanding their distribution in the Mediterranean (reaching Libya) without human intervention as a result of climate change. Libya, and globally the strait of Sicily, has a special geographical position in the Mediterranean. For approximately 44 of the 63 alien species listed here, the first Libyan record was also the first for the central Mediterranean. This highlights the importance of the region for monitoring introductions in the Mediterranean. This sector is regarded as an area of biogeographical transition (Bianchi, 2007) and it is located at the crossroads between different Mediterranean basins and hydrographic conditions. In this

area, significant changes have occurred in the biological communities and many alien species have appeared in the last few years. Lessepsian migration through the Suez Canal probably has a conspicuous influence on Libyan marine biodiversity, but Libya also seems to harbour alien species arriving from the West (Atlantic). In both cases, a warm temperate or subtropical affinity of most species is evident.

Compared to other Mediterranean regions, the number of alien species recorded in Libya is low: 63 species versus 107 in Tunisia and 232 in Egypt (Katsanevakis *et al.*, 2012; UNEP-MAP-RAC/SPA, 2013). However, this number is certainly underestimated for various reasons. The scientific effort of investigation, including taxonomic expertise availability is limited; small-sized Lessepsian immigrants (macrophytes and invertebrates) have been poorly studied and many species probably remain undiscovered. Moreover, the deep benthic assemblages were not investigated. As an example, the divergence between Tunisia and Libya in numbers is more obvious in Decapoda (17 in Tunisia vs. 4 in Libya), Bivalvia (7 in Tunisia vs. 3 in Libya), Gastropoda (15 in Tunisia vs 9 in Libya) (UNEP-MAP-RAC/SPA, 2013). The difference is also due to the lack of data for several groups in Libya such as Phaeophyceae, Polychaeta, Copepoda, Isopoda, Cephalopoda, Bryozoa, Cnidaria, Hydrozoa, Ascidiacea, Echinodermata, Platyhelminthes and Elasmobranchii.

## Conclusion

Marine protected areas in the Mediterranean have not escaped biological invasions and most of them have been affected by introduced invasive alien species that threaten marine biodiversity (Otero *et al.*, 2013). In El-Kouf National Park, among the nine alien species recorded, *C. racemosa* var. *cylindracea*, *Siganus* spp. and *Percnon gibbesi* are classified among the major invaders of the Mediterranean. However, no data are available on the environmental impact of these species on the El-Kouf NP.

An in-depth study of marine organisms would substantially increase the number of alien species occurring in Libya, especially the proportion of Indo-Pacific species arriving via the Suez Canal. Monitoring marine assemblages of MPAs provides a valuable opportunity for further in-depth knowledge of native and introduced species.

## Acknowledgements

The surveys in the north-eastern part of Libya (the Cyrenaican coast) were conducted within the framework of the MedMPAnet Project implemented by UNEP/MAP-RAC/SPA and funded by RAC/SPA, the European Commission (EC), the Spanish Agency for International Cooperation to Development (AECID), and the French Global Environment Facility (FFEM).

Argyro Zenetos was supported from the Euro-

pean Community's Seventh Framework Programme ([FP7/2007-2013]) under grant agreement n° 287600 - PERSEUS project (Policy-oriented marine Environmental Research for the Southern European Seas).

The MedMPAnet Project activities implemented in Libya were carried out thanks to the valuable support of the Environment General Authority (EGA). We thank Mr. El Maki Ayad Elagil and the EGA staff for their valuable help during the field studies in the El-Kouf NP region.

The authors are very grateful to four anonymous reviewers for their kind and helpful contributions.

## References

- Akyol, O., Ünal, V., Ceyhan, T., Bilecenoglu, M., 2005. First confirmed record of *Lagocephalus sceleratus* (Gmelin, 1789) in the Mediterranean Sea. *Journal of Fish Biology*, 66, 1183-1186.
- Al-Hassan, L.A.J., El-Silini, O.A., 1999. Check-list of bony fishes collected from the Mediterranean coast of Benghazi, Libya. *Revista de Biología Marina & Oceanografía*, 34 (2), 291-301.
- Andreakis, N., Procaccini, G., Maggs, C., Kooistra, W.H.C.F., 2007. Phylogeography of the invasive seaweed *Asparagopsis* (Bonnemaisoniales, Rhodophyta) reveals cryptic diversity. *Molecular Ecology*, 16, 2285-2299.
- Armsby, J.K., 1980. *Kouf National Park marine survey, final report: April-July 1980*, ACSAD. (published in Arabic).
- Azzurro, E., Soto, S., Garofalo, G., Maynou, F., 2012. *Fistularia commersonii* in the Mediterranean Sea: invasion history and distribution modeling based on presence-only records. *Biological Invasions*, 15 (5), 977-990.
- Babbini, L., Bressan, G., 1997. Recensement de Corallinacées de la Mer Méditerranée et considérations phytogéographiques. *Bibliotheca Phycologica*, 103, 1-421.
- Ben Abdallah, A.R., Al-Turky A., Fituri, A.A., 2003. Commercially exploited exotic fishes in Libya. *Bulletin de l'Institut National des Sciences et Technologies de la Mer (Salammbô) n°spécial*, (8), Actes des 6<sup>èmes</sup> Journées de l'ATSMer, Tunis (Tunisie) 28-30 Novembre 2003, 53-55.
- Ben Abdallah, A.R., Al-Turky A., Fituri, A.A., 2004. New records of invasive fishes in the western Libyan coast. *Bulletin de l'Institut National des Sciences et Technologies de la Mer (Salammbô) n°spécial*, (9), Actes des 7<sup>èmes</sup> Journées de l'ATSMer, Zarzis (Tunisie) 18-22 décembre 2004, 117-118.
- Ben Abdallah, A.R., Al-Turky A., Fituri, A.A., 2005. Exotic fishes along the coast of Libya (Arabic with English abstract) *Libyan Journal of Marine Science*, (10), 48-56.
- Ben Abdallah, A., Ben Souissi J., Méjri H., Capapé C., Golani D., 2007. First record of *Cephalopholis Taeniops* (Valenciennes) in the Mediterranean Sea. *Journal of Fish Biology* (71), 610-614.
- Ben Abdallah, A.R., Al-Turky, A., Nafti, A., Shakman, E., 2011. A new record of a Lessepsian fish, *Lagocephalus suzezensis* (Actinopterygii: Tetraodontiformes: Tetraodontidae), in the South Mediterranean (Libyan coast). *Acta Ichthyologica & Piscatoria*, 41 (1), 71-72.
- Ben Souissi, J., Zaouali, J., Ben-Abdallah, A., 2007. Sur la présence de nouvelles espèces exotiques dans la Grande et la Petite Syrte. *Rapport de la Commission internationale pour*

- l'Exploration Scientifique de la Mer Méditerranée*, 38, 435.
- Ben-Tuvia, A., 1964. Two siganid fishes of Red Sea origin in the Eastern Mediterranean. *Bulletin of Sea Fishery Research of Haifa Station*, 37, 3-9.
- Ben-Tuvia, A. 1966. Red Sea fishes recently found in the Mediterranean. *Copeia*, 2, 254-275.
- Bianchi, C.N., 2007. Biodiversity issues for the forthcoming tropical Mediterranean Sea. *Hydrobiologia*, 580, 7-21.
- Bianchi, C.N., Morri, C., 2003. Global sea warming and "tropicalization" of the Mediterranean Sea: biogeographic and ecological aspects. *Biogeographia*, 24, 319-327.
- Bilecenoglu, M., Kaya, M., Akalin, S., 2006. Range expansion of silverstripe blasp, *Lagocephalus sceleratus* (Gmelin, 1789) to the northern Aegean Sea. *Aquatic Invasions*, 1(4), 289-291.
- Blanc-Vernet, L., Clairefond, P., Orsolini, P., 1979. Les Foraminifères. *Annales de l'Université de Provence*, 6, 171-209.
- Boudouresque, C.F., Verlaque, M., 2010. Is global warming involved in the success of seaweed introductions in the Mediterranean Sea? In: *Seaweeds and their role in globally changing environments*. Israel, A., Einav, R. & Seckbach, J. (Eds), pp. 31-50, Springer publ., Dordrecht.
- Corsini, M., Margies, P., Kondilatos, G., Economidis, P.S., 2006. Three new exotic fish records from the SE Aegean Greek waters. *Scientia Marina*, 70, 319-323.
- DAISIE: Delivering Alien Species In Europe: <http://www.europe-aliens.org>
- De Toni, G.B., 1895. Phyceae japonicae novae addita enumeratione algarum in ditone maritima Japoniae hucusque collectarum. *Memorie del Reale Istituto Veneto di Scienze, Lettere ed Arti*, 25 (5), 78, II plates.
- De Toni, G.B., Levi, D., 1888. Pugillo di alghe tripolitane. *Rendiconti della Reale Accademia Nazionale dei Lincei*, Roma, Serie 4, 4, 240-250.
- Dukes, J.S., Mooney, H., 1999. Does global change increase the success of biological invaders? *Trends in Ecology and Evolution*, 14, 135-139.
- Elbaraasi, H., Bograra, O., Elsilini, O., Bojwari, J., 2013. First record of the Bermuda sea chub, *Kyphosus saltatrix* (Actinopterygii: Perciformes: Kyphosidae), in the coastal waters of Libya. *Acta ichthyologica & Piscatoria*,
- Elkrwe, H.M., Elhawaj, H., Galil, B., Ben Abdallah, A., 2008. The first record of *Percnon gibbesi* (H. Milne-Edwards, 1853) (Crustacea: Decapoda: Plagusidae) from the southern rim of the Mediterranean. *Aquatic Invasions*, 3, 243-245.
- EU, 2008. Directive of the European Parliament and the Council Establishing a Framework for Community Action in the Field of Marine Environmental Policy (Marine Strategy Framework Directive). European Commission. Directive 2008/56/EC, OJ L 164.
- EU, 2011. Our Life Insurance, Our Natural Capital: an EU Biodiversity Strategy to 2020. European Commission. COM/2011/244.
- Flagella, M.M., Guala, I., Lorenti, M., Buia, M.C., 2003. *Asparagopsis taxiformis* and *Caulerpa racemosa* interaction with algal community and ecophysiological traits. In: *Proceedings of the third European Phycological Congress*, Belfast (UK), 21-26 July 2003, p. 37.
- Gallardo, T., Gómez Garreta, A., Ribera, M.A., Cormaci, M., Furnari, G. *et al.*, 1993. Check-list of Mediterranean Seaweeds, II. Chlorophyceae Wille S.I. *Botanica Marina*, 36, 399-421.
- Giannuzzi-Savelli, R., Pusateri, F., Palmeri A., Ebreo, C., 1994. Atlante delle Conchiglie marine del mediterraneo (1). *La Conchiglia*, Roma.
- Giannuzzi-Savelli, R., Pusateri, F., Palmeri A., Ebreo, C., 1997. Atlante delle Conchiglie marine del mediterraneo. (2) Caenogastropoda. *La Conchiglia*, Roma, 258p.
- Giannuzzi-Savelli, R., Pusateri, F., Palmeri, A., Ebreo, C., Coppini, M. *et al.*, 2001. *Atlante delle conchiglie marine del Mediterraneo. Vol. 7: Bivalvia Protobranchia- Pteromorphia*. Evolver, Roma, 246 p.
- Godeh, M.M., Nizamuddin, M., Elmenifi F.A., 1992. Marine algae from eastern coast of Libya (Cyrenaica). *Pakistan Journal of Botany*, 24 (1), 11-21.
- Godeh, M.M., El-Menifi, F.O., Said A.A., 2009. Marine algae of Tobruk and Ain Ghazal coasts, Libya. *Journal of Science and its Applications*, Garyounis University Press, 3, 42-55.
- Gofas, S., Zenetos, A., 2003. Exotic molluscs in the Mediterranean basin: current status and perspectives. *Oceanography & Marine Biology: an annual review*, 41, 237-277.
- Golani, D., Orsi-Relini, L., Massuti, E., Quignard, J.P., 2002. Vol. 1. Fishes. In: *CIESM Atlas of Exotic Species in the Mediterranean*. Briand, F. (Ed.), CIESM Publishers, Monaco, 256 pp.
- Gómez Garreta, A., Gallardo, T., Ribera, M.A., Cormaci, M., Furnari, G. *et al.*, 2001. Checklist of the Mediterranean seaweeds. III. Rhodophyceae Rabenh. 1. Ceramiales Oltm. *Botanica Marina*, 44, 425-460.
- Halim, Y., 1990. On the potential migration of Indo-Pacific plankton through the Suez Canal *Bulletin de l'Institut Océanographique de Monaco*, n° special (7), 11-27.
- Kacem-Snoussi, A., Ben Abdallah, R. A., El Turkey, A. A., Ben Moussa, N. M., 2009. Guide to bony fishes in Libyan waters (Arabic) (ed) Marine Research Center 237p.
- Katsanevakis, S., Poursanidis, D., Yokes, B., Mačić, V., Beqiraj, S. *et al.*, 2011. Twelve years after the first report of the crab *Percnon gibbesi* (H. Milne-Edwards, 1853) in the Mediterranean: current distribution and invasion rates. *Journal of Biological Research-Thessaloniki*, 16, 224-236.
- Katsanevakis, S., Bogucarskis, K., Gatto, F., Vandekerkhove, J., Deriu, I. *et al.*, 2012. Building the European Alien Species Information Network (EASIN): a novel approach for the exploration of distributed alien species data. *BioInvasions Records*, 1, 235-245.
- Katsanevakis, S., Zenetos, A., Belchior, C., Cardoso, A.C., 2013a. Invading European Seas: assessing pathways of introduction of marine aliens. *Ocean & Coastal Management*, 76, 64-74.
- Katsanevakis, S., Gatto, F., Zenetos, A., Cardoso, A.C., 2013b. How many marine aliens in Europe? *Management of Biological Invasions*, 4, 1, 37-42.
- Klein, J., Verlaque, M., 2008. The *Caulerpa racemosa* invasion: a critical review. *Marine Pollution Bulletin*, 56, 205-225.
- Langer, M.R., 2008. Foraminifera from the Mediterranean and the Red Sea. Aqaba-Eilat, the improbable Gulf. In: *Environment, Biodiversity and Preservation*. Por, F.D.), pp. 399-417, Magnes Press.
- Leliaert, F., Coppejans, E., 2006. A revision of *Cladophoropsis* Børgesen (Siphonocladales, Chlorophyta). *Phycologia*, 45, 657-679.
- Maccagno, T.P., 1939. Contributo alla conoscenza dei crostacei della Tripolitania. *Bollettino dei Musei di Zoologia e di Anatomia Comparata della Reale Università di Torino*, 47, 437-452.
- Manning, R.B., Holthuis, L.B., 1981. West African brachyuran

- crabs. *Smithsonian Contributions to Zoology*, 306: xii, 1-379.
- Milazzo, M., Azzurro, E., Badalamenti, F., 2012. On the occurrence of the silverstripe blaasop *Lagocephalus sceleratus* (Gmelin, 1789) along the Libyan coast. *BioInvasions Records*, 1 (2), 135-137.
- Monterosato T. A. (di), 1917. Molluschi viventi e quaternari raccolti lungo le coste della Tripolitania dall'Ing. Camillo Crema. *Bollettino Zoologico Italiano Ser. 3, 4*: 1-28, pl. 1 (fig. 1-35).
- Mouneimne, N., 1979. Poissons nouveaux pour les côtes libanaises. *Cybium* (6), 105-110.
- Muschler, R. 1910. Algae. In: *Catalogue Raisonné des Plantes de Tripolitaine. Florae Libycae Prodromus*. Durand, E. & Barratte, J.F.G. (Eds), pp. 293-313, Geneve.
- Negoescu, I., 1981. Contribution à l'étude des Isopodes Anthuridés (Isopoda, Anthuridae) de la Mer Méditerranée. *Rapports de la Commission Internationale pour l'Exploration Scientifique de la Mer Méditerranée*, 27 (2), 219-220.
- Ni Chualáin, F., Maggs, C.A., Saunders, G.W., Guiry, M.D., 2004. The invasive genus *Asparagopsis* (Bonnemaisoniaceae, Rhodophyta): Molecular systematics, morphology, and ecophysiology of *Falkenbergia* isolates. *Journal of Phycology*, 40, 1112-1126.
- Nizamuddin, M., 1981. Contribution to the marine algae of Libya. Dictyotales. *Bibliotheca Phycologica*, 54, 1-122, XXXIX plates, 1 map.
- Nizamuddin, M., 1985. A new species of *Cystoseira* C. Ag. (Phaeophyta) from the Eastern part of Libya. *Nova Hedwigia*, 42, 119-122.
- Nizamuddin, M., 1988. *Cladophoropsis gerloffii*, a new siphonocladaceous alga from the coast of Libya. *Willdenowia*, 17, 229-234.
- Nizamuddin, M., 1991. *The green marine algae of Libya*. Bern, Elga Publishers, 230 pp.
- Nizamuddin, M., 1995. A caespitose-tophulose *Cystoseira* species from Tripoli, Libya. *Pakistan Journal of Botany*, 27, 49-54
- Nizamuddin, M., El Menifi Fathalla, A., 1993. A new species of the genus *Codium* (Chlorophyta-Codiales) from the eastern coast of Libya. *Pakistan Journal of Botany*, 25, 208-214.
- Nizamuddin, M., Godeh, M., 1989. *Stypopodium tubruquense* (Phaeophyta, Dictyotales) a new species from the Mediterranean Sea. *Willdenowia*, 18, 603-608.
- Nizamuddin, M., West, J.A., Meñez, E.G., 1979. A list of marine algae from Libya. *Botanica Marina*, 24, 465-476.
- Nizamuddin, M., El-Menifi, F., Godeh, M., 1994. Re-establishment of the genus *Poropsis* Kützing. *Pakistan Journal of Botany*, 26, 215-219.
- Norman, J.R., 1929. Notes on fishes of the Suez Canal. *Proceedings of the Zoological Society* (2), 616 p, London.
- Occhipinti-Ambrogi, A., 2007. Global change and marine communities: Alien species and climate change. *Marine Pollution Bulletin*, 55, 342-352.
- Ortiz, M., Petrescu, I., 2007. The marine Amphipoda (Crustacea: Gammaridea) of the Republic of Libya, SE Mediterranean. *Travaux du Muséum National d'Histoire Naturelle "Grigore Antipa"*, 1, 11-23.
- Otero, M., Cebrian E., Francour, P., Galil, B., Savini, D., 2013. *Monitoring Marine Invasive Species in Mediterranean Marine Protected Areas (MPAs): A strategy and practical guide for managers*. Malaga, Spain: IUCN. 136 pages.
- Petersen, H.E., 1918. Algae (excl. calcareous algae). In: *Report on the Danish Oceanographical Expeditions 1908-1910 to the Mediterranean and adjacent seas. Vol. II. Biology*. Schmidt, J. (Ed.) Vol. [Art.] K.3, 20 pp. København.
- Petrou, A., Isaias, E., Chrysanthou, K., Scarcella, G., 2011. Fishing yields of *Lagocephalus sceleratus* in Cyprus. *Proceedings of the Tenth International Conference on the Mediterranean Coastal Environment, MEDCOAST 11*, E. Özhan (Ed.), 25-29 October 2011, Rhodes, Greece, 445-452.
- Pimentel, D., Lach, L., Zuniga, R., Morrison, D., 2000. Environmental and economic costs of non-indigenous species in the United States. *BioScience*, 50, 53-65.
- Relini, M., Orsi-Relini, L., Puccio, V., Azzurro, E., 2000. The exotic crab *Percnon gibbesi* (H. Milne Edwards, 1853) (Decapoda, Grapsidae) in the Central Mediterranean. *Scientia Marina*, 64, 337-340.
- Ribera, M.A., Gómez-Garreta, A., Gallardo, T., Cormaci, M., Furnari, G. et al., 1992. Check-list of Mediterranean Seaweeds. I. Fucophyceae (Warming 1884). *Botanica Marina*, 35, 109-130.
- Röckel, D., 1986. Sensational find in the Mediterranean. *La Conchiglia*, 18 (210-211), 12.
- Ruiz, G.M., Fofonoff, P.W., Hines, A.H., Grosholz, E.D., 1999. Non-indigenous species as stressors in estuarine and marine communities: assessing invasion impacts and interactions. *Limnology and Oceanography*, 44, 950-972.
- Sala, E., Kizilkaya, Z., Yildirim, D., Ballesteros, E., 2011. Alien marine fishes deplete algal biomass in the Eastern Mediterranean. *PLoS ONE* 6 (2), e17356. doi:10.1371/journal.pone.0017356.
- Sghaier, Y.R., Zakhama-Sraieb, R., Benamer, I., Charfi-Cheikhrouha, F., 2011. Occurrence of the seagrass *Halophila stipulacea* (Hydrocharitaceae) in the southern Mediterranean Sea. *Botanica Marina*, 54, 575-582.
- Shakman, E., 2008. *Lessepsian migrant fish species of the coastal waters of Libya: Status, biology, ecology*. PhD Thesis, Universität Rostock, 125s Rostock.
- Shakman, E., Kinzelbach, R., 2006. The Halfbeak fish *Hemiramphus far* (Forsskål, 1775) in the coastal waters of Libya. *Zoology in the Middle East (Heidelberg)*, 39, 111-112.
- Shakman, E., Kinzelbach, R., 2007a. First record of the Sweeper fish *Pempheris vanicolensis* Cuvier, 1821 on the eastern Libyan coast. *Rostocker Meeresbiologische Beiträge*, 18, S. 79-81.
- Shakman, E., Kinzelbach, R., 2007b. Commercial fishery and fish species composition in coastal waters of Libya. *Rostocker Meeresbiologische Beiträge*, 18, S. 63-78.
- Shakman, E., Kinzelbach, R., 2007c. Distribution and characterization of Lessepsian migrant fishes along the coast of Libya. *Acta Ichthyologica et Piscatoria*, 37 (1), 7-15.
- Shakman, E., Winkler, H.M., Oeberst, R., Kinzelbach, R., 2008. Morphometry, age and growth of *Siganus luridus* Rüppell, 1828 and *Siganus rivulatus* Forsskål, 1775 (Siganidae) in the central Mediterranean (Libyan coast). *Revista de Biologia Marina & Oceanografia*, 43 (3), 521-529.
- Shameel, M., 1983. Notes on the seaweeds of Tripoli, Libya. *Pakistan Journal of Botany*, 15, 79-83.
- Štirn, J., 1970. Some notes on western trends of Lessepsian migration. In: *Journées Ichthyologiques*, Rome, 30 Nov- 1 Dec., pp. 187-190, CIESM, Monaco.
- Streftaris, N., Zenetos, A., 2006. Alien marine species in the Mediterranean - the 100 'worst invasives' and their impact. *Mediterranean Marine Science*, 7 (1), 87-118.
- Taskin, E., Jahn, R., Öztürk, M., Furnari, G., Cormaci, M.,

2012. *The Mediterranean Cystoseira* (with photographs). pp. [2] 1-75. Manisa, Turkey: Celal Bayar University.
- UNEP-MAP-RAC/SPA. 2005. Action Plan concerning species introductions and invasive species in the Mediterranean Sea. RAC/SPA (Ed.), Tunis, 30 pp.
- UNEP/MAP-RAC/SPA. 2009. *Report of the MedPosidonia Project-Libya (Project to develop the inventorying, mapping and monitoring of Posidonia meadows in four Mediterranean countries: Algeria, Libya, Tunisia and Turkey)*. RAC/SPA edition, Tunis. 29 pp.
- UNEP-MAP-RAC/SPA, 2013. Marine Mediterranean Invasive Alien Species (MAMIAS).
- Vitousek, P.M., D'Antonio, C.M., Loope, L.L., Rejm Nek, M., Westbrook, S.R., 1997. Introduced species: a significant component of human-caused global change. *New Zealand Journal of Ecology*, 21 (1), 1-16.
- WoRMS Editorial Board, 2013. World Register of Marine Species. Available from <http://www.marinespecies.org> at VLIZ (Accessed May 2013)
- Zaouali, J., Ben Souissi, J., Galil, B.S., D'Udekem D'Acoz, C., Ben Abdellah, A., 2007a. Grapsoid crabs (Crustacea: Decapoda: Brachyura) new to the Sirte Basin, southern Mediterranean Sea - the roles of vessel traffic and climate change. *JMBA2, Biodiversity Records*, 5770.
- Zaouali, J., Ben Souissi, J., Galil, B.S., D'Udekem D'Acoz, C., Ben Abdellah, A., 2007b. Contribution à l'étude des peuplements actuels des substrats solides de l'étage médiolittoral de la Méditerranée méridionale. *Rapports de la Commission Internationale pour l'Exploration Scientifique de la Mer Méditerranée*, 38, 639.
- Zarmouth, M.M., Nizamuddin, M., 1991. *Sargassum rametaceum* (Phaeophyta, Fucales), a new species from eastern coast of Libya. *Willdenowia*, 21, 269-273.
- Zenetos, A., Gofas, S., Verlaque, M., Çinar, M.E., Garcia Raso, J.E. *et al.*, 2010. Alien species in the Mediterranean Sea by 2010. A contribution to the application of European Union's Marine Strategy Framework Directive (MSFD). Part I. Spatial distribution. *Mediterranean Marine Science*, 11 (2), 318-493.
- Zenetos, A., Gofas, S., Verlaque, M., Çinar, M.E., Garcia Raso, J.E. *et al.*, 2011. Errata to the review article (Medit. Mar. Sci., 11 (2): 318-493). Alien species in the Mediterranean Sea by 2010. A contribution to the application of European Union's Marine Strategy Framework Directive (MSFD). Part I. Spatial distribution. *Mediterranean Marine Science*, 12 (2), 509-514.
- Zenetos, A., Gofas, S., Morri, C., Rosso, A., Violanti, D., *et al.*, 2012. Alien species in the Mediterranean Sea by 2012. A contribution to the application of European Union's Marine Strategy Framework Directive (MSFD). Part 2. Introduction trends and pathways. *Mediterranean Marine Science*, 13 (2), 328-352.
- Zenetos, A., Koutsogiannopoulos, D., Ovalis, P., Poursanidis, D., 2013. The role played by citizen scientists in monitoring marine alien species in Greece. *Cahier de Biologie Marine*, 54, 3, 419-426.
- Zgozi, S.W, Haddoud, D.A., Rough, A., 2002. Influence of environmental factors on distribution and abundance of macrobenthic organisms at Al Gazala lagoon (Libya) (in Arabic) *Technical report of Marine Research center of Tajura*, 23-27.
- Zupanovic, S., El-Buni, A.A., 1982. A contribution to demersal fish studies off the Libyan coast. *Bulletin of the Marine Research Centre*, Tripoli, Libya, 3, 78-122.
- Websites visited
- AlgaeBase: <http://www.algaebase.org>
- DAISIE: <http://www.europe-aliens.org>
- FishBase: <http://www.fishbase.org>