

Mediterranean Marine Science

Vol 15, No 2 (2014)



Marine fishes in the Black Sea: recent conservation status

M. H. YANKOVA, D. PAVLOV, P. IVANOVA, E. KARPOVA, A. BOLTACHEV, B. ÖZTÜRK, L. BAT, M. ORAL, M. MGELADZE

doi: [10.12681/mms.700](https://doi.org/10.12681/mms.700)

To cite this article:

YANKOVA, M. H., PAVLOV, D., IVANOVA, P., KARPOVA, E., BOLTACHEV, A., ÖZTÜRK, B., BAT, L., ORAL, M., & MGELADZE, M. (2014). Marine fishes in the Black Sea: recent conservation status. *Mediterranean Marine Science*, 15(2), 366–379. <https://doi.org/10.12681/mms.700>

Marine fishes in the Black Sea: recent conservation status

M. YANKOVA¹, D. PAVLOV², P. IVANOVA¹, E. KARPOVA³, A. BOLTACHEV³, B. ÖZTÜRK⁴,
L. BAT⁵, M. ORAL⁴ and M. MGELADZE⁶

¹Institute of Oceanology, BAS, First of May St., P. O. Box: 152, 9000, Varna, Bulgaria

²Society of Innovative Ecologists in Bulgaria, Dr. Bassanovich St., No. 10, 9010, Varna, Bulgaria

³Institute of Biology of Southern Seas, Sevastopol, Nakhimov av., 99011, Sevastopol, Ukraine

⁴Faculty of Fisheries, Istanbul University, Ordu St., No: 200, 34470, Laleli, Istanbul, Turkey

⁵Faculty of Fisheries, Sinop University, 57000, Sinop, Turkey

⁶Water Ecology and Fisheries Laboratory, National Environmental Agency, Agmashenebeli, Tbilisi, 0112, Georgia

Corresponding author: maria_y@abv.bg

Handling Editor: Costas Papaconstantinou

Received: 18 November 2013; Accepted: 13 March 2014; Published on line: 30 May 2014

Abstract

A revised checklist of the fish fauna of the Black Sea is reported. This paper is the first attempt to present an actual Check List of the fishes in the Black Sea according to the data available in Black Sea countries, as well as their current conservation status, enlisted in IUCN. The total number of Black Sea fish species is 189. Concerning the conservation status, only two species (1.06 %) are extinct (*Acipenser nudipectus* and *A. sturio*), 3.70 % are critically endangered, 16.40 % are vulnerable, 1.06% are endangered; for 10.58 % there is a lack of data, 26.46% has been classified in the category “Least concern”, 2.65 % are “Near threatened” and 38.10% are “Not evaluated”.

Keywords: Keywords: Black Sea, Ichthyofauna, list of species, IUCN.

Introduction

The first description of the Black Sea fishes has been performed by Pallas in 1811. The most complete information about the species composition of fishes in the Black Sea is given by Slastenenko (1955-1956), who listed 189 species. Data on fish species diversity in the Black Sea can be also found in the works of Berg (1949), Rass (1949, 1965), Svetovidov (1964) and Vasil'eva (2007). Svetovidov (1964) has presented 150 marine, diadromous fish species inhabited water of the Black Sea. Vasil'eva (2007) enlisted 176 species for the Black Sea. In the Black Sea Red Data book (Dumont *et al.*, 1999) 41 species for the Bulgarian Black Sea Coast were included. Chichkof (1912) described 51 fish species. Drensky (1921) reported 87 fish species for the Bulgarian Black Sea. Stoyanov *et al.* (1963) and Georgiev & Kolemanova (1978) reported 140 fish species for the Bulgarian Black Sea fish fauna.

Zhivkov *et al.* (2005) describe 218 species (107 freshwater and brackish and 111 marine species) for Bulgarian waters. Karapetkova & Zhivkov (2010) have presented 210 species of Bulgarian Ichthyofauna.

Literature reviews of the fish fauna of the Turkish Black Sea waters have been conducted by a number of scientists (Kocataş *et al.*, 1987; Mater & Meriç 1996;

Öztürk 1999; Bilecenoğlu *et al.*, 2002; Bat *et al.*, 2005; Fricke *et al.*, 2007; Keskin 2010). According to these authors, there are different numbers of fish species in the Turkish Black Sea waters: Kocataş *et al.* (described 150 species, 1987), Öztürk (described 140 fish, 1999), Bilecenoğlu *et al.* (151 fish, 2002), Bat *et al.* (identified 94 fish, 2005), Fricke *et al.* (recorded 157 species, 2007) and Keskin (161 fish, 2010).

The total number of fish species found in the Ukrainian Black Sea waters is 141; among them, 4 species are freshwater inhabitants according to CIESM (2010). Boltachev and Karpova (2012) recorded 135 species in the Black Sea along the Crimean Peninsula.

The fishes of Georgia have been studied by Cuvier (1829), Günther (1866) and Sharvashidze (1982). Presently, the fish checklist of Georgia contains 167 species of which 61 are freshwater inhabitants, 76 marine and 30 species are migratory (Ninua & Japoshvili, 2008).

The aim of the study is to present the actual status of the Black Sea Fish Check List, combining the data from all Black Sea countries.

Materials and Methods

The fish species checklist for the Black Sea was prepared following an exhaustive bibliographic review and

analysis of published articles. The present compilation provides a review of data obtained from the literature. The checklist is based not only on recent and relevant publications, but also on older records. The Latin names of fishes are presented according to their taxonomic priority. Current and old names were analyzed and compared with the available data, in order to clarify the taxonomic status.

The nomenclature follows (Eschmeyer & Fongr, 2014). Fish diversity in the Black Sea was examined by country (including Black Sea region): Bulgaria, Georgia, Romania, Turkey, Ukraine and Russia. For Black Sea countries, research has not been carried out for all listed species regarding their IUCN status, and they have been scientifically determined and categorized. On account of this, possible categorization of the species according to IUCN criteria is not relevant at the moment for most of the listed species. The conservation status of Black Sea marine fishes was completed using the IUCN Red List categories and criteria (version 2 of 2013). All these assessments are available on the IUCN Red List website: <http://www.IUCNredlist.org>.

Results and Discussion

Table 1 presents the annotated list of 189 fish species in the Black Sea along with their distribution.

The nomenclature follows Eschmeyer & Fongr, (2014) catalogue with exception of the following species:

- *Atherina boyeri* Risso, 1810, listed as *Atherina pontica* (Eichwald, 1831). Based on genetic analysis, the authors have a lot of evidence that, in Black Sea waters, the correct name is *Atherina pontica* (Eichwald, 1831). (Dobrovolov & Georgiev, 1995; Dobrovolov & Ivanova, 1999; Dobrovolov *et al.*, 2003).
- *Trachurus mediterraneus* Steindachner, 1868, listed as *Trachurus mediterraneus ponticus* Aleev, 1956. *Trachurus mediterraneus ponticus* Aleev, distinct from the Mediterranean one, i.e. *Tr. mediterraneus* Steindachner (1868), (Dobrovolov, 2000).
- *Scophthalmus maximus* (Linnaeus, 1758) listed as *Psetta maxima* (Linnaeus, 1758). Currently, the genus affiliation of turbot in the Black Sea is not clearly defined. Intensive research (genetic and morphological study) is ongoing to establish genera and species status.

Species reported as alien in the Black Sea (Alexandrov *et al.*, 2007), such as *Oncorhynchus gorboscha*, *Lateolabrax japonicus*, *Oncorhynchus mykiss* and *Plecoglossus altivelis*, *Salvelinus fontinalis*, *Salmo salar*, which were intentionally imported for aquaculture, are not included in our annotated list as they are mostly freshwater species.

The ichthyofauna of the Black Sea includes genera that are very different as regards their origin and ecology. Forty-two species were considered to be endangered to

various extents: seven species were classified as “Critically endangered” (3.70 %), two as “Extinct” (1.06%) and two as “Endangered” (1.06%), thirty one as “Vulnerable” (16.40 %). Fifty species have been classified in the category “Least concern” (26.46%). Concerning the conservation status, 10.58 % are “Data Deficient” and 2.65 % “Near threatened”. More information is needed, as many of these Data Deficient species may in fact be threatened. It is important to focus conservation efforts and research on these species. 38.10% of the species were considered to be “Not Evaluated” (Fig. 1, Table 1).

There is substantial lack of information on the conservation status of nearly 50% of Black Sea marine species (which are assessed as DD and NE). The high percentage of data deficient and not evaluated fish species in the Black Sea indicates the necessity for additional careful study of these species, because some of them could be threatened species. The diagram presents the species (percentage) for which no information is available (Fig. 1).

For these species, it is necessary to take urgent conservation and protection measures. It is also important to implement further monitoring and appropriate managements and recovery plans for these species.

Some commercial fish species are already under threat, such as Bluefin tuna, Sturgeons and Mackerel. The frequency of appearance of *Acipenser gueldenstaedtii*, *A. stellatus* and *Huso huso* throughout the Southern Black Sea coasts is the highest in the Yeşilirmak-Kizilirmak Basin and Sakarya Basin, Turkish coast (Ustaoğlu *et al.*, 2011). *A. persicus* was newly recorded from the Sinop-Samsun coast of the Black Sea (Bat *et al.*, 2005). This species, like *A. nudiventris* and *A. sturio*, should to be marked as extinct (EX), in view of the literature data (Tsekov *et al.*, 2008a). On the other hand, changes in Black sea ichthyofauna diversity are due to natural permanent penetration of alien species through the Bosphorus strait – “mediterrization” or economic activity – artificial introduction, shipping and other factors (Alexandrov *et al.*, 2007; Yankova *et al.*, 2013). Some of the new




Fig.1: Regional Red List status of all marine fish species in the Black Sea. Categories are abbreviated as: EN-Endangered; CR-Critically Endangered; EX-Extinct; VU-Vulnerable; LC-Least Concern; DD-Data Deficient; NE-Not Evaluated; NT-Near Threatened.

Table 1. Fish species in the Black Sea. I. Origin: A (M) Atlanto-Mediterranean species, (C): cosmopolitans, E(M): Mediterranean endemic, E(B): Black Sea endemic, (IP): Indo-Pacific species, (Eu W): European Waters and rare species (*); II. Conservation status: extinct (EX); Critically endangered (CR), Vulnerable (VU); least concern (LC); Not evaluated (NE), No available information (DD), - Near Threatened (NT); III. Distribution: Bulgaria (1); Georgia (2); Romania (3); Russian Federation (4); Turkey (5); Ukraine (6).

Families / species	Origin	IUCN Status	Country of occupancy	References
Acipenseridae				
<i>Acipenser gueldenstaedtii</i> Brandt & Ratzeburg, 1833	E(B)	CR	1, 2, 3, 4, 5, 6	Bat <i>et al.</i> , 2005; Dobrovlov <i>et al.</i> , 2005 ; Karapetkova & Zhivkov, 2010; Oral <i>et al.</i> , 2013; Tsekov <i>et al.</i> , 2008a; Zivkov <i>et al.</i> , 2005,
<i>Acipenser stellatus</i> Pallas, 1771	E(B)	CR	1, 2, 3, 4, 5, 6	Bat <i>et al.</i> , 2005; Dobrovlov <i>et al.</i> , 2005 ; Karapetkova & Zhivkov, 2010; Oral <i>et al.</i> , 2013; Tsekov <i>et al.</i> , 2008a; Zivkov <i>et al.</i> , 2005,
<i>Acipenser nudiiventris</i> Lovetsky, 1828	E(B)	EX	1, 2, 3, 4, 5, 6	Bat <i>et al.</i> , 2005; Fricke <i>et al.</i> , 2007; Karapetkova & Zhivkov, 2010; Oral <i>et al.</i> , 2013;
<i>Acipenser ruthenus</i> Linnaeus, 1758	EuW	VU	1, 5, 6	Dobrovlov <i>et al.</i> , 2005; Karapetkova & Zhivkov, 2010; Oral <i>et al.</i> , 2013; Tsekov <i>et al.</i> , 2008a; Zivkov <i>et al.</i> , 2005;
<i>Acipenser sturio</i> Linnaeus, 1758	A(M)	EX	1, 2, 3, 4, 5, 6	Dobrovlov <i>et al.</i> , 2005, Fricke <i>et al.</i> , 2007; Karapetkova & Zhivkov, 2010; Oral <i>et al.</i> , 2013;
* <i>Acipenser persicus</i> Borodin, 1897	E(B)	CR	2, 4, 5	Bat <i>et al.</i> , 2005; Fricke <i>et al.</i> , 2007; Oral <i>et al.</i> , 2013;
<i>Huso huso</i> (Linnaeus, 1758)	E(B)	CR	1, 2, 3, 4, 5, 6	Dobrovlov <i>et al.</i> 2004, Dobrovlov <i>et al.</i> , 2005, Fricke <i>et al.</i> , 2007; Karapetkova & Zhivkov, 2010; Oral <i>et al.</i> , 2013; Tsekov <i>et al.</i> , 2008; Zivkov <i>et al.</i> 2005;
Alopiidae				
* <i>Alopias vulpinus</i> (Bonnaterre, 1788)	C	VU	5	Fricke <i>et al.</i> , 2007; Geldiay, 1969; Kabasakal, 1998; Vasil'eva, 2007;
Ammodytidae				
<i>Gymnammodytes cicereus</i> (Rafinesque, 1810)	E(M)	NE	1, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005 ;
Anguillidae				
<i>Anguilla anguilla</i> (Linnaeus, 1758)	A(M)	CR	1, 2, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Karapetkova & Zhivkov, 2010; Oral <i>et al.</i> , 2013; Zivkov <i>et al.</i> , 2005;
Atherinidae				
<i>Atherina pontica</i> (Eichwald, 1831)	A(M)	LC	1, 2, 3, 4, 5, 6	Dobrovlov & Georgiev, 1995; Dobrovlov & Ivanova, 1999; Dobrovlov <i>et al.</i> , 2003;
* <i>Atherina hepsetus</i> Linnaeus, 1758	A(M)	NE	1, 3, 4, 5, 6	Boltachev & Karpova, 2012; Dobrovlov & Georgiev, 1995; Erazi, 1942; Fischer <i>et al.</i> , 1987; Geldiay, 1969; Karapetkova & Zhivkov, 2010; Mater & Meriç, 1996; Mater & Bilecenoglu, 1999; Öztürk, 1999; Slastenenko, 1955-1956; Whitehead <i>et al.</i> , 1984-1986;
Balistidae				
* <i>Balistes caprisus</i> Gmelin, 1789	A(M)	NE	5, 6	Bilecenoglu <i>et al.</i> , 2002; Fricke <i>et al.</i> , 2007; Oven & Salekhova, 1969; Svetovidov, 1964;
Belonidae				
<i>Belone belone</i> (Linnaeus, 1761)	E(M)	NE	1, 2, 3, 4, 5, 6	Bilecenoglu <i>et al.</i> , 2002; Eschmeyer & Fongr, 2014; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005;
Blenniidae				
<i>Aidablennius (Blennius) sphyinx</i> (Valenciennes, 1836)	E(M)	NE	1, 2, 3, 4, 5, 6	Karapetkova & Zivkov, 2006, 2010 ; Fricke <i>et al.</i> , 2007; Oral <i>et al.</i> , 2013; Zivkov <i>et al.</i> , 2005 ;
* <i>Blennius ocellaris</i> Linnaeus, 1758	A(M)	NE	2, 5, 6	Fricke <i>et al.</i> , 2007;
<i>Coryphoblennius galerita</i> (Linnaeus, 1758)	E(M)	NE	1, 2, 3, 4, 5, 6	Bat <i>et al.</i> , 2006; Fricke <i>et al.</i> , 2007; Karapetkova & Zhivkov, 2010; Oral <i>et al.</i> , 2013; Zivkov <i>et al.</i> , 2005;
* <i>Microlophrys adriaticus</i> (Steindachner & KolomBatovic, 1883)	A(M)	DD	5, 6	Fricke <i>et al.</i> , 2007; Eschmeyer & Fongr, 2014;
<i>Parablennius (Blennius) zvonimiri</i> (KolomBatovic, 1892)	A(M)	DD	1, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005,
<i>Parablennius (Blennius) sanguinolentus</i> (Pallas, 1814)	A(M)	NE	1, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005 ;
<i>Parablennius (Blennius) tentacularis</i> (Brunnich, 1768)	A(M)	NE	1, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005;
<i>Parablennius gattorugine</i> (Linnaeus, 1758)	A(M)	LC	5	Bilecenoglu <i>et al.</i> , 2002; Vasil'eva, 2007;
<i>Parablennius incognitus</i> (Bath, 1968)	A(M)	NE	2, 4, 5, 6	Boltachev <i>et al.</i> , 2009; Bilecenoglu <i>et al.</i> , Fricke <i>et al.</i> , 2007; 2002; Vasil'eva, 2007;

(continued)

Table 1 (continued)

Families / species	Origin	IUCN Status	Country of occupancy	References
<i>Salaria (Lipophrys) pavo</i> (Risso, 1810)	A(M)	NE	1, 2, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Karapetkova & Zhivkov, 2010; Oral <i>et al.</i> , 2013; Zivkov <i>et al.</i> , 2005;
Bothidae				
<i>Arnoglossus laterna</i> (Walbaum, 1792)	A(M)	NE	5	Bat <i>et al.</i> , 2005; Fricke <i>et al.</i> , 2007;
<i>Arnoglossus thori</i> Kyle, 1913	A(M)	NE	5	Fricke <i>et al.</i> , 2007;
<i>Arnoglossus kessleri</i> Schmidt, 1915	E(M)	DD	1, 2, 3, 4, 5, 6	Boltachev & Karpova, 2012; Karapetkova & Zhivkov, 2010; Svetovidov, 1964; Zivkov <i>et al.</i> , 2005;
Callionymidae				
<i>Callionymus pusillus</i> Delaroche, 1809	A(M)	NE	1, 2, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Karapetkova & Zhivkov, 2010; Oral <i>et al.</i> , 2013; Zivkov <i>et al.</i> , 2005 ;
<i>Callionymus risso</i> Le Sueur, 1814 = (<i>C. belemus</i>)	A(M)	NE	1, 2, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Karapetkova & Zhivkov, 2010; Oral <i>et al.</i> , 2013; Zivkov <i>et al.</i> , 2005 ;
<i>Callionymus fasciatus</i> (Valenciennes in Cuvier & Valenciennes, 1837)	E(M)	LC	4, 5	Fischer <i>et al.</i> , 1987; Fricke <i>et al.</i> , 2007; Mater & Meriç, 1996; Mater & Bilecenoglu, 1999; Öztürk, 1999;
<i>Callionymus lyra</i> Linnaeus, 1758	A(M)	NE	2, 4, 5	Fischer <i>et al.</i> , 1987; Fricke <i>et al.</i> , 2007; Mater & Meriç, 1996; Mater & Bilecenoglu, 1999; Öztürk, 1999; Svetovidov, 1964; Whitehead <i>et al.</i> , 1984-1986;
Carangidae				
<i>Lichia amia</i> (Linnaeus, 1758)	A(M)	DD	1, 5	Bilecenoglu <i>et al.</i> , 2002; Fricke <i>et al.</i> , 2007; Keskin, 2010; Svetovidov, 1964; Karapetkova & Zhivkov, 2010; Stefanov, 2007; Vasil'eva, 2007; Zivkov <i>et al.</i> , 2005;
<i>Naucrates ductor</i> (Linnaeus, 1758)	C	VU	1, 5, 6	Bilecenoglu <i>et al.</i> , 2002; Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Svetovidov, 1964; Stefanov, 2007; Vasil'eva, 2007; Zivkov <i>et al.</i> , 2005;
<i>Trachurus mediterraneus ponticus</i> Aleev, 1956	A(M)	NE	1, 2, 3, 4, 5, 6	Aleev, 1956, 1957, 1959; Banarescu, 1964; Cautis, 1979; Cautis & Jonescu, 1979; Dobrovolov 1977; Dobrovolov 1981; Dobrovolov & Dobrovolova, 1983; Dobrovolov, 1986; Dobrovolov 1988; Dobrovolov & Terzieva, 1995; Dobrovolov 2000; Georgiev & Kolarov, 1962; Karapetkova & Zivkov 2006, 2010; Stoyanov <i>et al.</i> , 1963; Vasil'eva, 2007;
<i>Trachurus trachurus</i> (Linnaeus, 1758)	A(M)	DD	5, 6	Fricke <i>et al.</i> , 2007; Vasil'eva, 2007;
Centracanthidae				
<i>Centracanthus cirrus</i> Rafinesque, 1810	A(M)	DD	3, 5	Abaza <i>et al.</i> , 2006; Fricke <i>et al.</i> , 2007;
<i>Spicara smaris</i> (Linnaeus, 1758)	A(M)	NE	1, 2, 3, 5	Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005;
<i>Spicara maena</i> (Linnaeus, 1758)	A(M)	NE	1, 2, 4, 5, 6	Boltachev & Karpova, 2012; Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005;
<i>Spicara flexuosa</i> Rafinesque, 1810	A(M)	NE	1, 2, 3, 4, 5, 6	Boltachev & Karpova, 2012; Karapetkova & Zhivkov, 2010; Salekhova, 1979; Vasil'eva, 2007; Zivkov <i>et al.</i> , 2005;
Clupeidae				
<i>Alosa immaculata</i> Bennett, 1835	E(B)	VU	1, 2, 3, 4, 5, 6	Dobrovolov <i>et al.</i> , 2012; Fricke <i>et al.</i> , 2007; Freyhof & Kottelat 2008; Keskin, 2010; Karapetkova & Zhivkov, 2010; Mezhzherin <i>et al.</i> , 2009; Turan <i>et al.</i> , 2010; Vasil'eva, 2007;
<i>Alosa caspia</i> (Eichwald, 1838)	E(B)	LC	1, 2, 3, 4, 6	Dobrovolov <i>et al.</i> , 2012; Karapetkova & Zhivkov, 2010; Mezhzherin <i>et al.</i> , 2009; Vasil'eva, 2007; Zivkov <i>et al.</i> , 2005;
<i>Alosa tanaica</i> (Grimm 1901)	E(B)	LC	2, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Keskin, 2010;
<i>Alosa fallax</i> (Lacepede, 1803)	A(M)	LC	1, 3, 4, 5, 6	Dobrovolov <i>et al.</i> , 2012; Georgiev & Kolarov 1958; Karapetkova & Zhivkov, 2010;
<i>Alosa maeotica</i> (Grimm, 1901)	E(B)	LC	1, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Keskin, 2010;
<i>Clupeonella cultriventris</i> (Nordmann, 1840)	E(B)	LC	1, 2, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005 ;
<i>Sardina pilchardus</i> (Walbaum, 1792)	A(M)	NE	1, 2, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005 ;
<i>Sprattus sprattus</i> (Linnaeus, 1758)	A(M)	NE	1, 2, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005 ;
<i>Sardinella aurita</i> Valenciennes, 1847	C	NE	1, 2, 3, 5, 6	Boltachev & Karpova, 2012; Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005;
Chaetodontidae				
* <i>Heniochus acuminatus</i> (Linnaeus, 1758)	A(M)	LC	6	Boltachev & Astakhov, 2004;
Congridae				
* <i>Conger conger</i> (Linnaeus, 1758)	A(M)	DD	2, 5, 6	Fricke <i>et al.</i> , 2007; Keskin, 2010;

(continued)

Table 1 (continued)

Families / species	Origin	IUCN Status	Country of occupancy	References
Dactylopteridae				
* <i>Dactylopterus volitans</i> (Linnaeus, 1758)	A(M)	NE	6	Boltachev <i>et al.</i> , 2013; Movchan, 2011;
Dasyatidae				
<i>Dasyatis pastinaca</i> (Linnaeus, 1758)	A(M)	DD	1, 2, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005;
Echeneididae				
* <i>Echeneis naucrates</i> Linnaeus, 1758	C	NE	1	Valkanov, 1961;
Engraulidae				
<i>Engraulis encrasicolus</i> (Linnaeus, 1758)	E(B)	NE	1, 2, 3, 4, 5, 6	Chashchin, 1998; Dobrovolov, 1976; Fricke <i>et al.</i> , 2007; Grant, 2005; Ivanova & Dobrovolov, 2006; Ivanova <i>et al.</i> , 2013a; Keskin, 2010; Karapetkova & Zhivkov, 2010; Magoulas <i>et al.</i> , 2006; Samsun <i>et al.</i> , 2006; Turan <i>et al.</i> , 2004;
Gadidae				
<i>Merlangius merlangus</i> (Linnaeus, 1758)	E(M)	NE	1, 2, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Vasil'eva, 2007; Zivkov <i>et al.</i> , 2005;
* <i>Micromesistius poutassou</i> (Risso, 1827)	A(M)	NE	6	Boltachev <i>et al.</i> , 1999; Boltachev & Karpova, 2012;
Gasterosteidae				
<i>Gasterosteus aculeatus</i> Linnaeus, 1758	A(M)	LC	1, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005;
<i>Pungitius platygaster</i> (Kessler, 1859)	E(B)	LC	1, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005 ;
Gobiesocidae				
<i>Apletodon dentatus</i> (Facciola, 1887)	A(M)	NE	3, 5	Bat <i>et al.</i> , 2006;
<i>Apletodon microcephalus</i> (Brook, 1890)	A(M)	NE	1	Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005;
<i>Diplecogaster bimaculata bimaculata</i> (Bonnaterre, 1788)	A(M)	NE	1, 3, 4, 5, 6	Boltachev & Karpova, 2012; Erazi, 1942; Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Slastenko, 1955-1956; Zivkov <i>et al.</i> , 2005;
<i>Lepadogaster candolii</i> Risso, 1810	A(M)	NE	1, 2, 3, 4, 5, 6	Boltachev & Karpova, 2012; Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005;
<i>Lepadogaster lepadogaster</i> (Bonnaterre, 1788)	A(M)	LC	1, 2, 3, 4, 5, 6	Boltachev & Karpova, 2012; Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005;
Gobiidae				
<i>Aphia minuta</i> (Risso, 1810)	A(M)	NE	1, 2, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Vassilev <i>et al.</i> , 2010, 2012; Zivkov <i>et al.</i> , 2005;
<i>Benthophiloides brauneri</i> Beling & Iljin, 1927	E(B)	DD	1, 3, 6	Karapetkova & Zhivkov, 2010; Vassilev <i>et al.</i> , 2010, 2012 ; Zivkov <i>et al.</i> , 2005;
<i>Benthophilus stellatus</i> (Sauvage, 1874)	E(B)	LC	1, 3, 6	Karapetkova & Zhivkov, 2010; Vassilev <i>et al.</i> , 2010, 2012; Zivkov <i>et al.</i> , 2005;
<i>Caspiosoma caspium</i> (Kessler, 1877)	E(B)	LC	6	Eschmeyer & Fongr, 2014;
<i>Chromogobius quadrivittatus</i> (Steindachner, 1863)	A(M)	LC	1, 4, 6	Karapetkova & Zhivkov, 2010; Kovtun, 2013; Vassilev <i>et al.</i> , 2010, 2012, Vasil'eva, 2007; Zivkov <i>et al.</i> , 2005 ;
* <i>Chromogobius zebratus</i> (KolomBatovic, 1891)	E(M)	LC	6	Kovtun & Karpova, 2014;
* <i>Gammogobius steinitzi</i> Bath, 1971	E(M)	DD	6	Kovtun & Manilo, 2013;
<i>Gobius niger</i> Linnaeus, 1758	A(M)	NE	1, 2, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Ivanova <i>et al.</i> , 2013b ; Vassilev <i>et al.</i> , 2010, 2012 ; Zivkov <i>et al.</i> , 2005 ;
<i>Gobius cobitis</i> Pallas, 1814	A(M)	LC	1, 2, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Oral <i>et al.</i> , 2013; Vassilev <i>et al.</i> , 2010, 2012 ; Zivkov <i>et al.</i> , 2005 ;
<i>Gobius bucchichi</i> Steindachner, 1870	E(M)	NE	1, 3, 4, 5, 6	Boltachev & Karpova, 2012; Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Oral <i>et al.</i> , 2013; Vassilev <i>et al.</i> , 2010, 2012 ; Zivkov <i>et al.</i> , 2005 ;
<i>Zosterisessor ophiocephalus</i> (Pallas, 1814)	E(M)	DD	1, 5, 6	Boltachev & Karpova, 2012; Dobrovolov <i>et al.</i> , 1995; Fricke <i>et al.</i> , 2007; Ivanova <i>et al.</i> , 2013b; Keskin, 2010; Karapetkova & Zhivkov, 2010; Oral <i>et al.</i> , 2013;
<i>Gobius paganellus</i> Linnaeus, 1758	A(M)	NE	1, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Vassilev <i>et al.</i> , 2010, 2012 ; Zivkov <i>et al.</i> , 2005 ;
<i>Gobius cruentatus</i> Gmelin, 1789	A(M)	NE	5, 6	Boltachev <i>et al.</i> , 2009; Engin <i>et al.</i> , 2007; Fricke <i>et al.</i> , 2007; Keskin, 2010; Slastenenko 1955-1956;

(continued)

Table 1 (continued)

Families / species	Origin	IUCN Status	Country of occupancy	References
<i>Gobius xanthocephalus</i> Heymer et Zander, 1992	E(M)	LC	4, 6	Boltachev <i>et al.</i> , 2009; Boltachev & Karpova, 2012; Vasil'eva, 2007;
<i>Knipowitschia cameliae</i> Nalbant et Otel, 1995	E(B)	CR	3	Vasil'eva, 2007;
<i>Knipowitschia caucasica</i> (Berg, 1916)	E(B)	LC	1, 2, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Ivanova <i>et al.</i> , 2013b; Keskin, 2010; Karapetkova & Zhivkov, 2010; Vassilev <i>et al.</i> , 2010, 2012, Zivkov <i>et al.</i> , 2005;
<i>Knipowitschia longecaudata</i> (Kessler, 1877)	E(B)	LC	1, 2, 3, 5, 6	Fricke <i>et al.</i> , 2007; Ivanova <i>et al.</i> , 2013b ; Keskin, 2010 ; Karapetkova & Zhivkov, 2010; Vassilev <i>et al.</i> , 2010, 2012 , Zivkov <i>et al.</i> , 2005 ;
<i>Mesogobius Batrachocephalus</i> (Pallas, 1814)	E(B)	LC	1, 2, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Vassilev <i>et al.</i> , 2010, 2012; Zivkov <i>et al.</i> , 2005 , Boltachev & Karpova, 2012;
<i>Millerigobius macrocephalus</i> (Kolombatovič, 1891)	E(M)	NE	6	Boltachev & Karpova, 2012;
<i>Neogobius (Ponticola) eurycephalus</i> (Kessler, 1874)	E(B)	LC	1, 3, 4, 5, 6	Keskin, 2010; Vassilev <i>et al.</i> , 2010, 2012
<i>Neogobius (Ponticola) platyrostris</i> (Pallas, 1814)	E(B)	LC	1, 4, 5, 6	Keskin, 2010; Karapetkova & Zhivkov, 2010; Vassilev <i>et al.</i> , 2010, 2012 ; Zivkov <i>et al.</i> , 2005;
<i>Neogobius (Ponticola) cephalargoides</i> (Pinchuk, 1976)	E(B)	NE	1, 3, 4, 6	Karapetkova & Zhivkov, 2010; Vassilev <i>et al.</i> , 2010, 2012; Zivkov <i>et al.</i> , 2005;
<i>Neogobius (Ponticola) kessleri</i> (Günther, 1861)	E(B)	LC	1, 3, 5, 6	Keskin, 2010; Karapetkova & Zhivkov, 2010; Vassilev <i>et al.</i> , 2010, 2012 ; Zivkov <i>et al.</i> , 2005;
<i>Apolonia fluviatilis</i> (Pallas, 1814)	E(B)	LC	1, 2, 3, 4, 5, 6	Brown & Stepien, 2008; Dobrovolov <i>et al.</i> , 1995; Fricke <i>et al.</i> , 2007; Ivanova <i>et al.</i> , 2013b; Keskin, 2010; Neilson & Stepein, 2009; Stepien & Tumeo, 2006;
<i>Apolonia melanostomus</i> (Pallas, 1814)	E(B)	LC	1, 2, 3, 4, 5, 6	Brown & Stepien, 2008; Dobrovolov <i>et al.</i> , 1995; Fricke <i>et al.</i> , 2007; Ivanova <i>et al.</i> , 2013b; Keskin, 2010; Neilson & Stepein, 2009; Stepien & Tumeo, 2006;
<i>Neogobius (Ponticola) ratan</i> (Nordmann, 1840)	E(B)	NE	1, 2, 3, 4, 5, 6	Keskin, 2010; Vassilev <i>et al.</i> , 2010, 2012 Zivkov <i>et al.</i> , 2005;
<i>Neogobius (Ponticola) syrman</i> (Nordmann, 1840)	E(B)	LC	1, 3, 4, 5, 6	Ivanova <i>et al.</i> , 2013b; Keskin, 2010; Karapetkova & Zhivkov, 2010; Vassilev <i>et al.</i> , 2010, 2012; Zivkov <i>et al.</i> , 2005;
<i>Babka gymnotrachelus</i> (Kessler, 1857)	E(B)	LC	1, 2, 3, 4, 5, 6	Neilson & Stepein, 2009;
<i>Pomatoschistus marmoratus</i> (Risso, 1810)	A(M)	NE	1, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Ivanova <i>et al.</i> , 2013b ;Keskin, 2010; Karapetkova & Zhivkov, 2010; Vassilev <i>et al.</i> , 2010, 2012 , Zivkov <i>et al.</i> , 2005;
<i>Pomatoschistus minutus</i> (Pallas, 1770)	A(M)	NE	1, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Vassilev <i>et al.</i> , 2010, 2012 ; Zivkov <i>et al.</i> , 2005 ;
<i>Proterorhinus marmoratus</i> (Pallas, 1811)	E(B)	VU	1, 2, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Vassilev <i>et al.</i> , 2010, 2012; Zivkov <i>et al.</i> , 2005 ;
<i>Pomatoschistus Bathi</i> Miller, 1982	E(M)	DD	1, 2, 4, 6	Boltachev & Karpova, 2012; Vasil'eva, 2007; Vassilev <i>et al.</i> , 2010; 2012;
<i>Pomatoschistus pictus</i> (Malm, 1865)	E(M)	NE	4	Vasil'eva, 2007;
<i>Tridentiger trigonocephalus</i> (Gill, 1859)	IP	NE	6	Boltachev & Karpova, 2010;
* <i>Zebrus zebrus</i> (Risso, 1827)	E(M)	LC	5, 6	Boltachev & Karpova, 2013; Kovačić & Engin, 2009; Keskin, 2010;
Gymnuridae				
* <i>Gymnura altavela</i> (Linnaeus, 1758)	A(M)	VU	5	Bilecenoglu <i>et al.</i> , 2002; Fricke <i>et al.</i> , 2007; Keskin, 2010;
Hexanchidae				
<i>Hexanchus griseus</i> Bonnaterre, 1788	C	NT	5	Kabasakal, 2013;
Labridae				
* <i>Coris julis</i> (Linnaeus, 1758)	A(M)	LC	1, 3, 5	Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005,
<i>Ctenolabrus rupestris</i> (Linnaeus, 1758)	A(M)	NE	1, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005;
<i>Labrus viridis</i> Linnaeus, 1758	A(M)	VU	1, 3, 4, 5, 6	Boltachev & Karpova, 2012; Erazi, 1942; Fischer <i>et al.</i> , 1987; Fricke <i>et al.</i> , 2007; Geldiay, 1969; Keskin, 2010; Karapetkova & Zhivkov, 2010; Mater & Meriç, 1996; Mater & Bilecenoglu, 1999; Öztürk, 1999; Slastenenko, 1955-1956; Svetovidov, 1964; Whitehead <i>et al.</i> , 1984-1986; Zivkov <i>et al.</i> , 2005;
<i>Symphodus cinereus</i> (Bonnaterre, 1788)	A(M)	LC	1, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010;

(continued)

Table 1 (continued)

Families / species	Origin	IUCN Status	Country of occupancy	References
<i>Symphodus roissali</i> (Risso, 1810)	A(M)	LC	1, 2, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010;
<i>Symphodus tinca</i> (Linnaeus, 1758)	A(M)	LC	1, 2, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Oral <i>et al.</i> , 2013
<i>Symphodus ocellatus</i> (Linnaeus, 1758)	E(M)	VU	1, 2, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010;
<i>Symphodus scina</i> (Forsskål, 1775)	E(M)	VU	3, 4, 5, 6	Boltachev & Karpova, 2012; Vasil'eva, 2007;
Lophiidae				
<i>Lophius piscatorius</i> Linnaeus, 1758	A(M)	NE	1, 2, 3, 4, 5, 6	Bat <i>et al.</i> , 2005; Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005 ;
* <i>Lophius budegassa</i> Spinola, 1807	A(M)	VU	5	Fricke <i>et al.</i> , 2007; Keskin, 2010; Mater & Bilecenoglu, 1999; Öztürk, 1999;
Moronidae				
<i>Dicentrarchus labrax</i> (Linnaeus, 1758)	A(M)	VU	1, 2, 3, 4, 5, 6	Bat <i>et al.</i> , 2005; Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005;
* <i>Morone saxatilis</i> (Walbaum, 1792)	A(M)	NE	2, 4, 6	Aleev <i>et al.</i> , 2007; Black Sea Transboundary Diagnostic Analysis 2007; Vasil'eva, 2007; Zaitsev and Öztürk, 2001;
Mugilidae				
<i>Chelon labrosus</i> (Risso, 1827)	A(M)	LC	1, 5, 6	Antović, 2013; Boltachev & Yurakhno, 2002; Boltachev & Karpova, 2012; Dobrovolov <i>et al.</i> , 2003; Erguden <i>et al.</i> , 2010; Fricke <i>et al.</i> , 2007; Keskin, 2010; Turan <i>et al.</i> , 2005;
<i>Liza aurata</i> (Risso, 1810)	A(M)	LC	1, 2, 3, 4, 5, 6	Bat <i>et al.</i> , 2005; Fricke <i>et al.</i> , 2007; Keskin, 2010 ; Karapetkova & Zhivkov, 2010;
<i>Liza ramada</i> (Risso, 1827)	A(M)	LC	1, 3, 5, 6	Boltachev & Karpova, 2012; Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010;
<i>Liza saliens</i> (Risso, 1810)	A(M)	LC	1, 2, 3, 4, 5, 6	Bat <i>et al.</i> , 2005; Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010;
<i>Mugil cephalus</i> Linnaeus, 1758	A(M)	LC	1, 2, 3, 4, 5, 6	Bat <i>et al.</i> , 2005; Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010;
<i>Liza haematocheila</i> (Temminck & Schlegel 1845)	IP	NE	1, 2, 3, 4, 5, 6	Karapetkova & Zivkov, 2006, 2010; Dobrovolov <i>et al.</i> , 2003; Zivkov <i>et al.</i> , 2005;
Mullidae				
<i>Mullus barbatus</i> (Linnaeus, 1758)	A(M)	LC	1, 2, 3, 4, 5, 6	Eschmeyer & Fongr, 2014; Fricke <i>et al.</i> , 2007; Karapetkova & Zivkov, 2006, 2010; Keskin, 2010; Zivkov <i>et al.</i> , 2005 ;
<i>Mullus surmuletus</i> Linnaeus, 1758	A(M)	NE	2, 4, 5	Fricke <i>et al.</i> , 2007; Ivanova <i>et al.</i> , 2014 ; Keskin, 2010;
Merlucciidae				
* <i>Merluccius merluccius</i> (Linnaeus, 1758)	A(M)	NE	5	Fricke <i>et al.</i> , 2007; Keskin, 2010;
Ophidiidae				
<i>Ophidion rochei</i> Müller, 1845	E(M)	DD	1, 2, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005;
Percidae				
<i>Percarina demidoffii</i> Nordmann 1840	E(B)	NT	3, 6	Vasil'eva, 2007;
<i>Sander marinus</i> (Cuvier in Cuvier & Valenciennes, 1828)	E(B)	DD	1, 6	Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005
Pleuronectidae				
<i>Platichthys flesus</i> (Linnaeus, 1758)	A(M)	LC	1, 2, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007 ; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005 ;
Pomacentridae				
<i>Chromis chromis</i> Linnaeus, 1758	A(M)	NE	1, 2, 3, 4, 5, 6	Bat <i>et al.</i> , 2005; Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005 ;
Pomatomidae				
<i>Pomatomus saltatrix</i> (Linnaeus, 1766)	C	NE	1, 2, 3, 4, 5, 6	Bat <i>et al.</i> , 2005; Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005 ;
Phycidae				
<i>Gaidropsarus mediterraneus</i> (Linnaeus, 1758)	A(M)	LC	2, 5	Bat <i>et al.</i> , 2005; Fricke <i>et al.</i> , 2007; Keskin, 2010;
Rajidae				
<i>Raja clavata</i> Linnaeus, 1758	C	VU	1, 2, 3, 4, 5, 6	Bat <i>et al.</i> , 2005; Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005;
Salmonidae				

(continued)

Table 1 (continued)

Families / species	Origin	IUCN Status	Country of occupancy	References
<i>Salmo labrax</i> Pallas, 1814	E(B)	LC	1, 2, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005 ;
<i>Salmo coruhensis</i> Turan, Kottelat & Engin, 2010	E(M)	DD	5	Oral <i>et al.</i> , 2013; Turan <i>et al.</i> , 2009;
<i>Salmo rizeensis</i> Turan, Kottelat & Engin, 2010	E(M)	DD	5	Oral <i>et al.</i> , 2013; Turan <i>et al.</i> , 2009;
Sciaenidae				
* <i>Argyrosomus regius</i> (Asso, 1801)	A(M)	DD	5	Fricke <i>et al.</i> , 2007; Keskin, 2010;
<i>Sciaena umbra</i> Linnaeus, 1758	A(M)	VU	1, 2, 3, 4, 5, 6	Bat <i>et al.</i> , 2005; Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005;
<i>Umbrina cirrosa</i> (Linnaeus, 1758)	A(M)	NE	1, 2, 3, 4, 5, 6	Bat <i>et al.</i> , 2005; Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005 ;
Scombridae				
* <i>Auxis rochei rochei</i> (Risso, 1810)	C	LC	5	Fricke <i>et al.</i> , 2007; Keskin, 2010;
* <i>Euthynnus alletteratus</i> (Rafinesque-Schmaltz, 1810)	A(M)	LC	1, 5	Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005;
<i>Sarda sarda</i> (Bloch, 1793)	C	LC	1, 2, 3, 4, 5, 6	Bat <i>et al.</i> , 2005; Boltachev & Karpova, 2012; Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Oral <i>et al.</i> , 2013; Vasil'eva, 2007; Zivkov <i>et al.</i> , 2005 ;
<i>Scomber japonicus</i> Houttuyn, 1782	A(M)	LC	1, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005;
<i>Scomber scombrus</i> Linnaeus, 1758	A(M)	LC	1, 2, 3, 4, 5, 6	Eschmeyer & Fongr, 2014, Karapetkova & Zhivkov, 2010; Vasil'eva, 2007; Zivkov <i>et al.</i> , 2005;
<i>Thunnus thynnus</i> (Linnaeus, 1758)	A(M)	EN	1, 2, 3, 4, 5, 6	Bat <i>et al.</i> , 2005; Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Oral <i>et al.</i> , 2013; Vasil'eva, 2007; Zivkov <i>et al.</i> , 2005;
Scophthalmidae				
<i>Psetta maxima</i> Linnaeus (1758)	A(M)	NE	1, 3, 4, 5, 6	Atanasov <i>et al.</i> , 2011; Karapetkova & Zhivkov, 2006, 2010; Tsekov <i>et al.</i> , 2008b; Zivkov <i>et al.</i> , 2005,.
<i>Scophthalmus maeoticus</i> (Pallas 1814)	A(M)	NE	1, 2, 3, 4, 5, 6	Vasil'eva, 2007;
<i>Scophthalmus rhombus</i> (Linnaeus, 1758)	A(M)	VU	1, 3, 4, 5, 6	Eschmeyer & Fongr, 2014; Fricke <i>et al.</i> , 2007; Vasil'eva, 2007;
Scorpaenidae				
<i>Scorpaena porcus</i> Linnaeus, 1758	A(M)	NE	1, 2, 3, 5, 6	Bat <i>et al.</i> , 2005; Fricke <i>et al.</i> , 2007; Karapetkova & Zhivkov, 2006, 2010 ; Keskin, 2010; Oral <i>et al.</i> , 2013; Zivkov <i>et al.</i> , 2005;
<i>Scorpaena notata</i> Rafinesque, 1810	A(M)	VU	3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Keskin, 2010;
Scyliorhinidae				
<i>Scyliorhinus canicula</i> (Linnaeus, 1758)	A(M)	VU	4, 5	Aksiray, 1987; Erazi, 1942; Fischer, 1973; Geldiay, 1969; Slastenenko, 1955-1956; Mater & Meriç, 1996; Mater & Bilecenoglu, 1999; Öztürk, 1999; Svetovidov, 1964; Vasil'eva, 2007;
Serranidae				
<i>Serranus cabrilla</i> (Linnaeus, 1758)	A(M)	NE	1, 2, 6	Boltachev & Karpova, 2013, Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005;
* <i>Serranus hepatus</i> (Linnaeus, 1758)	A(M)	NT	3, 5	Dalgıç <i>et al.</i> , 2013; Fricke <i>et al.</i> , 2007;
<i>Serranus scriba</i> (Linnaeus, 1758)	A(M)	NT	1, 2, 5	Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005; Zivkov <i>et al.</i> , 2005;
Soleidae				
<i>Buglossidium luteum</i> (Risso, 1810)	A(M)	VU	4, 5	Fricke <i>et al.</i> , 2007; Keskin, 2010; Vasil'eva, 2007;
<i>Microchirus variegatus</i> (Donovan, 1808)	A(M)	VU	5	Fricke <i>et al.</i> , 2007; Keskin, 2010; Vasil'eva, 2007;
<i>Solea nasuta</i> (Pallas, 1814)	E(M)	NE	1, 2, 5	Karapetkova & Zhivkov, 2010; Vasil'eva, 2007; Zivkov <i>et al.</i> , 2005;
<i>Solea solea</i> (Linnaeus, 1758)	A(M)	NE	4, 5, 6	Bat <i>et al.</i> , 2005; Fricke <i>et al.</i> , 2007; Keskin, 2010;
Sparidae				
<i>Boops boops</i> (Linnaeus, 1758)	A(M)	LC	1, 2, 3, 4, 5, 6	Bat <i>et al.</i> , 2005; Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005;
<i>Dentex dentex</i> (Linnaeus, 1758)	A(M)	VU	1, 3, 4, 5, 6	Bat <i>et al.</i> , 2005; Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005;
<i>Diplodus annularis</i> (Linnaeus, 1758)	A(M)	VU	1, 2, 3, 5	Bat <i>et al.</i> , 2005; Fricke <i>et al.</i> , 2007; Keskin, 2010 ; Karapetkova & Zhivkov, 2010; ; Zivkov <i>et al.</i> , 2005;

(continued)

Table 1 (continued)

Families / species	Origin	IUCN Status	Country of occupancy	References
<i>Diplodus puntazzo</i> (Walbaum, 1792)	A(M)	VU	1, 2, 3, 4, 5, 6	Bat <i>et al.</i> , 2005; Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005;
<i>Diplodus sargus sargus</i> (Linnaeus, 1758)	E(M)	NE	1, 3, 4, 5, 6	Bat <i>et al.</i> , 2005; Fricke <i>et al.</i> , 2007; Keskin, 2010; Tkachenko, 2013; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005;
<i>Diplodus vulgaris</i> (Geoffroy Saint-Hilaire, 1817)	A(M)	NE	1, 5, 6	Bat <i>et al.</i> , 2005; Bilecenoglu <i>et al.</i> , 2002; Eschmeyer & Fongr, 2014; Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Svetovidov, 1964; Stefanov, 2007; Tkachenko, 2013; Vasil'eva, 2007; Zivkov <i>et al.</i> , 2005,
<i>Lithognathus mormyrus</i> (Linnaeus, 1758)	A(M)	NE	5, 6	Bilecenoglu <i>et al.</i> , 2002; Boltachev <i>et al.</i> , 2013; Stefanov, 2007; Vasil'eva, 2007; Zivkov <i>et al.</i> , 2005;
<i>Oblada melanura</i> (Linnaeus, 1758)	A(M)	NE	1, 3, 5	Karapetkova & Zhivkov, 2010; Stefanov, 2007; Vasil'eva, 2007; Zivkov <i>et al.</i> , 2005;
<i>Pagellus erythrinus</i> (Linnaeus, 1758)	A(M)	NE	1, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005;
<i>Sarpa salpa</i> (Linnaeus, 1758)	A(M)	NT	1, 2, 3, 4, 5, 6	Bat <i>et al.</i> , 2005; Boltachev & Yurakhno, 2002; Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005;
<i>Spondyliosoma cantharus</i> (Linnaeus, 1758)	A(M)	NE	1, 4, 5, 6	Bat <i>et al.</i> , 2005; Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Svetovidov, 1964; Vasil'eva, 2007; Zivkov <i>et al.</i> , 2005;
<i>Sparus aurata</i> Linnaeus, 1758	A(M)	VU	1, 2, 3, 5, 6	Bat <i>et al.</i> , 2005; Boltachev & Yurakhno, 2002; Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005;
Sphyraenidae				
<i>Sphyraena sphyraena</i> (Linnaeus, 1758)	A(M)	VU	1, 3, 4, 5, 6	Boltachev <i>et al.</i> , 2009; Boltachev & Karpova, 2012; Fricke <i>et al.</i> , 2007; Karapetkova & Zhivkov, 2010; Keskin, 2010; Svetovidov, 1964; Vasil'eva, 2007; Zivkov <i>et al.</i> , 2005;
<i>Sphyraena pinguis</i> Günther, 1874	IP	NE	6	Boltachev, 2009;
Sphyrnidae				
<i>Sphyrna zygaena</i> (Linnaeus, 1758)	C	VU	3	Vasil'eva, 2007;
Squalidae				
<i>Squalus acanthias</i> Linnaeus, 1758	C	EN	1, 2, 3, 4, 5, 6	Bat <i>et al.</i> , 2005; Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005;
<i>Squalus blainville</i> (Risso, 1827)	C	DD	1, 5	Bat <i>et al.</i> , 2005; Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005;
Squatinae				
* <i>Squatina squatina</i> (Linnaeus, 1758)	A(M)	CR	5	Bilecenoglu <i>et al.</i> , 2002; Fricke <i>et al.</i> , 2007; Keskin, 2010;
Syngnathidae				
<i>Hippocampus hippocampus</i> (Linnaeus, 1758)	A(M)	DD	1, 2, 3, 4, 5, 6	Bat <i>et al.</i> , 2005; Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Oral <i>et al.</i> , 2013; Zivkov <i>et al.</i> , 2005;
<i>Nerophis ophidion</i> (Linnaeus, 1758)	A(M)	VU	1, 2, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Oral <i>et al.</i> , 2013; Zivkov <i>et al.</i> , 2005 ;
<i>Syngnathus abaster</i> Risso, 1827	A(M)	VU	1, 2, 3, 4, 5, 6	Bat <i>et al.</i> , 2005; Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Oral <i>et al.</i> , 2013; Zivkov <i>et al.</i> , 2005;
* <i>Syngnathus acus</i> Linnaeus, 1758	A(M)	NE	1, 3, 4, 5, 6	Bat <i>et al.</i> , 2005; Boltachev <i>et al.</i> , 2009; Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Oral <i>et al.</i> , 2013; Zivkov <i>et al.</i> , 2005 ;
<i>Syngnathus schmidtii</i> Popov, 1927	E(B)	NE	1, 5, 6	Bilecenoglu <i>et al.</i> , 2002; Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005;
<i>Syngnathus tenuirostris</i> Rathke, 1837	E(M)	DD	1, 2, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Oral <i>et al.</i> , 2013; Zivkov <i>et al.</i> , 2005 ;
<i>Syngnathus typhle</i> Linnaeus, 1758	A(M)	LC	1, 2, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Oral <i>et al.</i> , 2013; Zivkov <i>et al.</i> , 2005;
<i>Syngnathus variegatus</i> Pallas, 1814	E(B)	VU	1, 2, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005 ;
Trachinidae				
<i>Trachinus draco</i> Linnaeus, 1758	A(M)	NE	1, 2, 3, 4, 5, 6	Bat <i>et al.</i> , 2005; Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Oral <i>et al.</i> , 2013; Zivkov <i>et al.</i> , 2005;

(continued)

Table 1 (continued)

Families / species	Origin	IUCN Status	Country of occupancy	References
Triglidae				
<i>Chelidonichthys (Aspitrigla) cuculus</i> (Linnaeus, 1758)	A(M)	NE	5, 6	Fricke <i>et al.</i> , 2007; Keskin, 2010;
<i>Chelidonichthys lucerna</i> (Linnaeus, 1758)	A(M)	VU	1, 2, 3, 4, 5, 6	Bat <i>et al.</i> , 2005; Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Oral <i>et al.</i> , 2013; Svetovidov, 1964; Vasil'eva, 2007; Zivkov <i>et al.</i> , 2005;
<i>Eutrigla gurnardus</i> (Linnaeus, 1758)	A(M)	VU	1, 5	Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005 ;
Trichiuridae				
<i>Lepidopus caudatus</i> (Euphasen, 1788)	A(M)	NE	5	Fricke <i>et al.</i> , 2007; Keskin, 2010;
Tripterygiidae				
<i>Tripterygion tripteronotum</i> (Risso, 1810)	E(M)	VU	3, 4, 5, 6	Boltachev & Karpova, 2012; Erazi, 1942; Geldiay, 1969; Keskin, 2010; Mater & Meriç, 1996; Mater & Bilecenoglu, 1999; Öztürk, 1999; Svetovidov, 1964; Slastenenko, 1955-1956; Vasil'eva, 2007; Whitehead <i>et al.</i> , 1984-1986;
Uranoscopidae				
<i>Uranoscopus scaber</i> Linnaeus, 1758	A(M)	VU	1, 2, 3, 4, 5, 6	Bat <i>et al.</i> , 2005; Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov, 2010; Oral <i>et al.</i> , 2013; Zivkov <i>et al.</i> , 2005;
Xiphiidae				
<i>Xiphias gladius</i> Linnaeus, 1758	C	LC	1, 2, 3, 4, 5, 6	Karapetkova & Zhivkov, 2010; Zivkov <i>et al.</i> , 2005;
Zeidae				
<i>Zeus faber</i> Linnaeus, 1758	C	NE	1, 2, 3, 4, 5, 6	Fricke <i>et al.</i> , 2007; Keskin, 2010; Karapetkova & Zhivkov BS TDA Zhivkov, 2010; Zivkov <i>et al.</i> , 2005;

species form their own populations and become constant inhabitants of the Black Sea; others remain rare or known from single findings (Boltachev *et al.*, 2009).

Analyzing the geographic origin of marine species, most of them (59.26%) have Atlanto-Mediterranean origin, 7.41% are cosmopolitans, 31.22% are endemics (Mediterranean and Black Sea) and 2.12% are introduced species (Indo-Pacific and European Waters) (Fig. 2, Table). The recent increasing species diversity in the Black Sea region is due to Mediterranean immigrants.

Marine fishes in the Black Sea are affected by a number of major threats as:

- 1) Over-exploitation of commercial fish stocks (for targeted species).
- 2) Industrial and domestic pollution.

- 3) Habitat loss - land use changes, river barriers, drainage.
- 4) Human disturbance fish farming, fishery management and the introduction of new species.

Conservation measures in the Black Sea, which should apply:

- Fishing restriction and management
- Marine Protected Areas are an important conservation tool. MPAs allow for the conservation of species and their biophysical environments and are an effective way to safeguard ecosystem services. When carefully designed and managed, MPAs can increase fish species richness (Tunesi & Molinari, 2005).
- Several international conventions and national initiatives exist to protect fish species. – Four International conventions such as CITES, the Bern Convention on the conservation of European wildlife and Natural Habitats and the Convention on the Conservation of Migratory species of wild animals (CMS or Bern Convention), are relevant to the conservation and management of Black Sea marine fish fauna.

The effective implementation and control of existing measures (national, international and regular) is very important for the survival of Black Sea fish species.

At national level, regulations are in place for the protection of red listed species. At the international level, however, no universal agreements between Black Sea countries exist extending protection to Red Listed species, with the exception of mammals.

Although some species are protected under national, regional or international conventions, the vast majority of




Fig. 2: Origin of marine species along the Black Sea Coast.

threatened species are not protected or subject to effective management plans. The creation of effective marine protected areas should be implementing in order to reduce pressure on fish populations and safeguard fish habitats. Further funding and research for endemic fish species is high priority.

Conclusions

This paper presents the actual and complete Black Sea fish check list as well as the conservation status of marine fish species. For most of the species (about 50% of all marine fish species), the conservation status was evaluated as data deficient and not evaluated. These groups may be included in the large proportion of threatened species

Sound fishery management and common conservation measures should be applied in all Black Sea countries. Priority areas for further research should be specified. Furthermore, monitoring of fish species and populations and genetical identification are needed for sustainable use of commercial fish species as well as protection of biodiversity. Scientific collaboration between Black Sea countries should continue in order to update the Check List and conservation status of marine fish species when new available information is obtained.

References

- Abaza, V., Boicenco, L., Moldoveanu, M., Timofte, F., Bologna, A. *et al.*, 2006. Evolution of marine biodiversity status at the Romanian Black Sea coast as result of anthropogenic modifications in the last decades. p. 50-51. In: *1st Biannual Scientific Conference- Black Sea Ecosystem 2005 and Beyond, 9-12 May, 2006*, Istanbul, Turkey.
- Aleev, Yu. G., 1956. On the taxonomy of the Black-Sea scad. *Voprosy Ikhtyologii*, 7, 175-184 (In Russian).
- Aleev, Yu. G., 1957. The scads (Trachurus) in the USSR seas. *Trud Sevastopolskii biologicheskoi stantsii.*, 9, 167-242 (In Russian).
- Aksiray, F., 1987. *Türkiye deniz balıkları ve tayin anahtarı* [Marine fishes of Turkey and a key to species]. Istanbul Üniversitesi Rektörlüğü Yayınları, Istanbul, 811 pp. (In Turkish).
- Aleev, Yu. G., 1959. On the reproduction of the southern population of the horse mackerel in the western areas of Black Sea. *Trud Sevastopolskii biologicheskoi stantsii.* 12, 270-285. (In Russian).
- Aleev, B., Boltachev, A., Kharchenko, T., Liashenko, A., Son, M. *et al.*, 2007. Trends of aquatic alien species invasion in Ukraine. Aquatic Invasions, *The European Journal of Applied Research on Biological Invasions in Aquatic Ecosystems*, 2 (3), 215-242.
- Antović, I., 2013. Continuous Osteological Characters in the Reconstruction of Phylogenetic Relationships of the Six Euro-Mediterranean Mullet Species (Mugilidae). *Zoological Science*, 30 (9), 754-759.
- Atanassov, I., Ivanova, P., Panayotova, M., Tsekov, A., Rusanov, K., 2011. Mitochondrial control region DNA variation in turbot populations from the Bulgarian and Romanian Black Sea coasts. *Biotechnology & Biotechnological Equipment*, 25/4, ISSN: 2627-2633.
- Banarescu, P. (Ed.), 1964. *The fauna of the popular Republic of Romania. Pisces – Osteichthyes XIII Academici Republicii Populare*. Romine Press, Romania, 959 pp. (In Romanian).
- Bat, L., Erdem, Y., Ustaoglu, S., Yardim, Ö., Satilmis, H., 2005. A Study on the Fishes of the Central Black Sea Coast of Turkey. *Journal of the Black Sea/Mediterranean Environment*, 11 (3), 287-302.
- Bat, L., Demirci, G.G., Öztürk, M., 2006. Occurrence of *Apletodon dentatus bacescui* (Murgoci, 1940) (Gobiesocidae) and *Coryphoblennius galerita* (Linnaeus, 1758) (Blenniidae) at the central Black Sea coast of Turkey. *Journal of the Black Sea/Mediterranean Environment*, 12 (1), 59-65.
- Berg, L.S., 1949. *Ryby presnykh vod SSSR i sopredelnykh stran (Fish of the freshwater bodies of the USSR and adjacent states)*. 4th ed. Nauka Publishers, Moscow, Leningrad, 1382 pp. (In Russian).
- Bilecenoglu, M., Taskavak, E., Mater, S., Kaya, M., 2002. *Checklist of the marine fishes of Turkey*. Zootaxa 113. Magnolia Publishers, Auckland, New Zealand, 194 pp.
- Black Sea Transboundary Diagnostic Analysis (BS TDA). May, 2007, 227pp.
- Boltachev, A., Gaevskaya, A., Zuev, G., Yurakhno, V., 1999. The blue whiting, *Micromesistius poutassou* (Risso, 1826) (Pisces: Gadidae), the new species for the Black Sea fauna. *Ecologiya Morya*, 48, 79-82.
- Boltachev, A., Yurakhno, V., 2002. New evidences of ongoing mediterraneanization of the Black Sea ichthyofauna. *Journal Ichthyology*, 42 (9), 713-719.
- Boltachev, A., 2004. Specifying species belonging of Barracuda of group *Sphyraena obtusata* (Pisces: Sphyraenidae) found in the Black Sea. *Journal Ichthyology*, 49 (1), 128-131.
- Boltachev, A., Astakhov, D., 2004. An unusual finding of pennant coralfish *Heniochus acuminatus* (Chaetodontidae) in Balaklava Bay (Sevastopol, Southwestern Crimea). *Journal of Ichthyology*, 44 (9), 798-799.
- Boltachev, A.R., 2009. Specifying Species Belonging of Barracuda of Group *Sphyraena obtusata* (Pisces: Sphyraenidae) Found in the Black Sea. *Journal of Ichthyology*, 2009, 49 (1), 128-131.
- Boltachev, A., Karpova, E., Danilyuk, O., 2009. Findings of new and rare fish species in the coastal zone of the Crimea (the Black Sea). *Journal of Ichthyology*, 49 (4), 277-291.
- Boltachev, A.R., Karpova, E.P., 2010. Naturalization of the Pacific Chameleon Goby *Tridentiger trigonocephalus* (Perciformes, Gobiidae) in the Black Sea (Crimea, Sevastopol Galf). *Journal of Ichthyology*, 50 (2), 188-296.
- Boltachev, A.R., Karpova, E.P., 2012. *Marine fishes of the Crimean peninsula*. Business Informs Publishers, Simferopol, Ukraine, 120 pp., ISBN: 978 966 648 319 8. (In Russian).
- Boltachev, A.R., Karpova, E.P., Kirin, M.P., 2013. The first find of the sand steenbras *Lithognathus mormyrus* (L., 1758) (Osteichthyes, Sparidae) in the Black Sea coastal zone of the Crimea. *Marine Ecological Journal*, 12 (4), 96.
- Boltachev, A.R., Karpova, E.P., 2013. Faunistic revision of alien fish species in the Black Sea. *Russian Journal of Biological Invasions*, (In press).
- Brown, J., Stepien, C., 2008. Ancient divisions, recent expansions: phylogeography and population genetics of the round goby *Apollonia melanostoma*. *Molecular Ecology*, 17, 2598-2615.
- Cautis, I., 1966. Cerc. Provect. *Bulletin Institute Pisc*, 252, 31-40. (In Romanian).
- Cautis, I., 1979. Biologie de la reproduction du chin chard de la Mer Noire. p. 461-478. In: *Le Chinchard de la Mer Noire*

- (*Trachurus mediterraneus ponticus*), *etude monographique*. Pora, E. (Ed.). Deuxieme Partie, Publishers, Romania.
- Chashchin, A., 1998. The anchovy and other pelagic fish stocks transformations in the Azov-Black Sea Basin under environmental and fisheries impact. *The Proceeding of the First Symposium on Fisheries and Ecology*, 1-10.
- Chichkof, G., 1912. Contribution à l'étude de la faune de la Mer Noire. Animaux recoltés sur les cotes Bulgares. *Archives of Zoological Experimental Genetics Notes Review*, 10, 39-39.
- CIESM, 2010. *Climate forcing and its impacts on the Black Sea Marine Biota*. p. 105–111. In: 39th CIESM Workshop Monographs, Briand, F. (Eds). CIESM Publishers, Monaco.
- Cuvier, G., 1829. *Le règne animal*. 2nd ed. Déterville et Crochard. Paris, France, 406 pp. (In French).
- Dalgıç, G., Gümüş, A., Zengin, M., 2013. First record of Brown comber *Serranus hepatus* (Linnaeus, 1758) for the Black Sea. *Turkish Journal of Zoology*, 37, 523-524.
- Dobrovolo, I., 1976. Multiple forms of lactate dehydrogenase in anchovy (*Engraulis encrasicolus* L.) from the Black Sea, Sea of Azov and the Atlantic Ocean. *Comptes rendus de l'Academie bulgare des Sciences*, 29, 877-880.
- Dobrovolo, I., 1977. Electrophoretic investigations of proteins from some Carangidae representatives in relation to interspecies differentiation. *Proceeding of Institute of Fisheries Varna*, 15, 35-46. (In Bulgarian.)
- Dobrovolo, I., S., 1981. Biochemic-genetic investigations on some Black Sea fishes. *Scientific reports*. Black Sea in the 1300 years history of Bulgaria, 339-346.
- Dobrovolo, I., Dobrovolo, S., 1983. Biochemical polymorphism of the Black Sea and Mediterranean scads. *Proc. of the Institute of Fisheries Varna*, XX, 101-107.
- Dobrovolo, I., 1986. Study of population structure of the Black Sea horse mackerel with gen-markers. p. 29-31. In: *Anniversary National Conference of Biology, 29-31 May 1986*. Pleven, Bulgaria.
- Dobrovolo, I., 1988. *Biochemical and population genetic investigations of industrial fish species in the waters of Bulgaria and the World Ocean*. MSc. Thesis. Institute of Zoology, Bulgaria, 533 p.
- Dobrovolo, I. S., 2000. Genetic divergence between the scad subspecies *Trachurus mediterraneus* (Carangidae, Pisces) from the Black Sea and the Mediterranean. *Mediterranean Marine Science*, 1(1), 133-141.
- Dobrovolo, I., Georgiev, Zh., Dobrovolo, S., 1995. Comparative electrophoretic investigations of the species of the family Gobiidae (Pisces) in the Bulgarian sector of the Black Sea. *Proceeding of the Institute of Fisheries-Varna*, 23, 48-68.
- Dobrovolo, I., Terzieva, P., 1995. Biochemical-genetic monitoring on the populations of three commercially important fish species in the Black Sea. p. 7-13. In: *4th Scientific Conference with International Participation "Ecology, economics and environment of the Black Sea area"*, 1-2 June, Varna, Bulgaria.
- Dobrovolo, I., Georgiev, Zh., 1995. Protein taxonomy of some atherinids (Pisces, Atherinidae) from the Black Sea and the Mediterranean. *Proceedings of the Institute of Fisheries-Varna*, 23, 36-47.
- Dobrovolo, I., Ivanova, P., 1999. Biochemical genetic comparison of the *Atherina boyeri* and *Atherina mochon pontica* (Pisces, Atherinidae). *Folia zoologica*, 48 (1), 55-60.
- Dobrovolo, I., Ivanova, P., Vasilev, V.P., Jonkov, J.I., 2003. Genetic divergence of mugilid fishes (Genus Mugilidae, Pisces) in the Bulgarian Black Sea coastal waters. In: p. 389-396. *Proceeding of the 30th International Conference Pacem in Maribus, a year after Johannesburg. Ocean Governance and Sustainable Development: Ocean and Coasts – a Climpse into the Future*, 27-30 October, 2003, Kiev, Ukraina.
- Dobrovolo, I., Ivanova, P., Apostolu, A., Manolov, Zh., 2003. Systematic of the Atherinidae species in the Black Sea and Mediterranean basins based on biochemical-genetic data. p. 721-728. In: *6th International Conference on the Mediterranean Coastal Environment, MEDCOAST 03, 7-11 October*. Ravenna, Italy.
- Dobrovolo, I., Cekov, A., Ivanova, P., Yankova, M., 2004. Identification on the black cavier of great sturgeon *Huso huso* and sterlet *Acipenser ruthenus* (Acipenseridae, Pisces), *National Center for Agrarian Science, Journal of Animal Science*, XLI (3), 30-32.
- Dobrovolo, I., Ivanova, P., Tzekov, A., 2005. Genetic-biochemical identification of some sturgeons and their hybrids (Pisces, Acipenseridae). *Verhandlungen Internationale Vereinigung für Theoretische und Angewandte Limnologie* 29 (2), 917-921.
- Dobrovolo, I., Ivanova, P., Georgiev, Zh., Panayotova, M., Raykov *et al.*, 2012. Allozyme variation and genetic identification of shad species (Pisces: Clupeidae, Genus Alosa) along the Bulgarian Black Sea coast. *Acta zoologica bulgarica* 64 (2), 175-183.
- Drensky, P., 1921. Contribution à l'étude de la faune de la Mer Noire, recoltés sur les cotes bulgares. *Annals of the Academy of Sciences*, (25), 6-12. (In French).
- Dumont, J.H., Mamaev, V.O., Zaitsev, Y.P., 1999. *Black Sea Red Data Book*. UNOPS Publishers, New York. 413 pp.
- Engin, S., Turan, D., Kovacic, M., 2007. First record of the red mouthed goby, *Gobius cruentatus* (Gobiidae), in the Black Sea. *Cybius*, 31, 87-88.
- Erazi, R.A.R., 1942. Marine fishes found in the Sea of Marmara and in the Bosphorus. *Revue de la Faculte des Sciences de l'universite D'Istanbul*, 7(1/2), 103-114.
- Erguden, D., Gurlek, M., Yaglioglu, D., Turan, C., 2010. Genetic Identification and Taxonomic Relationship of Mediterranean Mugilid Species Based on Mitochondrial 16S rDNA Sequence Data. *Journal of Animal and Veterinary Advances*, 9 (2), 336-341.
- Eschmeyer, W.N., Fongr, J.D., 2014. Species by family/subfamily. <http://research.calacademy.org/research/ichthyology/catalog/SpeciesByFamily.asp>. Electronic version accessed dd mmm 2014.
- Fischer, W. (Ed.), 1973. *Fiches FAO d'identification des espèces pour les besoins de la pêche*. Méditerranée et mer Noire (Zone de pêche 37). FAO, Rome, pag.var.
- Fischer, W., Bauchot, M.L., Schneider, M., 1987. *Fiches FAO d'identification des espèces pour les besoins de la pêche*. Méditerranée et mer Noire. Zone de pêche 37. FAO and EEC, Rome, 761-1530.
- Freyhof J., Kottelat, M., 2008. *Alosa immaculata*. In: IUCN 2010. IUCN Red List of Threatened Species.
- Fricke, R., Bilecenoglu, M., Sari, H.M., 2007. Annotated checklist of fish and lamprey species (Gnathostomata and Petromyzontomorpha) of Turkey, including a Red List of threatened and declining Species. *Stuttgarter Beiträge zur Naturkunde*, 706, 1-168.
- Geldiay, R., 1969. Izmir Körfezinin baslica balıkları ve muhtemel invasyonları [Important fishes found in the Bay of Izmir

- and their possible invasions]. Ege Üniversitesi Fen Fakültesi Monografileri, İzmir, 135 pp. (In Turkish).
- Georgiev, Zh., Kolarov, P., 1958. The twaite shad *Alosa fallax nilotica* Geoffroy (Clupeidae, Pisces) – a new fish for Bulgarian Fauna. *Proceeding of Institute of Zoology*, 7, 351-354.
- Georgiev, Z.M., Kolarov, P., 1962. On the migration and distribution of horse mackerel (*Trachurus mediterraneus ponticus*, Aleev) in the western part of Black Sea. *Arbeiten des Zentralen Forschungsinstitutes für Fischzucht und Fischerei –Varna*, 2, 148-172.
- Georgiev, G., Kolemanova, K.A., 1978. Ribite v Cherno more. p. 124-173 In: *Cherno more*. Valkanov, V., Marinov, H., Danov, H., Vladev, P. (Eds). Cherno more. George Bakalov Publishers, Varna, Bulgaria. (In Bulgarian).
- Grant, W., 2005. A second look at mitochondrial DNA variability in European anchovy (*Engraulis encrasicolus*): assessing models of population structure and the Black Sea isolation hypothesis. *Genetica*, 125, 293-309.
- Günther, A., 1866. *Catalogue of the fishes in the British Museum*. Vol.6. Order of the Trustees. London, UK, 512 pp.
- IUCN 2013. The IUCN Red List of Threatened Species. Version 2013.2. <http://www.IUCNredlist.org>. Downloaded on 21 November 2013.
- Ivanova, P., Dobrovolov, I., 2006. Population-genetic structure on European anchovy (*Engraulis encrasicolus* L. 1758) (Osteichthyes: Engraulidae) from Mediterranean basin and Atlantic Ocean. *Acta Adriatica*, 47 (1), 13-22.
- Ivanova, P., Dobrovolov, I., Bat, L., Kideys, A., Nikolsky, V. et al., 2013a. Application of esterase polymorphism to specify population genetic structure of *Engraulis encrasicolus* (Pisces: Engraulidae) in the Black and Azov Seas. *Marine Ecological Journals*, 12 (4), 45-52.
- Ivanova, P., Dobrovolov, I., Apostolou, A., Vasilev, M., Velkov, B., et al., 2013b. Protein Biomarkers for Identification of Some Gobiid Species (Actinopterygii: Gobiidae) along the Bulgarian Black Sea Coast *Acta Zoologica Bulgarica*, 65 (4), 429-438.
- Ivanova, P., Dobrovolov, I., Tsekov, A., 2014. Genetic differentiation between *Mullus barbatus* from north-western part of the Black Sea and *Mullus surmuletus* (Pisces, Mullidae) from Mediterranean. *Ecologia Balkanica*, in press Online ISSN: 1313-9940, Print ISSN: 1314-0213.
- Kabasakal, H., 1998. A note on the occurrence of the thresher shark, *Alopias vulpinus* from south-western Black Sea. *Journal of the Marine Biological Association of the United Kingdom*, 78 (2), 685-686.
- Kabasakal, H., 2013. Bluntnose sixgill shark, *Hexanchus griseus* (Chondrichthyes: Hexanchidae), caught by commercial fishing vessels in the seas of Turkey between 1967 and 2013. *Annales, Series Historia Naturalis*, 23 (1), 33-48.
- Karapetkova, M., Zhivkov, M., 2006. *Ribite v Bulgaria*. Geya Libris Publishers, Sofia, 190 pp. (In Bulgarian).
- Karapetkova, M., Zhivkov, M., 2010. *Ribite v Bulgaria*. Geya Libris Publishers, Sofia, 215 pp. (In Bulgarian).
- Keskin, Ç., 2010. A review of fish fauna in the Turkish Black Sea. *Journal Black Sea Mediterranean Environment*, 16 (2), 195-210.
- Kocataş, A., Ergen, Z., Mater, S., Özel, I., Katağan, T. et al., 1987. Marine Fauna. p. 141-161. In: *Biological diversity in Turkey*. Kence, A. (Eds). Environmental Problems Foundation of Turkey, Önder Matbaa Publishers, Ankara, Turkey.
- Kovačić, M., Engin, S., 2009. First record of the zebra goby, *Zeburus zeburus* (Gobiidae) in the Black Sea. *Cybius*, 33 (1), 83-84.
- Kovtun, O.A., 2013. The new finding of a rare goby *Chromogobius quadrivittatus* (Steindachner, 1863) (Actinopterygii, Perciformes, Gobiidae) in the marine underwater cave peninsula Tarhankut (Black Sea). *Marine Ecological Journal*, 12 (1), 18.
- Kovtun, O.A., Manilo, L.D., 2013. Mediterranean fish - *Gammogobius steinitzi* Bath, 1971 (Actinopterygii: Perciformes: Gobiidae) a new representative of the Black Sea ichthyofauna. *Acta ichthyologica et piscatoria*, 43 (4), 307-314.
- Kovtun, O.A., Karpova, E.P., 2014. *Chromogobius zebratus* (Kolombatovic, 1891) (Actinopterygii, Perciformes, Gobiidae) from marine underwater cave of Tarhankut peninsula (western Crimea), a species new for the Black Sea. *Marine Ecological Journal*, 13 (1), 72. (In Russian).
- Magoulas, A., Castilho, R., Caetano, S., Marcato, S., Patarnello, T., 2006. Mitochondrial DNA reveals a mosaic pattern of phylogeographical structure in Atlantic and Mediterranean populations of anchovy (*Engraulis encrasicolus*). *Molecular Phylogenetics and Evolution*, 39, 734-746.
- Mater, S., Meriç, N., 1996. Türkiye Omurgalıları Tür Listesi. p. 129-172. In: *Deniz Balıkları Pisces*. Kence, A., Bilgin, C., (Eds). Nural Matbaacılık A.Ş Publishers, Ankara, Turkey. (In Turkish).
- Mater, S., Bilecenoglu, M., 1999. Türkiye deniz balıkları [Marine fishes of Turkey]. p. 790-808. In: *Genel Zoocoğrafya ve Türkiye Zoocoğrafyası*. Demirsoy, A. (Ed), Meteksan Matbaası Publishers, Ankara, (In Turkish).
- Mezhzherin, S.V., Fedorenko, V.L., Verlatyi, B.D., 2009. Differentiation and allozyme variability of shads from the genus *Alosa* (Clupeiformes, Alosiinae) from Azov-Black Sea basin. *Cytology and Genetics*, 43 (2), 118-122.
- Movchan Yu.V., 2011. *Fishes of Ukraine*. Kiev Press “Zoloti vorota”, 420 pp. (In Ukrainian).
- Neilson, M.E., Stepien, C.A., 2009. Escape from the Ponto-Caspian: Evolution and biogeography of an endemic goby species flock (Benthophilinae: Gobiidae: Teleostei). *Molecular Phylogenetics and Evolution* 52, 84-102.
- Ninua, N.S., Japoshvili, B.O., 2008. Check list of fishes of Georgia. *Proceeding of Institute of Zoology- Georgian Academy of Sciences*, 23 (1), 163-176.
- Oral, M., Bat, L., Uysal, İ., 2013. Species sheets Part VI. Fish. pp. 118-283. In: Red Data Book Black Sea, Turkey. Öztürk, B., Oral, M., Topaloğlu, B., Bat, L., Okudan Arslan, E.Ş. et al., (Eds.). Published by Turkish Research Foundation (Tudav), Istanbul, TURKEY. Publication number 38, 323 pages, ISBN 978-975-8825-30-1.
- Oven, L., Salekhova, L., 1969. On the Black Sea ichthyofauna mediterraneanization. *Hydrobiological Journal*, 4, 124-127. (In Russian).
- Öztürk, B., 1999. *Black Sea Biological Diversity Turkey*. GEF Black Sea Environmental Programme. United Nations Publications, New York, USA, 228 pp.
- Rass, T.S., 1949. Ihtiofauna Chernogo morya i ee ispol'zovanie (Fish resources of the European seas of the USSR and possibilities of supplementing them by acclimatization). *Proceeding of Institute of Oceanology Academy of Sciences*, 4, 103-123. (In Russian).
- Rass, T.S., 1965. *Rybnye resursy Evropeiskikh morei SSSR i vozmozhnost' i ikh popolneniya akklimatizatsiei (Fish resources of the European seas of the USSR and opportunities to re-*

- plenish them by acclimatization). Nauka Publishers, Moscow, USSR, 108 pp. (In Russian).
- Samsun, O., Samsun, N., Kalayci, F., Bilgin, S., 2006. A study on recent variations in the population structure of European anchovy (*Engraulis encrasicolus* L., 1758) in the Southern Black Sea. E.U. *Journal of Fisheries and Aquatic Sciences*, 23 (3), 301–306.
- Sharvashidze, V., 1982. *Ribyi Gruzii (Fishes of Georgia)*. Ganatleba Publishers, Tbilisi, Georgia, 288 pp. (In Russian).
- Slastenenko, E., 1955-1956. *Karadeniz Havzasi Balıkları (The Fishes of the Black Sea Basin)*. Et Ve Balik Kurumu Publishers, Istanbul, Turkey, 711 pp. (In Turkish).
- Stepien, C., Tumeo, M., 2006. Invasion genetics of Ponto-Caspian gobies in the Great Lakes: a 'cryptic' species, absence of founder effects, and comparative risk analysis. *Biological Invasions*, 8, 61–78.
- Stefanov, T., 2007. Fauna and Distribution of Fishes in Bulgaria In: V. Fet, A. Popov A. (eds) *Biogeography and Ecology of Bulgaria*. Published by Springer, P.O. Box 17, 3300 AA Dordrecht, The Netherlands, 109-139.
- Stoyanov, S., Georgiev, Z., Ivanov, L., Hristov, D., Kolarov, P., Aleeva, K., Karapetkova, M., 1963. *Ribite v Cherno more (Fish in the Black Sea)*. Varna state Publishers, Sofia, Bulgaria, 98 pp. (In Bulgarian).
- Svetovidov, A.N., 1964. *Ribyi Chernogo moria (Fishes of the Black Sea)*. Nauka Publishers, Moscow, Leningrad, 525 pp. (In Russian).
- Tkachenko, P.V., 2013. First records of the *Diplodus sargus sargus* (Linnaeus, 1758) into the northwest part of the Black Sea. *Marine Ecological Journal*, 12 (3), 54. (In Russian).
- Tsekov, A., Ivanova, P., Angelov, M., Atanasova, S., Bloesch, J., 2008a. Natural sturgeon hybrids along the Bulgarian Black Sea coast and in the Danube River. *Acta Zoologica Bulgarica*, 60, 311–316.
- Tsekov, A., Ivanova, P., Dobrovolov, I. 2008b. First data for species identification of turbot along the Bulgarian Black Sea coast based on the genetic-biochemical analyses. *Acta Zoologica Bulgarica*, Suppl.2, 299-304.
- Tunesi, L., Molinari, A., 2005. Species richness and biogeographic outlines of the fish assemblage of the Portofino Marine Protected Area (Ligurian Sea). *Biologia Marina Mediteranea*, 12 (1), 116–123.
- Turan, C., Ergden, D., Gürlek, M., Başusta, N., Turan, F., 2004. Morphometric structuring of the anchovy (*Engraulis encrasicolus* L.) in the Black, Aegean and Northeastern Mediterranean Seas. *Turkish journal of veterinary and animal sciences*, 28, 865-871.
- Turan, D., Kottelat, M, Engin, S., 2009. Two new species of trouts, resident and migratory, sympatric in streams of northern Anatolia (Salmoniformes: Salmonidae). *Ichthyol. Explor. Freshwat.* 20 (4), 333-364.
- Turan, C., Caliskan, M., Kucuktas, H., 2005. Phylogenetic relationships of nine mullet species (Mugilidae) in the Mediterranean Sea. *Hydrobiologia*, 532, 45-51.
- Turan, C., Yağlıoğlu, D., Öztürk, B., Ergüden, D., Gürlek, M. et al., 2010. Morphometric and meristic analysis of Black Sea Shad (*Alosa immaculate* Bennett, 1835) populations in the Marmara Sea and Black Sea. p. 326-334. In: *Symposium the Marmara Sea, 25-26 September 2010*. Istanbul, Turkey.
- Ustaoglu Tiril, S., Zengin, M., Akbulut, B., Memis, D., Alagil, F. et al., 2011. A participatory approach to tagging and monitoring as an initial step in developing a sturgeon conservation strategy along the Turkish Black Sea Coast. *Journal of Applied Ichthyology*, 27 (2), 411-414.
- Valkanov, Al., 1961. Nature, Bulgarian academy of sciences, X, 1.
- Vasil'eva, E.D., 2007. *Ryby Chernogo morya*. Opredelitel' morskikh, solonovatovodnyh, evrigalinnyh i prohodnyh vidov s cvetnymi ilustraciami, sobrannymi Bogorodskim SV (*Fishes of the Black Sea*). Key to marine, brakish-water, eurihaline and anadromous species with color illustrations collected by Bogorodsky SV. VNIRO Publishers, Moscow, Russia, 238 pp. (In Russian).
- Vassilev, M., Apostolou, A., Velkov, B., Ivanova, P., Panayotova, M. et al., 2010. Status of Gobiid ichthyofauna (Gobiidae) in Bulgaria: Taxonomical, conservative, ecological and social aspects. In: *IV International symposium of ecologists of the Republic of Monte Negro, Budva, 6-9 October 2010*. Natura Montenegrina, Podgorica, 10 (2), 115-124.
- Vassilev, M., Apostolou, A., Velkov, B., Dobrev, D., Zarev, V., 2012. *Atlas of the gobies (Gobiidae) in Bulgaria*. Institute of Biodiversity and Ecosystem Research, Bulgarian Academy of Sciences, Bulgaria, 112 pp.
- Whitehead, P.J.P., Bauchot, M.L., Hureau, J.C., Nielsen, J., Tor-tonese, E. (Eds.) (1984 - 1986). *Fishes of the North-eastern Atlantic and the Mediterranean*. UNESCO, Paris, 1473 pp.
- Yankova, M., Pavlov, D., Ivanova, P., Karpova, E., Boltachev, A. et al., 2013. Annotated check list of the non-native fish species (Pisces) of the Black Sea. *Journal Black Sea/Mediterranean Environment*, 19 (2), 247-255.
- Zaitsev, Yu., Öztürk, B. (Ed.), 2001. *Exotic species in the Aegean, Marmara, Black, Azov and Caspian Seas*. Published by Turkish Marine Research Foundation. Istanbul, Turkey, 267 pp.
- Zivkov, M., Prodanov, K., Trichkova, T., Raikova-Petrova, G., Ivanova, P., 2005. Ribite v Balgariya - prouchenost, opazvane i ustojchivo izpolzvanie (Fishes in Bulgaria - research priorities, conservation and sustainable use). p. 247-282. In: *Current state of Bulgarian biodiversity – problems and perspectives*. Petrova, A. (Eds). Sofia; Bulgaria: Drakon Publishers, Sofia, Bulgaria. (In Bulgarian).