

Mediterranean Marine Science

Vol 15, No 3 (2014)

Vol 15, No 3 (2014)



Biology and population dynamics of by-catch fish species of the bottom trawl fishery in the western Mediterranean

F. ORDINES, M.T. FARRIOLS, J. LLEONART, B. GUIJARRO, A. QUETGLAS, E. MASSUTÍ

doi: [10.12681/mms.812](https://doi.org/10.12681/mms.812)

To cite this article:

ORDINES, F., FARRIOLS, M., LLEONART, J., GUIJARRO, B., QUETGLAS, A., & MASSUTÍ, E. (2014). Biology and population dynamics of by-catch fish species of the bottom trawl fishery in the western Mediterranean. *Mediterranean Marine Science*, 15(3), 613–625. <https://doi.org/10.12681/mms.812>

Supplementary Data

Biology and population dynamics of by-catch fish species of the bottom trawl fishery in the western Mediterranean

F. ORDINES, M. T. FARRIOLS, J.LLEONART, B. GUIJARRO,
A. QUETGLAS and E. MASSUTÍ

Mediterranean Marine Science, 2014, 15 (3), 613-625




Fig. S1: Species composition (in % of landed biomass) of the mixed fish category (MFC). “Other” are species representing less than 1%: *Trisopterus minutus*, *Zeus faber*, *Diplodus annularis*, *Trachinus radiatus*, *Helicolenus dactylopterus*, *Lepidotrigla dieuzeidei*, *Anthias anthias*, *Microchirus variegatus*, *Citharus linguatula*, *Microchirus ocellatus*, *Peristedion cataphractum*, *Callanthias ruber*, *Scorpaena spp.*, *Scorpaena elongata*, *Sparidae*, *Cepola macrophthalma*, *Phycis blennoides*, *Microchirus spp.*, *Serranus hepatus*, *Lophius piscatorius*, *Phycis phycis*, *Scorpaena lophei*, *Thalassoma pavo*, *Diplodus spp.*, *Gaidropsarus biscayensis*, *Coris julis*, *Illex coindetii*, *Solea spp.*, *Serranus scriba*, *Ophidion barbatum*, *Arnoglossus thori*, *Pagellus spp.*, *Bothus podas*, *Lophius budegassa*, *Lepidorhombus boscii*, *Scomber scombrus*, *Spicara smaris*, *Arnoglossus rueppelii*, *Synchiropus phaeton*, *Centracanthus cirrus*, *Pagellus bogaraveo*, *Merluccius merluccius*, *Arnoglossus imperialis*, *Engraulis encrasicolus*, *Liocarcinus depurator*.




Fig. S2: Quarterly evolution of the % of presence of opaque rings in the margin of the otoliths of the studied species. *Chelidonichthys cuculus*: continuous lines; *Trigloporus lastoviza*: dashed lines; *Serranus cabrilla*: dotted lines; *Trachinus draco*: dash-dotted lines.




Fig. S3: Fitting of von Bertalanffy growth function to age-length data of the studied species. *Chelidonichthys cuculus*: continuous lines; *Trigloporus lastoviza*: dashed lines; *Serranus cabrilla*: dotted lines; *Trachinus draco*: dash-dotted lines.

Table S1. Summary of the main biological parameters for each species. MED: Mediterranean, AT: Atlantic; TL range: total length range of the individuals studied; Age range: age range of the individuals studied; M: males, F: females, and T: both sexes.

| Species | Study Area | TL range (cm) | Age range (years) | | | L_{∞} | | | k | | | L_{50} | | | Reference |
|---------------------|----------------------|---------------|-------------------|----|----|--------------|------|------|------|---------|---------|----------|--------------------|----------------------------------|---------------------------------------|
| | | | T | F | M | T | F | M | T | F | M | T | F | M | |
| <i>C. cuculus</i> | Aegean Sea (MED) | | | 28 | 20 | | 0.22 | 0.51 | | | | | | Papaconstantinou, 1983 | |
| | Mediterranean | | | | | | | | | 19 | | | | Bauchot, 1987 | |
| | Gulf of Lion (MED) | 10-30 | 1-5 | 35 | | 0.28 | | | | | | | | Campillo, 1992 | |
| | Adriatic Sea (MED) | 10-36 | 1-5 | 29 | 29 | | 0.40 | 0.41 | | | | | | Marsan <i>et al.</i> , 1998 | |
| | Tyrrhenian Sea (MED) | 7-27 | 1-3 | 24 | 23 | | 0.74 | 0.59 | | | | 18 | | Colloca <i>et al.</i> , 2003 | |
| | Adriatic Sea (MED) | 10-26 | | | | | | | | | | 17 | 15 | Vallisneri <i>et al.</i> , 2012 | |
| | Douarnenez Bay (AT) | 14-45 | 1-21 | | 42 | 37 | | 0.46 | 0.52 | | | 28 | 27 | Baron, 1985a; 1985b | |
| | English Channel (AT) | 9-43 | 1-4 | 36 | 41 | 36 | 0.24 | 0.23 | 0.25 | | | | | Dorel, 1986; ICES, 2006 | |
| | Irish Sea (AT) | 10-43 | 1-7 | 42 | 41 | 41 | 0.21 | 0.24 | 0.21 | 0.28 | | 28 | 26 | Marriott <i>et al.</i> , 2010 | |
| | Gulf of Lion (MED) | | | | | | | | | | | | 19 < L_{50} < 21 | Kartas, 1971 | |
| <i>T. lastoviza</i> | Greek waters (MED) | | | | | | | | | | | 14 | | Papaconstantinou, 1986 | |
| | Egypt (MED) | | | | | | | | | 14 | | 15 | | Abdallah & Faltas, 1998 | |
| | Gulf of Gabès (MED) | | | | | | | | | | | 15 | | Boudaya, 2000 | |
| | Gulf of Tunis (MED) | 11-26 | | | | | | | | 16 | 16 | 16 | | Ben Jrad <i>et al.</i> , 2010 | |
| | Gulf of Gabès (MED) | | | | 33 | 30 | | 0.13 | 0.13 | | | | | Boudaya <i>et al.</i> , 2010 | |
| | Douarnenez Bay (AT) | | | | | | | | | | | | 29 | Baron, 1985 | |
| | Tunis (MED) | | | | | | | | | 10 (SL) | | | | Bouain, 1981 | |
| | Mediterranean | | | | | | | | | 15 (SL) | | | | Bauchot, 1987 | |
| | Aegean Sea (MED) | | | | 26 | 30 | | 0.30 | | | | | | | Papaconstantinou <i>et al.</i> , 1994 |
| | Greek waters (MED) | | 1-8 | 24 | | 0.30 | | | | | | | | | Politou & Papaconstantinou, 1995 |
| <i>S. cabrilla</i> | Cretan Shelf (MED) | 6-20 | 1-5 | 22 | | 0.39 | | | | | | | | Tserpes & Tsimenides, 2001 | |
| | Edremit Bay (MED) | 9-22 | 1-4 | 34 | | 0.11 | | | | | 15 (FL) | | | Torku-Koc <i>et al.</i> , 2004 | |
| | Aegean Sea (MED) | 7-23 | 1-6 | 24 | | 0.30 | | | | | 13 | | | İlhan <i>et al.</i> , 2010 | |
| | Canary Islands (AT) | | | | | | | | | 15 (SL) | | | | García-Díaz <i>et al.</i> , 1997 | |
| | Aegean Sea (MED) | 15-37 | | | | | | | | | 16 | 15 | | Kınacıgil <i>et al.</i> , 2008 | |
| | Black Sea (MED) | 9-26 | 1-6 | 29 | 33 | 29 | 0.28 | 0.18 | 0.17 | | | | | Ak & Genç, 2013 | |
| | Kattagat (AT) | 9-40 | 1-14 | 38 | 35 | | 0.15 | 0.16 | | | | | | Bagge, 2004 | |

References in Table S1

- Abdallah, M., Faltas, S.N., 1998. Reproductive biology of *Trigla lucerna* and *Trigloporus lastoviza* in the Egyptian Mediterranean water. *Bulletin of the National Institute of Oceanography and Fisheries*, 24, 285-303.
- Ak, O., Genç Y., 2013. Growth and reproduction of the greater weever (*Trachinus draco* L., 1758) along the eastern coast of the Black Sea. *Journal of the Black Sea/Mediterranean Environment*, 19, 95-110.
- Bagge, O., 2004. The biology of the greater weever (*Trachinus draco*) in the commercial fishery of the Kattegat. *ICES Journal of Marine Science*, 61, 933-943.
- Baron, J., 1985a. The Triglidae (Teleostei, Scorpaeniformes) of the Bay of Douarnenez. The growth of *E. gurnardus*, *T. lucerna*, *T. lastoviza* & *A. cuculus*. *Cybium*, 9, 127-144.
- Baron, J., 1985b. The Triglidae (Teleostei, Scorpaeniformes) of the Bay of Douarnenez. The reproduction of *E. gurnardus*, *T. lucerna*, *T. lastoviza* & *A. cuculus*. *Cybium*, 9, 252-281.
- Bauchot, M.L., 1987. Serranidae. P. 1317-1319. In: *Fiches FAO d'Identification des Espèces pour les Besoins de la Pêche Méditerranée et Mer Noire, Zone de Pêche 37 (Rev.1), Vol II, Vertébrés*. Fischer, W., Scheneider, M., Bauchot, M.L. (Eds.). FAO, Rome.
- Ben Jrad, L., Fehri-Bedoui, R., Ben Slama, S., Ben Hassine, O.K., 2010. Reproduction et régime alimentaire de *Trigloporus lastoviza* (Triglidae) dans le golfe de Tunis (Méditerranée occidentale). *Cybium*, 34, 353-365.
- Bouain, A., 1981. Les serrans (Téléostéens, Serranidés) des côtes sud de la Tunisie. Taille de première maturité, période de reproduction. *Cybium*, 4, 65-75.
- Boudaya, L., 2000. *Écobiologie de deux espèces de Triglidae (Pisces, Teleostei) du golfe de Gabès: Trigloporus lastoviza (Bonnaterre, 1788) et Aspitrigna obscura (Linnaeus, 1764)*. Mémoire de DEA. Université du Sud, Tunisie, 137pp.
- Campillo, A., 1992. *Les pêcheries françaises de Méditerranée: synthèse des connaissances*. Institut Français de Recherche pour L'exploitation de la Mer, France, 206 pp.
- Dorel, D., 1986. *Poissons de L' Atlantique nord est relations taille-poids*. Institut Français de Recherche pour L' exploitation de la Mer, France, 165 pp.
- García-Díaz, M.M., Tuset, V.M., González, J.A., Socorro, J., 1997. Sex and reproductive aspects in *Serranus cabrilla* (Osteichthyes: Serranidae): Macroscopic and histological approaches. *Marine Biology*, 127, 379-386.
- ICES, 2006. *Report of the working group on the assessment of new MOU species (WGNEW)*. ICES Headquarters, ICES Advisory Committee on Fisheries Management, CM2006 /ACFM: 11, 234 pp.
- Kartas F., 1971. *Les Triglidae de la mer Catalane, distribution, croissance et reproduction: genres Lepidotrigla, Trigloporus lastoviza, Eutrigla gurnardus*. MSc Thesis. University of Paris, France 173 pp.
- Kinacgil, H.T., İlkyaz, A.T., Metin, G., Ulaş, A., Soykan, O. et al., 2008. *Determining the first reproduction length, age and growth parameters of Aegean Sea demersal fish for the regulation of fisheries management*. Tübitak-Çaydag, 327 pp.
- Marriott, A.L., Latchford, J.W., McCarthy, I.D., 2010. Population biology of the red gurnard (*Aspitrigna cuculus* L.; Triglidae) in the inshore waters of Eastern Anglesey and North-west Wales. *Journal of Applied Ichthyology*, 26, 504-512.
- Marsan, R., Ungaro, N., Marzano, M.C., Martino, M., 1998. Growth of *Aspitrigna cuculus* (Osteichthyes, Triglidae) in the south-western Adriatic area: preliminary results. *Biologia Marina Mediterranea*, 5, 694-696. (in Italian).
- Papaconstantinou, C., 1983. Aspects of the biology of *Aspitrigna cuculus* (L., 1758) (Pisces, Scorpaeniformes) in the Gulf of Saronikos. *Thalassographica*, 6, 49-75.
- Papaconstantinou, C., 1986. The life history of rock gurnard (*Trigloporus lastoviza*, Brünn., 1768) in the Saronikos Gulf. *Journal of Applied Ichthyology*, 2, 75-86.
- Papaconstantinou, C., Politou, C.-Y., Caragitsou, E., Stergiou, K.I., Mytilineou, E. et al., 1994. *Investigations on the abundance and distribution of demersal stocks of primary importance in the Thermaikos Gulf and the Thracian Sea (Hellas)*. National Centre for Marine Research, Technical Report, North Aegean Sea Series 4/1994.
- Politou, C.-Y., Papaconstantinou, C., 1995. Age and growth of comber, *Serranus cabrilla* (L., 1785) in the Thracian Sea and the Thermaikos Gulf (Northern Greece). *Rapport de la Commission Internationale pour l'Exploration Scientifique de la Mer Méditerranée*, 34.
- Tserpes, G., Tsimenides, N., 2001. Age, growth and mortality of *Serranus cabrilla* (Linnaeus, 1758) on the Cretan shelf. *Fisheries Research*, 51, 27-34.
- Vallisneri, M., Montanini, S., Stagioni M., 2012. Size at maturity of triglid fishes in the Adriatic Sea, northeastern Mediterranean. *Journal of Applied Ichthyology*, 28, 123-125.