Mediterranean Marine Science

Vol 17, No 1 (2016)

VOL 17, No 1 (2016)

Diets of peracarid crustaceans associated with the orange coral *Astroides calycularis* in southern Spain

A. TERRÓN-SIGLER, D. LEÓN-MUEZ, P. PEÑALVER-DUQUE, F. ESPINOSA TORRE

doi: [10.12681/mms.1298](https://doi.org/10.12681/mms.1298)

To cite this article:

TERRÓN-SIGLER, A., LEÓN-MUEZ, D., PEÑALVER-DUQUE, P., & ESPINOSA TORRE, F. (2015). Diets of peracarid crustaceans associated with the orange coral *Astroides calycularis* in southern Spain. *Mediterranean Marine Science*, 17(1), 170–173. <https://doi.org/10.12681/mms.1298>

Diets of peracarid crustaceans associated with the orange coral *Astroides calycularis* in southern Spain

A. TERRÓN-SIGLER^{1,2}, D. LEÓN-MUEZ², P. PEÑALVER-DUQUE^{1,2} and F. ESPINOSA TORRE¹

¹ Departamento de Zoología, Facultad de Biología, Universidad de Sevilla, Avda. Reina Mercedes 6, 41012-Sevilla, Spain

² Asociación Hombre y Territorio. C/ Betania nº 13. CP. 41007 Sevilla, Spain

Corresponding author: terronsigler@hombreyterritorio.org

Handling Editor: Kostas Kapiris

Received: 6 March 2015; Accepted: 19 November 2015; Published on line: 22 February 2016

Abstract

The endangered and Mediterranean endemic orange coral (*Astroides calycularis*) hosts an important macrofaunal assemblage. The gut contents of the main peracarids associated with the orange coral were analysed. In total 161 specimens belonging to 11 species and 9 families were examined on the southern coast of the Iberian Peninsula. The gut content study was carried out introducing the specimens of each species in Hertwig's liquid. The analysis revealed that the peracarid species associated with *A. calycularis* had different feeding strategies and their main food source was detritus. The results highlight that peracarids may depend on the host and the detritus that the coral produces.

Keywords: *Astroides calycularis*, peracarid crustaceans, associated macrofauna, feeding, gut contents.

Introduction

Astroides calycularis (Pallas, 1766) is an azooxanthellate colonial scleractinian, with a calcium carbonate exo-skeleton (Zibrowius, 1980) that inhabits the rocky shores from the surface to a depth of 50 m (Rossi, 1971). It occupies both well-lit and dark habitats and appears to prefer environments with strong currents (Zibrowius, 1978; 1995; Kružić *et al.*, 2002). Its population density can be locally high, with colonies covering up to 90% of the sea bottom (Goffredo *et al.*, 2011). Populations of *A. calycularis* mainly have a limited geographical distribution in the Mediterranean Sea, restricted to the south-western area (Zibrowius, 1980; 1995; Bianchi, 2007). Though, the species has been also quoted in Adriatic Sea (Grubelić *et al.*, 2004). Andalusian rocky shores (southern Iberian Peninsula) host one of the highest densities of the orange coral populations (Terrón-Sigler *et al.*, 2015).

Terrón-Sigler *et al.* (2014) observed that peracarid crustaceans were the most prevalent species associated with the orange coral. Additionally, three of the peracarids associated with *A. calycularis* had a strong orange colour (i.e. *Janira maculosa*, *Sthenothoe cavimana*, and *Caprella penantis*) (Terrón-Sigler *et al.*, 2014). However, the nature of the association is unclear, whether it is obligate or facultative, and whether the coral is used as a substrate (e.g. refuge, spawning, brood care or protection by crypsis) or as a feeding source (partial or totally).

Amphipods are the most diverse group of crustaceans in terms of lifestyle, trophic types and habitats (De Broyer & Jazdzewski, 1996). Due to their population's

characteristics, amphipods may play major roles in the ecology of their habitats (Conlan, 1994). Amphipods are known to have versatile feeding strategies (i.e. Sarvala & Uitto, 1991) and show a wide range of feeding habits (Conradi & Cervera, 1995). They also constitute an important food source for a large variety of marine predators (Vázquez-Luis *et al.*, 2013); hence playing a key role in energy flow through food webs (Guerra-García *et al.*, 2014). Consequently, the main objective of the present study is to explore if peracarids associated with *A. calycularis* feed on the host coral's tissues, in an effort to highlight the nature of the association.

Material and Methods

161 specimens of peracarids were collected from 23 colonies of *A. calycularis* from Marina del Este beach (Granada coast; Andalusia; Spain) in July 2011 (Figure 1). Colonies were covered *in situ* with plastic bags before they were removed in order to keep all the associated fauna. The samples were fixed in 4% formalin and subsequently placed in 70% ethanol in the laboratory. The colonies were washed, the macrofauna sieved using a 0.5 mm size mesh, and the specimens collected were sorted and identified at the species level, when possible.

Diet study was analysed following the methodology proposed by Bello & Cabrera (1999) with slight variations. Recently, this method was successfully used in amphipods (i.e. Navarro-Barranco *et al.*, 2013; Vázquez-Luis *et al.*, 2013; Guerra-García *et al.*, 2014).

Fig. 1: Map of the study site showing Granada coast (Andalusia, Spain) and Marina del Este beach, where the colonies of *Astroides calycularis* were collected (* = 36°43'15" N; 3°43'43" W).

Specimens of each species were introduced in vials with Hertwig's liquid (consisting of 270 g of chloral hydrate, 19 ml of chloridric acid 1 N, 150 ml of distilled water and 60 ml of glycerine) and oven-heated at 65 °C for 4 to 6 hours depending on the cuticle thickness of the specimens. After this, samples were mounted on slides for microscope study. The percentage of absolute gut content (i.e. total area occupied by the content in the whole digestive tract) and the relative gut content (i.e. area occupied by each component within the total gut content) were estimated using a microscope equipped with an ocular micrometer (at 40× or 100×). Mean and standard error of the mean were calculated.

Results and Discussion

The collected peracarid specimens belonged to 11 species and distributed in 9 families (Table 1), representing the dominant peracarids associated with *A. calycularis* in the south Iberian Peninsula. Gut contents of the studied amphipod species included detritus, crustaceans and macroalgae (Table 1). The dominant component was detritus, followed by crustacean pieces and macroalgal tissues. In a diet analysis of marine amphipods around the Iberian Peninsula, Guerra-García *et al.* (2014) concluded that detritivorous species had fuller guts than

carnivorous species, where empty guts predominated. Nevertheless, trophic classifications may be subjective. In our study, three species namely *Liljeborgia psaltrica*, *S. cavimana* and *Eusiroides dellavallei*, presented empty guts. Although, these three species might be considered carnivorous, the amphipods associated with *A. calycularis* colonies presented the majority of the gut content occupied by detritus, highlighting the importance of detritus for the amphipods already noted in the Iberian Peninsula by Guerra-García *et al.* (2014).

Gut content ranged from 16% in *Leucothoe spinicarpa* to 64.3% in *J. maculosa*. In general, species with detritus had a higher area occupied by content in the digestive tract than other species.

Regarding caprellid species, these findings are in agreement with other studies, where the detritus was the dominant food source in seaweed-associated caprellids (Guerra-García & Tierno de Figueroa, 2009; Guerra-García *et al.*, 2014). In the present study, the two caprellids collected from orange coral colonies showed more than 98% detritus composition in their gut contents.

There is scarce information about the feeding habits of marine isopods (e.g. Navarro-Barranco *et al.*, 2013). In this study, *Paragnathia formica* showed 100% detritus gut content, while *J. maculosa* had a diverse gut content (Table 1).

Table 1. Gut contents of the studied peracarid species. N: number of specimens of each species examined, n: number of specimens with detected digestive contents. % Abs: total area occupied by the content in the whole digestive tract. Det: detritus, Crust: crustaceans, Malg: Macroalgae. Mean values with standard errors of the mean (in parentheses) are included.

PERACARIDS	N/n	% Abs	% Det	% Crust	% Malg	Feeding group
<i>Isopods</i>						
<i>Janira maculosa</i> Leach, 1814	22/22	64.3 (19.1)	93.9 (8.6)	5.7 (3.1)	0.5	Omnivorous
<i>Paragnathia formica</i> Hesse, 1864	12/10	27.5 (15.5)	100	--	--	Detritivorous
<i>Amphipods</i>						
<i>Lembos</i> sp. Bate, 1857	23/22	50.2 (30.7)	92.7 (6.3)	7.3 (5.2)	--	Omnivorous
<i>Autonoe spiniventris</i> Della Valle, 1893	9/8	59.4 (39)	100	--	--	Detritivorous
<i>Leucothoe spinicarpa</i> Abildgaard, 1789	19/17	16 (9.1)	98.1 (3.1)	1.9 (0.8)	--	Omnivorous
<i>Liljeborgia psaltrica</i> Krapp-Schickel, 1975	20/0	--	--	--	--	Carnivorous
<i>Sthenothoe cavimana</i> Chevreux, 1908	20/0	--	--	--	--	Carnivorous
<i>Eusiroides dellavallei</i> Chevreux, 1899	3/0	--	--	--	--	Carnivorous
<i>Caprella acanthifera</i> Leach, 1814	14/14	61.1 (3.7)	100	--	--	Detritivorous
<i>Phtisica marina</i> Slabber, 1769	11/8	19.4 (13.9)	98.7 (3.5)	1.3	--	Omnivorous
<i>Tanaids</i>						
<i>Tanais dulongii</i> Audouin, 1826	8/7	55.7 (28.7)	98.8 (3.7)	1.2	--	Omnivorous

The tanaid *Tanais dulongii* associated with *Fucus spiralis* in the south Iberian peninsula is herbivorous with fucoid algae its primary diet source (Torrecilla-Roca & Guerra-García, 2012). The specimens associated with *A. calycularis* did not show this feeding habit; more than 98% of the gut content was detritus and the remaining was composed of crustacean fragments. This finding highlights the plasticity in the feeding habits of peracarid species depending on the host or habitat where they inhabit.

Scleractinian corals function as structural engineers (Jones *et al.*, 1994), contributing to an increase in habitat complexity and surface topography, which promotes biodiversity by mediating competition or predation (Hixon & Menge, 1991; Coker *et al.*, 2009). Corals provide a large surface area where invertebrate assemblages can live, as well as refuges from predator, food source in the form of coral tissue, mucus and its associated detritus, and a hard skeleton used as a substratum by specialized burrowers (Castro, 1988). Taking all these factors into account, the peracarids associated with *A. calycularis* might use the coral skeleton as a refuge against potential predators (by hiding or as crypsis) or as food source of detritus that the colonies generate. Commensalisms between amphipods and cnidarians has been documented worldwide (e.g. Vader & Krapp-Schickel, 1996; Esquete *et al.*, 2014), but there is little information about this association with corals. Therefore, more studies are necessary in this regard with scleractinian corals. The analysis of gut contents of peracarids associated with the intertidal algae *F. spiralis* revealed that the assemblage is dominated by species feeding mainly on this algae (Torrecilla-Roca & Guerra-García, 2012), in contrast with other crustaceans inhabiting different seaweed assemblages which mainly feed on detritus (Guerra-García & Tierno de Figueroa, 2009; Alarcón-Ortega *et al.*, 2012). This could highlight that, depending upon the host with which the peracarids are associated, the relationship could be host-dependent

or not, but further research needs to be carried out in this regard.

Acknowledgements

Our research was supported by the Regional Government of Andalusia (General Office of Environment Management, Regional Ministry of Environment). We thank the General Office of Marine Protected Area from the Spanish state (National Ministry of Environment). Financial support for this work was provided by the Asociación Hombre y Territorio (<http://www.hombreyterritorio.org>) and the Laboratorio de Biología Marina (Departamento de Zoología, Facultad de Biología, Universidad de Sevilla). Special thanks to our colleague J. M. Guerra-García for his assistance with laboratory methods and taxonomic identifications, and Manuel González O'Sullivan and Paul Worrall for their assistance during the English review process. The experiments complied with current Spanish laws.

References

- Alarcón-Ortega, L.C., Guerra-García, J.M., Sánchez-Moyano, J.E., Cupul-Magaña, F.G., 2012. Feeding habits of caprellids (Crustacea: Amphipoda) from the west coast of Mexico. Do they feed on their hosting substrates? *Zoologica Baetica*, 23, 11-20.
- Bello, C.L., Cabrera, M.I., 1999. Uso de la técnica microhistológica de Cavender y Hansen en la identificación de insectos acuáticos. *Boletín de Entomología Venezolana*, 14, 77-79.
- Bianchi, C.N., 2007. Biodiversity issues for the forthcoming tropical Mediterranean Sea. *Hydrobiologia*, 580, 7-21.
- Castro, P., 1988. Animal symbioses in coral reef communities: a review. *Symbiosis*, 5, 161-184.
- Coker, D.J., Pratchett, M.S., Munday, P.L., 2009. Coral

- bleaching and habitat degradation increase susceptibility to predation for coral-dwelling fishes. *Behavioral Ecology*, 20, 1204-1210.
- Conlan, K.E., 1994. Amphipod crustaceans and environmental disturbance: a review. *Journal of Natural History*, 28, 519-554.
- Conradi, M., Cervera, L., 1995. Variability in trophic dominance of amphipods associated with the bryozoan *Bugula neritina* (L., 1758) in Algeciras Bay (Southern Iberian Peninsula). *Polskie Archiwum Hydrobiologii*, 42, 483-494.
- De Broyer, C., Jazdzewski, K., 1996. Biodiversity of the Southern Ocean: towards a new synthesis for the Amphipoda (Crustacea). *Bolletino del Museo Civico di Storia Naturale di Verona*, 20, 547-568.
- Esquete, P., Sibaja-Cordero, J., Troncoso, J.S., 2014. A new genus and species of Leptocheliidae (Crustacea: Peracarida: Tanaidacea) from Isla del Coco (Costa Rica). *Zootaxa*, 3741 (2), 228-242.
- Goffredo, S., Caroselli, E., Gasparini, G., Marconi, G., Putignano, M.T. *et al*, 2011. Colony and polyp biometry and size structure in the orange coral *Astroides calycularis* (Scleractinia: Dendrophylliidae). *Marine Biology Research*, 7 (3), 272-280.
- Grubelic, I., Antolic, B., Despalatovic, M., Grbec, B., Beg Paklar, G., 2004. Effect of climatic fluctuations on the distribution of warm-water coral *Astroides calycularis* in the Adriatic Sea new records and review. *Journal of the Marine Biological Association of the United Kingdom*, 84, 599-602.
- Guerra-García, J.M., Tierno de Figueroa, J.M., Navarro-Barranco, C., Ros, M., Sánchez-Moyano, J.E. *et al.*, 2014. Dietary analysis of the marine Amphipoda (Crustacea: Peracarida) from the Iberian Peninsula. *Journal of Sea Research*, 85, 508-517.
- Guerra-García, J.M., Tierno de Figueroa, J.M., 2009. What do caprellids feed on? *Marine Biology*, 156, 1881-1890.
- Hixon, M.A., Menge, B.A., 1991. Species diversity: prey refuges modify the interactive effects of predation and competition. *Theoretical Population Biology*, 39, 178-200.
- Jones, C.G., Lawton, J.H., Shachak, M., 1994. Organisms as ecosystem engineers. *Oikos*, 69, 373-386.
- Kruzic, P., Zibrowius, H., Pozar-Domac, A., 2002. Actiniaria and Scleractinia (Cnidaria, Anthozoa) from Adriatic Ser (Croatia): first records, confirmed occurrences and significant range extensions of certain species. *Italian Journal Zoology*, 69, 345-353.
- Navarro-Barranco, C., Tierno de Figueroa, J.M., Guerra-García, J.M., Sánchez Tocino, L., García-Gómez, J.C., 2013. Feeding habits of amphipods (Crustacea: Malacostraca) from shallow soft bottom communities; comparison between marine caves and open habitats. *Journal of Sea Research*, 78, 1-7.
- Rossi, L., 1971. Cnidari e Ctenofori d'Italia. *Cuaderni della Civica Stazione Idrobiologica di Milano*, 2, 77-86.
- Sarvala, J., Uitto, A., 1991. Production of the benthic amphipod *Pontoporeia affinis* and *P. femorata* in a Baltic archipelago. *Ophelia*, 34, 71-90.
- Terrón-Sigler, A., Peñalver, P., León-Muez, D., Espinosa, F., 2014. Spatio-temporal macrofaunal assemblages associated with the endangered orange coral *Astroides calycularis* (Scleractinia: Dendrophylliidae). *Aquatic Biology*, 21, 143-154. doi: 10.3354/ab00577.
- Terrón-Sigler, A., León-Muez, D., Peñalver, P., Gálvez-César, R., Espinosa Torre, F., 2015. Geographic distribution of *Astroides calycularis* (Scleractinia: Dendrophylliidae) as a baseline to assess future human impacts on the Southern Iberian Peninsula. *Journal of the Marine Biological Association of the United Kingdom*, 1-9. doi:10.1017/S0025315415001113.
- Torrecilla-Roca, I., Guerra-García, J.M., 2012. Feeding habits of the peracarid crustaceans associated with the alga *Fucus spiralis* in Tarifa Island, Cádiz (Southern Spain). *Zoologica Baetica*, 23, 39-47.
- Vader, W., Krapp-Schickel, G., 1996. Redescription and biology of *Sthenothoe brevicornis* Sars (Amphipoda: Crustacea), an obligate associate of the sea anemone *Actinostola callosa* (Verrill). *Journal of Natural History*, 30 (1), 51-66.
- Vázquez-Luis, M., Sánchez-Jerez, P., Bayle-Sempere, J.T., 2013. Does the invasion of *Caulerpa racemosa* var. *cylindracea* affect the feeding habits of amphipods (Crustacea: Amphipoda). *Journal of the Marine Biological Association of the United Kingdom*, 93, 87-94.
- Zibrowius, H., 1978. Les Scléractiniaires des grottes sous-marines en Méditerranée et dans l'Atlantique nord-oriental (Portugal, Madeïre, Canaries, Açores). *Pubblicazioni della Stazione zoologica di Napoli. Marine Ecology*, 40, 516-45.
- Zibrowius, H., 1980. Les Scléractiniaires de la Méditerranée et de l'Atlantique nord-orinetal. *Memories del'Institut Océanographique Monaco* 11, 198-201.
- Zibrowius, H., 1995. The Southern *Astroides calycularis* in the Pleistocene of the Northern Mediterranean – An indicator of climatic changes (Cnidaria, Scleractinia). *Geobios*, 28, 9-16.