Mediterranean Marine Science

Vol 17, No 2 (2016)

Bryozoa from the Mediterranean coast of Israel

N. SOKOLOVER, P. D. TAYLOR, M. ILAN

doi: [10.12681/mms.1390](https://doi.org/10.12681/mms.1390)

To cite this article:

SOKOLOVER, N., TAYLOR, P. D., & ILAN, M. (2016). Bryozoa from the Mediterranean coast of Israel. *Mediterranean Marine Science*, 17(2), 440–458. <https://doi.org/10.12681/mms.1390>

Supplementary Data
Bryozoa from the Mediterranean coast of Israel

N. SOKOLOVER, P.D. TAYLOR and M. ILAN
Mediterranean Marine Science, 2016, 17 (2)

Appendix

Suppl. Fig. 1: *Thalamoporella harmelini* (TAU-BR25039). **A.** Living colony in its natural habitat. **B.** Part of a colony. **C.** Edge of branch with marginal kenozooids (left). **D.** Group of autozooids and two avicularia. **E.** Three caliper spicules. **F.** Compass spicule. Scale bars: 1 mm (B); 500 μ m (C, D); 20 μ m (E); 10 μ m (F).

Suppl. Fig. 2: *Crisularia plumosa* (TAU-BR25023). **A.** Part of a colony. **B.** Pedunculate avicularium (left) and ovicell (right). **C.** Abfrontal view. **D.** Living colony in natural habitat. Scale bars: 500 μ m (A, C); 20 μ m (B).

Suppl. Fig. 3: *Drepanophora birbira*. **A.** Colony view (TAU-BR25055). **B.** Young colony, no secondary calcification (TAU-BR25056). **C.** Orifice with denticle and avicularium (TAU-BR25057). **D.** Ovicell (TAU-BR25055). Scale bars: 500 μ m (A, B); 20 μ m (C, D).

Suppl. Fig. 4: *Microporella harmeri*. **A.** Part of colony with ovicellate zooids (TAU-BR25080). **B.** Oblique view of infertile autozooids and avicularia (TAU-BR25080). **C.** Unbleached autozooid showing avicularium and open mandible, orifice with in-situ operculum, and 4 oral spines (TAU-BR25079). **D.** Unbleached autozooid showing avicularium with closed mandible (TAU-BR25079). Scale bars: 500 μ m.

Suppl. Fig. 5: *Mucropetraliella thenardii*. **A.** Part of a colony (TAU-BR25026). **B.** Ancestrula and periancestrular zooids (TAU-BR25026). **C.** Ovicell and mucro supporting five avicularia (TAU-BR25089). **D.** Orifice with two lateral avicularia, broken mucro (TAU-BR25089). Scale bars: 500 μ m (A, B, C); 20 μ m (D).

Suppl. Fig. 6: *Trematooeicia ligulata*. **A.** Three autozooids and an interzooidal avicularium (right) (TAU-BR25028). **B.** Orifice with suboral avicularium (TAU-BR25027). **C.** Part of a colony (TAU-BR25027). **D.** Living colony (stereoscope view) (TAU-BR25027). Scale bars: 100 μ m (A, B); 30 μ m (C).

Suppl. Fig. 7: *Schizoretepora hassi* (TAU-BR25082). **A.** Part of a colony with ovicellate zooids. **B.** Giant avicularium. **C.** Abfrontal view. **D.** Part of a living colony (stereoscope view). Scale bars: 100 μ m (A, B); 200 μ m (C).