



Mediterranean Marine Science

Vol 17, No 2 (2016)



High xenodiversity versus low native diversity in the south-eastern Mediterranean: bryozoans from the coastal zone of Lebanon

J. G. HARMELIN, G. BITAR, H. ZIBROWIUS

doi: 10.12681/mms.1429

To cite this article:

HARMELIN, J. G., BITAR, G., & ZIBROWIUS, H. (2016). High xenodiversity versus low native diversity in the southeastern Mediterranean: bryozoans from the coastal zone of Lebanon. *Mediterranean Marine Science*, *17*(2), 417–439. https://doi.org/10.12681/mms.1429

High xenodiversity versus low native diversity in the south-eastern Mediterranean: bryozoans from the coastal zone of Lebanon

J.G. HARMELIN¹, G. BITAR² and H. ZIBROWIUS³

 ¹Aix-Marseille Université, France, GIS Posidonie & Mediterranean Institute of Oceanography (MIO), OSU Pytheas, Station Marine d'Endoume, 13007 Marseille
 ²Department of Natural Sciences, Faculty of Sciences, Lebanese University, Hadath, Lebanon
 ³Le Corbusier 644, 280 Boulevard Michelet, 13008 Marseille, France Corresponding author: jean-georges.harmelin@univ-amu.fr

Handling Editor: Antonietta Rosso

Received: 18 July 2015; Accepted: 12 February 2016; Published on line: 21 Arpil 2016

Abstract

Because of its location in the warmest corner of the Mediterranean, its proximity to the northern entrance of the Suez Canal, i.e. the gateway for massive exotic biota introduction into the Mediterranean, the occurrence of intensive shipping activity and strong human pressure, the Lebanese coastal zone is an area that is exceptionally well-suited for investigating the effects of these extreme conditions on Mediterranean biodiversity.

Bryozoans, which are among the main components of sessile communities, dramatically reflect the impact of these particular conditions. Their assemblages, sampled by diving along the whole coast of Lebanon during a pluriannual programme, mainly between 1999 and 2003, consist of 93 species (12 Cyclostomata, 2 Ctenostomata, 79 Cheilostomata). The native part of this bryozoan fauna exhibits low diversity, with an unexpected absence of many taxa, from family to species level, which are very common in the rest of the Mediterranean. It is also characterized by a high proportion of endemic species, in contrast with the general eastward trend of decreasing endemicity observed in the Mediterranean, and by the strong presence of 'southern' thermophilic species.

With 27 non-indigenous species, the xenodiversity is exceptionally high, particularly in the cheilostome pool (26 species), but was probably undersampled. Moreover, one may assume that new non-indigenous bryozoans (NIB) are now established along the Levantine coasts. This trend is expected to increase in the near future with the intensification of surface water warming and the increase in shipping activity and propagule flux generated by the expansion of the Suez Canal.

Keywords: Biodiversity, biogeography, biological invasions, Bryozoa, climatic changes, endemism, Levantine basin, Suez Canal.

Introduction

The Mediterranean Sea is deemed to host a very high level of biodiversity for its size (Bianchi & Morri, 2000; Boudouresque, 2004; Coll et al., 2010; Bianchi et al., 2012), but with contrasting local situations and a decreasing gradient from the western to the eastern Mediterranean (Fredj, 1974; Boudouresque, 2004; Bianchi, 2007; Coll et al., 2010). Overall, Mediterranean species richness stems essentially from the combination of a complex paleogeographical history and a wide variety of subregional climatic conditions, resulting in the presence of numerous endemics and of biota of different biogeographical origins established during successive phases of the geological history of the Mediterranean (Pérès, 1967, 1985; Bianchi & Morri, 2000; Taviani, 2002). However, there is a wide disparity in knowledge of the biodiversity between Mediterranean subregions and taxonomic groups, with southern areas and some invertebrate groups, such as Bryozoa, remaining poorly investigated. Furthermore, the status of Mediterranean biodiversity needs constant re-assessment as its rate of change over time is increasing dramatically as a result of human stressors acting on coastal and offshore ecosystems, climate warming and introduction of non-indigenous species (Bianchi *et al.*, 2012).

The Mediterranean Sea is an exceptionally rich recipient area of non-indigenous species, mainly fuelled by three anthropogenic pathways of introduction of alien species, i.e. aquaculture, shipping and the Suez Canal (e.g. Zibrowius, 1992, Streftaris et al., 2005; Zenetos et al., 2012; Galil & Goren, 2014). The opening in 1869 of a connection between the Red Sea and the Mediterranean via the Suez Canal triggered an extensive colonization of the Mediterranean, particularly the eastern basin, by 'lessepsian' species belonging to most groups of flora and fauna (e.g. Por, 1990; Boudouresque, 1999a; Galil, 2007; Zenetos et al., 2012). This trend is accelerating as a result of hydrological changes in the Canal and the general warming of the Mediterranean that favours thermophilic species (Boudouresque, 1999a; Occhipinti-Ambrogi, 2007; Lejeusne et al., 2009; Bitar, 2008, 2010, Por. 2010).

The Levantine basin is the warmest corner of the Mediterranean (Abboud-Abi Saad et al. 2004; Taupier-

Letage, 2008), a condition exacerbated by global warming, which shapes the marine communities, entails the exclusion of shallow-water biota intolerant of warm temperatures, such as the sea-grass Posidonia oceanica (Celebi et al., 2006) and increases the invasibility of the coastal zone by thermophilic alien species (Occhipinti-Ambrogi, 2007; Lejeusne et al., 2009; Occhipinti-Ambrogi & Galil, 2010). The contiguity of the Levantine coasts with the northern entrance of the Suez Canal and the favourable circulation of surface waters (Hamad et al., 2006) are other fundamental drivers of the intense colonization by lessepsian species observed along these coasts (e.g., Por, 1990; Galil, 2007). This process, in constant progression, may controversially be viewed as a rejuvenation of the Mediterranean with a return of the tropical Tethys conditions (Por, 2010). The present status of coastal marine communities at sites of interest for conservation in Lebanon was recently surveyed (RAC/ SPA - UNEP/MAP, 2014) but, unfortunately, the baseline data on their past condition are very poor.

Rates and timing of colonization by alien species reported from different Mediterranean sub-regions in review papers are often biased by wide disparities in study effort and levels of expertise locally devoted to these issues (e.g. Nunes *et al.*, 2014). In addition, as stressed by Streftaris *et al.* (2005) and Por (2010), there is an imbalance in knowledge of the biological invasion of the Mediterranean Sea between taxonomic groups and some species are listed on the basis of unverified records (Zibrowius, 1992). The same remark also applies to the native elements of the same groups in the eastern Mediterranean, and particularly the Levantine basin.

Bryozoans are major components of benthic communities, displaying a wide variety of colony shapes, sizes and life history strategies. They also are among the main fouling organisms colonizing artificial substrata (Ryland, 1965; Gordon & Mawatari, 1992; Karlson & Osman, 2012). Because of particular traits of their morphology, life-history and ecology, many species are especially prone to dispersal outside their native geographical range by man-induced vectors (Gordon & Mawatari, 1992; Watts et al., 1998). Fouling on immerged surfaces of vessels is the most common mode of transport by shipping but larval transfer via ballast water is assumed to be effective for species producing long-lived cyphonautes larvae (Carlton, 1985). Translocation may also occur by means of rafting, i.e. settlement on drifting items, such as plastic debris (Winston et al., 1977; Barnes & Milner, 2005).

The actual magnitude of the non-indigenous bryozoan species in the Mediterranean as a whole is still imperfectly known and is underestimated (Zenetos *et al.*, 2012; Por, 2010). In addition, some old, poorly documented records need to be critically re-examined. According to a recent review (Zenetos *et al.* 2012), 31 alien bryozoan species have been recorded in the Mediterranean as a whole and 22 species in the eastern Mediterranean, the main pathways of their introduction being shipping and the Suez Canal.

Compared with the western Mediterranean, knowledge of the native bryozoan fauna of the eastern Mediterranean is deficient and more focused on the Aegean Sea (e.g. Harmelin, 1968-1969; Hayward, 1974, 1975; Ünsall, 1975; Ünsall & d'Hondt, 1978-1979; Morri et al., 1999; Koçak & Aydin Önen, 2014) than on the Levantine Basin (e.g. O'Donoghue & de Watteville, 1939; Gautier, 1956; d'Hondt, 1988; Koçak et al., 2002; Abdelsalam & Ramadan, 2008; Abdelsalam, 2014; Sokolover, 2014). Non-indigenous bryozoans present in the Levantine basin were recorded from Egypt (Hastings, 1927), Israel (Powell, 1969; Eitan, 1972a; Goren, 1980; d'Hondt, 1988; Sokolover, 2014), Turkey (Ünsal, 1975; Ünsal & d'Hondt, 1978-1979; Koçak, 2007), North Cyprus (Koçak et al, 2002) and Lebanon (Bitar & Kouli-Bitar, 2001 and references below).

This study provides an inventory of the bryozoan assemblages from the whole coastal zone of Lebanon collected in the context of field surveys spread over several years. The studied collection has first given rise to several taxonomic revisions (Harmelin, 2014 a-b; Harmelin et al., 2007, 2009, 2011, 2012). These previous works provided the basis for clarifying several taxonomic uncertainties, describing new taxa and highlighting the occurrence of many non-indigenous species never previously recorded in the Mediterranean. The present work provides new insights on the diversity and distribution of both native and exotic bryozoans in a region where knowledge on this group was very poor and which highlights the specificity of this corner of the Mediterranean, strongly impacted by the combined effects of non-indigenous species and the tropicalization of the Mediterranean.

Materials and Methods

The present bryozoan collection was provided by field surveys of the marine benthos carried out mainly between 1999 and 2003 within the framework of the French-Lebanese programme of scientific cooperation CEDRE. Sampling was performed by the authors by SCUBA diving between 2 m and 42 m depth from Tripoli (north) to Tyre (south), i.e. along ca. 80 % of the 225 km-long coast of Lebanon. Bryozoans were collected at 35 stations distributed in 18 localities (Table 1, Fig. 1). These localities are distributed to the north (loc. 1-9) and the south (loc. 10-18) of the 34° parallel, which delimits arbitrarily two regions in the present study. The northern region includes Tripoli, a major urban centre with a relatively active port and offshore uninhabited islands, and two towns, Batroun and Jbail (Byblos). The southern region comprises the city of Beirut, with satellite towns and a large port, and the towns of Saida and Tyre, with smaller harbours. For analysis of the species distribution, the northern and southern localities were grouped into two areas each: the Tripoli area (loc. 1-3), Batroun area



Fig. 1: Geographical distribution of the 18 localities sampled for bryozoans in four areas along the Lebanese coast. Number of non-indigenous bryozoan species recorded in these areas and their percentage in the local cheilostome pool.

(loc. 4-9), Beirut area (loc. 10-15) and Tyre area (loc. 16-18). Each species was coded according to four classes corresponding to the frequency of occurrence in the 18 sampled localities (class A: > 50 %; class B: 26-50 %; class C: 10-25 %; class D: < 10 %). Sampled stations were categorised *a posteriori* into four habitat types: (i) ports: piers, quays, jetty blocks; (ii) shaded habitats: cavities, overhangs on natural hard bottoms; (iii) exposed rock walls: upper and lateral surfaces of rocks; (iv) small substrata, biogenic concretions. This categorisation of habitat types reveals differences between the northern and southern batches of samples, with in particular a higher contribution of port habitats in the south (50.0 % vs. 10.5 % of stations). The depth range and mean depth $(\pm$ standard deviation) of the sampled stations, on the other hand, were not significantly different between the two regions (north: 3-35 m, 14.9 ± 9.9 m; south: 3-42 m, 12.9 ± 11.5 m). The distribution of sampled stations in four depth ranges shows that sampling effort was roughly equal at < 5 m, 16-25 m and > 25 m, while it was greater in the 5-15 m depth range. An approximate indication of the abundance of species in each sample was given by using a semi-quantitative ranking limited to three classes (1: one or a few tiny or small colonies, 2: several small or medium-sized colonies, 3: numerous small and mediumsized colonies or several large colonies). These data were used only to identify the most and the least abundant species. Specimens were examined routinely with a stereomicroscope and, in some cases, with a Hitachi S 570 scanning electron microscope (SEM) after being cleaned in a 7.5 % solution of sodium hypochlorite, rinsed, airdried and coated with gold. Taxa not identified at species level were not taken into account in the number of native species or when calculating proportions of non-indigenous species. The geographical distribution of the latter was categorized mainly according the classification in ecoregions by Spalding *et al.* (2007).

Results

General bryozoan diversity

Species richness for the whole collection amounts to 93 species (Tables 2, 3), including 12 cyclostomes, 2 ctenostomes and 79 cheilostomes. This assemblage includes 43 families, only eight of which comprise more than three species, while 25 (58 %) are monospecific. Seven taxa (5 cyclostomes, 2 cheilostomes) were not identified at species level (Tables 2, 3). Recorded diversity was higher in the northern region than in the southern region (82 vs. 66 species) and this N-S disparity in species richness was more pronounced for Cyclostomata (12 vs. 6 species) than for Cheilostomata (68 vs. 58 species). The Batroun area was the richest with 68 species, followed by the Tripoli area (57 species), while the two southern areas, Beirut and Tyre, accounted for 48 and 37 species, respectively. This bryozoan fauna included (i) a poorly diverse native component characterized by numerous 'southern' species, high endemicity, and unexpected absence of several taxa very common in the rest of the Mediterranean, and (ii) a high proportion of nonindigenous species.

Native bryozoan fauna: general features

With 59 taxa identified at species level, the native species richness was notably poor, especially in Cheilostomata (51 species), despite fairly intensive sampling in various habitats (Table 3). All native species were recorded in non-port habitats, except Cryptosula palliasana, recorded only once at Marina Villamar (loc. 15). In contrast, only 10 native species were recorded in ports. Among non-port habitats, the highest number of native species (56 species, including 7 cyclostomes and 1 ctenostome) were recorded on shaded rock walls (small semidark caves, overhangs), sampled between 6 and 42m depth. In this habitat, bryozoans often encrusted live and dead skeletons of three scleractinian species, Madracis pharensis (Heller, 1868), Phyllangia mouchezi (Lacaze-Duthiers, 1897) and Polycyathus muellerae (Abel, 1959), which were the most common corals on the sampled coastal bottoms.

The cheilostome assemblage comprises 24 species endemic to the Mediterranean (Table 2), i.e. 47 % of the native cheilostome pool. In contrast, only one species, *Copidozoum tenuirostre*, was identified in the category of

Table 1. List of 35 stations sampled at 18 localities along the coast of Leband	on.
---	-----

Stations	Coordinates	Depth	Habitat	Date
1A: Port	34.45472 N 35 81754 E	2-5m	Jetty blocks	19/IX/2002
1B: Port	34.45472 N	2-5m	Jetty blocks	08/VII/2003
2A: Ramkine Island	34.49639 N	14m	Overhang, cave	22/X/1999
2B: Lighthouse	35.76056 E 34.49444 N	12m	Pottery debris	22/V/1000
Island	35.77806.E 34.35722 N	14.24	Diana de la construcción de la c	22/1/1999
3A: Bay	35.75972 E	14-24m	Biogenic concretions	26/X/1999
4A: Chak el Hatab	35.66833 E	5-14m	Cave walls	18/IX/1999
4B: Hannouch	34.29417 N 35.66999 E	22m	Biogenic concretions	13/VII/2003
5A: Nearshore area	34.28611 N 35.66361 E	3-10m	Cave walls	18, 23/X/1999
5B: Nearshore area	34.28611 N 35.66361 F	10m	On <i>Spondylus spinosus</i> shell	02/V/2001
5C: Slope	34.28611 N	22m	Cave walls	14/IX/2002
5D: Slope	35.66361 E 34.28611 N	35m	Rocky wall, on	24/IX/2002
SE: Slope	35.66361 E 34.28611 N	25	Corallinaceae	06/3/11/2002
SE: Slope	35.66361 E 34 28722 N	35m	Rocky wall, on Adeonella	06/ V11/2003
6A: Bay	35.65555 E	6m	Rocks, on <i>Dictyopteris</i>	26/VI/1997
6B: 'Phoenician wall'	35.65555 E	9m	Rocky wall	16/X/1999
6C: Nearshore area	34.28722 N 35.65555 E	2-6m	Biogenic concretions	26/IX/2002
7A: Kafar Abida	34.22861 N 35.65333 E	10-12m	Rocky wall	06/VIII/1995
8A: Nearshore area	34.19556 N 35.62944 E	16m	Rocky wall	19/VIII/1993
8B: Nearshore area	34.19556 N 35.62944 E	28m	Biogenic concretions	08/VI/2000
8C: Small harbour	34.19250 N	1-2m	Quay wall	19/VIII/1993
9A · Tablieh shoal	35.63278 E 34.12417 N	17m	Shells & Cymodocea	08/VI/2000 17/X/1999
	35.63111 E 34.01417 N	20.20m	leaves	10/0/11/2002
	35.63139 E 33.91528 N	20-30111	Stones	10/ 11/2003
11A: Hart el Kelb	35.48780 E 33.91222 N	34m	Rocks, shells	21/X/1999
12A: Commercial port	35.52833 E	5m	Quay wall	02/VI/2000
12B: Commercial port	33.91222 N 35.52833 E	3-8m	On Thais shell	16/IX/2002
12C: Commercial port	33.91222 N 35.52833 E	5-8m	Pier, outside port	09/VII/2003
12D: Commercial port	33.90890 N 35.52280 F	14m	Pier, inside port	09/VII/2003
12E: Commercial port	33.90417 N	5-7m	Pier, inside port	16/IX/2002
13A: St Simon Islet	33.86250 N	3-4m	Rocks	about 1968
13B: Raoucheh coast	35.48139 E 33.88830 N	5-8m	Shells & Corallinaceae	IX/1992,
	35.46694 E 33.84306 N	2 11	Dienll	VI/2015
14А: Анроп	35.47555 E 33.77888 N	3-11m		10/ V11/2003
15A: Marina Villamar	35.46944 E	3m	Pier wall	07/VI/2000
16A: Harf el Rijmeh	35.36944 E	11m	On Spondylus shell	05/VI/2000
17A: Harf el Jouani	33.50861 N 35.34028 E	14m	Overhang	06/VI/2000
18A: Nearshore area	33.26567 N 35.19000 E	10m	Rocks	25/X/1999
18B: El Kasmieh	33.26528 N	36-42m	On <i>Phyllangia</i> ,	25/X/1999
	Stations 1A: Port 1B: Port 2A: Ramkine Island 2B: Lighthouse Island 3A: Bay 4A: Chak el Hatab 4B: Hannouch 5A: Nearshore area 5B: Nearshore area 5D: Slope 5D: Slope 6A: Bay 6B: 'Phoenician wall' 6C: Nearshore area 7A: Kafar Abida 8A: Nearshore area 8B: Nearshore area 8B: Nearshore area 8B: Nearshore area 8B: Nearshore area 8D: Nearshore area 10A: Aquamarina 11A: Harf el Kelb 12A: Commercial port 12B: Commercial port 12B: Commercial port 12D: Commercial port 13A: St Simon Islet 13B: Raoucheh coast 14A: Airport 15A: Marina Villamar 16A: Harf el Rijmeh 17A: Harf el Jouani 18B: El Kasmieh	StationsCoordinates1A: Port 34.45472 N1B: Port 35.81754 E2A: Ramkine Island 34.45472 N35.81754 E 34.45472 N2A: Ramkine Island 35.76056 E2B: Lighthouse 34.49444 NIsland 35.77806 E3A: Bay 34.35722 N3A: Bay 34.35722 N4A: Chak el Hatab 35.66999 E5A: Nearshore area 35.66999 E5A: Nearshore area 35.66361 E5D: Nearshore area 35.66361 E5D: Slope 34.28611 N35.66361 E 35.66361 E5D: Slope 35.66361 E5D: Slope 35.66361 E5E: Slope 34.28611 N35.66555 E 34.28611 N6C: Nearshore area 35.65555 E6B: 'Phoenician wall' 34.28722 N36.65555 E 34.28722 N7A: Kafar Abida 35.65555 E7A: Kafar Abida 35.62944 E8B: Nearshore area 35.6278 E9A: Tablich shoal 35.6139 E10A: Aquamarina 35.6139 E12B: Commercial port 35.91222 N33.90890 N 39.90890 N12D: Commercial port 35.2833 E12B: Commercial port 35.62944 E12E: Commercial port 35.62944 E12E: Commercial port 35.6139 E12B: Raoucheh coast 34.8830 N13A: St Simon Islet 33.88830 N13B: Raoucheh coast 35.46694 E13B: Raoucheh coast 35.46944 E13A: Nea	StationsCoordinatesDepth1A: Port 34.45472 N $2.5m$ 1B: Port 35.81754 E $2.5m$ 1B: Port 34.45472 N $2.5m$ 2A: Ramkine Island 35.76056 E $14m$ 2B: Lighthouse 34.49639 N $14m$ 2B: Lighthouse 34.49639 N $12m$ 1Sland $35.77806.E$ $12m$ 3A: Bay 35.75972 E $14.24m$ 4A: Chak el Hatab 35.66933 E $22m$ 5A: Nearshore area 35.66361 E $3-10m$ 35. Statison area 35.66361 E $3-10m$ 5B: Nearshore area 35.66361 E $32m$ 5D: Slope 34.28611 N $32m$ 5D: Slope 35.66361 E $35m$ 5E: Slope 34.28611 N $35m$ 5E: Slope 34.28722 N $6m$ 6A: Bay 34.28722 N $9m$ 6C: Nearshore area 35.65355 E $9m$ 6C: Nearshore area 35.65333 E $10-12m$ 8A: Nearshore area 35.62944 E $28m$ 8A: Nearshore area 35.63278 E $1-2m$ 9A: Tablieh shoal 35.4112 N $10m$ 35.4280 N 39.1528 N $11a$ 10A: Aquamarina 35.6313 E $10-12m$ 8A: Nearshore area 35.63278 E $1-2m$ 9A: Tablieh shoal 35.112 F 39.1528 N11A: Harf el Kelb 35.91528 N $31m$ 12B: Commercial port 35.2833 E $5m$ 12B: Commercial port 35.2833 E $5m$ 12B: Commercial por	StationsCoordinatesDepthHabitat1A: Port 34.45472 N2-5mJetty blocks1B: Port 34.45472 N2-5mJetty blocks2A: Ramkine Island 34.49639 N14mOverhang, cave2B: Lighthouse 34.49444 N12mPottery debrisJakad 35.77056 E14mOverhang, cave3A: Bay 34.35722 N14-24mBiogenic concretions3A: Chak el Hatab 35.66333 E5-14mCave walls4A: Chak el Hatab 35.66333 E5-14mCave walls5A: Nearshore area 35.66361 E3-10mCave walls5B: Nearshore area 35.66361 E22mBiogenic concretions5C: Slope 35.66361 E22mCave walls5D: Slope 34.28611 N35mCorallinaceae5D: Slope 34.28611 N35mRocky wall, on5E: Slope 34.28611 N35mRocky wall, on5E: Slope 34.28611 N35mRocky wall6A: Bay 34.28722 N9mRocky wall6A: Bay 34.28722 N9mRocky wall7A: Kafar Abida 35.65355 E9mRocky wall8A: Nearshore area 35.62361 L10-12mRocky wall8A: Nearshore area 35.62361 L10mRocky wall8A: Nearshore area 35.62361 L10mRocky wall8A: Nearshore area 35.62344 L10mRocky wall8A: Nearshore area 35.62378 L12mQuay wall

temperate-boreal Atlantic species. Taxonomic uncertainties (possibility of cryptic species) concerning several socalled 'cosmopolitan' or 'circumtropical' species detract from the reliability of other biogeographical categories.

Mediterranean 'southern' species

The assemblage of native cheilostomes recorded in Lebanon comprised several species known for their particular abundance in warm areas of the Mediterranean, such as *Reptadeonella violacea* and *Schizoporella errata*, both in class A of occurrence frequency (Table 2). Among these thermophilic cheilostomes widely distributed in Lebanon, four species are remarkable for their limited geographical range in the Mediterranean (Figs. 2, 3). Three of them, *Cellepora posidoniae, Hippaliosina depressa* and *Monoporella bouchardi*, are endemic to the Mediterranean, while the actual taxonomic status of the fourth, *Exechonella* cf. *antillea*, needs to be clarified. In contrast, another endemic 'southern' species, *Adeonella pallasii*, is remarkable for its limited range observed in Lebanon.

Cellepora posidoniae was represented by numerous colonies collected at 12 localities from Tripoli islands to Zahrani (3-34m depth). This material was ascribed with some hesitation to *C. posidoniae*, a species originally described from Chios (Aegean Sea) by Hayward (1975) as *Rhamphostomellina posidoniae*, type species of a new genus, *Rhamphostomellina* Hayward, 1975. This species was subsequently placed in the genus *Cellepora* Linnaeus, 1767 by Hayward & McKinney (2002) and redescribed with SEM pictures by Chimenz Gusso *et al.* (2014). The wall structure and some morphological details of this species were described by Gordon & Grischenko (1994) from Hayward's material. The colonies examined by

Hayward (1975), Unsal & d'Hondt (1978-1979) and Chimenz Gusso *et al.* (2014) were small, poorly calcified, growing on *Posidonia* leaves. In contrast, the Lebanese colonies were robust, well calcified, with rounded ramifications, growing on shaded hard substrata (Fig. 3A). Despite these zoarial and habitat differences, the Lebanese colonies were considered to be conspecific with *C. posidoniae* in having significant morphological similarities at zooidal level: primary orifice with a wide, low sinus, peristomial complex bearing a proximo-lateral avicularium with a proximal process delimiting a pseudosinus and a triangular, slightly hooked rostrum, and ovicell with two ectooecial foramina (Figs. 3B-C). However, a taxonomic reassessment of all *Cellepora* species from the Atlanto-Mediterranean region would be advisable.

Hippaliosina depressa (Fig. 4A) was recorded at 12 localities from Tripoli Islands to Tyre (5-42m depth). Lepralia depressa Busk, 1854 was originally described from the Aegean Sea, where its frequency was noticed by Harmelin (1968-1969) and Hayward (1974). It was recorded from the same area by Ünsal (1975) and Morri et al. (1999), and from the Levantine Basin, in Israel (d'Hondt, 1988; Sokolover, 2014), Egypt (Abdelsalam, 2014), Northern Cyprus (Koçak et al., 2002), and Lebanon (Bitar & Kouli-Bitar, 2001). Records of this species beyond the Eastern Mediterranean are scarce: Tunisia, Gulf of Gabes (David & Pouyet, 1979), SE Sicilia (Rosso et al., 2012), Corsica (Calvet, 1902; Gautier, 1962). The record by Powell (1969) at Acre, Haifa Bay of H. acutirostris Canu & Bassler, 1929, a species from Philippines, is doubtful and most likely concerns H. depressa.

Monoporella bouchardi (Fig. 4B) was particularly common along the coasts of Lebanon, as already pointed out by Harmelin (2014a). It was found at 13 localities (2-



Fig. 2: Geographical range of four 'southern' species, *Exechonella cf. antillea, Hippaliosina depressa, Cellepora posidoniae and Monoporella bouchardii* according to records by Calvet (1902), O'Donoghue & De Watteville (1939), Harmelin (1968-1969, 2014a + unpublished data), Hayward (1974, 1975), Ünsall (1975), David & Pouyet (1979), d'Hondt (1988), Morri et al. (1999), Koçak *et al.*, (2002), Rosso *et al.* (2012).



Fig. 3: Cellepora posidoniae, Selaata, 8 m. A: nodular colony. B: non ovicellate autozooids. C: ovicellate zooid. Scale bars, A: 5 mm, B: 200 μm, C: 100 μm

36m depth) from Tripoli Islands to Tyre. The geographical range of this Mediterranean endemic mostly covers the Eastern Mediterranean, although a few records are available from the Central Mediterranean (Harmelin, 2014a).

Exechonella cf. *antillea* (Fig. 4C) was present at 10 localities from 3 to 36m depth and in all samples from shaded walls from Selaata (3-35m depth). Its frequency in Lebanon contrasts with the scarcity of previous records, which are restricted to a few places in the Eastern Mediterranean (Fig. 2): Chios (Hayward, 1974: a single colony, first Mediterranean record), Cyprus (Koçak *et al.*, 2002; on *Posidonia* rhizomes), Israel, Akhzov canyon (Sokolover, 2014). This distribution pattern raises the question of the biogeographical status of this species: is it actually native to the Mediterranean or a cryptogenic species (*sensu* Carlton, 1996)? This species belongs to a worldwide "*E. antillea*" complex including, inter alia, the Caribbean *E. antillea* and species from W Africa (Cook & Bock, 2004). Several *Exechonella* species were

recorded from the Red Sea at Safaga Bay (Ostrovsky *et al.*, 2011) including *E. tuberculata* (MacGillivray, 1883), *E. brasiliensis* Canu & Bassler, 1928 and three unnamed species.

Adeonella pallasii is another typical 'southern' species endemic to the Mediterranean. Unlike the preceding species, it was recorded from a single locality, Selaata, where colonies were aggregated on overhangs and cave walls at 20-35m depth. Its occurrence is worth noting as it is the only species in this collection with large, rigid, erect colonies. This species is common in the Aegean Sea, also present in the Ionian Sea (Rosso, 2003; Rosso & Novosel, 2010), and particularly abundant in the Adriatic Sea on shaded walls and cave entrances (Hayward & McKinney, 2002; Novosel *et al.*, 2004; Rosso & Novosel, 2010; JGH, pers. observ.), where it replaces the 'northern' *A. calveti* (Canu & Bassler, 1930) in the same ecological niche.



Fig. 4: Three 'southern' cheilostomes common in Lebanon. A: *Hippaliosina depressa*, B: *Monoporella bouchardii*, C: *Exechonel- la* cf. *antillea*. Scale bars: 200 μm.

,,, _,		~r · · · ·					
CVCLOSTOMATA	TR	BA	BE	TY	Oc.	NIB	En.
CYCLUSIUMAIA Annectocymidae							
Prohoscing horvi (Audonin 1826)	x	X			в	?	?
Annectocyma sp. 1. creeping		X	x	x	B	•	•
Annectocyma sp. 2, erect	•	X		•	Ċ		•
Entalophoroecia sp.		Χ			D		
Tubuliporidae							
Idmidronea atlantica (Forbes) in Johnston, 1847		Х	•	•	С	•	•
Diastoporidae							
<i>Liripora amphorae</i> (Harmelin, 1976)	X	•	•	•	D	•	•
Frondiporidae	v	v		v	n		
Frondipora verrucosa (Lamouroux, 1821)	Х	Х	•	Х	В	•	•
Crisia regursa Heller 1867	v		v	v	C		v
Crisia sp	A X	•	Λ	A V	Č	•	Л
Lichenonoridae	Δ	•	•	Λ	C	·	•
Patinella radiata (Audouin 1826)	X	X	X		в		
Disporella hispida (Flemming, 1828)	X	X	X	x	Ă	:	:
Disporella sp.	X	X			C		
CTENOSTOMATA							
Pherusellidae							
Pherusella tubulosa (Ellis & Solander, 1786)	Χ	Х	Х	Х	В		
Vesiculariidae							
Amathia verticillata (delle Chiaje, 1822)	•	Х	Х	•	С	Х	•
CHEILOSTOMATA							
Acteidae	37	37	N 7		n		
Actea sica (Couch, 1844)	Х	Х	Х	•	В	•	•
Canidozoum tonuinostuo (Hinaka, 1990)		v			n		
Crassimarginatella sp	· x	Λ	· x	•	C	·	·
Retevirgula akdenizae Chimenz et al 1997	21	x	1	•	D	•	x
Anthroporidae	•	1	•	•	D	•	
Akatopora leucocypha (Marcus, 1937)	Χ			Х	С	Χ	
Bugulidae							
Bugula neritina (Linnaeus, 1758)	Χ	•	Х		С	Х	
Beaniidae							
Beania cf. hirtissima (Heller, 1867)	Х	•	•	Х	С	•	•
Beania magellanica (Busk, 1852)	•	X	Х	•	С	•	•
					n		
Licornia jolloisii (Audouin, 1826)	•	X	X	•	B	X	•
Cradoscrupocellaria bertholletii (Audouin, 1826)	·v	Х	Х	•	C	•	÷
Cradoscrupocellaria af alligi Vioiro & Sponger Jones 2012		•	•	•	D D	•	Λ
Microporidae	Λ	•	•	•	D	•	•
Calpensia nobilis (Esper 1796)	x			X	C		
Mollia circumcincta (Heller 1867)	x	x	•		č	•	x
Monoporellidae	28	1	•	•	C	•	
Monoporella bouchardi (Audouin, 1826)	Χ	Χ	Χ	Х	Α		Х
Onychocellidae							
Onychocella marioni Jullien, 1881	Χ	Χ	Х	Х	Α		
Onychocella vibraculifera Neviani, 1895	X	Х	Х	•	В		Х
Rectonychocella disjuncta (Canu & Bassler, 1930)	Х	•	•	•	D	•	•
Thalamoporellidae					~		
Thalamoporella harmelini Soule et al., 1999	•	•	Х	•	C	X*	•
Thalamoporella rozieri (Audouin, 1826)	•	Х	•	•	D	Х	•
Chlidoniidae Chlidonii muifamii (Dertelini 1910)	v				р		
Cribyilinidaa	Λ	•	•	•	D	•	•
Collaring balzagi (Audouin, 1826)		v			n		
Puelling podunculata Gautier 1956	· v	X	•	•	C	•	· v
Puelling nicardi Harmelin, 1988	X	X	x	•	Ă	·	X
Puellina radiata (Moll 1803)			X	•	D	•	X
Puelling setosa Waters, 1899	x	•	-	•	Ď	•	-
Hippothoidae		•	•	•		•	•
Hippothoa flagellum Manzoni, 1870	Х	Х		Х	С		
Ĉĥorizoporidae							
Chorizopora brongnarti (Audouin, 1826)		Χ			С		

Table 2. List of bryozoan species and families recorded in Lebanon in two northern (Tripoli: TR, Batroun: BA) and two southern (Beirut: BE, Tyre: TY) areas. Oc.: rank of occurrence frequency in 18 localities (A >50%, B: 26-50%, C: 10-25%, D <10%). NIB: non-indigenous species, including two cryptogenic species: *). En.: Mediterranean endemic species.

(continued)

Table 2 (continued)							
Trypostegidae							
Trypostega venusta (Norman, 1864)	Х	Х	•	Х	В	•	•
Exochonella ef antillea (Osburn 1927)	X	x	x	x	Δ		
Adeonidae	2	2	21	2	7	•	•
Reptadeonella violacea (Johnston, 1847)	Х	Х	Х	Х	Α		
Adeonellidae							
Adeonella pallasii (Heller, 1867)	•	Х	•	•	D	•	Х
Cellenoraria labelligera Harmer 1957	x	x	X		в	x	
Celleporaria cf. sherryae Winston, 2009	•	X	X		Č	X	
Celleporaria cf. brunnea (Hincks, 1884)		•	Х		D	Х	
Celleporaria vermiformis (Waters, 1909)	X	•	•		D	X	•
Drepanophora birbira Powell, 1967	X	X	•	X	С	X	•
Parasmittina egyntiaca (Waters 1909)	x	x	X	x	A	x	
Parasmittina protecta (Thornely, 1905)		•	X	•	Ĉ	X	:
Parasmittina raigii (Audouin, 1826)		Х			D		
Parasmittina rouvillei (Calvet, 1902)	X	Х	Χ	Х	C	·	Х
Parasmittina spondylicola Harmelin et al., 2009	X	X	X	•	C	X	•
Parasmittina serruloides Harmelin et al., 2009 Smitting nitidissima (Hincks, 1880)		X V	X	$\dot{\mathbf{v}}$	B B	X	•
Watersinoridae	Λ	Л	Λ	Λ	D	Λ	·
Watersipora subtorquata (d'Orbigny, 1852)	Х	Х	Х		Α	Х	
Schizoporellidae							
Schizoporella errata (Waters, 1878)	Х	Х	X	Х	Α	•	·
Schizoporella cf. tetragona (Reuss, 1848)	• V	·	X	·	C	•	X
Schizoprachielia sanguinea (Norman, 1808) Margarettidae	Λ	Λ	•	Λ	В	•	•
Margaretta cereoides (Ellis & Solander, 1786)	Х	Х			В		X
Hippaliosinidae							
Hippaliosina depressa (Busk, 1854)	Х	Х	Х	Х	Α	•	•
Hippopodinidae		v			D		v
Cheiloporinidae	•	Λ	•	•	υ	•	λ
Hagiosynodos hadros Hayward & McKinney, 2002				Х	D		Х
Hagiosynodos kirchenpaueri (Heller, 1867)	Х	Х	Х	•	В		Х
Cryptosulidae					_		
Cryptosula pallasiana (Moll, 1803)	•	•	X	•	D	•	•
Microporella hrowni Harmelin et al. 2011			x		D	x	
Microporella cf. ciliata (Pallas, 1766)	•	x	X	:	Č	•	•
Microporella coronata (Audouin, 1826)	Х	Х		Х	В	Х	
Microporella genisii (Audouin, 1826)	Х	Х	Х	Х	В	Х	•
Microporella harmeri Hayward, 1988	•	Х	•	•	D	Х	•
Petraliellidae Micropatraliella theorardii (Audouin, 1826)				v	D	v	
Escharinidae	•	•	•	Δ	ν	Λ	•
Escharina vulgaris (Moll, 1803)	Х	Х	Χ	Х	Α		
Therenia rosei Berning et al., 2008		Х	Х	•	В	•	Х
Colatooeciidae					C		
Trematooecia ligulata Ayari & Taylor, 2008	•	Х	Х	Х	C	•	Х
Hippomenella mucronelliformis (Waters 1899)	х	X			С		x
Celleporidae	18	1	•	•	U	•	
Buskea cf. dichotoma (Hincks, 1862)		Х		Х	С		
Cellepora posidoniae (Hayward, 1974)	Х	X	Х	Х	Α	•	Х
Celleporina bitari Harmelin, 2014	· v	X	·v	$\dot{\mathbf{v}}$	D	X	•
Celleporina cunariensis Alistegui, 1989 Celleporina mangnevillana (Lamouroux, 1816)	Λ	A X	Λ	Λ	A C	•	×
Celleporina sp.	X	X	•	:	č	•	
Predanophora longiuscula (Harmer, 1957)			X	X	С	X	•
Scorpiodinipora costulata (Canu & Bassler, 1929)	Х	Х	Х	Х	В	Х	•
Phidoloporidae					D		
Dentiporella sardonica (Waters, 18/9) Dissionalaidochasma maditarrangum Chimony Guesso & Soula, 2002	·v	Х	•	$\dot{\mathbf{v}}$	D D	•	$\dot{\mathbf{v}}$
Retenorella grimaldii (Jullien 1903)		x	•	Λ	C	•	<u>л</u>
Rhynchozoon neapolitanum Gautier, 1962	X	X	X		B		x
Rhynchozoon pseudodigitatum Zabala & Maluquer, 1988	•	Х		Х	С		Х
Schizoretepora hassi Harmelin et al., 2007	X	Х	Х	Х	В	X*	•
N spacios	E7	60	10	27		27	75
		Uð	40	31		41	

Missing native taxa

An obvious peculiarity of the sampled assemblages was the absence of certain families, genera, or of some large, conspicuous species which are among the most common bryozoans in coastal habitats of other parts of the Mediterranean. These missing taxa include cyclostomes such as *Tubulipora spp.*, *Diplosolen obelia*, *Plagioecia spp.*. Among the Cheilostomata, the most obvious absences concern families such as Cellariidae (*Cellaria*), Bitectiporidae (*Hippoporina*, *Metroperiella*, *Schizomavella*, *Pentapora*), Romancheinidae (*Escharella*, *Escharoides*, *Hemicyclopora*, *Hippopleurifera*), Myriaporidae (*Myriapora*), Bugulidae (only represented by the nonindigenous *Bugula neritina*). Genera such as *Callopora*, *Hincksina*, *Fenestrulina*, *Smittoidea*, *Turbicellepora* are also missing.

The absence of *Schizomavella* species is all the more surprising in that this genus is one of the most species-rich in the Mediterranean (Rosso, 2003; Reverter-Gil *et al.*, 2016) and that several species [e.g. *S. mamillata* (Hincks, 1880), *S. cornuta* (Heller, 1867)] are major components of the landscape of coastal rocky bottoms in the western Mediterranean. This genus was not recorded from Israel by Sokolover (2014), but is not entirely excluded from the Levantine Basin. In Egypt, a *Schizomavella* species was recorded from Alexandria as *S. auriculata* (probably misidentified) by O'Donoghue & de Watteville (1939), and recently, *S. triangularis* Reverter & Fernandez-Pulpeiro was found in the same area by Abdelsalm (2014).

Other large bryozoans which contribute significantly to the Mediterranean coastal landscapes were also missing, such as *Schizoporella dunkeri* (Reuss, 1848), normally very common with *Reptadeonella violacea* on shallow hard substrata, and the large erect species *Myriapora truncata* (Pallas, 1766), *Smittina cervicornis* (Pallas, 1766), *Pentapora fascialis* (Pallas, 1766) and *Turbicellepora avicularis* (Hincks, 1860). Some other species playing the same landscaping role in other parts of the Mediterranean, such as *Dentiporella sardonica* and *Parasmittina rouvillei*, were present but with few, small colonies.

Non-indigenous bryozoans

All non-indigenous bryozoans (NIB) identified in the present collection, except *Bugula neritina* s.l. and *Amathia verticillata*, were recorded previously and described in taxonomic works (Harmelin *et al.*, 2007, 2009, 2011, 2012; Harmelin, 2014b). With the addition of *B. neritina* s.l. and *A. verticillata* (see below), this NIB pool comprises 27 species, including two cryptogenic species (*Schizoretepora hassi, Thalamoporella harmelini*), all belonging to the order Cheilostomata (Table 2) except for *A. verticillata* (Ctenostomata). This pool might also include *Exechonella* cf. *antillea* considering that it is closely related with a species group from tropical waters (see above) and its range in the Mediterranean is restricted to the eastern basin and is very limited (Fig. 2). However, pending a taxonomic re-assessment of the whole species group, this species has been provisionally left in the native group. Among the 16 cheilostome genera that composed the NIB pool, eight occurred in the Mediterranean Sea only with alien species: *Akatopora* Davis, 1934, *Drepanophora* Harmer, 1957, *Licornia* van Beneden, 1850, *Mucropetraliella* Stach 1936, *Predanophora* Tilbrook, 2006, *Scorpiodinipora* Balavoine, 1959, *Thalamoporella* Hincks, 1887 and *Celleporaria* Lamouroux, 1821. The latter and two other genera, *Parasmittina* Osburn, 1952 and *Microporella* Hincks, 1877, contributed particularly to the NIB pool, each with four species. No Cyclostomata species was considered as NIB.

The proportion of NIB reached 31.4 % in the bryozoan fauna as a whole and 33.8 % in the cheilostome assemblage (Table 3). The contribution of NIB was variable in the four areas sampled along the coast of Lebanon (Fig. 1). It was higher in the two southern areas, particularly in the Beirut area where the proportion of alien species reached 43 % of the cheilostome assemblage. However, the number of non-indigenous cheilostome species in this area was only marginally higher than in the adjacent Batroun area (18 vs. 17 species), while the number of native cheilostomes was clearly lower in the Beirut area than in the Batroun area (24 vs. 39 species). Therefore, the higher percentage of NIB observed in the Beirut area is explained by the lower diversity in native cheilostomes. In the Tyre area, which was the focus of a lower sampling effort than the other areas, species richness was relatively low (37 species in total), but the proportion of NIB (35.5 % of the cheilostome assemblage) was also higher than in the two northern areas.

Table 3. Number of bryozoan species recorded in the northern and southern regions and along the whole coast of Lebanon within the following categories: cyclostomes, ctenostomes, cheilostomes, species identified at species level, native cheilostomes, endemic cheilostomes and non-indigenous species (NIB); percentage of endemic cheilostomes (relative to native cheilostomes); percentage of NIB species relative to (1) whole assemblage identified at species level, (2) whole cheilostome assemblage identified at species level.

	Ν	N S	
	Lebanon	Lebanon	coast
Whole assemblage	82	66	93
Cyclostomes	12	6	12
Ctenostomes	2	2	2
Cheilostomes	68	58	79
Identified species	75	63	86
Native cheilostomes	46	35	51
Endemic cheilostomes	21	15	24
% Endemic cheilost.	45.7	42.9	47.1
NIB species	21	23	27
% NIB (1)	28.0	36.5	31.4
% NIB (2)	30.3	38.6	33.8



Fig. 5: Number of native (NAB) and non-indigenous (NIB) cheilostome species recorded in four habitat categories (histogram bars), total number of cheilostome species in each habitat and corresponding percentage of NIB species.

The distribution of the cheilostome species richness in the four habitat categories (Fig. 5) shows that the highest number of NIB species was found on shaded walls and the lowest in ports. On the other hand, in terms of the ratio between indigenous and non-indigenous species, the highest proportions in NIB were found in ports and small substrata. The great majority (81 %) of the 27 NIB species were reported from the 5-15 m depth range, as a result, at least in part, of the higher sampling effort made in this depth range (Table 4). The other depth categories harboured similar NIB diversity.

Eleven NIB species, i.e. 40.7 % of the NIB pool, were recorded in class A (7.4 %) and class B (33.3 %) of occurrence frequency (Table 2). These widespread species occurred predominantly in natural habitats on various substrates (rock walls, bioconcretions, shells, coral

and erect bryozoan skeletons), from shallow (3-9 m) to deep (34-42 m) stations (Table 4). In particular, Celleporaria labelligera, Microporella genisii, Parasmittina egyptiaca (Fig. 6), Smittina nitidissima, Schizoretepora hassi and Watersipora subtorquata were among the most frequent bryozoans recorded in Lebanon and were distributed across a wide depth range (Table 4). In constrast, 9 NIB species were recorded exclusively in a single area and, in some cases, at a single station. Four of these species, Thalamoporella rozieri, Microporella harmeri, Celleporina bitari and Mucropetraliella thenardii, were found in moderately anthropized localities (Batroun, Selaata and El Kasmieh-Tyre) without large ports. The five other species (Thalamoporella harmelini, Celleporaria cf. brunnea, C. vermiformis, Parasmittina protecta, Microporella browni) were collected off Beirut city or in



Fig. 6: Three non-indigenous bryozoans common in Lebanon. A: Celleporaria labelligera. B: Parasmittina egyptiaca. C: Microporella genisii. Scale bars: 200 µm.

ports and harbours (Tripoli and Beirut main ports, Beirut airport marine infrastructures, Khalde Villamar marina).

Discussion

Diversity of the native bryozoan assemblages

Paucispecificity, high endemism and strong presence of southern species are the most striking features of the native bryozoan assemblages of the Lebanese coastal bottoms.

Species richness of the native cheilostomes recorded in Lebanon represents only 14 % of the diversity of this group in the Mediterranean as a whole, as compiled by Rosso (2003). This deficit is less pronounced at generic (27 %) and family (48 %) levels. The paucispecificity of this native assemblage is also evident when compared with other Mediterranean regional assemblages from the same depth range (Tab. 5). Therefore, the bryozoans of Lebanon clearly illustrate the extreme end of the wellknown eastward depletion of marine biodiversity in the Mediterranean (Fredj, 1974; Bianchi & Morri, 2000; Boudouresque, 2004; Coll *et al.*, 2010). The exclusion of conspicuous components of Mediterranean hard substrata communities is quite surprising, especially since these missing taxa have been recorded in the Aegean Sea (Harmelin, 1968-1969; Hayward, 1974; Ünsal, 1975; Morri *et al.*, 1999). Similarly, the temperate-boreal species are almost entirely excluded while this category is well represented in the Aegean Sea (23 %: Hayward, 1974). In assemblages of hydroids from Lebanon, Atlanto-Mediterranean species were also lacking (Morri *et al.*, 2009). This phenomenon of regional exclusion raises the question of whether it reflects a natural, biogeographical distinctive feature of the Lebanese coastal area, recent environmental degradation or cases of species displacement by invasive aliens.

<u>Biogeographical particularity hypothesis</u>. A most obvious bionomic feature of the coastal communities in this part of the Levantine coasts, which includes Lebanon and Israel, is the absence of the endemic sea-grass *Posidonia oceanica*, which is widespread throughout the rest of the Mediterranean. The absence of this major ecosystem engineer species in this area (Celebi *et al.*, 2006; Pergent *et al.*, 2014) may be relatively recent as it was recorded in 1977 (Bitar, 2010). However, the ab-

Table 4. Depth distribution of non-indigenous bryozoan species recorded in Lebanese waters according to four depth categories.

 N-Re: number of records. In bold, the 11 most frequent species. *: cryptogenic species.



 Table 5. Species richness of native bryozoan regional pools recorded from different parts of the Mediterranean. Data from Harmelin (1976, part): Table I, species from Precoralligenous, Coralligenous and semi-dark caves. n.r.: not recorded.

Geographical area (source)	Cyclostomata	Cheilostomata
Lebanon (present collection)	7	51
Aegean Sea (Harmelin, 1968-1969)	14	85
Aegean Sea, Chios (Hayward, 1974)	n.r.	101
Aegean Sea, Milos (Morri et al., 1999)	12	70
Tunisia (Ben Ismail et al., 2012)	13	73
N Adriatic (Hayward & McKinney, 2002)	19	79
Adriatic Sea (Novosel et al., 2004)	9	68
Balearic Islands, Cabrera (Zabala, 1993)	26	109
NW Mediterranean, Provence (Harmelin, 1976, part)	23	83

sence of remaining dead matte, a remarkably persistent component of Posidonia beds (Pergent et al., 2014), in recently surveyed sites (this study and RAC/SPA-UNEP/ MAP, 2014) suggests that P. oceanica was restricted to very small areas. Our field surveys also showed that several conspicuous and normally common invertebrates were lacking, such as gorgonians and zoanthids among the Cnidaria, and many large echinoderms (unpublished data). The subtropical thermal regime occurring in the SE Mediterranean (Abboud-Abi Saad et al., 2004; Nykjaer, 2009) enhanced by global warming, is the most likely ecological driver (Celebi et al., 2006) of this regional exclusion process. The exploration of deep-water habitats below 50-60 m depth, where thermal conditions are cooler all year round (Abboud-Abi Saad et al., 2004), might perhaps reveal the occurrence of several taxa missing at shallower depths, including bryozoans.

Habitat degradation hypothesis. The Lebanese coast is highly urbanized and exposed to multiple human stressors (e.g. Nakhlé, 2003; Shaban, 2008; Bitar, 2008, 2010). Unfortunately, references to the past status of the benthos of the Levantine area are very limited, except for some taxa, such as bivalves (Crocetta *et al.*, 2013a). Some investigations suggest that the quality of coastal habitats has severely declined over the last decades, in parallel to alien invasions (Basson *et al.*, 1976; Bitar & Kouli-Bitar, 1995). The marked difference in cyclostome diversity observed between the northern and the southern regions may conceivably be attributed to differences in environmental quality, as suggested by the decline of cyclostome diversity observed along a gradient of increasing pollution in Provence (Harmelin & Capo, 2002).

Displacement hypothesis. The establishment of highly competitive exotic species is one of the major threats suffered by marine communities, with a dramatic impact on interspecific relationships (e.g. Occhipinti Ambrogi, 2001; Galil, 2007). Profound changes caused by exotic species due to competitive exclusion were observed in Lebanese coastal communites. Two obvious examples are provided by remarkably successful Lessepsian species. Firstly, in the guild of herbivorous fishes, the na-

tive sparid Sarpa salpa was driven out by two invasive lessepsian rabbit-fishes, Siganus luridus and S. rivulatus (Bariche et al., 2004). Secondly, in the mediolittoral zone of the same area, the native mussel Mytilus galloprovincialis was replaced by the lessepsian species Brachidontes pharaonis (Bitar, 2010; Crocetta et al., 2013a). In the present bryozoan collection, the native Watersipora cucullata (Busk, 1854), originally described from the Aegean Sea, was not recorded while the alien W. subtorquata was frequent (Harmelin, 2014b). However, the hypothesis that the latter is responsible for the exclusion of the former would appear to be doubtful. No invasive trend was observed among the non-indigenous bryozoans collected in Lebanon (see below) and there was no evidence of a displacement phenomenon by other alien sessile organisms in the sampled habitats.

The particular occurrence of 'southern' species is another spectacular feature of the native assemblage. These species confirm the observation by Rosso (2003) that the native Mediterranean bryozoan fauna includes species which are predominantly or exclusively distributed in the eastern basin, or even only in the Levant Sea. The geographical distribution pattern of these species is most likely related to their thermophilic character and shaped by particular winter thermal thresholds (e.g. February surface isotherm at 16°C), as suggested by the distribution of other biota (Bianchi, 2007; Coll et al., 2010; Bianchi et al., 2012). These thermophilic species might be remnants of the tropical-subtropical fauna present during the Pleistocene interglaciations (Taviani, 2002). The same assumption was made for lithistid sponges from a Lebanese submarine cave (Pérez et al., 2004). 'Southern' cheilostomes could thus become good indicators of the effects of global warming on the geographical range of species.

The high degree of endemicity noted within the assemblage of native cheilostomes (47 %) contrasts with lower rates observed for this group in the western Mediterranean (32 %: Gautier, 1962 in Hayward, 1974) and the Aegean Sea (36 %: Hayward, 1974). This high endemicity was unexpected as it is not consistent with the general pattern of endemicity in the Mediterranean, which shows an eastward decreasing trend (e.g. Fredj, 1974; Coll *et al.*, 2010). Consistently with this trend, a low proportion of endemic species was noted by Morri *et al.* (2009) among hydroids collected at the same Lebanese localities as the bryozoans. The paradoxically high proportion of bryozoan endemics probably originates from the climatic discrimination which particularly affects the Atlanto-Mediterranean species not adapted to the warm conditions of this corner of the Mediterranean, and thus increases the proportion of endemic species, including the 'southern' ones restricted to the eastern Mediterranean.

Non-indigenous species: Lebanon, a hot-spot of bryozoan xenodiversity

The present results confirm that cheilostomes have a remarkable capacity for dispersal using human-mediated vectors, and attest that the coastal zone of Lebanon is a bioinvasion hotspot. A recent survey of bryozoans along the Mediterranean coast of Israel also showed a high, albeit lower, proportion of NIB (Sokolover, 2014: about 25%). The strong presence of NIB in Levantine coastal waters contrasts with low numbers of alien bryozoans recorded in other parts of the eastern Mediterranean (e.g. only two species reported from Turkey: Kocak & Aydin Önen, 2014). However, the report of a low occurrence of alien bryozoans may result from insufficient sampling and a lack of taxonomic expertise. This bias is not specific to bryozoans and may affect the validity of comparisons of regional invasion rates. If the exotic bryozoans recorded in Lebanon are added to the list compiled by Zenetos et al. (2010, 2012) for the whole of the Mediterranean, the amended list comprises 40 species, including 5 Ctenostomata and 35 Cheilostomata. Thus, the Lebanese NIB species represent 74 % of the whole Mediterranean pool of exotic cheilostomes. Moreover, the actual occurrence of several species of this pool needs to be reassessed as they were recorded without explanation despite serious taxonomic problems [e.g. Celleporaria fusca (Busk, 1854) in d'Hondt, 1988]. The list of NIB recorded in Lebanon itself, will probably be adjusted by future taxonomic revisions. In particular, the actual geographical range of the two cryptogenic species is still a pending issue. A cause of underestimation of the bioinvasion by bryozoans was the sampling design of the studied collection. A more intensive sampling effort on microhabitats particularly rich in bryozoans, such as empty shells and other small substrates, and in deep-water habitats, would probably have provided more records of NIB species. Moreover, the present list was based on relatively old samples, the most recent being from 2003. Therefore, considering the intensification of the colonization of the Mediterranean by non-indigenous species during the last decades (e.g. Ben Rais Lasram & Mouillot, 2009; Boudouresque & Verlaque, 2012; Galil & Goren, 2014), it is highly probable that the NIB fauna established in the Lebanese waters is now significantly richer than what is presented here.

The proportion of NIB observed in Lebanon exceeds the highest proportions of non-indigenous species recorded in other groups from the same area, except for the opisthobranch fauna (37% of exotic or cryptogenic species: Crocetta et al., 2013b). According to Lakkis (2012), the highest percentages of alien species in Lebanese waters was observed among crustaceans (less copepods: 30 %) and zooplankton (18%). Among bivalves from Lebanon, alien species account for ca. 16% (Crocetta et al., 2013a). The same proportion (15.8 %) of exotic species was found among hydroids sampled together with the bryozoans studied here (Morri et al., 2009). At Lebanese coastal localities similar to those sampled for bryozoans, the bony fish assemblages censused by diving comprised 13 % of introduced species (Harmelin-Vivien et al., 2005). The highest proportion of alien species recorded in Levantine waters (56%) was found in an assemblage of 16 serpulid species from SE Turkey (Cinar, 2006).

Among the most abundant and widespread NIB in Lebanon (class A), only the worldwide invader Watersipora subtorquata was known from other parts of the Mediterranean Sea (Harmelin, 2014b), while Smittina nitidissima was the only one already known from the Levantine coasts (recorded as S. malleolus: d'Hondt, 1988). Other particularly frequent NIB species were never recorded before in the Mediterranean, such as *P. egyptiaca*, although it was an early colonizer of artificial structures in the southern part of the Suez Canal (Hastings, 1927; Ghobashy et al., 1980). The lack of previous records of these frequent exotic species may be due to misleading identifications and more generally to poor knowledge of the bryozoan fauna of the SE Mediterranean and also of the Red Sea, despite recent efforts (Ostrovsky et al., 2011). The cryptogenic status (Carlton, 1996) attributed to T. harmelini and S. hassi, two species described from Lebanese material and afterwards only recorded in Israel (Sokolover, 2014), is another example of gaps in our knowledge. Their extremely limited known range may also be a case of stenoendemicity or an indication that these species are native to poorly known non-Mediterranean areas. The remarkable diversity of Parasmittina, Microporella and Celleporaria in the NIB pool may be explained by the fact that they are among the most species-rich cheilostome genera in warm waters (cf. species lists in Bock, 2015).

The absence of cyclostomes in the NIB pool is worth noting. This lack may result from the poor dispersal capacity of larvae in this group (e.g. Hayward & Ryland, 1985) and probably also from a low tolerance of conditions of vessel fouling and shipping transfer. It is noticeable that cyclostomes were similarly absent from a pool of alien bryozoans recorded in Australia (Campbell *et al.*, 2003) and only one species was recorded by Gordon & Mawatari (1992) among fouling bryozoans in New Zealand ports. However, obvious taxonomic difficulties and poor knowledge of the cyclostome fauna from both the eastern Mediterranean and the Red Sea may also explain in part this absence. For instance, *Proboscina boryi*, to which some small specimens were provisionally ascribed, was considered here as native although this species is very poorly known and its origin, Mediterranean or Red Sea, was not indicated by Audouin (1826).

The wide spatial and habitat distribution of many NIB species in Lebanon (Table 4, Fig. 1) is a strong indication of a relatively old colonization and good acclimatation to the Levantine environment. On the other hand, the restricted range of species recorded in a single area or at a single station can be diversely interpreted. The record of a species apparently well established in a single port (e.g. *C. vermiformis* at Tripoli) may correspond to a primary recipient area directly supplied by shipping. In contrast, single records of inconspicuous colonies from moderate-ly anthropized localities (e.g. *M. thenardii* and *C. bitari*) are more likely attributable to deficient sampling.

Geographical range of the Lebanese NIB and inferred pathways

The geographical range of the NIB species recorded in Lebanon was compiled from recent relevant sources and categorized according to the biogeographical classification proposed by Spalding *et al.* (2007), with slight modifications (Table 6). As noted in previous descriptions of the Lebanese NIB species (Harmelin *et al.*, 2007, 2009, 2011, 2012; Harmelin, 2014b), knowledge of the actual range of many of them is hampered by taxonomic uncertainties involving sibling species. However, some clear trends can be evidenced.

Historical introductions. The Lebanese pool of nonindigenous cheilostomes includes three of the most widespread aliens in tropical to temperate seas, whose occurrence in anthropized sites in the Mediterranean has been known for a long time: *Amathia verticillata* (= *Zoobotryon verticillatum*: cf. Waeschenbach *et al.*, 2015 for generic change), *Bugula neritina* and *Watersipora subtorquata*.

The non-indigenous status of A. verticillata in the European waters has been recognized only recently (Galil & Gevili, 2014; Ferrario et al., 2014; Marchini et al., 2015). This ctenostome, often confused with an algae, is widely distributed in tropical and subtropical regions, whereas in the Mediterranean it is confined to ports and marinas (e.g. Prenant & Bobin, 1956). A recent taxonomic revision (Vieira et al., 2014) of W. subtorquata, originally described from Brazil and often confused with other Watersipora species, has clarified its actual range, which includes the Mediterranean. Its occurrence in this sea only at anthropized sites (Chimenz Gusso et al., 2004; Harmelin, 2014b) strongly suggests a man-induced introduction. For the same reasons, we consider that B. neritina s.l. is not indigenous to the Mediterranean. This Bugula is widespread in Mediterranean ports and marinas, fouling artificial structures and ship hulls (e.g. Gautier, 1962; Prenant & Bobin, 1966), most probably since several centuries ago, as indicated by its first record in this sea (Linnaeus, 1758 in Ryland *et al.*, 2011). Molecular data (Mackie *et al.*, 2006; Fehlauer-Ale *et al.*, 2014) have shown that *B. neritina* is a complex of three cryptic biological species but, unfortunately, Mediterranean populations were not considered in these genetic studies. The current data do not allow the designation with any confidence of the native region of type S, the most widely distributed haplotype, which occurs in the English Channel and Galicia, but the south-western Atlantic and California may be two possible alternatives (Fehlauer-Ale *et al.*, 2014).

It is conceivable that transoceanic transportation on ship hulls of these three non-indigenous species may have started with the first voyages of discovery and trade between Europe and the rest of the world, an historical pathway that should not be underestimated (Bianchi & Morri, 2000). It is thus quite likely that their Mediterranean populations reflect very complex introduction histories with multiple imports from different geographical sources.

Suez Canal corridor. Most NIB recorded in Lebanon also occur in the Red Sea (85% with two disputable records: Table 6) and many of them were recorded in the Gulf of Suez and northern Red Sea (Balavoine, 1959; Ostrovsky et al., 2011) and the Suez Canal (Hastings, 1927; Eitan, 1972b; Ghobashy et al., 1980). The Indo-Pacific realm is also well represented, particularly the western Indo-Pacific region (Table 6). About 15 NIB species were also recorded in other areas of the Levantine coast, particularly Israel (Powell, 1969; d' Hondt, 1988; Sokolover, 2014). This distribution pattern stresses the major contribution of the Suez Canal as a pathway to the colonisation of the Levantine coasts by non-indigenous bryozoans, and is consistent with results for other groups (reviews in Streftaris et al., 2005; Galil, 2007; Çinar et al. 2011; Zenetos et al., 2012). Bryozoan translocation through the Suez Canal involves both transportation (i.e. shipping and rafting) and migration (see below). The Suez Canal is not only a two-way transit route for thousands of vessels each month, but also a place where barges and tugs are working all year round, in contact with fouled quays and immersed structures. As stressed by Fox (1926), Galil (2007) and Mineur et al. (2012) when fouled, these craft are probably important vectors of introduction, functioning as moving relays from spot to spot along the Canal. For instance, P. egyptiaca was recorded on barges at Port Taufiq (southern end of the Suez Canal) by Hastings (1927).

Gibraltar gateway. Long-distance vessels coming from the Atlantic via Gibraltar Strait can be vectors of introduction of alien species ("herculean" introductions) in both the Levantine Basin and the Red Sea. This scenario probably concerns first Atlantic species native to tropical and warm-temperate areas, such as *Smittina nitidissima*,

Table 6. Geographical distribution of non-indigenous bryozoan species (NIB) recorded in Lebanon (*: cryptogenic species). 1: Other Levant localities (a: from N. Sokolover, 2014, with some taxonomic changes). 2: Rest of the Mediterranean. 3: Red Sea, including the Gulf of Aden. 4: W Indo-Pacific. 5: Central Indo-Pacific. 6: Warm-temperate NE Pacific. 7: Tropical W Atlantic. 8: Warm-temperate W Atlantic. 9: Temperate W Atlantic. 10: Tropical E Atlantic. 11: Warm-temperate E Atlantic (including Canaries Is. and S Lusitanian region). 12: Temperate E Atlantic. (x): uncertain.

	1	2	3	4	5	6	7	8	9	10	11	12
Amathia verticillata	Х	Х	Х	Х	X		Х				Х	
Akatopora leucocypha	(x)		(x)				Х	Х	Х			
Bugula neritina s.l.	Χ	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х	Х
Licornia jolloisii	Х		Х				Х	Х				
Thalamoporella harmelini *	X (a)											
Thalamoporella rozieri	X (a)		Х	Х			(x)					
Watersipora subtorquata	Χ	Х	Х	Х	Х	Х	Х	Х	Х	Х		
Celleporaria cf. brunnea	Χ	Х				Х					Х	Х
C. labelligera			Х		(x)							
C. vermiformis			Х		(x)							
C. cf. sherryae	X (a)		(x)				(x)	Х				
Drepanophora birbira	X (a)		Х									
Parasmittina egyptiaca	X (a)		Х									
P. protecta	X (a)		Х	Х						Х	Х	
P. serruloides				Х			(x)					
P. spondylicola			Х	Х								
Smittina nitidissima	Х		Х				Х			Х	Х	
Microporella browni			Х	Х								
M. coronata	(x)	(x)	Х									
M. genisii			Х	Х							Х	
M. harmeri	X (a)		Х	Х	Х						Х	
Mucropetraliella thenardii	X(a)		Х	(x)								
Celleporina bitari			Х		(x)							
Predanophora longiuscula			Х	(x)								
Scorpiodinipora costulata			Х	Х	Х		Х			Х		
Rhynchozoon larreyi	Х		Х		(x)							
Schizoretepora hassi *	X (a)											

Akatopora leucocypha and Celleporaria cf. sherryae, but also species from other seas, such as Celleporaria cf. brunnea. The invasive range of this species, possibly native to California, now includes the NE Atlantic (Portugal, Cascais: Canning-Clode *et al.*, 2013; France, Arcachon: André *et al.*, 2014), the western Mediterranean (Italy, several ports: Lodola *et al.*, 2015), the central Mediterranean (Gulf of Tarento, fouling near aquaculture farm: Lezzi *et al.*, 2015) and the eastern Mediterranean (Turkey: Koçak, 2007; Lebanon: Harmelin, 2014b).

Means of translocation and spreading processes

The colonisation of Levantine waters by NIB species results from two modes of translocation at different spatial scales, functioning both through the Suez Canal and within the Mediterranean, i.e. transportation and natural spread (Fig. 7). Transportation of colonies is realized by attachment to moving substrata, mainly vessels and, occasionally, drifting items. Natural spread consists in an alternation between short-distance dispersal of propagules in the flow (essentially larvae), and attached stages (sexually mature colonies established on natural or artificial substrata). The distribution of NIB along the Levantine coasts most likely results from the linkage of transportation with natural spread using stepping stone habitats. Other vectors, such as release from shellfish aquaculture and aquaria, appear to be negligible in this regional context.

Shipping is a major but heterogeneous vector of species introduction, which includes both transcontinental exchanges by large vessels and coastal navigation by smaller merchant vessels, fishing boats or recreational craft. The maritime traffic across the Mediterranean is among the most intensive in the world (Dobler, 2002; Flagella *et al.*, 2006; Abdulla & Linden, 2008) and is a major source of alien introduction (Zibrowius, 2002; Galil *et al.*, 2008, Nunes *et al.*, 2014). It is well known to be the most powerful vector of introduction of bryozoans, (Ryland, 1965; Gordon & Mawatari, 1992; Watts *et al.*, 1998), chiefly by hull fouling. However, settlement of colonies in less exposed niche areas of vessels, such as 'sea-chests' (Jackson, 2008), may be an even more ef-



Fig. 7: Introduction processes and spread of alien bryozoans in the Levantine basin. Large arrows: main shipping routes; small double arrows: coastal shipping and working boats in the Suez Canal; black triangles: step by step spread of alien bryozoans by larval dispersal and colony settlement. 1: colony transfer by shipping; 2: colony transfer by rafting; 3: larval dispersal.

ficient introduction pathway for bryozoans. Moreover, if ballast waters cannot be vectors of direct transfer of bryozoan lecithotrophic larvae, given their short life duration and the length of the voyage, it might be possible that some larvae loaded with ballast water may occasionally settle on the walls of ballast tanks. This might concern especially sciaphilic species with short life cycles, such as several NIB recorded in Lebanon. If this occurs despite ballast water management (e.g. Endresen *et al.* 2004), this 'interior hull fouling' (Drake *et al.*, 2005) would lead to the development of fertile colonies and the possible production of larvae released into the sea during the ballast discharge.

Rafting on drifting substrata, either natural (e.g. algae: pelagic *Sargassum*) or artificial (e.g. plastic debris), allows sessile organisms, including bryozoans, to be transported over long distances (Winston *et al.*, 1977; Barnes *et al.*, 2009). The Mediterranean is severely polluted by vast amounts of plastic debris (Galgani *et al.*, 1995; Galil *et al.*, 1995; Suaria & Aliani, 2014). The abundance and nature of the debris along the Lebanese coasts, both of local and distant origin (pers. observ.), suggest that their role as vector of the dispersal of NIB along the Levantine coast should be considered.

Apart from the rare transfer of colony fragments of light erect species, propagules responsible for the spread of bryozoan local populations are free larvae released by fertile colonies. Most gymnolaemate bryozoans (all species recorded here) produce short-lived lecithotrophic larvae (Ryland, 1981; Jackson, 1986) which settle short distances from the parents, as confirmed by population genetics (Goldson et al., 2001). However, exceptional hydrological events may occasionally extend the dispersal range of these larvae. In any case, the geographical spread of a bryozoan population depends for the most part on the intensity of larvae production and the availability of habitats suitable for the settlement of larvae. Species whose life-history is characterized by an early and abundant production of larvae are particularly well suited to active step-by-step spread, including an opportunistic occupation of ephemeral and/or artificial substrata. This is in particular the case of Celleporina bitari, Smittina nitidissima, Parasmittina serruloides and P. spondylicola, small colonies of which are formed almost entirely of ovicellate zooids (Harmelin, 2014b; Harmelin et al., 2009).

Why are there so many alien bryozoans in Lebanese waters?

The conditions that foster the establishment of exotic species in native marine communities and their successful spread have been widely discussed in the literature (for the Mediterranean, e.g. Zibrowius, 1992; Boudour-esque, 1999b; Occhipinti-Ambrogi, 2001; Streftaris *et al.*, 2005; Galil, 2007). Clearly, the exceptional success of NIB in Lebanese coastal waters mainly results from

the combination of two conditions: a highly favourable local context and strong propagule pressure from species well-adapted to this environment.

The geographical position of Lebanon in the Mediterranean is the key feature of the particular suitability of this area for colonisation by NIB species as it combines proximity to the Suez Canal, permanent circulation of surface waters favouring linkages with the Canal and warm climate. Other local features reinforce the invasibility of the Lebanese coasts, such as intensive shipping activity converging on the ports of Beirut and Tripoli and proximity to routes to the Canal entrance, Turkish ports and the Black Sea, and benthic communities combining naturally low diversity and poor conditions.

Oceanographical context. The permanent flow direction of the Atlantic surface waters flowing off the eastern Levantine coasts is northerly (Hamad *et al.*, 2006). This surface circulation can thus be a powerful vector of supply of alien propagules (essentially free stages of larvae), released at or near the Mediterranean mouth of the Suez Canal, i.e. about 400-600 km upstream of Lebanon. However, this dispersal route is subject to fluctuations due to mesoscale anticyclonic eddies (Hamad *et al.*, 2006) and, as in any coastal system, local factors (e.g. wind, shoreline geography, underwater topography, etc.) increase the stochasticity of dispersal and recruitment. Currents are also vectors of transfer of drifting objects that may be colonized by non-indigenous sessile organisms.

Climatic context of Lebanese coastal waters. The marine water bodies of the south-east Levantine basin are the warmest of the Mediterranean Sea, as clearly seen on satellite thermal images of surface water (e.g. Hamad *et al.*, 2006; Taupier-Letage, 2008). Temperature profiles recorded in Lebanon (Abboud-Abi Saad *et al*, 2004) show that within the 0-40m depth range, the mean temperature does not fall below 16.9°C in winter and reaches 30°C in summer. These thermal conditions, intensified by the effect of global climate change on the Mediterranean Sea water temperature (Nykjaer, 2009), are thus particularly favourable to subtropical and tropical invaders (Occhipinti-Ambrogi, 2007; Lejeusne *et al.*, 2009; Por, 2010; Raitsos *et al.*, 2010).

Condition of benthic communities. The naturally poor diversity of the native bryozoan assemblages, and more generally of the native sessile benthos in the southeastern Mediterranean, might be favourable to the colonisation of NIB species able to occupy poorly exploited niches. The assumption that the naturally species poor Levantine area serves as a 'Godot' basin waiting for colonizers (Taviani, 2002; Oliviero & Taviani, 2003), has often been argued to explain the massive biotic invasion of the Levantine Basin (e.g. Galil, 2000). According to Boudouresque & Verlaque (2012), this hypothesis remains questionable, as counter-examples exist in the Mediterranean. The decline of Lebanese benthic communities due to pollution (Nakhlé, 2003; Shaban, 2008), responsible in part for their low diversity, may favour the settlement of resistant NIB species. Experiments with fouling assemblages in New-Zealand ports involving native and non-native bryozoans have shown that increased pollution resulted in the decline of native species diversity and percentage cover while non-native species were not affected (Piola & Johnston, 2008). The occurrence of *Celleporaria* cf. *brunnea* in the port of Beirut (Harmelin, 2014b) confirms the aptitude of this species to occupy port and anthropized habitats, already noted in California by Soule & Soule (1981) and indicated by records in the Mediterranean (Turkey: Koçak, 2007; Italy: Lezzi *et al.*, 2015, Lodola *et al.*, 2015) and eastern Atlantic (Portugal: Canning-Clode *et al.*, 2013).

Status of alien bryozoans in Lebanese communities

According to Zenetos et al. (2005, 2010), alien species can be classified as *established* (naturalized species with self-perpetuating populations), casual (recorded only once, presumed to be non-established), questionable (for taxonomic or other reasons) and cryptogenic. How may these categories be applied to the NIB recorded here, and more generally to bryozoans? Galil (2007) noted that it may be difficult to judge from a single record whether the occurrence of a new alien species reveals an on-going colonization or only an ephemeral intrusion. This doubt is reasonable, particularly when this record involves a species of which the individuals are able to move over long distances at the adult stage, such as fishes. In the present collection, only two NIB, Celleporaria vermiformis and C. cf. brunnea, were represented by a single colony. However, for C. vermiformis, the available sample was a fragment of a large fertile colony growing among many others on a jetty block (H. Z. observation), while C. cf. brunnea, also collected on a jetty, also occurs in other Mediterranean ports (see above). Therefore, these species are considered as established in Lebanese ports. More generally, as already stressed (Harmelin, 2014b), the actual spatial distribution and local abundance of bryozoan species are all the more likely to be underestimated as field sampling is not focused on bryozoans. This was the case for the studied collection, the initial purpose of which was a general overview of the Lebanese benthos. Therefore, all NIB recorded in Lebanon are considered to be established with self-reproducing populations. Moreover, considering that these NIB species have apparently no adverse impact on the recipient communities, they are considered to be non-invasive. This assessment refers to the concept of 'invasive' as implying the occurrence of harmful changes and biodiversity damage in the recipient ecosystems caused by the spread of alien species (e.g. IUCN, 1999, Boudouresque & Verlague, 2002).

Expected trends in the Levantine basin

We have shown from a relatively old series of samples that the bryozoan assemblages of Lebanon reflect particularly well the peculiarities of this region, i.e. poorly diverse communities, warm climate favouring thermophilic species and high incidence of alien species illustrated by an unexpectedly high proportion of non-indigenous species among cheilostomes. The current situation, 12 years after the last sampling survey, is most likely worse, with an increased number of NIB species established in the coastal zone. In the near future, the intensification of surface water warming and increased shipping activity through the Suez Canal and the Mediterranean will boost the introduction of new alien species, leading to drastic changes in the structure of bryozoan assemblages. In particular, the new expansion of the Suez Canal, inaugurated in August 2015, that aims to significantly increase the shipping activity, will result in the intensification of both the direct transfer of propagules from the Red Sea and the ship-mediated import of non-indigenous warm-water colonizers, with catastrophic consequences for the Mediterranean ecosystems well beyond the Levantine basin. Bryozoans thus provide a fascinating model of future trends with regard to the marine fauna in other regions under the impact of global and local changes.

Ackowledgements

Field surveys were supported by the French-Lebanese Cooperation CEDRE. Thanks are expressed to the staff of the Centre CNRS of Batroun, Lebanon, particularly Dr G. Khalaf, director, to the persons who helped us during field trips, to M. Paul, who improved the English writing, to A. Rosso and two anonymous reviewers for their precious comments on the manuscript.

References

- Abboud-Abi Saad, M., Romano, J.C., Bensoussan, N., Fakhi, M., 2004. Suivis temporels comparés de la structure thermique d'eaux côtières libanaises (Batroun) et françaises (Marseille) entre juin 1999 et octobre 2002. Comptes Rendus Geoscience, 336, 1379-1390.
- Abdelsalam, K.M., 2014. Benthic bryozoan fauna from the Northern Egyptian coast. *Egyptian Journal of Aquatic Research*, 40, 269-282.
- Abdelsalam, K.M., Ramadan, S.E., 2008. Fouling Bryozoa from some Alexandria harbours, Egypt. (II) Encrusting species. *Mediterranean Marine Science*, 9 (2), 5-20.
- Abdulla, A., Linden, O. (Eds). 2008. Maritime traffic effects on biodiversity in the Mediterranean Sea: Review of impacts, priority areas and mitigation measures. Malaga, Spain: IUCN Centre for Mediterranean Cooperation. 184 pp.
- André, F., Corolla, J.P., Lanza, B., Rochefort, G., 2014. Bryozoaires d'Europe. Les carnets du plongeur. Editions Neptune Plongée, Gargas, 255 pp.

- Audouin, V., 1826. Explication sommaire des planches de polypes de l'Egypte et de la Syrie, publiées par Jules-César Savigny, Membre de l'Institut; offrant un exposé des caractères naturels des genres avec la distinction des espèces. p. 225-244. In: *Description de l'Egypte. Histoire Naturelle*. Jomard, E.F. (Ed.). Imprimerie Nationale, Paris.
- Balavoine, P., 1959. Bryozoaires. Mission Robert Ph. Dollfus en Egypte (Décembre 1927 – Mars 1929). S.S. "Al Sayad". Résultats Scientifiques 3^e partie, 34, 257-280.
- Bariche, M., Letourneur, Y., Harmelin-Vivien, M., 2004. Temporal fluctuations and settlement patterns of native and Lessepsian herbivorous fishes on the Lebanese coast (eastern Mediterranean). *Environmental Biology of Fishes*, 70, 81-90.
- Barnes, D.K., Galgani, F., Thompson, R.C., Barlaz, M., 2009. Accumulation and fragmentation of plastic debris in global environments. *Philosophical Transactions of the Royal Society. Series B, Biological Sciences*, 364 (1526), 1985-1998.
- Barnes, D.K.A., Milner, P., 2005. Drifting plastic and its consequences for sessile organism dispersal in the Atlantic Ocean. *Marine Biology*, 146 (4), 815-825.
- Basson, P.W., Hardy, J.T., Lakkis, V., 1976. Ecology of marine macroalgae in relation to pollution along the coast of Lebanon. *Acta Adriatica*, 18, 19, 307-325.
- Ben Ismail, D., Rabaoui, L., Diawara, M., Ben Hassine, O.K., 2012. The Bryozoan assemblages and their relationship with certain environmental factors along the shallow and subtidal Tunisian coasts. *Cahiers de Biologie Marine*, 53, 231-242.
- Ben Rais Lasram, F., Mouillot, D., 2009. Increasing southern invasion enhances congruence between endemic and exotic Mediterranean fish fauna. *Biological Invasions*, 11, 697-711.
- Bianchi, C.N., 2007. Biodiversity issues for the forthcoming tropical Mediterranean Sea. *Hydrobiologia*, 580, 7-21.
- Bianchi, C.N., Morri, C., 2000. Marine biodiversity of the Mediterranean Sea: Situation, problems and prospects for future research. *Marine Pollution Bulletin*, 40, 5, 367-376.
- Bianchi, C.N., Morri, C., Chiantore, M., Montefalcone, M., Parravicini, V. *et al.*, 2012. Mediterranean Sea biodiversity between the legacy from the past and a future of change. p. 1-55. In: *Life in the Mediterranean Sea: A look at habitat changes*. Sambler, N. (Ed.). Nova Science Publishers, New York.
- Bitar, G., 2008. National overview on vulnerability and impacts of climate change on marine and coastal biodiversity in Lebanon. Contract RAC/SPA, N° 16: 41 pp.
- Bitar, G., 2010. Impact des changements climatiques et des espèces exotiques sur la biodiversité et les habitats marins au Liban. *Rapport Commission Internationale pour l'Exploration scientifique en Mer Méditerranée*, 39, 452.

- Bitar, G., Kouli-Bitar, S., 1995. Impact de la pollution sur la répartition des peuplements de substrat dur à Beyrouth (Liban – Méditerranée Orientale). *Rapport Commission Internationale pour l'Exploration scientifique en Mer Méditerranée*, 34, 19.
- Bitar, G., Kouli-Bitar, S. 2001. Nouvelles données sur la faune et la flore benthique de la côte Libanaise. Migration lessepsienne. *Thalassia Salentina*, 25, 71-74.
- Bock, P.E., 2015. *The Bryozoa home page*. http://bryozoa.net/ (Accessed 15 June 2015)
- Boudouresque, C.F., 1999a. The Red Sea-Mediterranean link: unwanted effects of canals. p. 213-228. In: *Invasive Species and Biodiversity Management*. Sandlund, O.T., Schei, P.J., Viken, Å. (Eds). Kluwer Academic Publishers, Dordrecht.
- Boudouresque, C.F., 1999b. Introduced species ithe Mediterranean: routes, kinetics and consequences. p. 51-72. In: *Proceedings of the Workshop on Invasive* Caulerpa *species in the Mediterranean*, Heraklion, Crete, Greece, 18-20 March 1998. MTS N° 125, UNEP, Athens.
- Boudouresque, C.F., 2004. Marine biodiversity in the Mediterranean: status of species, populations and communities. *Scientific Report of the Port-Cros National Park, Fr.*, 20, 97-146.
- Boudouresque, C.F., Verlaque, M., 2002. Biological pollution in the Mediterranean Sea: invasive versus introduced macrophytes. *Marine Pollution Bulletin*, 44, 32-38.
- Boudouresque, C.F., Verlaque, M., 2012. An overview of species introduction and invasion processes in marine and coastal lagoon habitats. *Cahiers de Biologie Marine*, 53, 309-317.
- Calvet, L., 1902. Matériaux pour servir à l'histoire de la faune des bryozoaires marins des côtes françaises. II. Bryozoaires marins des côtes de Corse. *Travaux de l'Institut de Zoologie de l'Université de Montpellier et de la Station Zoologique de Cette*, (2) 12, 1-52.
- Campbell, M.L., Nair, R., Hewitt, C.L., 2003. Xenodiversity of the marine bryozoan fauna in Australian waters. *Records of the South Australian Museum Monograph Series*, 7, 31-37.
- Canning-Clode, J., Souto, J., McCann, L., 2013. First record of *Celleporaria brunnea* (Bryozoa: Lepraliellidae) in Portugal and in the East Atlantic. *Marine Biodiversity Records*, 6, e108. doi: 10.1017/ S1755267213000821.
- Carlton, J.T., 1985. Transoceanic and interoceanic dispersal of coastal marine organisms: the biology of ballast water. *Oceanography and Marine Biology Annual Review*, 23, 313–371.
- Carlton, J.T., 1996. Biological invasions and cryptogenic species. *Ecology*, 77 (6), 1653-1655.
- Celebi, B., Cemal Gucu, A., Ok, M., Sakinan, S., Akoglu, E., 2006. Hydrographic indications to understand the absence of *Posidonia oceanica* in the Levant Sea

(Eastern Mediterranean). *Biologia Marina Mediterranea*, 13, 34-38.

- Chimenz Gusso, C., Boccia, P., Giovannini, N., 2004. Importance of faunistic and taxonomical studies for a correct analysis of the zoogeography of Mediterranean Bryozoa. *Biogeographia*, 25, 93-108.
- Chimenz Gusso, C., Nicoletti, L., Bondanese, C., 2014. Briozoi. *Biologia Marina Mediterranea*, 21 (suppl. 1), 1-336.
- Çinar, M.E., 2006. Serpulid species (Polychaeta: Serpulidae) from the Levantine coast of Turkey (eastern Mediterranean), with special emphasis on alien species. *Aquatic Invasions*, 1 (4), 223-240.
- Çinar, M.E., Bilecenoğlu, M., Öztürk, B., Katagan, T., Yokes, M.B. *et al.*, 2011. An updated review of alien species on the coast of Turkey. *Mediterranean Marine Science*, 12, 2, 257-315.
- Coll, M., Piroddi, C., Steenbeek, J., Kaschner, K., Ben Rais Lasram, F. *et al.*, 2010. The biodiversity of the Mediterranean Sea: estimates, patterns, and threats. *PLoS ONE* 5 (8): e11842.
- Cook, P.L., Bock, P.E., 2004. *Exechonella* (Exechonellidae Bryozoa, Cheilostomata) from the Recent and Miocene of southern Australia. *Proceedings of the Royal Society of Victoria*, 116 (2), 265-282.
- Crocetta, F., Bitar, G., Zibrowius, H., Oliverio, M., 2013a. Biogeographical homogeneity in the eastern Mediterranean Sea. II. Temporal variation in Lebanese bivalve biota. *Aquatic Biology*, 19, 75-84.
- Crocetta, F., Zibrowius, H., Bitar, G., Templado, J., Oliverio, M., 2013b. Biogeographical homogeneity in the eastern Mediterranean Sea - I: the opisthobranchs (Mollusca: Gastropoda) from Lebanon. *Mediterranean Marine Science*, 14 (2), 403-408.
- David, L., Pouyet, S., 1979. Etude sédimentologique et écologique du plateau tunisien et du golfe de Gabès.7: la mer Pélagienne. *Géologie Méditerranéenne*, 6 (1), 265-270.
- Dobler, J.P., 2002. Analysis of shipping patterns in the Mediterranean and Black seas. In: CIESM, Alien marine organisms introduced by ships in the Mediterranean and Black seas. *CIESM Workshop Monographs*, 20, 19-28.
- Drake, L.A., Meyer, A.E., Forsberg, R.L., Baier, R.E., Doblin, M.A. *et al.*, 2005. Potential invasion of microorganisms and pathogens via 'interior hull fouling': biofilms inside ballast water tanks. *Biological Invasions*, 7, 969-982.
- Endresen, Ø., Behrens, H.L., Brynestad, S., Andersen, A.B., Skjong, R., 2004. Challenges in global ballast water management. *Marine Pollution Bulletin*, 48, 615-623.
- Eitan, G., 1972a. Types of metamorphosis and early astogeny in *Hippopodina feegeensis* (Busk) (Bryozoa– Ascophora). *Journal of Experimental Marine Biology and Ecology*, 8, 27-30.

- Eitan, G., 1972b. Contributions to the knowledge of Suez Canal migration. Additions to the bryozoan fauna of the Suez Canal. *Israel Journal of Zoology*, 21, 377-384.
- Fehlauer-Ale, K.H., Mackie, J.A., Lim-Fong, G.E., Ale, E., Pie, M.R. *et al.*, 2014. Cryptic species in the cosmopolitan *Bugula neritina* complex (Bryozoa, Cheilostomata). *Zoologica Scripta*, 43, 193-205.
- Ferrario, J., Marchini, A., Lodola, A., Occhipinti Ambrogi, A., 2014. The pseudoindigenous bryozoan Zoobotryon verticillatum along the Mediterranean and European Atlantic coasts. Biologia Marina Mediterranea, 21 (1), 117-118.
- Flagella, M.M., Soria, A., Buia, M.C., 2006. Shipping traffic and introduction of non-indigenous species: study case in the Mediterranean Sea. *Ocean and Coastal Management*, 49 (12), 947-960.
- Fox, H.M., 1926. Cambridge Expedition to the Suez Canal, 1924. I. General Part. *Transactions of the Zoological Society of London*, 22, 1, 1-64.
- Fredj, G. 1974. Stockage et exploitation des données en écologie marine. C. Considérations biogéographiques sur le peuplement benthique de la Méditerranée. Mémoires de l'Institut Océanographique, Monaco, 7, 1-88.
- Galgani, F., Jaunet, S., Campillo, A., Guenegen, X., His, E., 1995. Distribution and abundance of debris on the continental shelf of the north-western Mediterranean Sea. *Marine Pollution Bulletin*, 30, 11, 713-717.
- Galil, B.S., 2000. A sea under siege alien species in the Mediterranean. *Biological Invasions*, 2, 177-186.
- Galil, B.S., 2007. Seeing red: Alien species along the Mediterranean coast of Israel. *Aquatic Invasions*, 2, 4, 281-312.
- Galil, B.S., Gevili, R., 2014. *Zoobotryon verticillatum* (Bryozoa, Ctenostomatida, Vesiculariidae), a new occurrence on the Mediterranean coast of Israel. *Marine Biodiversity Records*, 7, e17.
- Galil, B.S., Goren, M., 2014. Metamorphoses: Bioinvasions in the Mediterranean Sea. p. 463-478. In *The Mediterranean Sea: It is history and present challenges.* Goffredo, S., Dubinsky, Z. (Eds), DOI 10.1007/978-94-007-6704-1_27. Springer Science+Business Media, Dordrecht.
- Galil, B.S., Golik, K., Türkay, M., 1995. Liter at the bottom of the sea: a sea bed survey in the Eastern Mediterranean. *Marine Pollution Bulletin*, 30, 1, 22-24.
- Galil, B.S., Occhipinti-Ambrogi, A., Gollasch, S., 2008.
 Biodiversity impacts of species introductions via marine vessels. p. 117-158. In: *Maritime traffic effects on biodiversity in the Mediterranean Sea: Review of impacts, priority areas and mitigation measures*.
 Abdulla, A., Linden, O. (Eds). IUCN Centre for Mediterranean Cooperation, Malaga, Spain.
- Gautier, Y.V., 1956. Première faunule des Bryozoaires des côtes syriennes. *Vie Milieu*, 7, 4, 554-561.

- Gautier, Y.V., 1962. Recherches écologiques sur les bryozoaires chilostomes en Méditerranée occidentale. *Recueil des Travaux de la Station Marine d'Endoume*, 38 (Bull. 24), 1-434.
- Ghobashy, A.F.A., El-Komi, M.M., Ramadan, S.E., 1980. Fouling in the Suez Canal. p. 75-92. In: *Proceedings of the 5th Congress on Marine Corrosion and Fouling*, 19-23 May 1980, Spain.
- Goldson, A.J., Hughes, R.N., Gliddon, C.J., 2001. Population genetic consequences of larval dispersal mode and hydrography: a case study with bryozoans. *Marine Biology*, 138, 1037-1042.
- Gordon, D.P., Grischenko, A.V., 1994. Bryozoan frontal shields: the type species of *Desmacystis*, *Rhamphostomella*, *Rhamphostomittina*, *Rhamphostomellina*, and new genus *Arctonula*. *Zoologica Scripta*, 23 (1), 61-72.
- Gordon, D.P., Mawatari, S.F. 1992. Atlas of marinefouling Bryozoa of New Zealand ports and harbours. *Miscellaneous Publications. N.Z. Oceanographic Institute*, 107, 1-52.
- Goren, M., 1980. Development of benthic community on artificial substratum at Ashdod (Eastern Mediterranean). *Oceanologica Acta*, 3, 275-283.
- Hamad, N., Millot, C., Taupier-Letage, I., 2006. The surface circulation in the eastern basin of the Mediterranean Sea. *Scientia Marina*, 70, 3, 457-503.
- Harmelin, J.G., 1968-1969. Bryozoaires récoltés au cours de la campagne du «Jean Charcot» en Méditerranée orientale (août-septembre 1967). - I. Dragages. *Bulletin du Muséum national d'Histoire naturelle*, Paris, (2), 40, 6, 1178-1208. Ibid. (2) 41, 1, 295-311.
- Harmelin, J.G., 1976. Le sous-ordre des Tubuliporina (Bryozoaires Cyclostomes) en Méditerranée. Ecologie et systématique. Mémoires de l'Institut Océanographique, 10, 1-326.
- Harmelin, J.G., 2014a. Monoporella bouchardii (Audouin & Savigny, 1826) (Bryozoa, Cheilostomata):
 A forgotten taxon redescribed from Eastern Mediterranean material. Cahiers de Biologie Marine, 55, 91-99.
- Harmelin, J.G., 2014b. Alien bryozoans in the eastern Mediterranean Sea — new records from the coasts of Lebanon. *Zootaxa*, 3893 (3), 301-338.
- Harmelin, J.G., Capo, S., 2002. Effects of sewage on bryozoan diversity in Mediterranean rocky bottoms.
 p. 151-158. In: *Bryozoan Studies 2001: Proceedings* of the 12th International Bryozoology Association Conference. Wyse Jackson, P., Buttler, P. Wyse Jackson, Buttler, C., Spencer Jones, M. (Eds). Swets & Zeitlinger Publishers, Sassenheim.
- Harmelin, J.G., Bitar, G., Zibrowius, H., 2007. Schizoretepora hassi n. sp. (Bryozoa, Phidoloporidae) from Lebanon (Eastern Mediterranean), and reappraisal of Schizotheca serratimargo (Hincks, 1886). Cahier de Biologie Marine 48, 179-186.

- Harmelin, J.G., Bitar, G., Zibrowius, H., 2009. Smittinidae (Bryozoa, Cheilostomata) from coastal habitats of Lebanon (Mediterranean Sea), including new and non-indigenous species. *Zoosystema*, 31 (1), 163-187.
- Harmelin, J.G., Ostrovsky, A.N., Caceres-Chamizo, J.P., Sanner, J., 2011. Bryodiversity in the tropics: taxonomy of *Microporella* species (Bryozoa, Cheilostomata) with personate maternal zooids from Indian Ocean, Red Sea and southeast Mediterranean. *Zootaxa*, 2798, 1-30.
- Harmelin, J.G., Vieira, L.M., Ostrovsky, A.N., Cáceres-Chamizo, J.P., Sanner, J., 2012. Scorpiodinipora costulata (Canu & Bassler, 1929) (Bryozoa, Cheilostomata), a taxonomic and biogeographic dilemma: complex of cryptic species or human-mediated cosmopolitan colonizer? Zoosystema, 34 (1), 123-138.
- Harmelin-Vivien, M.L., Bitar, G., Harmelin, J.G., Monestiez, P., 2005. The littoral fish community of the Lebanese rocky coast (eastern Mediterranean Sea) with emphasis on Red Sea immigrants. *Biological Invasions*, 7, 625-637.
- Hastings, A.B., 1927. Cambridge expedition to the Suez Canal, 1924. Pt. 20. Report on the Polyzoa. *Transactions of the Zoological Society, London*, 22, 331-353.
- Hayward, P.J., 1974. Studies on the cheilostome bryozoan fauna of the Aegean island of Chios. *Journal of Natural History*, 8, 369-402.
- Hayward, P.J., 1975. A new species of epiphytic bryozoan from the Aegean Sea. *Cahiers de Biologie Marine*, 16, 207-212.
- Hayward, P.J., McKinney, F.K., 2002. Northern Adriatic Bryozoa from the vicinity of Rovinj, Croatia. *Bulletin of the American Museum of Natural History*, 270, 1-139.
- Hayward, P.J., Ryland J.S., 1985. *Cyclostome Bryozoans*. Synopses of the British Fauna (New Series), 34, published for the Linnean Society of London and the Estuarine and Brackish-Water Sciences Association by E.J. Brill/Dr. W. Backhuys, London, 147 pp.
- Hondt, J.L. d', 1988. Bryozoa from the coast of Israel. *Bolletino Zoologico*, 3, 191-203.
- IUCN (World Conservation Union), 1999. IUCN guidelines for the prevention of biodiversity loss due to biological invasion. *Species*, 31-32, 28-42.
- Jackson, J.B.C., 1986. Modes of dispersal of clonal benthic invertebrates: consequences for species distributions ans genetic structure of local populations. *Bulletin of Marine Science*, 39, 588-606.
- Jackson, L., 2008. *Marine biofouling: An assessment of risks and management initiatives*. Global Invasive Species Programme (GISP) and the UNEP Regional Seas Programme. 68 pp.
- Karlson, R.H., Osman, R.W., 2012. Species composition and geographic distribution of invertebrates in fouling communities along the east coast of the USA: a regional perspective. *Marine Ecology Progress Series*, 458, 255-268.

- Koçak, F., 2007. A new alien bryozoan *Celleporaria brunnea* (Hincks, 1884) in the Aegean Sea (eastern Mediterranean). *Scientia Marina*, 71 (1), 191-195.
- Koçak, F., Aydin Önen, S., 2014. Checklist of Bryozoa on the coasts of Turkey. *Turkish Journal of Zoology*, 38, 1-12.
- Koçak, F., Balduzzi, A., Benli, H.A., 2002. Epiphytic bryozoan community of *Posidonia oceanica* (L.) Delile meadow in the northern Cyprus (Eastern Mediterranean). *Indian Journal of Marine Science*, 3, 235-238.
- Lakkis, S., 2012. Introduction of Indo-Pacific and Eritrean species into Lebanese seawaters (Levantine Basin, East Mediterranean). *Cahiers de Biologie Marine*, 53, 381-390.
- Lejeusne, C., Chevaldonné, P., Pergent-Martini, C., Boudouresque, C.F., Pérez, T., 2009. Climate change effects on a miniature ocean: the highly diverse, highly impacted Mediterranean Sea. *Trends in Ecology & Evolution*, 25, 250-260.
- Lezzi, M., Pierri, C., Cardone, F., 2015. Presence of *Celleporaria brunnea* (Bryozoa: Lepraliellidae) in the Central Mediterranean: first occurrence in the Gulf of Tarento. *Marine Biodiversity Records*, 8, e137.
- Lodola, A., Ferrario, J., Occhipinti-Ambrogi, A., 2015. Further Mediterranean expansion of the non-indigenous bryozoan *Celleporaria brunnea*: multiple records along the Italian coasts. *Scientia Marina*, 79 (2), 263-274.
- Mackie, J.A., Keough, M.J., Christidis, L., 2006. Invasion patterns inferred from cytochrome oxidase I sequences in three bryozoans, *Bugula neritina*, *Watersipora subtorquata*, and *Watersipora arcuata*. *Marine Biology*, 149, 285-295.
- Marchini, A., Ferrario, J., Minchin, D., 2015. Marinas may act as hubs for the spread of the pseudo-indigenous bryozoan *Amathia verticillata* (Delle Chiaje, 1822). *Scientia Marina*, 79 (3), 355-365.
- Mineur, F., Cook, E.J., Minchin, D., Bohn, K., Macleod, A. et al., 2012. Changing coasts: marine aliens and artificial structures. Oceanography and Marine Biology: An Annual Review, 50, 189-234.
- Morri, C., Bianchi, C.N., Cocito, S., Peirano, A., De Biase, A. M. *et al.*, 1999. Biodiversity of marine sessile epifauna at an Aegean island subject to hydrothermal activity: Milos, eastern Mediterranean Sea. *Marine Biology*, 135, 729-739.
- Morri, C., Puce, S., Bianchi, C.N., Bitar, G., Zibrowius, H. *et al.*, 2009. Hydroids (Cnidaria: Hydrozoa) from the Levant Sea (mainly Lebanon), with emphasis on alien species. *Journal of the Marine Biological Association of the United Kingdom*, 89, 49-62.
- Nakhlé, K.F., 2003. Le mercure, le cadmium et le plomb dans les eaux littorales libanaises: apports et suivi au moyen de bioindicateurs quantitatifs (éponges, bivalves et gastéropodes). PhD Thesis Université Paris 7, 246 pp.

Novosel, M., Požar-Domac, A., Pasarić, M., 2004. Diversity and distribution of the Bryozoa along underwater cliffs in the Adriatic Sea with special reference to thermal regime. *P.S.Z.N.: Marine Ecology*, 25 (2), 155-170.

Nunes, A.L., Katsanevakis, S., Zenetos, A., Cardoso, A.C., 2014. Gateways to alien invasions in the European seas. *Aquatic Invasions*, 9, 2, 133-144.

- Nykjaer, L., 2009. Mediterranean Sea surface warming 1985-2006. *Climate Research*, 39, 11-17.
- Occhipinti Ambrogi, A., 2001. Transfer of marine organisms: a challenge to the conservation of coastal biocoenoses. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 11, 243-251.
- Occhipinti-Ambrogi, A., 2007. Global change and marine communities: Alien species and climate change. *Marine Pollution Bulletin*, 55, 342-352.
- Occhipinti-Ambrogi, A., Galil, B., 2010. Marine alien species as an aspect of global change. *Advances in Oceanography and Limnology*, 1, 199-218.
- O'Donoghue, C. H., de Watteville, D., 1939. The fishery grounds near Alexandria. XX – Bryozoa. Fouad I Institute of Hydrobiology & Fisheries. Notes & Memoirs, 34, 1-58.
- Oliverio, M., Taviani, M., 2003. The Eastern Mediterranean Sea: tropical invasions and niche opportunities in a "Godot Basin". *Biogeographia*, 24, 313-318.
- Ostrovsky, A.N., Cáceres-Chamizo, J., Vávra, N., Berning, B., 2011. Bryozoa of the Red Sea: history and current state of research. p. 67-97. In: *Annals of Bryozoology* 3. Wyse Jackson, P.N., Spencer Jones, M.E. (Eds). International Bryozoology Association Publ., Dublin.
- Pergent, G., Bazairi, H., Bianchi, C.N., Boudouresque, C.-F., Buia, M.-C. *et al.*, 2014. Climate change and Mediterranean seagrass meadows: a synopsis for environmental managers. *Mediterranean Marine Science*, 15/2, 462-473.
- Pérès, J.M., 1967. The Mediterranean benthos. Oceanography and Marine Biology Annual Review, 5, 449-533.
- Pérès, J.M., 1985. History of the Mediterranean biota and colonization of the depths. pp. 198-232. In: Western Mediterranean. Margalef, R. (Ed.). Pergamon Press.
- Pérez, T., Vacelet, J., Bitar, G., Zibrowius, H., 2004. Two new lithistids (Porifera: Demospongiae) from a shallow eastern Mediterranean cave (Lebanon). *Journal* of the Marine Biological Association, U.K., 84, 15-24.
- Piola, R.F., Johnston, E.L., 2008. Pollution reduces native diversity and increases invader dominance in marine hard-substrate communities. *Diversity and Distributions*, 14, 2, 329-342.
- Por, F.P., 1990. Lessepsian migration. An appraisal and new data. *Bulletin de l'Institut océanographique, Monaco*, n° special, 7, 1-10.

- Por, F.P., 2010. Climate optimum rejuvenates the Mediterranean marine world. *Integrative Zoology*, 5, 112-121.
- Powell, N.A., 1969. Indo-Pacific Bryozoa new to the Mediterranean coast of Israel. *Israel Journal of Zoology*, 18, 157-168.
- Prenant, M., Bobin, G., 1956. Bryozoaires, première partie: Entoproctes, Phylactolèmes, Cténostomes. *Faune de France*, 60, 1-398.
- Prenant, M., Bobin, G., 1966. Bryozoaires, deuxième partie: Chilostomes Anasca. *Faune de France*, 68, 1-647.
- RAC/SPA UNEP/MAP, 2014. Ecological characterization of sites of interest for conservation in Lebanon: Enfeh Peninsula, Ras Chekaa cliffs, Roucheh, Saida, Tyre and Nakoura. By Ramos-Esplá, A.A., Bitar, G., Khalaf, G., El Shaer, H., Forcada, A., Limam, A., Ocaña, O., Sghaier, Y.R., Valle, C., RAC/SPA Publ., MedMPAnet Project, Tunis. 168 pp.
- Raitsos, D.E., Beaugrand, G., Georgopoulos, D., Zenetos, A., Pancucci-Papadopulou, M.A. *et al.*, 2010.
 Global climate change amplifies the entry of tropical species into the Mediterranean Sea. *Limnology & Oceanography*, 55 (4), 1478-1484.
- Reverter-Gil, O., Souto, J., Novosel, M., Tilbrook, K.J., 2016. Adriatic species of *Schizomavella* (Bryozoa: Cheilostomata). *Journal of Natural History*, 50, 5-6, 281-321.
- Rosso, A., 2003. Bryozoan diversity in the Mediterranean Sea. *Biogeographia*, 24, 227-250.
- Rosso, A., Novosel, M., 2010. The genus Adeonella (Bryozoa, Ascophora) in the Mediterranean, with description of two new living species and rediscovery of a fossil one. Journal of Natural History, 44 (25-28), 1697-1727.
- Rosso, A., Di Martino, E., Sanfilippo, R., Di Martino, V., 2012. Bryozoan Communities and Thanatocoenoses from Submarine Caves in the Plemmirio Marine Protected Area (SE Sicily). p. 251-269. In: *Bryozoan Studies 2010. Proceedings of the 15th IBA Conference, 2010 Kiel, Germany.* Ernst, A., Schäfer, P., Scholz, J. (Eds.). Lecture Notes in Earth System Sciences, 143. Springer, Berlin, Heidelberg.
- Ryland, J.S., 1965. *Catalogue of main marine fouling organisms (found on ships coming into European waters)*. 2. *Polyzoa*. Paris, OECD
- Ryland, J.S., 1981. Colonies; growth and reproduction. p. 221-226. In: *Recent and fossil Bryozoa*. Larwood, G.P., Nielsen, C. (Eds). Olsen & Olsen, Fredensborg.
- Ryland, J.S., Bishop, J.D.D., De Blauwe, H., El Nagar, A., Minchin, D. *et al.*, 2011. Alien species of *Bug-ula* (Bryozoa) along the Atlantic coasts of Europe. *Aquatic Invasions*, 6, 17-31.
- Shaban, A., 2008. Use of satellite images to identify marine pollution along the Lebanese coast. *Environmental Forensics*, 9, 2-3, 205-214.

- Sokolover, N., 2014. *The Bryozoa community along the Israeli Mediterranean*. PhD Thesis, Tel Aviv University, 153 p.
- Soule, J.D., Soule, D.F., 1981. Heavy metals uptake in bryozoans. p. 227-233. In: *Recent and fossil Bryozoa. Papers presented at the 5th International Conference on Bryozoa, Durham 1980.* Larwood, G.P., Nielsen, C. (Eds.). Olsen & Olsen, Fredensborg.
- Spalding, M.D., Fox, H.E., Allen, G.R., Davidson, N., Ferdaña, Z.A. *et al.*, 2007. Marine ecoregions of the world: a bioregionalization of coastal and shelf areas. *Bioscience*, 57, 7, 573-583.
- Streftaris, N., Zenetos, A., Papathanassiou, E., 2005. Globalisation in marine ecosystems: the story of non-indigenous marine species across European seas. *Oceanography and Marine Biology: An Annual Review*, 43, 419-453.
- Suaria, G., Aliani, S., 2014. Floating debris in the Mediterranean Sea. *Marine Pollution Bulletin*, 86, 1-2, 494-504.
- Taupier-Letage, I., 2008. On the use of thermal images for circulation studies: Applications to the Eastern Mediterranean Basin. p. 153-164. In: *Remote sensing of the European seas*. Barale, V., Gade, M. (Eds), Springer Science and Business Media B.V.
- Taviani, M., 2002. The Mediterranean benthos from late Miocene up to Present: ten million years of dramatic climatic and geologic vicissitudes. *Biologia Marina Mediterranea*, 9(1), 445-463.
- Ünsal, I., 1975. Bryozoaires marins de Turquie. Istanbul Üniversitesi Fen Fakültesi Mecm. B, 40 (1-4), 37-54.
- Ünsal, I., d'Hondt, J.L., 1978-1979. Contribution à la connaissance des bryozoaires marins de Turquie (Eurystomata et Cyclostomata). *Vie et Milieu*, 28-29, 4 AB, 613-634.
- Vieira, L.M., Spencer Jones, M.E., Taylor, P.D., 2014. The identity of the invasive fouling bryozoan *Water-sipora subtorquata* (d'Orbigny) and some other congeneric species. *Zootaxa*, 3857 (2), 151-182.
- Waeschenbach, A., Vieira, L.M., Reverter-Gil, O., Souto-Derungs, J., Nascimento, K.B. *et al.*, 2015. A phylogeny of Vesiculariidae (Bryozoa, Ctenostomata) supports synonymization of three genera and reveals

possible cryptic diversity. *Zoologica Scripta*, 44 (6), 587-702.

- Watts, P.C., Thorpe, J.P., Taylor, P.D., 1998. Natural and anthropogenic dispersal mechanisms in the marine environment: a study using cheilostome Bryozoa. *Philosophical Transactions of the Royal Society of London* B, 353, 453-464.
- Winston, J.E., Gregory, M.R., Stevens, L.M., 1977. Encrusters, epibionts and other biota associated with pelagic plastics: a review of biogeographical, environmental and conservation issues. p. 81-97I. In: *Marine debris: sources, impact and solutions*. Coe, J.M., Rogers, D.B. (Eds), Springer, Berlin, Heidelberg, New York.
- Zabala, M., 1993. Els briozous. In: *Historia Natural de l'Arxipèlag de Cabrera*. Alcover, J.A., Ballesteros, E., Fornós, J.J. (Eds). *Monografies Societat d'Història Natural de les Balears*, 2, 561-577.
- Zenetos, A., Çinar, M.E., Pancucci-Papadopoulou, M.A., Harmelin, J.G., Furnari, G. *et al.*, 2005. Annotated list of marine alien species in the Mediterranean with records of the worst invasive species. *Mediterranean Marine Science*, 6 (2), 63-118.
- Zenetos, A., Gofas, S., Morri, C., Rosso, A., Violanti, D. et al., 2012. Alien species in the Mediterranean Sea by 2012. A contribution to the application of European Union's Marine Strategy Framework Directive (MSFD). Part 2. Introduction trends and pathways. *Mediterranean Marine Science*, 13 (2), 328-352.
- Zenetos, A., Gofas, S., Verlaque, M., Çinar, M.E., Garcia Raso, J.E. *et al.*, 2010. Alien species in the Mediterranean Sea by 2012. A contribution to the application of European Union's Marine Strategy Framework Directive (MSFD). Part I. Spatial distribution. *Mediterranean Marine Science*, 11 (2), 318-493.
- Zibrowius, H., 1992. Ongoing modification of the Mediterranean marine fauna and flora by the establishment of exotic species. *Mésogée*, 51, 83-107.
- Zibrowius, H., 2002. Assessing scale and impact of shiptransported alien fauna in the Mediterranean? In: CIESM, Alien marine organisms introduced by ships in the Mediterranean and Black Seas. *CIESM Workshop Monographs*, 20, 63-68.