

Mediterranean Marine Science

Vol 17, No 1 (2016)

VOL 17, No 1 (2016)



Haplotype variation in the *Physa acuta* group (Basommatophora): genetic diversity and distribution in Serbia

M. J. RAKOVIĆ, M. B. RAKOVIĆ, A. M. PETROVIĆ, N. Z. POPOVIĆ, J. A. ĐUKNIĆ, Z. Z. NAUNOVIC, M. M. PAUNOVIĆ

doi: [10.12681/mms.1453](https://doi.org/10.12681/mms.1453)

To cite this article:

RAKOVIĆ, M. J., RAKOVIĆ, M. B., PETROVIĆ, A. M., POPOVIĆ, N. Z., ĐUKNIĆ, J. A., NAUNOVIC, Z. Z., & PAUNOVIĆ, M. M. (2016). Haplotype variation in the *Physa acuta* group (Basommatophora): genetic diversity and distribution in Serbia. *Mediterranean Marine Science*, 17(1), 292–301. <https://doi.org/10.12681/mms.1453>

Haplotype variation in the *Physa acuta* group (Basommatophora): genetic diversity and distribution in Serbia

M.J. RAKOVIĆ¹, M.B. RAKOVIĆ², A. M. PETROVIĆ³, N. Z. POPOVIĆ¹, J. A. ĐUKNIĆ¹, Z. Z. NAUNOVIĆ⁴
and M. M. PAUNOVIĆ¹

¹ Institute for Biological Research “Siniša Stanković”, University of Belgrade, Bulevar despota Stefana 142, Belgrade, Serbia

² Natural History Museum of Belgrade, Njegoševa 51, Serbia

³ Faculty of Biology, Institute of Zoology, University of Belgrade, Studentski trg 16, Belgrade, Serbia

⁴ University of Belgrade, Faculty of Civil Engineering, Bulevar kralja Aleksandra 73, Belgrade, Serbia

Corresponding author: mpaunovi@ibiss.bg.ac.rs

Handling Editor: Eleni Kalogianni

Received: 1 August 2015; Accepted: 2 March 2016; Published on line: 30 March 2016

Abstract

The genus *Physa* (= *Physella*) includes the most abundant and diverse freshwater gastropods native to North America. Due to their invasive nature, many species occur throughout the world. The most abundant species, *Physa acuta*, has been introduced to Europe, Africa, Asia and Australia by human commerce and migrating birds. This species is widely distributed throughout Serbia. The aim of this study was to explore the genetic diversity of *P. acuta* from Serbia, and to determine the evolutionary relationships among native Physidae populations from North America, Mexico and Cuba and populations from Europe, using sequences of the mitochondrial 16S rDNA gene. The ML (Maximum Likelihood) tree revealed two clades within Physidae, and two clades that correspond to the families Planorbidae and Lymnaeidae. In the Physidae clade there are two separate clades: one includes the species *Physa spelunca*, and the second includes samples of *P. acuta*. We determined three different haplotypes within specimens from Serbia. One haplotype is genetically closest to *Physa heterostrofa* (synonym of *P. acuta*) from Philadelphia, while the other two are very close to *P. acuta* specimens from New Mexico. Together with other samples, our findings corroborate the notion that we are dealing with one panmictic population of *P. acuta* and not with several separate species, despite the high genetic diversity between and among the populations. Our results indicate that in the same population in Serbia, there is high genetic distance between samples. Despite the small number of analyzed samples, our findings point to multiple introductions of *P. acuta* from different locations in America.

Keywords: 16S rDNA variation, Physidae, *Physa acuta*, alien species, Danube, Serbia.

Introduction

Physa (= *Physella*) are the most abundant and diverse freshwater gastropods native to North America, which due to their invasive nature occur throughout the world (Burch, 1989). They are hermaphrodites with the ability to self-fertilize and can be found in a wide variety of freshwater habitats worldwide. Dillon (2000) provides an excellent ecological overview of this group. Based primarily on shell characteristics over 30 species have been described (Te, 1980; Burch, 1989).

The Physidae family can be divided into six groups, i.e. *aplexa*, *marmorata*, *fontinalis*, *acuta*, *gyrina* and *pomili*, which are phylogenetically different, possess distinctive penial morphology, and to some extent distinctive shell morphology (Te, 1978; Wethington, 2004). The influence of environmental factors and genetic variability must also be taken into consideration when distinguishing members of the family, since the use of only shell morphology can lead to misidentification (DeWitt *et al.*, 2000). Taylor's (2003) classification scheme of the Phy-

sidae, was primarily based on the penial complex. The author assumed that changes in penial morphology were progressive, and classified the family into grades and clades based on whether groups possess primitive or specialized characteristics.

Members of the *acuta* group can be identified by the following characters: shell shape is oval and the spire is pointed (acute); the aperture is large and narrow at the top; the shell is dull and striated longitudinally (DeWitt *et al.*, 2000). This group is characterized by considerable genetic variation, both within species and within populations. This variation, along with the absence of reproductive isolation between nominal species, has led to many species within this group to be lumped together under the name *Physa acuta*, including *P. heterostropha*, *P. integra*, *P. cubensis*, and *P. virgata* (Dillon *et al.*, 2002; Wethington 2004). Taylor (2003) has transferred *Physa* (*Physella*) *acuta* to the genus *Haitia* Clench & Aguayo, 1932, but herein we maintain the name *P. acuta*, following the classification of Wethington (2004).

Physa acuta was recorded in most parts of Europe, up to Sweden to the north, the British Isles to the west, Ukraine to the east, and the Mediterranean to the south. It has been also introduced to Africa, Asia and Australia by human commerce (de Jong *et al.*, 2014). In Serbia, according to Novaković (2014), *P. acuta* was recorded in a total of 74 sampling sites situated at 52 watercourses and 4 reservoirs. In Vojvodina, the species is present mainly in the Danube–Tisa–Danube Canal (DTD), and in small and medium watercourses at altitudes up to 500 m (Novaković, 2014).

The aim of this study was to explore the genetic diversity of *P. acuta* from Serbia, and to determine the evolutionary relationships of the examined population with native Physidae populations from the United States of America, Canada, Mexico and Cuba, as well as with populations from Europe, using mitochondrial 16S rDNA gene sequences.

Materials and Methods

Fieldwork and species identification

Investigation of freshwater mollusks was performed during sampling of the faunal composition of the wetlands (backwater) in the Danube River (Fig. 1). The samples were collected along the right bank of the river at the following localities, the Čibuklija wetland (44.804719° N, 21.319505° W; river km 1082.70) and along the left bank of the river, the Žilova wetland (44.778011° N, 21.191754° W; river km 1090.50). Fieldwork was conducted in 2010, 2011, and 2014, using the Van-Veen type of grab with a sample area of 270 cm² and a “kick-netting” type of benthos network (mesh size 500 µm). After collection, samples were transferred to the laboratory where species were identified under a stereomicroscope (MBL2000, Krüss, Germany) and a binocular magnifier (MSZ5000, Krüss, Germany) according to Zhadin (1952), Pflieger (1999), Glöer (2002) and Glöer & Meier-Brook (2003). Table 1 provides a summary of the sampling locations, the number of individuals typed for the mitochondrial large subunit rDNA (16S) marker, and voucher specimen information for the Serbian *P. acuta* and sequences taken from GenBank database at the NCBI (<http://www.ncbi.nlm.nih.gov/>, National Center for Biotechnology Information). Additionally, seven specimens were included as representative of other freshwater Basommatophora families (Lymnaeidae, Planorbidae; see Table 1) as related species from the same habitat.

Isolation of soft tissue from snails

Individuals used for the molecular analyses were transferred to the laboratory alive, without use of a fixative. The soft tissues were isolated using the microwave oven (MWO) method, by treating individual specimens with microwaves in a microwave oven MO17DE, Gore-nje (output power of 700 W) at various intervals, depending on the length of the shell, and according to a previ-

ously published protocol (Galindo *et al.*, 2014), modified for freshwater snails (see Appendix I). The MWO method allowed for the preservation of undamaged snail shells that have been deposited in the collection of the Department of Hydroecology and Water Protection, Institute for Biological Sciences “Siniša Stanković”, University of Belgrade, Serbia. For molecular analyses, the soft parts of the body were stored in 96% ethanol.

DNA isolation, amplification and sequencing

DNA was extracted using the DNeasy Blood and Tissue Kit (QIAGEN, Valencia, CA, USA) according to the manufacturer’s instructions. For larger specimens, 20–30 mg of foot tissue was used. The target gene fragment was amplified using the following primers: 16Sar ‘5-CGCCT-GTTTATCAAAAACAT-3’ and 16Sbr ‘5-CCGGTCT-GAACTCAGATCACGT-3’ (Kessing *et al.*, 1989). DNA amplification was performed in a final volume of 25 µl, containing 1 µl of the extracted DNA, 12.5 µl of 1 x KAPA2G Robust HotStart ReadyMix (containing 2 mM MgCl₂ and 1.25 µl of 0.5 µM of each primer, and 9 µl of water. PCR was performed in an Eppendorf Mastercycler® (Hamburg, Germany) using the following thermal profile: initial denaturation at 94°C for 3 min, 38 cycles at 94°C for 30 s, at 56°C for 30 s, at 72°C for 45 s, and a final extension step at 72°C for 7 min. The PCR products were purified using the QIAquick PCR Purification Kit (QIAGEN, Valencia, CA, USA) according to the manufacturer’s instructions. DNA sequencing in both directions was performed by Macrogen Europe Inc. (Amsterdam, Netherlands).

Phylogenetic analysis

For the detection of defects and polymorphic sites at the ends of the sequences we used Sequencher 5.2.4. software (Trial free version). Comparison of the obtained sequences with sequences in the gene bank database was performed using the Basic Local Alignment Tool (BLAST), available at <http://www.ncbi.nlm.nih.gov>. Sequences were aligned using the program Clustal_W with the parameters provided in the software package MEGA (version 6.0; Tamura *et al.*, 2013). For calculation of average genetic distances, Kimura’s two-parameter method (K2P) of base substitution was used. The Maximum Likelihood (ML) tree was obtained using MEGA 6 software. The robustness of the tree was assessed using a bootstrap analysis with 1.000 replicates. The haplotype network was constructed with the TCS program (version 1.21; Clement *et al.*, 2000).

Results

P. acuta was found in the Čibuklija and Žilova wetlands of the Danube. Beside *P. acuta*, we recorded several other species from the order Basommatophora,




Fig. 1: Distribution of *P. acuta* group in Canada, North America, Mexico and Cuba (A), Europe (B) and Serbia (C). Distribution of *P. acuta* in North and Central America (white circles) was compiled from the data of Wethington *et al.* (2009), Wethington & Guralnick (2004) and Kraus *et al.* (2014). The European (white circles) range is based on Wethington & Lydeard (2007), data and distribution in Serbia is based on Novaković (2014, black circles), and on our field sampling data (white circles).

which were collected for molecular analysis. In total, six specimens of *P. acuta* and *Planorbarius corneus* and one specimen of *Lymnaea stagnalis* and *Stagnicola palustris* (Table 1) were collected. All collected specimens of *P. acuta* were treated as a single population, since both wetlands / marshes are a few kilometers apart and their habitat features are the same.

The ML tree inferred with the alignment of the matrix, including 51 specimens, revealed two clades within Physidae, and two clades corresponding to the families Planorbidae and Lymnaeidae (Fig. 2). Within the clade Physidae there are two separate clades: one contains the species *P. spelunca* (*acuta* group B), and the second includes the rest of the samples of *P. acuta* (*acuta* group A). We identified three different haplotypes in the specimens from Serbia. One haplotype was genetically closest to the species *P. heterostrofa* (now considered as synonym of *P. acuta*) from Philadelphia, while two others were very close to *P. acuta* specimens from New Mexico.

Haplotype networks of *P. acuta* samples, together with *P. spelunca* (Fig. 3) revealed that samples from Europe possess distant haplotypes in the network matrix. These results confirmed that in Serbia, in the same population, there is a high genetic variability between samples. K2P

distances varied between our samples from 3.5 to 3.7%. Moreover, samples from France, similarly to the Serbian samples, are clearly separated in two different clusters, with K2P distances of 2.8-3.5% (See Appendix II).

Discussion

Biodiversity of freshwater ecosystems is changing rapidly due to anthropogenic transoceanic biotic exchange, and water habitats are considerably affected by biological invasions (Pyšek *et al.*, 2010), as has already been described for the river Danube (Paunović *et al.*, 2007a,b; Arbaciauskas *et al.*, 2008; Sommerwerk *et al.*, 2009; Sommerwerk *et al.*, 2010; Paunović *et al.*, 2015). Invasive species can be introduced by intent or by chance. Introduced species compete with native inhabitants for resources, which can cause suppression of native species from their natural habitats. With regard to Serbia, it has been confirmed that the rate of introduction of alien species in some biotopes is growing, as they are suitable recipient areas (Zorić *et al.*, 2010a). For the largest number of alien species, the most suitable recipient areas are heavily modified and anthropo-

Table 1. Locality details, GenBank accession numbers and status of the freshwater snails typed for mt16S region. I = introduced (Europe); N = native (America).

Taxon	Locality	16S (N/hN)*	Specimen code	Status	GenBank Accession#
Physidae					
<i>Physa acuta</i>	France; Montpellier	1/1		I	AY651224
<i>Physa acuta</i>	France; Montpellier	1/1		I	AY651223
<i>Physa acuta</i>	France; Montpellier	1/1		I	EU038321
<i>Physa acuta</i>	Netherlands	1/1		I	EU038342
<i>Physa acuta</i>	Serbia; Danube	3/1	mr21; mr23; mr24	I	
<i>Physa acuta</i>	Serbia; Danube	2/1	mr19; mr20	I	
<i>Physa acuta</i>	Serbia; Danube	1/1	mr22	I	
<i>Physa acuta</i>	Nebraska	1/1		N	KF850479
<i>Physa acuta</i>	Montana	1/1		N	KF850476
<i>Physa acuta</i>	Wyoming	1/1		N	AY651241
<i>Physa acuta</i>	Missouri	1/1		N	AY651226
<i>Physa acuta</i>	Arizona	1/1		N	EU038308
<i>Physa acuta</i>	Niagara River, Canada	1/1		N	EU038309
<i>Physa acuta</i>	Colorado	1/1		N	AY651219
<i>Physa acuta</i>	Charleston; South Carolina	2/1		N	GQ415017; GQ415009
<i>Physa acuta</i>	Charleston; South Carolina	1/1		N	GQ415021
<i>Physa acuta</i>	Charleston; South Carolina	1/1		N	GQ415010
<i>Physa acuta</i>	Charleston; South Carolina	1/1		N	GQ415020
<i>Physa acuta</i>	Charleston; South Carolina	1/1		N	GQ415019
<i>Physa acuta</i>	Charleston; South Carolina	1/1		N	GQ415018
<i>Physa acuta</i>	Charleston; South Carolina	1/1		N	GQ415016
<i>Physa acuta</i>	Charleston; South Carolina	1/1		N	GQ415015
<i>Physa acuta</i>	Charleston; South Carolina	1/1		N	GQ415014
<i>Physa acuta</i>	Charleston; South Carolina	1/1		N	GQ415013
<i>Physa acuta</i>	Charleston; South Carolina	1/1		N	GQ415012
<i>Physa acuta</i>	New Mexico	1/1		N	EU833960
<i>Physa acuta</i>	New Mexico	1/1		N	EU833959
<i>Physa acuta</i>	New Harmony, Indiana	1/1		N	EU038325
<i>Physa acuta</i>	New Harmony, Indiana	1/1		N	EU038324
<i>Physa acuta</i>	Cuba	1/1		N	EU038319
<i>Physa acuta</i>	Cuba	2/1		N	EU038320; EU038318
<i>Physa acuta</i>	Mexico	1/1		I	HQ283247
<i>Physa heterostrophia</i>	Pennsylvania, Philadelphia	1/1		N	AY651231
<i>Physa heterostrophia</i>	Pennsylvania, Philadelphia	1/1		N	AY651230
<i>Physa spelunca</i>	Lower Kane Cave; Wyoming	1/1		N	AY651243
<i>Physa spelunca</i>	Lower Kane Cave; Wyoming	1/1		N	AY651242
<i>Physa virgata</i>	Gila River; Arizona	1/1		N	AY651210
<i>Physa virgata</i>	Gila River; Arizona	1/1		N	AY651209
Lymnaeidae					
<i>Lymnaea stagnalis</i>	Serbia; Danube	1/1	mr33	N	
<i>Stagnicola palustris</i>	Serbia; Danube	1/1	mr18	N	
Planorbidae					
<i>Planorbarius corneus</i>	Serbia; Danube	2/1	mr26; mr31	N	
<i>Planorbarius corneus</i>	Serbia; Danube	1/1	mr27	N	
<i>Planorbarius corneus</i>	Serbia; Danube	1/1	mr28	N	
<i>Planorbarius corneus</i>	Serbia; Danube	1/1	mr29	N	
<i>Planorbarius corneus</i>	Serbia; Danube	1/1	mr32	N	

* N = number of individuals sequenced; hN = number of unique haplotypes obtained.

genic aquatic habitats, which are also suitable for the spread of the alien populations of *P. acuta*. Moderate to high level of organic pollution, intensive inland water navigation and invasive species corridors in the vicinity also contribute to the degree of successful invasiveness of an area. In Europe, there are four main invasive cor-

ridors: the northern, central, western and southern corridors. The southern corridor links the Black and North Sea by the Danube, Rhine and Main rivers and is the most important corridor for the spread of alien species (Bij de Vaate *et al.*, 2002; Panov *et al.*, 2009; Panov *et al.*, 2010).




Fig. 2: Phylogenetic trees based on mt16S rDNA, obtained using the Maximum Likelihood (ML) method. Bootstrap values are indicated below the branches. Scale bar indicates the number of substitutions per site.

The Danube is one of the most interesting routes of introduction and spread of aquatic alien species, with the spreading continuing through its main tributaries, the Sava, the Tisa and the Morava. This corridor was most probably the main corridor through which *P. acuta* spread to Serbia. That is also supported by the high num-

ber of individuals recorded at the Sava River during the “GLOBAQUA” project surveys of 2014 and 2015 (unpublished data). We believe that wetland birds contribute to the spread of North American species along the Danube, as Malone (1965) linked the spread of North American species into Europe with wetland birds. The snail *P.*




Fig. 3: Haplotype networks from 43 *Physa acuta* group specimens, obtained using statistical parsimony (TCS). Circles represent specific haplotypes; the size of the circles reflects the number of individuals with a particular haplotype (not to scale); the dots between the circles represent mutational steps.

acuta has now attained worldwide distribution and can be considered as one of the most cosmopolitan recent snail species (Dillon *et al.*, 2002; Wethington & Lydeard, 2007). This species is known as a rapid (re)coloniser of freshwater systems of variable environmental conditions (Chlyeh *et al.*, 2006). Overall, in Serbian rivers, 21 alien species of aquatic macroinvertebrates have been identified and classified as white, black and grey, depending on their degree of invasiveness (Zorić *et al.*, 2010a,b; Raković, 2015).

The family Physidae is characterized by high intra-species genetic variability (Journey, 2014), which is corroborated by our results. The first clade, which includes *P. spelunca*, was observed to be phylogenetically distinct, as expected. This species lives in a hot spring in a cave, filled with toxic sulphuric gas and feeds primarily on bacteria (Turner & Clench, 1974). Differences in penial morphology of the *acuta* group have been observed, i.e. the eastern forms (*P. acuta*, *P. heterostrophia* and *P. integra*) have transparent, muscular penial sheaths, while the sheaths of the western forms (*P. virgata* and *P. spelunca*) are less transparent and more opaque (Wethington & Lydeard, 2007).

The second clade within the Physidae is characterized by relatively high genetic distance (up to 5.58%) for the analyzed 16S rDNA gene, and this result is consistent with previous findings, of samples from other regions (Wethington *et al.*, 2009). The samples from Serbia exhibit considerably distant positions in the haplotype network of the *P. acuta* group. One haplotype group clusters with samples from Canada and Pennsylvania, while the other group is closer to a population from New Mexico, corroborating previous claims that these are actually one panmictic population and not a separate species. The high genetic variability, determined in both the native and the introduced populations, makes it difficult to draw conclusions about the origin of the invasive stock in Europe and Serbia. In addition, despite the high genetic differences within *P. acuta*, there is no reproductive isolation, as reported by Dillon *et al.* (2002), who sampled six populations from diverse environments across North America and Europe. These authors also suggested that the North American populations appear to be relatively stable and morphologically diverse; their range is largely prehistoric and taxonomists have recognized numerous subspecies and forms. Elsewhere the species seems to be more morphologically uniform and biologically invasive, which is the case of the population in Serbia. The high genetic variability of alien populations of *P. acuta* most likely contributes to the success of this species in its establishment, maintenance and dispersal in new habitats (Hartl & Clark, 1989). The observed high genetic variability, despite the small number of analyzed samples, also suggests that there were potentially multiple introductions of the species from different locations. Further studies are required to further elucidate the dynamics of *P. acuta*.

Acknowledgements

This study was supported by the Ministry of Education, Science and Technological Development of the Republic of Serbia, Grants No. 176018, No. 173025, No. 43001 and the European Union's Seventh Framework Programme under grant agreement No 603629-ENV-2013-6.2.1-GLOBAQUA. We are grateful to Miloš Jović for his assistance in the construction of the distribution map. We also express our thanks to Dr. Goran Poznanović for his constructive comments and English proof reading.

References

- Arbaciauskas, K., Semenchenko, V., Grabowski, M., Leuven, R.E.S.W., Paunović, M. *et al.*, 2008. Assessment of biological contamination of benthic macroinvertebrate communities in European inland waterways. *Aquatic Invasions*, 3 (2), 206-224.
- Bij de Vaate, A., Jazdzewski, K., Ketelaars, H.A., Gollasch, S., Van der Velde, G., 2002. Geographical pattern in range

- extension of Ponto-Caspian macroinvertebrate species in Europe. *Canadian Journal of Fisheries and Aquatic Sciences*, 59 (7), 1159-1174.
- Burch, J.B., 1989. *North American freshwater snails*. Malacological Publications, Hamburg, Michigan, 365 pp.
- Chlyeh, G., Dodet, M., Delay, B., Khallaayoune, K., Jarne, P., 2006. Spatio-temporal distribution of freshwater snail species in relation to migration and environmental factors in an irrigated area from Morocco. *Hydrobiologia*, 553 (1), 129-142.
- Clement M., Posada D., Crandall K.A., 2000. TCS: a computer program to estimate gene genealogies. *Molecular Ecology*, 9 (10), 1657-1659.
- de Jong, Y., Verbeek, M., Michelsen, V., de Place Bjørn, P., Los, W., et al., 2014. Fauna Europaea—all European animal species on the web. *Biodiversity Data Journal*, 2, e4034.
- DeWitt, T.J., Robinson, B.W., Wilson, D.S., 2000. Functional diversity among predators of a freshwater snail imposes an adaptive trade-off for shell morphology. *Evolutionary Ecology Research*, 2, 129-148.
- Dillon, R.T., 2000. *The Ecology of Freshwater Molluscs*. Cambridge University Press, United Kingdom, 509 pp.
- Dillon, R.T., Wethington, A.R., Rhett, J.M., Smith, T.P., 2002. Populations of the European freshwater pulmonate *Physa acuta* are not reproductively isolated from American *Physa heterostropha* or *Physa integra*. *Invertebrate Biology*, 121 (3), 226-234.
- Galindo, L.A., Puillandre, N., Strong, E.E., Bouchet, P., 2014. Using microwaves to prepare gastropods for DNA barcoding. *Molecular Ecology Resources*, 14(4), 700-705.
- Glöer, P., 2002. *Süßwasser gastropoden Nord-und Mitteleuropas: Bestimmungsschlüssel. Lebensweise. Verbreitung*. Conch Books. Hackenheim, 327 pp.
- Glöer, P., Meier-Brook, C., 2003. *Süßwassermollusken: Ein Bestimmungsschlüssel für die Bundesrepublik Deutschland*. DJN, Hamburg, 138 pp.
- Hartl, D.L., Clark, A.G., 1989. *Principles of Population Genetics*. Sinauer Associates Inc.: Sunderland, MA, 635 pp.
- Journey, R. N., 2014. *Molecular characterization of the freshwater snail Physella acuta*. PhD Thesis, University of New Mexico, 454 pp.
- Kessing, B., H. Croom, A. Martin, C. McIntosh, W. O. McMillan et al., 1989. *The simple fool's guide to PCR, version 1.0*. Department of Zoology, University of Hawaii, Honolulu.
- Kraus, T., Brant, S. and Adema, C. 2014. Characterization of Trematode Cercariae from *Physella acuta* in the Middle Rio Grande. *Comparative Parasitology*, 81 (1), 105-109.
- Malone, C., 1965. Dispersal of aquatic gastropods via the intestinal tract of water birds. *Nautilus*, 78 (4), 135-139.
- Novaković, B., 2014. Distribution of the freshwater snail *Physella acuta* (Draparnaud 1805) in Serbia in the 2009-2013 period. *Zbornik radova Zaštita voda*. Tara, 77-81.
- Panov, V.E., Boris, G., Alexandrov, B.G., Arbaciauskas, K., Binimelis, R. et al., 2009. Assessing the Risks of Aquatic Species Invasions via European Inland Waterways: from Concepts to Environmental Indicators. *Integrated Environmental Assessment and Management*, 5 (1), 110-126.
- Panov, V., Alexandrov, B., Arbaciauskas, K., Binimelis, R., Copp, G.H. et al., 2010. Risk Assessment of Aquatic Invasive Species Introductions via European Inland Waterways. In: Settele, J., Penev, L., Georgiev, T., Grabaum, R., Grobelnik, V., Hammen, V., Klotz, S., Kotarac, M., Kuhn, I. (Eds). *Atlas of Biodiversity Risk*. Sofia, Pensoft Publishers, 140-3 pp.
- Paunović, M., Csányi, B., Knežević, S., Simić, V., Nenadić, D. et al., 2007a. Distribution of Asian clams *Corbicula fluminea* (Müller, 1774) and *C. fluminalis* (Müller, 1774) in Serbia. *Aquatic Invasions*, 2 (2), 105-112.
- Paunović, M., Jakovcev-Todorovic, D., Simic, V., Stojanovic, B., Cakic, P., 2007b. Macroinvertebrates along the Serbian section of the Danube River (stream km 1429-925). *Bio-logia*, 62 (2), 1-9.
- Paunović, M., Csányi, B., Simonović, P., Zorić, K., 2015. Invasive Alien Species in the Danube. In: Liška I. (Ed.) *The Handbook of Environmental Chemistry: The Danube River Basin*. Springer, Heidelberg, New York, Dordrecht, London, 389-410 pp.
- Pfleger, V., 1999. *A field guide in colour to Molluscs*. Blitz Editions. Leicester, UK, 216 pp.
- Pyšek, P., Bacher, S., Chytrý, M., Jarošík, V., Wild, J. et al., 2010. Contrasting patterns in the invasions of European terrestrial and freshwater habitats by alien plants, insects and vertebrates. *Global ecology and biogeography*, 19 (3), 317-33.
- Raković, M., 2015. *Diversity of Molluscs in the Danube River (1260 – 863.5 rkm) an taxonomic analysis of the genera Planorbarius, Radix, Physella and Ferrissia (Pulmonata: Basommatophora)*. PhD, University of Belgrade, Serbia, 147 pp.
- Sommerwerk, N., Hein, T., Schneider-Jakoby, M., Baumgartner, C., Ostojic, A. et al., 2009. Chapter 3. The Danube River Basin. In: *Rivers of Europe*. Tockner, K., Uehlinger, U., Robinson, C.T. (Eds). Elsevier, Academic Press, San Diego, 59-112 pp.
- Sommerwerk, N., Bloesch, J., Paunović M., Baumgartner C., Venohr, M. et al., 2010. Managing the world's most international river: the Danube River Basin. *Marine and Freshwater Research*, 61(7), 736-748.
- Tamura, K., Stecher, G., Peterson, D., Filipski, A., Kumar, S., 2013. MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Molecular Biology and Evolution*, 30, 2725-2729.
- Taylor, D.W., 2003. Introduction to Physidae (Gastropoda: Hygrophila) biogeography, classification, morphology. *Revista de Biologia Tropical*, Supplement, 51, 1-287.
- Te, G.A., 1978. *The systematics of the Family Physidae (Basommatophora: Pulmonata)*. PhD thesis, University of Michigan, 325 pp.
- Te, G.A., 1980. New classification system for the family Physidae (Pulmonata: Basommatophora). *Archiv für Molluskenkunde*, 110, 179-184.
- Turner, R.D., Clench, W.J., 1974. A new blind Physa from Wyoming with notes on its adaptation to the cave environment. *The Nautilus*, 88, 80-85.
- Wethington, A.R., 2004. *Phylogeny, taxonomy, and evolution of reproductive isolation in Physa (Pulmonata: Physidae)*. PhD, University of Alabama, Tuscaloosa, 119 pp.
- Wethington A.R., Guralnick R., 2004. Are populations of physids from different hot-springs distinctive lineages? *American Malacological Bulletin*, 19 (1-2), 135-144.
- Wethington, A.R., Lydeard C., 2007. A molecular phylogeny of Physidae (Gastropoda: Basommatophora) based on mito-

- chondrial DNA sequences. *Journal of Molluscan Studies*, 73 (3), 241-257.
- Wethington, A.R., Wise, J., Dillon, R.T., 2009. Genetic and morphological characterization of the Physidae of South Carolina (Gastropoda: Pulmonata: Basommatophora), with description of a new species. *The Nautilus*, 124 (4), 282-292.
- Zhadin, V.I., 1952. *Freshwater mollusks of Soviet Union*. Guide Series of Fauna Soviet Union, Russian, Zool. Instit. Akad. Nauk SSSR / ZI AN SSSR. Moskva- Leningrad. Academy of Sciences of the U.S.S.R, 368 pp.
- Zorić, K., Atanacković, A., Tomović, J., Vasiljević, B., Tubić, B., 2010a. Biological invasions of aquatic ecosystems in Serbia. Balkans regional young water professionals conference. 29-30 April 2010. Belgrade. Serbia, 69-71 pp.
- Zorić, K., Vranković, J., Cakić, P., Tomović, J., Vasiljević, B. *et al.*, 2010b. Chapter 15 Introduced species of aquatic macroinvertebrates. In: *Danube in Serbia – Joint Danube survey 2*. Paunović, M., Simonović, P., Simić, V., S. Simić (Eds). University of Belgrade, Institute for Biological Research “Siniša Stanković”, Belgrade, 267-280 pp.

Appendix I

A modified protocol for freshwater snails using MWO method:

I: Small snails – snails with shell length of 2-5 mm they were treated with electromagnetic waves in the microwave oven for 5-9 s: with 2 mm - 5 s / *Gyraulus crista* (Linnaeus, 1758); 4 mm - 6 s / *Bathyomphalus contortus* (L., 1758); 4 mm - 7 s / *Segmentina nitida* (M. 1774);

II: Medium snails – snails with shell length of 5-20 mm were treated with electromagnetic waves in the microwave oven for 8-20 s: with 8 mm - 10 s / *Anisus vortex* (L., 1758); 9 mm - 12 s (13 mm - 15 s) / *Physa acuta*

(D. 1805); 15 mm - 17 s (18 mm - 20 s) / *Planorbis* (L., 1758); 19 mm - 20 s / *Radix labiata* (R. 1835);

III: Large snails – snails with shell length of 20-40 mm were treated with electromagnetic waves in the microwave oven for a period of 20-40 s: with 25 mm - 30 s (35 mm - 40 s) / *Planorbarius corneus* (L., 1758);

IV: The largest snails – snails with shell length greater than 40 mm were treated with electromagnetic waves in the microwave for over 40 s: 47 mm - 50 s / *Lymnaea stagnalis* (L., 1758).

Appendix II

Estimates of evolutionary divergence between sequences *P. acuta* group.

Locality*	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42								
1 Nebraska																																																		
2 Pennsylvania	0.026																																																	
3 Pennsylvania	0.011	0.028																																																
4 Montana	0.000	0.026	0.011																																															
5 Arizona	0.007	0.028	0.013	0.007																																														
6 Arizona	0.013	0.030	0.015	0.013	0.015																																													
7 Mexico	0.011	0.022	0.013	0.011	0.013	0.015																																												
8 South Carolina	0.011	0.022	0.013	0.011	0.013	0.015	0.000																																											
9 South Carolina	0.013	0.024	0.015	0.013	0.015	0.017	0.007	0.007																																										
10 South Carolina	0.013	0.024	0.015	0.013	0.015	0.017	0.002	0.002	0.004																																									
11 South Carolina	0.009	0.020	0.011	0.009	0.011	0.013	0.002	0.002	0.004	0.004																																								
12 South Carolina	0.009	0.020	0.011	0.009	0.011	0.013	0.002	0.002	0.004	0.004	0.000																																							
13 South Carolina	0.011	0.022	0.013	0.011	0.013	0.015	0.004	0.004	0.007	0.007	0.002	0.002																																						
14 Wyoming	0.069	0.080	0.072	0.069	0.075	0.074	0.072	0.072	0.074	0.074	0.069	0.069	0.072																																					
15 Wyoming	0.069	0.085	0.072	0.069	0.075	0.069	0.076	0.076	0.078	0.078	0.074	0.074	0.076	0.013																																				
16 South Carolina	0.009	0.020	0.011	0.009	0.011	0.013	0.002	0.002	0.004	0.004	0.000	0.000	0.002	0.069	0.074																																			
17 South Carolina	0.013	0.024	0.015	0.013	0.015	0.017	0.007	0.007	0.004	0.009	0.004	0.004	0.007	0.074	0.078	0.004																																		
18 South Carolina	0.013	0.024	0.015	0.013	0.015	0.011	0.007	0.007	0.009	0.009	0.004	0.004	0.007	0.076	0.078	0.004	0.009																																	
19 South Carolina	0.011	0.022	0.013	0.011	0.013	0.015	0.004	0.004	0.007	0.007	0.002	0.002	0.004	0.072	0.076	0.002	0.007	0.007																																
20 South Carolina	0.013	0.024	0.015	0.013	0.015	0.017	0.002	0.002	0.004	0.004	0.004	0.004	0.007	0.074	0.078	0.004	0.004	0.009	0.007																															
21 South Carolina	0.009	0.020	0.011	0.009	0.011	0.013	0.002	0.002	0.004	0.004	0.000	0.000	0.002	0.069	0.074	0.000	0.004	0.004	0.002	0.004																														
22 New Mexico	0.017	0.035	0.020	0.017	0.020	0.013	0.020	0.020	0.022	0.017	0.017	0.020	0.080	0.076	0.017	0.020	0.017	0.020	0.020	0.017																														
23 New Mexico	0.020	0.037	0.022	0.020	0.022	0.015	0.022	0.022	0.022	0.024	0.020	0.020	0.022	0.082	0.078	0.020	0.022	0.020	0.022	0.020	0.002																													
24 Netherlands	0.015	0.033	0.017	0.015	0.017	0.009	0.017	0.017	0.020	0.020	0.015	0.015	0.017	0.076	0.072	0.015	0.020	0.015	0.017	0.020	0.015	0.011	0.013																											
25 Indiana	0.013	0.024	0.015	0.013	0.015	0.017	0.007	0.007	0.009	0.009	0.004	0.004	0.007	0.074	0.078	0.004	0.009	0.009	0.007	0.009	0.004	0.022	0.024	0.020																										
26 Indiana	0.009	0.026	0.011	0.009	0.011	0.013	0.011	0.011	0.009	0.009	0.009	0.009	0.011	0.074	0.074	0.009	0.013	0.013	0.011	0.013	0.009	0.017	0.020	0.015	0.013																									
27 Serbia	0.022	0.039	0.024	0.022	0.024	0.017	0.024	0.024	0.022	0.026	0.022	0.022	0.024	0.085	0.080	0.022	0.022	0.022	0.024	0.022	0.022	0.011	0.013	0.015	0.026	0.022																								
28 Serbia	0.022	0.039	0.024	0.022	0.024	0.017	0.024	0.024	0.022	0.026	0.022	0.022	0.024	0.085	0.080	0.022	0.022	0.022	0.024	0.022	0.022	0.011	0.013	0.015	0.026	0.022	0.000																							
29 Serbia	0.022	0.004	0.024	0.022	0.024	0.026	0.017	0.017	0.020	0.020	0.015	0.015	0.017	0.078	0.082	0.015	0.020	0.020	0.017	0.020	0.015	0.030	0.033	0.028	0.020	0.022	0.035	0.035																						
30 Serbia	0.024	0.041	0.026	0.024	0.026	0.020	0.026	0.026	0.024	0.028	0.024	0.024	0.026	0.087	0.082	0.024	0.024	0.026	0.024	0.024	0.013	0.015	0.017	0.028	0.024	0.002	0.002	0.037																						
31 Serbia	0.022	0.004	0.024	0.022	0.024	0.026	0.017	0.017	0.020	0.020	0.015	0.015	0.017	0.078	0.082	0.015	0.020	0.020	0.017	0.020	0.015	0.030	0.033	0.028	0.020	0.022	0.035	0.035	0.000	0.037																				
32 Serbia	0.022	0.004	0.024	0.022	0.024	0.026	0.017	0.017	0.020	0.020	0.015	0.015	0.017	0.078	0.082	0.015	0.020	0.020	0.017	0.020	0.015	0.030	0.033	0.028	0.020	0.022	0.035	0.035	0.000	0.037	0.000																			
33 France	0.024	0.039	0.024	0.024	0.022	0.024	0.026	0.026	0.028	0.028	0.024	0.024	0.026	0.082	0.082	0.024	0.028	0.028	0.026	0.028	0.024	0.033	0.035	0.028	0.028	0.024	0.037	0.037	0.035	0.039	0.035	0.035																		
34 Cuba	0.009	0.020	0.011	0.009	0.011	0.013	0.002	0.002	0.004	0.004	0.000	0.000	0.002	0.069	0.074	0.000	0.004	0.004	0.002	0.004	0.000	0.017	0.020	0.015	0.004	0.009	0.022	0.022	0.015	0.024	0.015	0.015	0.015	0.015	0.015	0.015	0.015	0.015	0.015	0.015	0.015	0.015	0.015	0.015	0.015	0.015	0.015			
35 Cuba	0.009	0.026	0.011	0.009	0.011	0.013	0.011	0.011	0.013	0.013	0.009	0.009	0.011	0.074	0.074	0.009	0.013	0.013	0.011	0.013	0.009	0.017	0.020	0.015	0.013	0.009	0.022	0.022	0.015	0.024	0.015	0.015	0.015	0.015	0.015	0.015	0.015	0.015	0.015	0.015	0.015	0.015	0.015	0.015	0.015	0.015	0.015	0.015		
36 Cuba	0.009	0.020	0.011	0.009	0.011	0.013	0.002	0.002	0.004	0.004	0.000	0.000	0.002	0.069	0.074	0.000	0.004	0.004	0.002	0.004	0.000	0.017	0.020	0.015	0.004	0.009	0.022	0.022	0.015	0.024	0.015	0.015	0.015	0.015	0.015	0.015	0.015	0.015	0.015	0.015	0.015	0.015	0.015	0.015	0.015	0.015	0.015	0.015		
37 Canada	0.017	0.009	0.020	0.017	0.020	0.022	0.013	0.013	0.015	0.015	0.011	0.011	0.013	0.074	0.078	0.011	0.015	0.015	0.013	0.015	0.011	0.026	0.028	0.024	0.015	0.017	0.030	0.030	0.004	0.033	0.004	0.004	0.030	0.011	0.017	0.011														
38 Arizona	0.024	0.046	0.028	0.024	0.030	0.030	0.030	0.030	0.033	0.033	0.028	0.028	0.030	0.067	0.067	0.028	0.033	0.033	0.030	0.033	0.02																													