

Mediterranean Marine Science

Vol 17, No 2 (2016)



Meta-analysis review of fish trophic level at marine protected areas based on stable isotopes data

J. J. de LOPE ARIAS, G. MATEU-VICENS, M. DEUDERO COMPANY

doi: [10.12681/mms.1628](https://doi.org/10.12681/mms.1628)

To cite this article:

de LOPE ARIAS, J. J., MATEU-VICENS, G., & DEUDERO COMPANY, M. (2016). Meta-analysis review of fish trophic level at marine protected areas based on stable isotopes data. *Mediterranean Marine Science*, 17(2), 496–507. <https://doi.org/10.12681/mms.1628>

Meta-analysis review of fish trophic levels in marine protected areas based on stable isotope data

J.J. DE LOPE ARIAS¹, G. MATEU-VICENS² and S. DEUDERO COMPANY¹

¹ Instituto Español de Oceanografía, Centre Oceanogràfic de les Balears, Moll de Ponent s/n, Palma de Mallorca, 07015, Spain

² Universitat de les Illes Balears, Palma de Mallorca, Spain

Corresponding author: salud.deudero@ba.ieo.es

Handling Editor: Paraskevi Karachle

Received: 21 December 2015 ; Accepted: 19 April 2016; Published on line: 19 July 2016

Abstract

Stable isotopes ($\delta^{15}\text{N}$) have been used to determine trophic levels in marine food webs. This study assessed if Marine Protected Areas (MPAs) affect the trophic levels of fishes based on stable isotopes in the western Mediterranean. A total of 22 studies including 600 observations were found and the final dataset consisted of 11 fish species and 146 observations comparing trophic levels inside and outside MPAs. The database was analysed by meta-analysis and the covariate selected was the level of protection (inside vs. outside MPAs). The results indicate significant differences between trophic levels inside and outside the MPAs; however, results differ from expectations since the trophic level was lower inside than outside MPAs. Three habitats were analysed (coastal lagoons, demersal, and littoral) and significant statistical differences were found among them: trophic level was higher in demersal habitats than in coastal lagoons and littoral areas. No significant differences were found in species classified by trophic functional groups. Several hypotheses are considered as possible explanations linked to protection level, time since protection, and MPA size. We debate the suitability of using the stable isotope $\delta^{15}\text{N}$ as a direct indicator of trophic level in evaluating MPA effects on food webs.

Keywords: Diet, fisheries effects, marine reserves, Mediterranean Sea.

Introduction

Marine Protected Areas (MPAs) have been established to prevent or reduce impacts exerted by humans in marine ecosystems. The benefits of the establishment of MPAs have been described both inside (García-Rubies & Zabala, 1990; Francour, 1994; Guidetti, 2002) and outside no-take areas (Russ & Alcala, 1996; Ojeda-Martinez *et al.*, 2007).

The trophic level (TL) of a species is an ecological trait that might provide valuable information on the ecological processes deriving from spatial regulations on fish catches in marine protected areas. TL could be determined by fish stomach contents analyses or by quantifying stable isotopes ratios of nitrogen (N). Stomach contents analysis provides information on prey consumption and is based on the transfer of organic matter through food chains. Species position within the food chain is determined by their diet and weight (Polis & Strong, 1996). There are several limitations to this methodology: it only provides a snapshot of the ingestion at a particular time, there exists the difficulty of distinguishing between assimilate and non-assimilated food sources, the amount that has been assimilated, and that which has been the contribution of primary producers.

Stable isotope signatures rely on the assumption of ‘we are what we eat’ and are based on the transformation of the carbon isotope (^{12}C) and nitrogen isotope (^{14}N) within an organism to its most stable isotopes, ^{13}C and ^{15}N , respectively. This process takes place because when an organism feeds, the food tends to stabilize. The ratio of stable isotopes of proteins in consumer species reflects those proteins in the diet in a predictable way (Hobson & Clark, 1992; Hobson, 1999).

This study focuses on $\delta^{15}\text{N}$ since that isotope is used to estimate the trophic level of marine organisms, especially when the base level is the primary producers (Cabana & Rasmussen, 1996). The abundance of $\delta^{15}\text{N}$ in the tissues of the consumer species is enhanced, more or less, by 3.4 ‰ relative to their prey (Vander Zanden *et al.*, 1997; Post, 2002). This type of analysis overcomes the limitations of analysis by stomach content. The results reflect the actual assimilation of nutrients and can be used for studies over a longer period of time because assimilation is reflected in the tissue analysis.

Meta-analyses have been widely applied as a tool to assess processes in MPAs so as to demonstrate the variations in ecological traits and fish life histories (Claudet *et al.*, 2008), effects of MPAs on fish populations (Coté *et al.*, 2001) as well as the interactions between cod and shrimp in oceanic food webs (Worm & Myers, 2003).

Although some meta-analyses on fisheries research have been performed since the 1990s, to our knowledge our study is the first to assess trophic level changes linked to Marine Protected Areas through the evaluation of stable isotopes in fishes.

In this context, the main aim of the present study was to investigate changes in trophic levels based on stable isotopes for fish species both inside Marine Protected Areas and outside MPAs. We have considered only data derived from stable isotopes in our analysis, not taking into consideration shifts in biomass, abundance or density related to fisheries effects. The initial hypothesis is that the fishing pressure exerted outside protected areas determines lower trophic levels of fishes with respect to those measured inside the protected areas. For that reason, we aimed to analyse how fish trophic level is influenced by protection. Additionally, we wanted to investigate differences in trophic level (TL) by habitat types, and assess differences in TL regarding trophic functional groups.

Material and Methods

This study has focused on the western Mediterranean, considering large number of observations of fish species that cover a vast array of ecosystems, coastal lagoons, demersal, and littoral areas and several functional trophic groups, carnivorous, omnivorous and piscivorous (according to FISHBASE¹).

TL for a given species can be modelled as a function of the $\delta^{15}N$, according to the equation:

$$TL_i = \frac{\delta^{15}N_i - \delta^{15}N_{ref}}{TE} + TL_{ref}$$

where TE represents the trophic enrichment and TL_{ref} refers to base trophic level in the studied ecosystem. For the Mediterranean, the final equation is:

$$TL_i = \frac{\delta^{15}N_i - \delta^{15}N_{ref}}{3.4} + 2$$

where 3.4 is the enrichment in each trophic level increase (Post, 2002) and 2 is the trophic level base at the Mediterranean Sea.

Database

Data have been obtained from a search in the ISI Web of Science using “trophic”, “stable isotopes”, and “Mediterranean” as keywords. A total of 22 studies were found until 2008, which corresponds to 600 observations.

1. Diet based on fishbase.org:

Piscivorous: Feed mainly on fishes and invertebrates.

Carnivorous: Feeds on crustaceans, decapods, mysids, worms, and molluscs

Omnivorous: Feed on seaweeds and small invertebrates, algae, crustaceans, etc

Subsequently, the information was summarised in a database in which an identity code was provided for each observation including different fields: study author, fish species and family, fish size, depth average, stable isotope ratios (carbon and nitrogen data), standard deviation, location, geographical coordinates, country, type of habitat, diet, protection effect (inside vs. outside), protection status, trophic level obtained from our calculations from stable isotope data, trophic level recorded from bibliographical cites at www.fishbase.org, and trophic level based on stomach contents reported in Stergiou & Karpouzi (2002).

From the original pool of studies, the observations lacking sample size and therefore without standard deviation estimates were rejected. The observations retained were those concerning species with information from both inside and outside the Marine Protected Area. Thus, the final dataset consists of 11 species and 146 observations. The covariate selected is the level of protection (inside vs. outside the marine protected area) (Table 1).

Analysis

To test the differences between trophic levels inside and outside the marine protected areas, the size effect (e_i) was utilised, with log-response ratio (Hedges *et al.*, 1999):

$$e_i = \ln\left(\frac{TL_{iin}}{TL_{iout}}\right)$$

where TL_{iin} and TL_{iout} are the trophic levels inside and outside the marine protected area, respectively. Subsequently, variance of the size effect (V_i) was calculated. V_i is used to derive weights in the meta-analysis to increase both the precision and the power of the test (Gurevitch & Hedges, 1999; Osenberg *et al.*, 1999). Thus, studies with more observations provide a greater weight. Variance of the size effect was calculated according to the equation:

$$V_i = \frac{(S_{iin})^2}{N_{iin}(\overline{TL}_{iin})^2} + \frac{(S_{iout})^2}{N_{iout}(\overline{TL}_{iout})^2}$$

where V_i is the variance associated to e_i , N_{iin} and N_{iout} corresponds to the number of observations inside and outside the protected area respectively, \overline{TL}_{iin} and \overline{TL}_{iout} are the mean trophic levels inside and outside the protected area respectively, and S_{iin} and S_{iout} are the standard deviations associated with the trophic levels. The variance is applied to calculate the weight for meta-analysis:

$$w_i = \frac{1}{V_i}$$

where w_i is the weight associated to the size effect e_i ; V_i is the variance defined above. Weighted cumulative effect was then calculated as:

Table 1. Author, species, stable isotope ratios trophic level (TL), standard deviation, number of samples, habitat and diet (fishbase).

Author	Species	No-take areas			Take areas			Habitat	Diet (fishbase)
		TL $\delta^{15}\text{N}$	SD	n samples	TL $\delta^{15}\text{N}$	SD	n samples		
Vizzini & Mazzola, 2008	<i>Boops boops</i>	2.90	0.10	52.00	-	-	-	Littoral	Omnivorous
Carlier <i>et al.</i> , 2007	<i>Boops boops</i>	-	-	-	3.63	0.70	3.00	Littoral	Omnivorous
Vizzini & Mazzola, 2008	<i>Boops boops</i>	2.96	0.10	27.00	-	-	-	Littoral	Omnivorous
Pinnegar <i>et al.</i> , 2000	<i>Boops boops</i>	-	-	-	2.63	0.36	3.00	Littoral	Omnivorous
Vizzini & Mazzola, 2008	<i>Boops boops</i>	2.96	0.10	27.00	-	-	-	Littoral	Omnivorous
-	<i>Boops boops</i>	-	-	-	3.13	0.53	3.00	-	-
-	<i>Boops boops</i>	2.96	0.10	27.00	-	-	-	Littoral	Omnivorous
-	<i>Boops boops</i>	-	-	-	3.13	0.53	3.00	-	-
Vizzini & Mazzola, 2008	<i>Coris julis</i>	3.01	0.20	62.00	-	-	-	Littoral	Carnivorous
Jennings <i>et al.</i> , 1997	<i>Coris julis</i>	-	-	-	3.81	0.11	4.00	Littoral	Carnivorous
Vizzini & Mazzola, 2008	<i>Coris julis</i>	3.10	0.20	60.00	-	-	-	Littoral	Carnivorous
Cardona <i>et al.</i> , 2007	<i>Coris julis</i>	-	-	-	4.10	0.70	10.00	Littoral	Carnivorous
Pinnegar <i>et al.</i> , 2000	<i>Coris julis</i>	3.19	0.50	58.00	-	-	-	Littoral	Carnivorous
Jennings <i>et al.</i> , 1997	<i>Coris julis</i>	-	-	-	4.22	0.27	6.00	Littoral	Carnivorous
Vizzini & Mazzola, 2008	<i>Coris julis</i>	3.22	0.20	70.00	-	-	-	Littoral	Carnivorous
Jennings <i>et al.</i> , 1997	<i>Coris julis</i>	-	-	-	4.25	0.11	5.00	Littoral	Carnivorous
-	<i>Coris julis</i>	3.04	0.19	47.88	-	-	-	-	-
Vizzini & Mazzola, 2008	<i>Coris julis</i>	-	-	-	3.19	0.50	58.00	Littoral	Carnivorous
-	<i>Coris julis</i>	3.04	0.19	47.88	-	-	-	-	-
Lepoint <i>et al.</i> , 2000	<i>Diplodus annularis</i>	-	-	-	3.34	0.40	2.00	Littoral	Carnivorous
Vizzini & Mazzola, 2008	<i>Diplodus annularis</i>	2.99	0.60	10.00	-	-	-	Littoral	Carnivorous
Jennings <i>et al.</i> , 1997	<i>Diplodus annularis</i>	-	-	-	3.93	0.13	5.00	Littoral	Carnivorous
Vizzini & Mazzola, 2008	<i>Diplodus annularis</i>	3.13	0.20	13.00	-	-	-	Littoral	Carnivorous
Cardona <i>et al.</i> , 2007	<i>Diplodus annularis</i>	-	-	-	4.25	0.50	10.00	Littoral	Carnivorous
Vizzini & Mazzola, 2008	<i>Diplodus annularis</i>	3.16	0.30	18.00	-	-	-	Littoral	Carnivorous
Jennings <i>et al.</i> , 1997	<i>Diplodus annularis</i>	-	-	-	4.40	0.20	5.00	Littoral	Carnivorous
-	<i>Diplodus annularis</i>	3.09	0.37	13.67	-	-	-	-	-
Jennings <i>et al.</i> , 1997	<i>Diplodus annularis</i>	-	-	-	4.40	0.20	5.00	Littoral	Carnivorous
-	<i>Diplodus annularis</i>	3.09	0.37	13.67	-	-	-	-	-
Lepoint <i>et al.</i> , 2000	<i>Diplodus annularis</i>	-	-	-	3.07	0.40	2.00	Littoral	Carnivorous
-	<i>Diplodus annularis</i>	3.09	0.37	13.67	-	-	-	-	-
Pinnegar <i>et al.</i> , 2000	<i>Diplodus annularis</i>	-	-	-	3.13	0.27	3.00	Littoral	Carnivorous
-	<i>Diplodus annularis</i>	3.09	0.37	13.67	-	-	-	-	-
Deudero <i>et al.</i> , 2004	<i>Diplodus annularis</i>	-	-	-	3.44	0.18	10.00	Littoral	Carnivorous
Badalamenti <i>et al.</i> , 2008	<i>Lophius budegassa</i>	3.90	0.15	6.00	-	-	-	Demersal	Piscivorous
Badalamenti <i>et al.</i> , 2008	<i>Lophius budegassa</i>	-	-	-	3.86	0.16	6.00	Demersal	Piscivorous
Badalamenti <i>et al.</i> , 2008	<i>Merluccius merluccius</i>	3.60	0.07	6.00	-	-	-	Demersal	Piscivorous
Badalamenti <i>et al.</i> , 2008	<i>Merluccius merluccius</i>	-	-	-	3.34	0.11	6.00	Demersal	Piscivorous
Badalamenti <i>et al.</i> , 2002	<i>Merluccius merluccius</i>	3.43	0.10	80.00	-	-	-	Demersal	Piscivorous
Badalamenti <i>et al.</i> , 2008	<i>Merluccius merluccius</i>	-	-	-	3.36	0.08	6.00	Demersal	Piscivorous
Badalamenti <i>et al.</i> , 2008	<i>Merluccius merluccius</i>	3.46	0.10	80.00	-	-	-	Demersal	Piscivorous
Badalamenti <i>et al.</i> , 2008	<i>Merluccius merluccius</i>	-	-	-	3.41	0.12	6.00	Demersal	Piscivorous
Badalamenti <i>et al.</i> , 2008	<i>Merluccius merluccius</i>	3.48	0.11	6.00	-	-	-	Demersal	Piscivorous
Badalamenti <i>et al.</i> , 2008	<i>Merluccius merluccius</i>	-	-	-	3.42	0.17	6.00	Demersal	Piscivorous
Badalamenti <i>et al.</i> , 2008	<i>Merluccius merluccius</i>	3.49	0.19	6.00	-	-	-	Demersal	Piscivorous
Badalamenti <i>et al.</i> , 2008	<i>Merluccius merluccius</i>	-	-	-	3.45	0.10	6.00	Demersal	Piscivorous
Badalamenti <i>et al.</i> , 2008	<i>Merluccius merluccius</i>	3.49	0.19	6.00	-	-	-	Demersal	Piscivorous
Badalamenti <i>et al.</i> , 2002	<i>Merluccius merluccius</i>	-	-	-	3.46	0.10	80.00	Demersal	Piscivorous
Badalamenti <i>et al.</i> , 2008	<i>Merluccius merluccius</i>	3.49	0.09	6.00	-	-	-	Demersal	Piscivorous

(continued)

Table 1 (continued)

Author	Species	No-take areas			Take areas			Habitat	Diet (fishbase)
		TL $\delta^{15}\text{N}$	SD	n samples	TL $\delta^{15}\text{N}$	SD	n samples		
Badalamenti <i>et al.</i> , 2008	<i>Merluccius merluccius</i>	-	-	-	3.48	0.12	6.00	Demersal	Piscivorous
Badalamenti <i>et al.</i> , 2008	<i>Merluccius merluccius</i>	3.50	0.12	6.00	-	-	-	Demersal	Piscivorous
Badalamenti <i>et al.</i> , 2002	<i>Merluccius merluccius</i>	-	-	-	3.49	0.10	80.00	Demersal	Piscivorous
Badalamenti <i>et al.</i> , 2008	<i>Merluccius merluccius</i>	3.51	0.12	6.00	-	-	-	Demersal	Piscivorous
Badalamenti <i>et al.</i> , 2008	<i>Merluccius merluccius</i>	3.53	0.14	6.00	3.50	0.18	6.00	Demersal	Piscivorous
Badalamenti <i>et al.</i> , 2008	<i>Merluccius merluccius</i>	-	-	-	3.55	0.09	6.00	Demersal	Piscivorous
Badalamenti <i>et al.</i> , 2008	<i>Merluccius merluccius</i>	3.56	0.10	6.00	-	-	-	Demersal	Piscivorous
Badalamenti <i>et al.</i> , 2008	<i>Merluccius merluccius</i>	-	-	-	3.55	0.17	6.00	Demersal	Piscivorous
-	<i>Merluccius merluccius</i>	3.50	0.12	19.45	-	-	-	-	-
Polunin <i>et al.</i> , 2001	<i>Merluccius merluccius</i>	-	-	-	3.59	0.21	3.00	Demersal	Piscivorous
Badalamenti <i>et al.</i> , 2008	<i>Mullus barbatus</i>	3.60	0.39	6.00	-	-	-	Demersal	Carnivorous
Badalamenti <i>et al.</i> , 2002	<i>Mullus barbatus</i>	-	-	-	3.61	0.41	6.00	Demersal	Carnivorous
Badalamenti <i>et al.</i> , 2008	<i>Mullus barbatus</i>	3.82	0.16	6.00	-	-	-	Demersal	Carnivorous
Badalamenti <i>et al.</i> , 2008	<i>Mullus barbatus</i>	3.84	0.51	6.00	3.62	0.19	6.00	Demersal	Carnivorous
Badalamenti <i>et al.</i> , 2008	<i>Mullus barbatus</i>	-	-	-	3.67	0.19	6.00	Demersal	Carnivorous
Badalamenti <i>et al.</i> , 2008	<i>Mullus barbatus</i>	3.88	0.08	6.00	-	-	-	Demersal	Carnivorous
Badalamenti <i>et al.</i> , 2008	<i>Mullus barbatus</i>	-	-	-	3.70	0.57	6.00	Demersal	Carnivorous
Badalamenti <i>et al.</i> , 2008	<i>Mullus barbatus</i>	3.49	0.38	6.00	-	-	-	Demersal	Carnivorous
Badalamenti <i>et al.</i> , 2002	<i>Mullus barbatus</i>	-	-	-	3.72	0.10	74.00	Demersal	Carnivorous
Badalamenti <i>et al.</i> , 2008	<i>Mullus barbatus</i>	3.58	0.26	6.00	-	-	-	Demersal	Carnivorous
Badalamenti <i>et al.</i> , 2002	<i>Mullus barbatus</i>	-	-	-	3.72	0.10	74.00	Demersal	Carnivorous
-	<i>Mullus barbatus</i>	3.70	0.30	6.00	-	-	-	-	-
Badalamenti <i>et al.</i> , 2008	<i>Mullus barbatus</i>	-	-	-	3.76	0.35	6.00	Demersal	Carnivorous
-	<i>Mullus barbatus</i>	3.70	0.30	6.00	-	-	-	-	-
Bautista-Vega <i>et al.</i> , 2008	<i>Mullus barbatus</i>	-	-	-	3.82	0.62	62.00	Littoral	Carnivorous
-	<i>Mullus barbatus</i>	3.70	0.30	6.00	-	-	-	-	-
Bautista-Vega <i>et al.</i> , 2008	<i>Mullus barbatus</i>	-	-	-	3.83	0.81	78.00	Littoral	Carnivorous
-	<i>Mullus barbatus</i>	3.70	0.30	6.00	-	-	-	-	-
Bautista-Vega <i>et al.</i> , 2008	<i>Mullus barbatus</i>	-	-	-	3.87	1.53	46.00	Littoral	Carnivorous
-	<i>Mullus barbatus</i>	3.70	0.30	6.00	-	-	-	-	-
Bautista-Vega <i>et al.</i> , 2008	<i>Mullus barbatus</i>	-	-	-	3.88	1.01	47.00	Littoral	Carnivorous
-	<i>Mullus barbatus</i>	3.70	0.30	6.00	-	-	-	-	-
Carlier <i>et al.</i> , 2007	<i>Mullus barbatus</i>	-	-	-	3.96	0.50	3.00	Littoral	Carnivorous
-	<i>Mullus barbatus</i>	3.70	0.30	6.00	-	-	-	-	-
Badalamenti <i>et al.</i> , 2008	<i>Mullus barbatus</i>	-	-	-	3.16	0.07	6.00	Coastal Lagoon	Carnivorous
Vizzini & Mazzola, 2008	<i>Mullus surmuletus</i>	2.93	0.40	19.00	-	-	-	Littoral	Carnivorous
Bautista-Vega <i>et al.</i> , 2008	<i>Mullus surmuletus</i>	-	-	-	3.77	0.39	44.00	Littoral	Carnivorous
Vizzini & Mazzola, 2008	<i>Mullus surmuletus</i>	2.96	0.20	38.00	-	-	-	Littoral	Carnivorous
Bautista-Vega <i>et al.</i> , 2008	<i>Mullus surmuletus</i>	-	-	-	3.83	0.96	35.00	Littoral	Carnivorous
Vizzini & Mazzola, 2008	<i>Mullus surmuletus</i>	3.10	0.30	33.00	-	-	-	Littoral	Carnivorous
Bautista-Vega <i>et al.</i> , 2008	<i>Mullus surmuletus</i>	-	-	-	3.88	0.70	37.00	Littoral	Carnivorous
-	<i>Mullus surmuletus</i>	3.00	0.30	30.00	-	-	-	-	-
Cardona <i>et al.</i> , 2007	<i>Mullus surmuletus</i>	-	-	-	3.90	1.20	10.00	Littoral	Carnivorous
-	<i>Mullus surmuletus</i>	3.00	0.30	30.00	-	-	-	-	-
Bautista-Vega <i>et al.</i> , 2008	<i>Mullus surmuletus</i>	-	-	-	3.91	1.06	35.00	Littoral	Carnivorous
-	<i>Mullus surmuletus</i>	3.00	0.30	30.00	-	-	-	-	-
Magni <i>et al.</i> , 2008	<i>Mullus surmuletus</i>	-	-	-	4.46	0.20	4.00	Coastal Lagoon	Carnivorous
-	<i>Mullus surmuletus</i>	3.00	0.30	30.00	-	-	-	-	-
Pinnegar <i>et al.</i> , 2000	<i>Mullus surmuletus</i>	-	-	-	2.73	0.17	3.00	Littoral	Carnivorous

(continued)

Table 1 (continued)

Author	Species	No-take areas			Take areas			Habitat	Diet (fishbase)
		TL $\delta^{15}\text{N}$	SD	n samples	TL $\delta^{15}\text{N}$	SD	n samples		
- Deudero <i>et al.</i> , 2004	<i>Mullus surmuletus</i>	3.00	0.30	30.00	-	0.03	-	Littoral	- Carnivorous
- Deudero <i>et al.</i> , 2004	<i>Mullus surmuletus</i>	3.00	0.30	30.00	-	-	-	-	- Carnivorous
Vizzini & Mazzola, 2008	<i>Mullus surmuletus</i>	2.99	0.30	43.00	3.59	0.14	1.00	Littoral	Piscivorous
Vizzini & Mazzola, 2006	<i>Serranus cabrilla</i>	3.13	0.20	41.00	3.60	0.30	1.00	Littoral	Piscivorous
Vizzini & Mazzola, 2006	<i>Serranus cabrilla</i>	3.16	0.30	4.00	3.13	0.20	41.00	Littoral	Piscivorous
Vizzini & Mazzola, 2008	<i>Serranus cabrilla</i>	3.25	0.30	35.00	3.51	0.40	1.00	Littoral	Piscivorous
Vizzini & Mazzola, 2006	<i>Serranus cabrilla</i>	3.10	0.20	8.00	3.42	0.30	14.33	Littoral	-
Vizzini & Mazzola, 2008	<i>Serranus scriba</i>	3.25	0.20	9.00	4.04	0.29	6.00	Littoral	Piscivorous
Jennings <i>et al.</i> , 1997	<i>Serranus scriba</i>	3.28	0.10	35.00	4.37	0.40	10.00	Littoral	Piscivorous
Vizzini & Mazzola, 2008	<i>Serranus scriba</i>	3.31	0.20	13.00	4.51	0.23	6.00	Littoral	Piscivorous
Jennings <i>et al.</i> , 1997	<i>Serranus scriba</i>	3.24	0.18	16.25	4.57	0.17	6.00	Littoral	Piscivorous
Pinnegar <i>et al.</i> , 2000	<i>Serranus scriba</i>	3.24	0.18	16.25	3.31	0.20	3.00	Littoral	Piscivorous
Jennings <i>et al.</i> , 1997	<i>Serranus scriba</i>	3.24	0.18	16.25	3.44	0.17	10.00	-	-
- Vizzini & Mazzola, 2008	<i>Serranus scriba</i>	2.78	0.10	39.00	3.46	0.33	1.00	Littoral	Piscivorous
- Deudero <i>et al.</i> , 2004	<i>Spicara maena</i>	2.78	0.10	39.00	2.73	0.32	3.00	Littoral	Carnivorous
Deudero <i>et al.</i> , 2004	<i>Spicara maena</i>	2.84	0.10	18.00	2.73	0.32	3.00	Littoral	Carnivorous
Vizzini & Mazzola, 2008	<i>Spicara maena</i>	3.22	0.60	6.00	2.73	0.32	3.00	-	-
Vizzini & Mazzola, 2008	<i>Symphodus ocellatus</i>	3.22	0.60	6.00	2.54	0.10	8.00	Littoral	Omnivorous
Vizzini <i>et al.</i> , 2002	<i>Symphodus ocellatus</i>	2.96	0.40	68.00	3.29	0.23	3.00	Coastal Lagoon	Omnivorous
Pinnegar <i>et al.</i> , 2000	<i>Symphodus tinca</i>	2.99	0.50	38.00	3.84	0.86	6.00	Littoral	Omnivorous
Vizzini & Mazzola, 2008	<i>Symphodus tinca</i>	3.01	0.40	10.00	3.99	0.40	10.00	Littoral	Carnivorous
Jennings <i>et al.</i> , 1997	<i>Symphodus tinca</i>	3.40	0.30	88.00	4.04	0.14	6.00	Littoral	Carnivorous
Vizzini & Mazzola, 2008	<i>Symphodus tinca</i>	3.09	0.40	51.00	4.10	1.42	6.00	Littoral	Carnivorous
Jennings <i>et al.</i> , 1997	<i>Symphodus tinca</i>	-	-	-	3.13	0.40	3.00	Littoral	-
Pinnegar <i>et al.</i> , 2000	<i>Symphodus tinca</i>	-	-	-	-	-	-	-	Carnivorous

$$\bar{e}_k = \frac{\sum_{i=1}^{n_k} w_i e_i}{\sum_{i=1}^{n_k} w_i}$$

where n_k is the number of studies for consideration and e_i and w_i are defined above. The fixed effect model was used since, among variance studies, it is lower than 0 and therefore, relative weights for the fixed effect model are more balanced than those for the random effect model. Total heterogeneity was obtained according to Hedges & Olkin (1985):

$$Q_{T_k} = \sum_{i=1}^{n_k} w_i (e_i - \bar{e}_k)^2$$

and the corresponding significance is tested against χ^2 distributions with $n_k - 1$ degrees of freedom. The meta-analysis was performed using R and Excel software packages.

The equation was proposed as a method to quantify heterogeneity and it is expressed in percentage of the total variability in a set of size effects due to true heterogeneity, that is, between-studies variability (Higgins & Thompson, 2002).

$$I^2 = \left(\frac{(Q - df)}{Q} \right) \times 100$$

where Q stands for “heterogeneity” and stands for “degrees of freedom”.

Results

The results indicate significant differences between fish trophic levels inside and outside MPAs. However, those differences differ from initial expectations, with higher TL outside MPAs despite the protection level (Fig. 1).

Total heterogeneity was obtained according to Hedges & Olkin (1985) by applying Q test measures heterogeneity among studies ($Q_t = 1638.93$, $p < 0.001$, $df = 145$).

The result points out the corresponding size effects were heterogeneous among the studies.

The heterogeneity (I^2) calculated in this meta-analysis was 98.71%. This result suggests that mean differences of $\delta^{15}\text{N}$ -fish varied more between protected areas and non-protected areas than could be expected by random error alone.

The final dataset used consists of eleven species and the covariate selected was the level of protection (inside vs. outside MPAs). The meta-analysis provided the following results on the carnivorous species: *Coris julis* ($e_i = -0.055 \pm 0.019$, 95% CI, $n = 145$), *Diplodus annularis* ($e_i = -0.195 \pm 0.023$, 95% CI, $n = 135$), *Mullus barbatus* ($e_i = -0.002 \pm 0.004$, 95% CI, $n = 498$), *Mullus surmuletus* ($e_i = -0.207 \pm 0.018$, 95% CI, $n = 440$), *Spicara maena* ($e_i = 0.026 \pm 0.000$, 95% CI $n=105$), *Symphodus tinca* ($e_i = -0.209 \pm 0.013$, 95% CI, $n=286$); omnivorous species: *Boops boops* ($e_i = -0.055 \pm 0.019$, 95% CI, $n=145$), *Symphodus ocellatus* ($e_i = 0.107 \pm 0.032$, 95% CI, $n=23$); piscivorous species: *Merluccius merluccius* ($e_i = 0.010 \pm 0.000$, 95% CI, $n=50$), *Serranus cabrilla* ($e_i = -0.085 \pm 0.006$, 95% CI, $n=180$) and *Serranus scriba* ($e_i = -0.193 \pm 0.018$, 95% CI, $n=155$). Significant size effect was found in all the fish species. In order to test habitat effects on the trophic level, the species were classified by habitat according to literature (Table 1). The meta-analysis showed significant size effect at all the habitats considered: coastal lagoons ($e_i = 0.037 \pm 0.004$, 95% CI, $n = 5$), demersal ($e_i = 0.006 \pm 0.001$, 95% CI, $n=41$) and littoral ($e_i = -0.144 \pm 0.004$, 95% CI, $n=100$). The results exhibit higher trophic level in demersal areas than in littoral and coastal lagoons (Fig. 2).

Similarly, to evaluate trophic changes at MPAs, the species were classified by trophic functional groups (Table 1) (Fig.3): carnivorous ($e_i = -0.126 \pm 0.003$, 95% CI, $n= 86$), omnivorous ($e_i = -0.001 \pm 0.008$, 95% CI, $n= 12$) and piscivorous ($e_i = -0.065 \pm 0.001$, 95% CI, $n= 48$). No significant effect was found in omnivorous species.




Fig. 1: Mean fish trophic level based on stable isotopes (\pm SD) inside and outside Marine Protected Areas.




Fig. 2: Mean fish trophic level (\pm SD) calculated by habitat derived from fish stable isotope data.




Fig. 3: Size effect average (\pm SD) by trophic functional groups of the fish species. No significant size effect has been found.

Discussion

The size effect for protection effects exhibits higher trophic levels outside Marine Protection Areas in spite of protection (Fig. 1). The initial hypothesis holds that inside MPAs with full protection, species have food resources available to achieve the maximum trophic level within the food web for each of the fish species considered. However, outside MPAs, species do not have plentiful food resources available because of fishing and overfishing effects so their trophic level is lower. The increased mortality caused by fishing changes both diet and trophic level (Pope & Knights, 1982). Also, a reduction in the average fish size associated with fishing could increase the apparent decline in the mean trophic level (Pauly *et al.*, 2000) and even change the trophic structure globally (Christensen *et al.*, 2014).

There is extensive literature about effects inside MPAs documenting increased density, biomass, individual size, and diversity in all functional groups (Dufour *et al.*, 1995; Halpern & Warner, 2002; Halpern, 2003), increased abundance and richness of fishes (Rowley, 1994; Bohnsack, 1998; Barrett *et al.*, 2007), improvement in the conservation and restoration of population and marine habitats (Fox *et al.*, 2012), avoidance of depletion of high trophic level populations due to fishing, and recovery of natural ecological structures (Jones, 2014) and management, education, and research issues (Angulo-Valdés & Hatcher, 2010). However, there is not much literature about how the establishment of MPAs modifies the species trophic level based on stable isotopes. Badalamenti *et al.* (2002) reported an increase of trophic level (based on $\delta^{15}\text{N}$) of some fish species (*M. merluccius*, *M. barbatus* and *Lophius budegassa*) after nine years of no-trawling at the Gulf of Castellammare.

Our overall results indicate that the trophic level does not increase with the presence of the protection itself. This conclusion is in line with previous studies indicating that trophic level does not necessarily increase with protection and in some cases it can even decrease (Deudero *et al.*, 2004; Vizzini & Mazzola, 2009; Sinopoli *et al.*, 2012).

The establishment of MPAs provides benefits within MPAs borders but also to adjacent no-take areas. For example, there are increases of fish abundance (Ojeda-Martinez *et al.*, 2007), density and species richness (Russ & Alcala, 1996) linked to the “spill-over effect” (Harmelin-Vivien *et al.*, 2008; Stobart *et al.*, 2009; Chateau & Wantiez, 2009), and/or a surplus of fish available in adjacent areas (Roberts & Polunin, 1991; DeMartini, 1993; McClanahan & Magni, 2000). However, no studies have been found to support the fact that trophic levels could be higher outside MPAs than inside in spite of protection, as this study concludes.

Protection Level

In this study, there is a wide array of protection status at the studied MPAs: fishing is prohibited in Columbretes and some parts of “Egadi Islands”, trawling is prohibited in some parts of “Egadi Island” and Gulf of Castellammare, and professional fishing and recreational marine fishing are under regulation in Cala Ratjada. The different fishing types developed inside the MPAs could have indirect effects on trophic structures (Pauly *et al.*, 1998) and alter the sources of primary production (Kaiser & Spencer, 1994) with implications for prey (Pinnegar *et al.*, 2000). Each type of fishing is aimed at capturing a specific type of organism, which occupies a different position within the food chain (Stergiou *et al.*, 2004). For example, angling aims at fishing for larger species of a higher trophic level, while trawling, purse or dredging are directed towards catching smaller species placed at lower trophic levels, such as shellfish and crustaceans. These types of fishing have increased in the last 50 years, with a consequent reduction in both sizes and trophic level of the catches (Watson *et al.*, 2006).

In areas where fishing is restricted the target species increase both their size and biomass (Roberts & Polunin, 1991). For instance, piscivorous species (predators) prey on small fishes, decreasing both their numbers and sizes and, this in turn, causes an increase in invertebrate species at the base of the food chain. Therefore, one would expect a higher trophic level where fishing is restricted than in fishing areas (Pauly *et al.*, 2001; Pinnegar *et al.*, 2002). In the case of Mallorca Island, sport fishing impacts coastal fish communities: Morales-Nin *et al.* (2005) estimated that recreational fisheries exerted an impact on the majority of species within trophic level 4 (31%), representing at least 1209.25 tons/year catches. This might be a feasible explanation of the results obtained in this study, related to a low protection level in the MPAs analysed.

Establishment of MPAs

Other factors regarding processes at MPAs affecting life history and ecology of marine species are dependent on time since establishment of the MPAs and the size of the no-take zone (Claudet *et al.*, 2008). There are several analyses where the “reserve effect” emerges during different periods of time. For example, after the first year an increase in biomass was observed (Russ & Alcala, 1996); during the first three years of protection, increases in density, biomass, and diversity occur (Halpern & Warner, 2002) with a significant increase in density, species richness, and size of the target (Seytre & Francour, 2008). Six years after MPA establishment, significant differences were found in total abundance, species richness, and diversity on species such as *S. cabrilla* and *C. julis* (Claudet *et al.*, 2006); 5-10 years after protection, the “reserve effect” emerges (Gell & Roberts, 2003) and MPAs are more effective when maintained for at least 15 years following establishment (Molloy *et al.*, 2009).

In our study, several temporal scenarios in MPA designation are included. Thus, “Egadi Islands” were established in 1991, Gulf of Castellammare was established in 1990, Columbretes were established in 1990, and Cala Ratjada was established in 2007. Some studies reported increase abundance, biomass, mean body size, and trophic level (based on stomach content) after 8 - 16 years of no fishing at Columbretes (Stobart *et al.*, 2009). Therefore, contrasting responses at temporal scales seems to be relevant from studies including 15 years of effective protection.

The trophic level responses of fish at MPAs are in agreement with previous research from Claudet *et al.* (2008) that exhibited no effect on economic value, body size, habitat, depth range and schooling behaviour of several species at MPAs in 12 locations of the Mediterranean Sea. Similarly, Tunesi *et al.* (2006) did not record a clear “reserve effect” after five years of protection in spite of significant differences in the size of the target species. That lack of results could be due to the critical lack of empirical knowledge about MPAs and the unsuccessful conservation measures made (Planes *et al.*, 2000; Fraschetti *et al.*, 2002; Claudet & Fraschetti, 2010), or a lack of knowledge in important scientific facts such as knowledge about dispersal schemes, the geographical range of the population, an appropriate experimental design, and environmental evaluation (Claudet & Pelletier, 2004). Establishment of appropriate MPA designation within the proper context such as a large-scale marine spatial planning has to be made (Agardy *et al.*, 2011).

Food sources

Habitat effects on the trophic levels have been analysed. Our results showed significant size effect in all the habitats considered (Fig.2). However, in our analyses, demersal habitats have a trophic level higher than coastal lagoons and littoral areas, which have similar trophic levels.

The species found in demersal areas are *M. merluccius* and *M. barbatus*. *Mullus barbatus* is carnivorous, feeding mainly on benthic invertebrates (Labropoulou &

Eleftheriou, 1997) such as the shrimp *Leptochela pugnax* (Golani & Galil, 1991). During its life cycle an increase in consumption of polychaetes and shrimps with predator size occurs along with a decrease in the smaller crustaceans consumption (Bautista-Vega *et al.*, 2008). *Merluccius merluccius* is piscivorous but juveniles feed on small crustaceans such as euphausiids and mysids (Bozzano *et al.*, 1997), shifting to fish prey when older as *M. merluccius*, *Argentina sphyraena*, *Spicara flexuosa*, *Centracanthidae*, *M. barbatus* (Carpentieri *et al.*, 2005). Both fish species *Merluccius* and *Mullus* feed on organisms with high nitrogen content (Bautista-Vega *et al.*, 2008; Froese & Pauly, 2014), explaining the higher trophic level than species that live in littoral habitats and coastal lagoons.

Littoral habitats and coastal lagoons exhibit high fish trophic levels (Fig. 2) since both habitats are highly influenced by many human activities impacting the functioning of coastal marine ecosystems. Due to human influences the inputs of nutrients with $\delta^{15}\text{N}$ from sewage and aquaculture facilities are variable (Vizzini & Mazzola, 2004). Studies of the largest sources of nitrogen (air, fertilizer, soil, and waste water) show that waste water has the higher content $\delta^{15}\text{N}$ (the modal value of $>15\text{‰}$) compared to other sources with lower values of $\delta^{15}\text{N}$ (between -5‰ and 5‰). All these nitrogen loads combined with fraction occurring during the ammonification processes (Cabana & Rasmussen, 1996) increase nutrients and organic matter in the ecosystem, resulting in eutrophication. Eutrophication may cause a reduction in species richness and diversity and indirectly change the trophic level occupied by each species in the system. Enrichment of $\delta^{15}\text{N}$ is significantly higher in coastal areas than in marine ecosystems due to anthropogenic influences (McClelland & Valiela, 1998; McGhie *et al.*, 2000; Umezawa *et al.*, 2002). The rise in available nitrogen makes $\delta^{15}\text{N}$ values increase in all benthic primary producers occurring during the ammonification processes and in the main herbivorous species, such as *Paracentrotus lividus* and *Sarpa salpa* (Prado *et al.*, 2009).

The trophic level can be influenced both by species diet and by human-derived activities. The lack of knowledge about the appropriate determination of the food sources represents limitations of trophic level estimation by stable isotope analysis. The uncertainty about species diet could limit trophic level estimation due to the fact that diet is a direct reflection on the trophic level. For this reason, the combined use of stable isotopes and stomach analysis could decrease this uncertainty (Ramsvatn, 2013; Shannon *et al.*, 2014).

Among the aims of this study the size effect change of trophic functional groups was assessed (Fig.3). Omnivorous species did not show significant size effect (*B. boops* and *S. ocellatus*), linked to the generalist diet of omnivorous species. Accordingly, the diet of *B. boops* is composed of zooplanktonic crustaceans (copepods), fish eggs, gastropods, macroalgae, *Posidonia oceanica* (Cresson *et al.*, 2014), and *S. ocellatus* feeds on zoobenthos (gastropods, polychaetes, copepods), plants, and zooplankton (mainly fish eggs and larvae) (Froese &

Pauly, 2014). Due to their diet, higher wide trophic levels could be expected in these species than in piscivorous species. Resource limitation outside the MPAs due to fishing and overfishing along with reserve effect within MPA boundaries could explain the lower trophic level recorded outside the MPA.

Life cycle

In this meta-analysis study, different trophic levels associated with fish species have been analysed, including different fish sizes. Size is a factor that might influence the species trophic level, linked to ontogenetic changes in fish diets and increasing TL with fish size (du Buit, 1995; Pinnegar *et al.*, 2003; Sarà & Sarà, 2007). The body mass of fish can increase by five times during their life cycle, and the trophic level increases with body size during ontogeny (Jennings *et al.*, 2002). Also, mean trophic level measured over time and space and its high variability may mask the potential of trophic level associated with the dynamics of the food chain (Greenstreet *et al.*, 1997) and ontogenetic change. Moreover, the trophic level of a species can be altered by the top-down and bottom-up dynamics within the trophic structure of an ecosystem (Devaraj, 2004; Shackell *et al.*, 2010).

$\delta^{15}\text{N}$ as direct indicator of trophic level

In this investigation, $\delta^{15}\text{N}$ has been considered as a direct indicator of trophic level; however, based on our results combined with published studies (Vanderklint & Ponsard, 2003; Schmidt *et al.*, 2004; Mancinelli *et al.*, 2013), other factors may bias trophic level interpretation. Thus, the isotopic fractionation should be considered. For example, with isotopic fractionation associated with herbivorous species, some studies show an increase higher than 3.4‰ relative to their prey, due to differences found in enzymatic and digestive systems of herbivorous species (Mill *et al.*, 2007; Wyatt *et al.*, 2010). Olive *et al.* (2003) indicated trophic level may be distorted by the feeding ratio, the nitrogen content of food, excretion rate, and assimilation efficiency.

The present study has showed differences in fish trophic level from stable isotope data obtained from the literature. Higher fish trophic levels are observed outside MPAs, indicating that many factors are involved in trophic responses and that protection by itself cannot guarantee the increase in food resources for fish species. The suitability of using the stable isotope ($\delta^{15}\text{N}$) as a direct indicator of trophic level in evaluating MPA effects on food webs should be considered carefully.

Acknowledgements

This research was partly funded by the European project COCONET 'Towards Coast to Coast Network of marine protected areas (from the shore to the high sea), coupled with sea-based wind energy potential' grant agreement 287844.

References

- Angulo-Valdés, J., Hatcher, B., 2010. A new typology of benefits derived from marine protected areas. *Marine Policy*, 34 (3), 635-644.
- Agardy, T., di Sciara, G., Christie, P., 2011. Mind the gap: addressing the shortcomings of marine protected areas through large scale marine spatial planning. *Marine Policy*, 35, 226-232.
- Badalamenti, F., D'Anna, G., Pinnegar, J., Polunin, N., 2002. Size-related trophodynamic changes in three target fish species recovering from intensive trawling. *Marine Biology*, 141 (3), 561-570.
- Badalamenti, F., Sweeting, C., Polunin, N., Pinnegar, J., D'Anna, G. et al., 2008. Limited trophodynamics effects of trawling on three Mediterranean fishes. *Marine Biology*, 154, 765-773.
- Barrett, S., Edgar, J., Buxton, D., Haddon, M., 2007. Changes in fish assemblages following 10 years of protection in Tasmanian marine protected areas. *Journal of Experimental Marine Biology and Ecology*, 345 (2), 141-157.
- Bautista-Vega, A., Letourneur, Y., Harmelin-Vivien, M., Salen-Picard, C., 2008. Difference in diet and size-related trophic level in two sympatric fish species, the red mullets *Mullus barbatus* and *Mullus surmuletus*, in the Gulf of Lions (north-west Mediterranean Sea). *Journal of Fish Biology*, 73, 2402-2420.
- Bell, D., Harmelin-Vivien, M., 1983. Fish fauna of French Mediterranean *Posidonia oceanica* seagrass meadows. 2. *Feeding habits*. *Tethys*, 11, 1-4.
- Bohnsack, J., 1998. Applications of marine reserves to reef fisheries management. *Australian Journal of Ecology*, 23, 298-304.
- Bozzano, A., Recasens, L., Sartor, P., 1997. Diet of the european hake *Merluccius merluccius* (Pisces: Merlucciidae) in the Western Mediterranean (Gulf of Lions). *Scientia Marina*, 61 (1), 1-8.
- Cabana, G., Rasmussen, J., 1996. Comparison of aquatic food chains using nitrogen isotopes. *Proceedings of the National Academy of Sciences of the United States of America*, 93, 10844-10847.
- Cardona, L., Revelles, M., Sales, M., Aguilar, A., Borrel, A., 2007. Meadows of the seagrass *Posidonia oceanica* are a significant source of organic matter for adjoining ecosystems. *Marine Ecology Progress Series*, 335, 123-131.
- Carlier, A., Riera, P., Amouroux, J., Bodiou, J.-Y., Gremare, J., 2007. Benthic trophic network in the Bay of Banyuls-sur-Mer (northwest Mediterranean, France): an assessment based on stable carbon and nitrogen isotopes analysis. *Estuarine Coastal and Shelf Science*, 72, 1-15.
- Carpentieri, P., Colloca, F., Cardinale, M., Belluscio, A., Ardizzone, G., 2005. Feeding habits of European hake (*Merluccius merluccius*) in the central Mediterranean Sea. *Fishery Bulletin*, 103 (2), 411-416.
- Chateau, O., Wantiez, L., 2009. Movement patterns of four coral reef fish species in a fragmented habitat in New Caledonia: implications for the design of marine protected area networks. *ICES Journal of Marine Science*, 66, 50-55.
- Christensen, V., Coll, M., Piroddi, C., Steenbeek, J., Buszowski, J. et al., 2014. A century of fish biomass decline in the ocean. *Marine Ecology Progress Series*, 512, 155-166.
- Claudet, J., Fraschetti, S., 2010. Human-driven impacts on marine habitats: A regional meta-analysis in the Mediterranean Sea. *Biological Conservation*, 134, 2195-2206.
- Claudet, J., Pelletier, D., 2004. Marine protected areas and artificial reefs: A review of the interactions between management and scientific studies. *Aquatic Living Resources*, 17, 129-138.
- Claudet, J., Osenberg, C., Benedetti-Cecchi, L., Domenici, P., García-Charton, J. et al., 2008. Marine reserves: size and age do matter. *Ecology Letters*, 11, 481-489.
- Claudet, J., Osenberg, P., Domenici, F., Badalamenti, M., Milazzo, J. et al., 2010. Marine reserves: Fish life history and ecological traits matter. *Ecological Applications*, 20, 830-839.
- Claudet, J., Pelletier, D., Jouvenel, Y., Bacher, F., Galzin, R., 2006. Assessing the effects of marine protected area (MPA) on a reef fish assemblage in a northwestern Mediterranean marine reserve: identifying community-based indicators. *Biological Conservation*, 130 (3), 349-369.
- Côté, I., Mosqueira, I., Reynolds, J., 2001. Effects of marine reserve characteristics on the protection of fish populations: a meta-analysis. *Journal of Fish Biology*, 59, 178-189.
- Cresson, P., Ruitton, S., Ourgaud, M., Harmelin-Vivien, M., 2014. Contrasting perception of fish trophic level from stomach content and stable isotope analyses: A Mediterranean artificial reef experience. *Journal of Experimental Marine Biology and Ecology*, 452, 54-62.
- du Buit, M., 1995. Food and feeding of cod (*Gadus morhua* L.) in the Celtic Sea. *Fisheries Research*, 22, 227-241.
- Devaraj, M., 2004. Three tank model. A top down or bottom up dominance analysis. *Journal of Theoretical*, 6, 5.
- DeMartini, E., 1993. Modeling the potential of fishery reserves for managing Pacific coral reef fishes. *Fishery Bulletin*, 91, 414-427.
- Deudero, S., Pinnegar, J., Polunin, N., Morey, G., Morales-Nin, B., 2004. Spatial variation and ontogenic shifts in the isotopic composition of Mediterranean littoral fishes. *Marine Biology*, 145, 971-981.
- Dufour, V., Jouvenel, J., Galzin, R., 1995. Study of a Mediterranean reef fish assemblage. Comparisons of population distributions between depths in protected and unprotected areas over one decade. *Aquatic Living Resources*, 8, 17-25.
- Fox, H., Mascia, M., Basurto, X., Costa, A., Glew, L. et al., 2012. Reexamining the science of marine protected areas: linking knowledge to action. *Conservation Letters*, 5, 1-10.
- Francour, P., 1994. Pluriannual analysis of the reserve effect on ichthyofauna in the Scandola natural reserve (Corsica, north-western Mediterranean). *Oceanologica Acta*, 17, 309-317.
- Fraschetti, S., Terlizzi, A., Micheli, F., Benedetti-Cecchi, L., Boero, F., 2002. Marine Protected Areas in the Mediterranean Sea: Objectives, Effectiveness and Monitoring. *Marine Ecology*, 23, 190-200.
- Froese, R., Pauly, D., 2014 (Eds). *Fish Base. World Wide Web Electronic Publication*. <http://www.fishbase.org> (version 11/2014) (Accessed 15 November 2014).
- García-Rubies, A., Zabala, M., 1990. Effects of total fishing prohibition on the rocky fish assemblages of Medes Island marine reserve (NW Mediterranean). *Scientia Marina*, 54, 317-328.
- Gell, F., Roberts, C., 2003. Benefits beyond boundaries: the fishery effects of marine reserves. *Trends in Ecology and Evolution*, 18, 448-455.
- Golani, D., Galil, B., 1991. Trophic relationship of colonizing and indigenous goatfishes (Mullidae) in the eastern Mediterranean with special emphasis on decapod crustaceans. *Hydrobiologia*, 218 (1), 27-33.
- Greenstreet, S., Bryant, A., Broekhuizen, N., Hall, S., Heath, M., 1997. Seasonal variation in the consumption of food by fish in

- the North Sea and implications for food web dynamics. *ICES Journal of Marine Science*, 54, 243-266.
- Guidetti, P., 2002. The importance of experimental design in detecting the effects of protection measures on fish in Mediterranean MPAs. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 12, 619-634.
- Gurevitch, J., Hedges, L., 1999. Statistical issues in ecological meta-analyses. *Ecology (USA)*, 80, 1142-1149.
- Harmelin-Vivien, M., Le Diréach, L., Bayle-Sempere, J., Charbonnel, E., García-Charton, J. *et al.*, 2008. Gradients of abundance and biomass across reserve boundaries in six Mediterranean marine protected areas: Evidence of fish spillover? *Biological Conservation*, 141 (7), 1829-1839.
- Halpern, S., 2003. The impact of marine reserves: do reserves work and does reserve size matter? *Ecological Applications*, 13 (1), S117-S137.
- Halpern, B., Warner, R., 2002. Marine reserves have rapid and lasting effects. *Ecology Letters*, 5, 361-366.
- Hedges, L., Olkin, I., 1985. *Statistical methods for meta-analysis*. Academic Press, Inc., USA, 369 pp.
- Hedges, L., Gurevitch, J., Curtis, P., 1999. The meta-analysis of response ratios in experimental ecology. *Ecology (USA)*, 80, 1150-1156.
- Higgins, J., Thompson, S., 2002. Quantifying heterogeneity in a meta-analysis. *Statistic in Medicine*, 21, 1539-1558.
- Hobson, K., 1999. Tracing origins and migration of wildlife using stable isotopes: a review. *Oecologia*, 120, 314-326.
- Hobson, K., Clark, R., 1992. Assessing avian diets using stable isotopes I: turnover of $\delta^{13}\text{C}$ in tissues. *Condor*, 94, 181-188.
- Jennings, S., Renones, O., Morales-Nin, B., Polunin, N., Moranta, J. *et al.*, 1997. Spatial variation in the ^{15}N and ^{13}C stable isotope composition of plants, invertebrates and fishes on Mediterranean reefs: implications for the study of trophic pathways. *Marine Ecology Progress Series*, 146, 109-116.
- Jennings, S., Warr, K., Mackinson, S., 2002. Use of size-based production and stable isotope analyses to predict trophic transfer efficiencies and predator-prey body mass ratios in food webs. *Marine Ecology Progress Series*, 240, 11-20.
- Jones, P., 2014. *Governing Marine Protected Areas. Resilience through diversity*. Routledge, USA, 239 pp.
- Kaiser, M., Spencer, B., 1994. Fish scavenging behavior in recently trawled areas. *Marine Ecology Progress Series*, 112, 41-49.
- Khuory, C., 1987. *Ichtyofauna des herbiers des Posidonies du parc national de Port-Croix: composition, éthologie, alimentaire et rôle dans le réseau trophique*. MSc thesis. Université Aix-Marseille, France, 230 pp.
- Labropoulou, M., Eleftheriou, A., 1997. The foraging ecology of two pairs of congeneric demersal fish species: importance of morphological characteristics in prey selection. *Journal of Fish Biology*, 50, 324-340.
- Lepoint, G., Nyssen, F., Gobert, S., Dauby, P., Bouqueneau, J., 2000. Relative impact of a seagrass bed and its adjacent epilithic algal community in consumer diets. *Marine Biology*, 136, 513-518.
- Magni, P., Rajagopal, S., Van der Velde, G., Fenzi, G., Kassenberg, J. *et al.*, 2008. Sediment features, macrozoobenthic assemblages and trophic relationships (^{13}C and ^{15}N analysis) following a dystrophic event with anoxia and sulphide development in the Santa Giusta lagoon (western Sardinia, Italy). *Marine Pollution Bulletin*, 57, 125-136.
- Mancinelli, G., Vizzini, S., Mazzola, A., Maci, S., Basset, A., 2013. Cross-validation of $\delta^{15}\text{N}$ and FishBase estimates of fish trophic position in a Mediterranean lagoon: the importance of isotopic baseline. *Estuarine Coastal and Shelf Science*, 135, 77-85.
- McClanahan, T., Mangi, S., 2000. Spill-over of exploitable fishes from a marine park and its effect on the adjacent fishery. *Ecological Application*, 10, 1792-1805.
- McClelland, J., Valiela, I., 1998. Changes in food web structure under the influence of increased anthropogenic nitrogen inputs to estuaries. *Marine Ecology Progress Series*, 168, 259-271.
- McGhie, T., Crawford, C., Mitchell, I., O'Brien, D., 2000. The degradation of fish-cage waste in sediments during fallowing. *Aquaculture*, 187, 351-366.
- Mill, A., Pinnegar, J., Polunin, N., 2007. Explaining isotope trophic-step fractionation: why herbivorous fish are different. *Functional Ecology*, 21, 1137-1145.
- Molloy, P., McLeant, B., Côté, M., 2009. Effects of marine reserves on fish populations: a global meta-analysis. *Journal of Applied Ecology*, 46, 743-751.
- Morales-Nin, B., Moranta, J., García, C., Tugores, M., Grau, A. *et al.*, 2005. The recreational fishery off Majorca Island (western Mediterranean): some implications for coastal resource management. *ICES Journal of Marine Science*, 62, 727-739.
- Ojeda-Martinez, C., Bayle-Sempere, J., Sánchez-Jerez, P., Forcada, A., Valle, C., 2007. Detecting conservation benefits in spatially protected fish populations with meta-analysis of long term monitoring data. *Marine Biology*, 151 (3), 1153-1161.
- Olive, P., Pinnegar, J., Polunin, N., Richards, G., Welch, R., 2003. Isotope trophic-step fractionation: a dynamic equilibrium model. *Journal of Animal Ecology*, 72, 608-617.
- Osenberg, C., Samelle, O., Goldberg, D., 1999. Meta-analysis in ecology: concepts, statistics, and applications. *Ecology (USA)*, 80, 1103-1104.
- Pauly, D., Christensen, V., Dalsgaard, J., Froese, R., Francisco, T., 1998. Fishing down marine food webs. *Science*, 279, 860-863.
- Pauly, D., Christensen, V., Froese, R., Palomares, M., 2000. Fishing down aquatic food webs. *American Scientist January-February*, 46-52.
- Pauly, D., Palomares, M., Froese, R., Sa-a, P., Vakily, M. *et al.*, 2001. Fishing down Canadian aquatic food webs. *Canadian Journal of Fisheries and Aquatic Sciences*, 58 (1), 51-62.
- Pinnegar, J., Polunin, N., Francour, P., Badalamenti, F., Chemello, R. *et al.*, 2000. Trophic cascades in fisheries and protected-area management of benthic marine ecosystems. *Environmental Conservation*, 27, 179-200.
- Pinnegar, J., Jennings, S., O'Brien, C., Polunin, N., 2002. Long-term changes in the trophic level of the Celtic Sea fish community and fish market price distribution. *Journal of Applied Ecology*, 39, 377-390.
- Pinnegar, J., Trenkel, V., Tidd, A., Dawson, W., du Buit, M., 2003. Does diet in Celtic Sea fishes reflect prey availability? *Journal of Fish Biology*, 63, 197-212.
- Polis, G., Strong, D., 1996. Food web complexity and community dynamics. *American Naturalist*, 147, 813-846.
- Polunin, N., Morales-Nin, B., Pawsey, W., Cartes, J., Pinnegar, J. *et al.*, 2001. Feeding relationships in Mediterranean bathyal assemblages elucidated by stable nitrogen and carbon isotope data. *Marine Ecology Progress Series*, 220, 13-23.
- Pope, J., Knights, J., 1982. Comparison of length distributions of combined catches of all demersal fishes in survey in the North Sea and at Faroe Bank. p. 116-118. In: *Multispecies ap-*

- proaches to fisheries management advice. Mercer MC (ed). Canadian Special Publication of Fisheries and Aquatic Sciences, Ottawa.
- Post, D., 2002. Using stable isotopes to estimate trophic position: models, methods and assumptions. *Ecology (USA)*, 83, 703-718.
- Planes, S., Galzin, R., García-Rubies, A., Goñi, R., Harmelin, J. *et al.*, 2000. Effects of marine protected areas on recruitment processes with special reference to Mediterranean littoral ecosystems. *Environmental Conservation*, 2, 126-143.
- Prado, P., Alcoverro, T., Romero, J., 2009. Influence of nutrients in the feeding ecology of seagrass (*Posidonia oceanica* L.) consumers: a stable isotopes approach. *Marine Biology*, 157, 715-724.
- Ramsvatn, S., 2013. *Investigating coastal ecosystem structure and dynamics using Ecopath mass-balance modeling and stable isotope data*. MSc thesis, University of Tromsø, Norway, 46 pp.
- Roberts, C., Polunin, N., 1991. Are marine reserves effects in management of reef fisheries? *Reviews in Fish Biology and Fisheries*, 1, 65-91.
- Rowley, R., 1994. Marine reserves in fisheries management. *Aquatic Conservation*, 4, 233-254.
- Russ, G., Alcala, A., 1996. Do marine reserve export adult fish biomass? Evidence from Apo Island, central Philippines. *Marine Ecology Progress Series*, 132, 1-9.
- Sarà, G., Sarà, R., 2007. Feeding habits and trophic levels of bluefin tuna *Thunnus thynnus* of different size classes in the Mediterranean Sea. *Journal of Applied Ichthyology*, 23, 122-127.
- Schmidt, K., McClelland, J., Mente, E., Montoya, J., Atkinson, A. *et al.*, 2004. Trophic-level interpretation based on $\delta^{15}\text{N}$ values: implication of tissue-specific fractionation and amino-acid composition. *Marine Ecology Progress Series*, 266, 43-58.
- Seytre, C., Francour, P., 2008. Is the Cape Roux marine protected area (Saint-Raphaël, Mediterranean Sea) an efficient tool to sustain artisanal fisheries? First indications from visual censuses and trammel net sampling. *Aquatic Living Resources*, 21, 297-305.
- Shackell, N., Frank, K., Fisher, J., Petrie, B., Leggett, W., 2010. Decline in top predator body size and changing climate alter trophic structure in an oceanic ecosystem. *Proceedings of the Royal Society. Series B. Biological Sciences*, 277, 1353-1360.
- Shannon, L., Coll, M., Bundy, A., Gascuel, D., Heymans, J. *et al.*, 2014. Trophic level-based indicators to track fishing impacts across marine ecosystems. *Marine Ecology Progress Series*, 512, 115-140.
- Sinopoli, M., Fanelli, E., D'Anna, G., Badalamenti, F., Pipitone, C., 2012. Assessing the effects of a trawling ban on diet and trophic level of hake, *Merluccius merluccius*, in the southern Tyrrhenian Sea. *Scientia Marina*, 76 (4), 677-690.
- Stergiou, K., Karpouzi, V., 2002. Feeding habits and trophic levels of Mediterranean fish. *Reviews in Fish Biology and Fisheries*, 11, 217-254.
- Stergiou, K., Moutopoulos, D., Krassas, G., 2004. Body size overlap in industrial and artisanal fisheries for five commercial fish species in the Mediterranean Sea. *Scientia Marina*, 68 (1), 179-188.
- Stoner, A., Zimmerman, J., 1988. Food pathways associated with penaeid shrimps in a mangrove-fringed estuary. *Fisheries Bulletin*, 86, 543-551.
- Stobart, B., Warwick, R., González, C., Mallol, S., Díaz, D. *et al.*, 2009. Long-term and spill-over effects of a marine protected area on an exploited fish community. *Marine Ecology Progress Series*, 384, 47-60.
- Tunesi, L., Molinari, A., Salvati, E., 2006. Fish assemblage of the marine protected area of Cinque Terre (NW Mediterranean Sea): First characterization and assessment by visual census. *Chemistry and Ecology*, 22, (Supplement 1), S245-S253.
- Umezawa, Y., Miyajima, T., Yamamuro, M., Kayanne, H., Koike, I., 2002. Fine-scale mapping of land-derived nitrogen in coral reefs by $\delta^{15}\text{N}$ in macroalgae. *Limnology and Oceanography*, 47, 1405-1416.
- Vanderklift, M., Ponsard, S., 2003. Sources of variation in consumer-diet $\delta^{15}\text{N}$ enrichment: a meta-analysis. *Oecologia*, 136 (2), 169-182.
- Vander Zanden, M., Cabana, G., Rasmussen, J., 1997. Comparing trophic position of freshwater fish calculated using stable nitrogen isotope ratios ($\delta^{15}\text{N}$) and literature dietary data. *Canadian Journal of Fisheries and Aquatic Sciences*, 54, 1142-1158.
- Verlaque, M., 1985. Note préliminaire sur le comportement alimentaire de *Sarpa salpa* (Sparidae) en Méditerranée. Rapports. *Commission Internationale Mer Méditerranée*, 29, 193-196.
- Vizzini, S., Sara, G., Michener, R., Mazzola, A., 2002. The role and contribution of the seagrass *Posidonia oceanica* (L.). Delile organic matter for secondary consumers as revealed by carbon and nitrogen stable isotope analysis. *Acta Oecologia*, 23, 277-285.
- Vizzini, S., Mazzola, A., 2004. Stable isotope evidence for the environmental impact of a land-based fish farm in the western Mediterranean. *Marine Pollution Bulletin*, 49, 61-70.
- Vizzini, S., Mazzola, A., 2006. The effects of anthropogenic organic matter inputs on stable carbon and nitrogen isotopes in organisms from different trophic levels in a southern Mediterranean coastal area. *Science of the Total Environment*, 368, 723-731.
- Vizzini, S., Mazzola, A., 2008. The fate of organic matter sources in coastal environments: a comparison of three Mediterranean lagoons. *Hydrobiology*, 611, 67-79.
- Vizzini, S., Mazzola, A., 2009. Stable isotopes and trophic positions of littoral fishes from a Mediterranean marine protected area. *Environmental Biology of Fishes*, 84 (1), 13-25.
- Watson, R., Revenga, C., Kura, Y., 2006. Fishing gear associated with global marine catches I. Database development. *Fisheries Research*, 79, 97-102.
- Worm, B., Myers, R., 2003. Meta-analysis of cod-shrimp interactions reveals top-down control in oceanic food webs. *Ecology (USA)*, 84, 162-173.
- Wyatt, A., Waite, A., Humphries, S., 2010. Variability in isotope discrimination factors in coral reef fishes: implications for diet and food web reconstruction. *PLoS ONE*, 5 (10), e13682.