

## Mediterranean Marine Science

Vol 17, No 3 (2016)



### Soft Bottom Molluscan Assemblages of the Bathyal Zone of the Sea of Marmara

A. DOĞAN, B. ÖZTÜRK, B. BİTLİS-BAKIR, N. TÜRKCÜ

doi: [10.12681/mms.1748](https://doi.org/10.12681/mms.1748)

#### To cite this article:

DOĞAN, A., ÖZTÜRK, B., BİTLİS-BAKIR, B., & TÜRKCÜ, N. (2016). Soft Bottom Molluscan Assemblages of the Bathyal Zone of the Sea of Marmara. *Mediterranean Marine Science*, 17(3), 678–691.  
<https://doi.org/10.12681/mms.1748>

## Soft Bottom Molluscan Assemblages of the Bathyal Zone of the Sea of Marmara

A. DOĞAN<sup>1,3</sup>, B. ÖZTÜRK<sup>1</sup>, B. BİTLİS-BAKIR<sup>2</sup> and N. TÜRKÇÜ<sup>1</sup>

<sup>1</sup> Ege University, Faculty of Fisheries, Department of Hydrobiology, 35100, Bornova, İzmir, Turkey

<sup>2</sup> Dokuz Eylül University, Institute of Marine Sciences and Technology, Inciraltı, 35340, İzmir, Turkey

Corresponding author: [alper.dogan@ege.edu.tr](mailto:alper.dogan@ege.edu.tr)

Handling Editor: Argyro Zenetos

Received: 19 April 2016; Accepted: 21 June 2016; Published on line: 23 September 2016

### Abstract

This study deals with the soft bottom molluscan species collected from the bathyal zone of the Sea of Marmara in 2013. Replicated samples were taken using a Box Core, sampling an area of 0.1 m<sup>2</sup> at 31 stations along two depth transects, 500 and 1000 m. A total of 1229 individuals belonging to 4 classes and 28 species were collected. Two species (*Akritogyra conspicua* and *Liostomia hansgei*) are new records for the marine molluscan fauna of Turkey and four species (*Benthonella tenella*, *Odostomia silesui*, *Syrnola minuta* and *Crenilabium exile*) are new records for the molluscan fauna of the Sea of Marmara. A relatively richer fauna was determined at a depth of 500 m (25 species) compared to 1000 m (17 species). The most dominant species at 500 m and 1000 m depths were *Crenilabium exile* and *Parthenina flexuosa*, respectively. Number of species and number of individuals varied significantly only between depths, while number of individuals changed significantly among basins (ANOVA test). A significant difference in species assemblages between the Tekirdağ and the Central Basins was detected (ANOSIM test). Multivariate analysis showed that depth was the main factor influencing the molluscan assemblages in the area.

**Keywords:** Mollusca, Bathyal zone, New records, Taxonomy, Ecology, Distribution, Sea of Marmara.

### Introduction

The Sea of Marmara, with a surface area of 11,500 km<sup>2</sup> and a maximum depth of 1390 m, is a semi-enclosed basin in the Mediterranean system located between the continents of Europe and Asia. It is connected to the Black Sea, the largest anoxic basin in the world, and the Aegean Sea through the İstanbul and Çanakkale Straits. The Sea of Marmara has two prominently different water masses; the upper layer originating from the Black Sea (from the surface to about 25 m), with brackish water (22–26‰ salinity) and a renewal time of about 5–6 months and the lower layer with more saline water (up to 38.5‰ salinity) originated from the Mediterranean Sea with a renewal time of about 6–7 years (Beşiktepe *et al.*, 1994). Because of the outflow of brackish Black Sea water, there is a strong halocline throughout the Sea of Marmara, leading to low-oxygen conditions below the thin well-mixed surface layer, at 25 m depth (Beşiktepe *et al.*, 1994; Kaminski *et al.*, 2002). On the other hand, the Sea of Marmara is subjected to high volumes of wastewater discharges from land-based sources. The basin receives a total of 1.9 x 10<sup>6</sup> tons of TOC (total organic carbon) and 2.7 x 10<sup>5</sup> tons of TN (total nitrogen) per year from the Black Sea inflow (Albayrak *et al.*, 2006), which is one of the important factors threatening biodiversity.

Studies on the molluscs of the Sea of Marmara date back to the end of 19<sup>th</sup> century. In that period, Colombo

(1885) recorded certain species of different systematic groups such as Coelenterata, Bryozoa, Polychaeta, Crustacea, Mollusca, Echinodermata and Pisces from the infra and the circalittoral zones. Sturany (1895) dealt only with molluscan species from the circalittoral and bathyal (one station only) zones. Ostroumoff (1896) conducted the most detailed study in the 19<sup>th</sup> century; he performed benthic and pelagic samplings at 61 stations and reported more than 700 macrobenthic and planktonic species. Marion (1898) recorded some species of Coelenterata, Crustacea, Mollusca and Echinodermata from gravelly mud bottoms with shell fragments, at a depth of 40–42 m. Later on, further studies on the molluscan fauna of the Sea of Marmara were conducted by different authors. Among them, Demir (1952) focused on the benthic invertebrates along the shores of the Bosphorus and Prince Islands. Tortonese (1959) worked on the soft bottoms of the infra and the circalittoral zones and reported some macrobenthic species of Porifera, Polychaeta, Mollusca, Echinodermata and Tunicata. Oberling (1969–1971) investigated littoral molluscan species between Tekirdağ and Avşa Island. Yüksek (1989) studied the littoral biota of the southern coast of the Sea of Marmara. Balkis (1992) determined the macrobenthic species of the littoral zone of Marmara Island. Albayrak & Balkis (1996a, b) and Albayrak *et al.* (2004) focused on the benthic prosobranch gastropods and the bivalvian species of the hard and the soft bottoms of the infra and the circalittoral zones of the

Bosphorus and the Sea of Marmara. Ritt *et al.* (2010) reported some macrobenthic species from the three micro habitats (bioturbated sediment, reduced sediment and carbonate crust) near a brackish-water cold seep of the bathyal zone on the North Anatolian Fault. Among the above mentioned studies, only Sturany (1895), Ostroumof (1896) and Ritt *et al.* (2010) identified molluscan specimens in some samples (8-13 samples) collected at depths exceeding 500 m.

The aim of this paper was to elucidate the composition and the structure of the molluscan assemblages occurring in the bathyal zone of the Sea of Marmara and their relationship with environmental variables.

## Materials and Methods

### Study Area

The Sea of Marmara is divided into three major sub-basins with bathyal zones, namely, Tekirdağ, Central and Çınarcık. They are about 1200 m deep and separated by saddles as shallow as 400–600 m (McHugh *et al.*, 2008). Among these, the Çınarcık Basin, which is located south of the Prince Islands, has a maximum depth of 1270 m. The Central Basin, which is located between Marmara Ereğlisi and Kapıdağ Peninsula, harbours the deepest point (1268 m) of the Sea of Marmara. The Tekirdağ Basin, which is situated in front of the Ganos Mountains, has a maximum depth of 1133 m (Gazioğlu *et al.*, 2002).

### Data collection

The benthic material was collected from 31 stations in the bathyal zone of the Sea of Marmara (Fig. 1; Table 1) by R/V Yunus-S in June 2013. All material was collected

using a Box Core, sampling an area of 0.1 m<sup>2</sup>. The first 30 cm of the sediment samples were evaluated. Some samples could not be taken at some stations (Table 1) due to the weather conditions and the bottom structure. No sample could be taken at 500 m depth of station 34, as the bottom was covered with rocks. At each station, three replicates were taken for benthic community analysis and an additional sample for granulometric and chemical analysis of the sediment. Samples were sieved with a 0.5 mm mesh on board the R/V Yunus-S and the retained material was placed in jars containing seawater with 4% formaldehyde solution. In addition to the box-core sampler, we also used a bottom-trawl to sample large-sized deep-sea faunal components of the Sea of Marmara. However, no molluscan specimens were gathered by this bottom-trawl hauling (one in each basin). At the laboratory, the material was washed with tap water and specimens were sorted according to taxonomic groups under a stereomicroscope. The sorted specimens were then preserved in 70% ethanol. The molluscan specimens were identified and counted under stereomicroscopes. Bottom-water samples were taken with a CTD bottle at each station during sampling. Temperature, salinity and dissolved oxygen concentrations were determined on board the R/V Yunus-S. Water samples for analysing nitrite, nitrate, ammonia, phosphate phosphorus and silicate were pre-filtered, frozen and immediately transferred to the laboratory.

### Data analysis

Community parameters such as number of species, number of individuals, Shannon–Wiener's diversity index ( $\log_2$  base) ( $H'$ ) and Pielou's evenness index ( $J'$ ) were calculated for the samples. In order to determine




Fig. 1: Map of the investigated area and sampling sites.

**Table 1.** Coordinates, sampling dates, number of sample (N) depth and bottom structure of the sampled stations.

Stations	Latitude	Longitude	Date	Depth (m)	N	Sediment type
Y13_500	40.773056° N	27.395278° E	10.06.2013	500	3	Clay
Y13_1000	40.796944° N	27.443056° E	10.06.2013	1000	2	Clay
Y17_500	40.711667° N	27.616389° E	09.06.2013	500	3	Clay
Y17_1000	40.764167° N	27.549444° E	09.06.2013	1000	3	Clay
Y18_500	40.892500° N	27.551111° E	10.06.2013	500	3	Clay with shell fragments
Y18_1000	40.873333° N	27.563611° E	10.06.2013	1000	2	Clay
Y19_500	40.926389° N	27.709722° E	11.06.2013	500	3	Gravelly clay
Y19_1000	40.883611° N	27.706944° E	11.06.2013	1000	2	Clay
Y20_500	40.918611° N	27.912500° E	13.06.2013	500	3	Clay
Y20_1000	40.881667° N	27.927222° E	12.06.2013	1000	2	Clay
Y21_500	40.703611° N	27.967778° E	15.06.2013	500	3	Clay
Y21_1000	40.756667° N	27.970000° E	15.06.2013	1100	2	Clay
Y23_500	40.716667° N	28.151111° E	15.06.2013	500	3	Clay
Y23_1000	40.775278° N	28.096944° E	15.06.2013	1000	2	Clay
Y24_500	40.927778° N	28.112500° E	13.06.2013	500	3	Clay
Y24_1000	40.881944° N	28.074722° E	13.06.2013	1190	1	Clay
Y29_500	40.733056° N	28.721389° E	17.06.2013	500	3	Clay
Y30_500	40.698611° N	28.871944° E	17.06.2013	500	3	Clay
Y30_1000	40.732222° N	28.880556° E	17.06.2013	1030	3	Clay
Y31_500	40.891667° N	28.458889° E	23.06.2013	500	3	Clay
Y32_500	40.892222° N	28.571111° E	23.06.2013	500	3	Gravelly clayey sand
Y33_500	40.901111° N	28.748611° E	23.06.2013	500	3	Clay
Y33_1000	40.861111° N	28.729167° E	23.06.2013	1000	1	Clay
Y34_1000	40.872222° N	28.836944° E	24.06.2013	1000	1	Clay
Y36_500	40.851667° N	28.967500° E	22.06.2013	500	2	Clay
Y36_1000	40.839722° N	28.958889° E	22.06.2013	1000	1	Gravelly sand with shell fragments
Y37_500	40.783611° N	29.132500° E	21.06.2013	500	3	Gravelly clay
Y37_1000	40.765556° N	29.120556° E	21.06.2013	1000	1	Clay
Y39_500	40.683611° N	29.149722° E	19.06.2013	500	3	Clay with shell fragments
Y39_1000	40.702778° N	29.152500° E	19.06.2013	1215	2	Clay
Y40_500	40.720000° N	29.292500° E	19.06.2013	500	3	Clay

species distribution patterns, the abundance data of all stations were analysed using cluster and multidimensional scaling (MDS) techniques, based on the Bray–Curtis similarity index (group average technique). The groups of stations (A, B and C) on the dendrogram were highlighted in grey. The relationship between environmental variables and patterns of similarity of the mollusc assemblages was assessed by the MDS plot with superimposed vectors whose correlation is higher than 0.3, using the PRIMER package (Clarke & Warwick, 2001). Prior to the cluster and multidimensional scaling analyses, the raw data (number of individuals) were transformed using they  $y_{ji} = \log(x_{ji} + 1)$  equation. Pearson's correlation analysis was used to assess relationships between environmental variables and community parameters.

The differences in the number of species, number of individuals, diversity and evenness index values were tested using ANOVA. Pairwise analysis of similarities (ANOSIM) was carried out to test significant differences regarding depths, basins and sediment types.

Nutrients and chlorophyll-a were analysed using a spectrophotometer (Parsons *et al.*, 1984). Total organic carbon concentration (TOC) in each sediment sample was estimated according to the modified Walkley Black titration method (Gaudette *et al.*, 1974). Granulometric analyses were carried out according to Erguvanli (1995). Three particle size fractions were determined including sand (2 mm-0.063 mm), silt (0.063 mm-0.002 mm) and clay (<0.002 mm). The physical & chemical properties of the stations will be presented at a later date by Aksu *et al.* (in preparation).

The specimens identified in this study were deposited at the Museum of the Faculty of Fisheries, Ege University (EFSM), İzmir, Turkey.

## Results

Examination of materials obtained from 31 stations revealed a total of 28 molluscan species and 1229 indi-



viduals belonging to 4 classes (Caudofoveata, Gastropoda, Bivalvia and Scaphopoda). The class Caudofoveata was represented by one species and one individual, Gastropoda by 22 species and 1162 individuals, Bivalvia by 4 species and 54 individuals and Scaphopoda by 1 species and 12 individuals (Table 2).

Among the classes, Gastropoda accounted for 91% and 96% of the total number of specimens collected at depths of 500 and 1000 m, respectively (Table 2). The dominant species at 500 m depth were *Crenilabium exile* (26.37% of total individuals), *Parthenina flexuosa* (15.02%), *Alvania cimicoides* (9.16%) and *Megastomia conoidea* (8.42%), all comprising 59% of total specimens (Table 2). *Parthenina flexuosa* was the most dominant (60.46%) species at 1000 m depth, followed by *C. exile* (22.28%), *B. tenella* (6.17%) and *Yoldiella philippiana* (3.66%).


The most frequent species at 500 m depth were *A. cimicoides* and *C. exile* (each present in 32% of the samples), *Megastomia conoidea* (18%), *P. flexuosa* (16%), *S. minuta* (16%) and *Cylichna cylindracea* (16%) (Table 2). *Parthenina flexuosa* and *C. exile* were the most frequent species (72% each) at 1000 m and deeper.

The mean number of species and number of individuals, diversity and evenness values at all stations are presented in Figure 2. The highest number of species was encountered at 1000 m, at station Y37 (7 species) and Y20 (6 species), and at 500 m depth, at station Y40 (6 species). Among 31 sampling stations, four stations (Y17\_500, Y21\_500, Y31\_500 and Y32\_500) did not have any molluscan species. Station Y20\_1000 had the highest molluscan density (1560 ind. m<sup>-2</sup>), followed by Y23\_1000, Y24\_1000 and Y39\_1000 (640 ind.m<sup>-2</sup> in

**Table 2.** List of species and number of individuals (N) recorded at two different sampling depths (500 and 1000 m) in the Sea of Marmara, with their dominancy (D%) and frequency values (F%).

Groups/Species	N (500 m)	N (1000 m)	F% (500m )	F% (1000 m)	D% (500 m)	D% (1000 m)
<b>CAUDOFOVEATA</b>						
<i>Falcidens guttuosus</i> (Kowalewsky, 1901)	1		2.00		0.37	
<b>GASTROPODA</b>						
<i>Putzeysia wiseri</i> (Calcara, 1842)	10	3	10.00	4.00	3.66	0.31
* <i>Akritogyra conspicua</i> (Monterosato, 1880)	5	7	6.00	12.00	1.83	0.73
<i>Bittium submamillatum</i> (de Rayneval & Ponzi, 1854)	3		4.00		1.10	
<i>Alvania cimicoides</i> (Forbes, 1844)	25	24	32.00	24.00	9.16	2.51
<i>Alvania punctura</i> (Montagu, 1803)	3		6.00		1.10	
** <i>Benthonella tenella</i> (Jeffreys, 1869)	6	59	6.00	36.00	2.20	6.17
<i>Pusillina inconspicua</i> (Alder, 1844)	3		2.00		1.10	
<i>Laeviphitus verduini</i> van Aartsen, Bogi & Giusti, 1989		3		8.00		0.31
<i>Ceratia proxima</i> (Forbes & Hanley, 1850)	2		4.00		0.73	
<i>Hyala vitrea</i> (Montagu, 1803)	5	1	6.00	4.00	1.83	0.10
* <i>Liostomia hansgei</i> Warén, 1991		4		12.00		0.42
<i>Mangelia nuperrima</i> (Tiberi, 1855)	2		4.00		0.73	
<i>Megastomia conoidea</i> (Brocchi, 1814)	23	11	18.00	36.00	8.42	1.15
** <i>Odostomia silesui</i> Nofroni, 1988	2		2.00		0.73	
<i>Odostomia unidentata</i> (Montagu, 1803)	1		2.00		0.37	
<i>Parthenina flexuosa</i> (Monterosato, 1874)	41	578	16.00	72.00	15.02	60.46
<i>Parthenina interstincta</i> (Adams, J., 1797)	1		2.00		0.37	
** <i>Syrnola minuta</i> H. Adams, 1869	20	5	16.00	12.00	7.33	0.52
<i>Turbonilla micans</i> (Monterosato, 1875)		2		8.00		0.21
** <i>Crenilabium exile</i> (Jeffreys, 1870)	72	213	32.00	72.00	26.37	22.28
<i>Roxania utriculus</i> (Brocchi, 1814)	16	3	12.00	4.00	5.86	0.31
<i>Cylichna cylindracea</i> (Pennant, 1777)	9		16.00		3.30	
<b>BIVALVIA</b>						
<i>Yoldiella philippiana</i> (Nyst, 1845)	5	35	2.00	44.00	1.83	3.66
<i>Lucinoma borealis</i> (Linnaeus, 1767)	2		2.00		0.73	
<i>Lucinoma kazani</i> Salas & Woodside, 2002	1		2.00		0.37	
<i>Myrtea amorpha</i> (Sturany, 1896)	8	3	2.00	4.00	2.93	0.31
<b>SCAPHOPODA</b>						
<i>Entalina tetragona</i> (Brocchi, 1814)	8	4	12.00	8.00	2.93	0.42

\*Species new records for the marine molluscan fauna of Turkey, \*\*Species new records for the Sea of Marmara.



**Fig. 2:** Mean number of species, faunal densities (number of individuals per m<sup>-2</sup>), diversity index and evenness index at each station, with  $\pm$  standard errors.

each). The highest diversity index value ( $H'=2.27$ ) was calculated at station Y37\_1000, followed by Y40\_500 ( $H'=2.18$ ) and Y36\_500 ( $H'=2.06$ ). The highest evenness index value ( $J'=0.95$ ) was found at station Y36\_500, followed by Y40\_500 ( $J'=0.89$ ) and Y33\_1000 ( $J'=0.87$ ).

The maximum values of community parameters in each sample, such as number of species, number of individuals, diversity index and evenness index at all stations are presented in Figure 3. The highest number of species was found at stations Y39\_1000 (11 species), Y39\_500 (9 species), Y24\_500 (8 species) and Y40\_500 (8 species). The highest molluscan density (430 ind.m<sup>-2</sup>) at 500 m depth was determined at stations Y39\_500 and Y40\_500 (290 ind. m<sup>-2</sup>). Stations Y20\_1000 (3020 ind.m<sup>-2</sup>) and Y39\_1000 (1240 ind.m<sup>-2</sup>) had the highest molluscan densities at 1000 m depth. The highest diversity index values were calculated at stations Y40\_500 ( $H'=2.8$ ), Y19\_500 ( $H'=2.62$ ) and Y37\_1000 ( $H'=2.27$ ). As stations Y13\_500, Y18\_500, Y20\_500, Y36\_500 and Y37\_500 had few species and specimens, the evenness index attained its maxima ( $J'=1$ ) at these stations. The highest evenness index values at 1000 m were found at stations Y20\_1000 ( $J'=0.94$ ), Y17\_1000 ( $J'=0.88$ ) and Y33\_1000 ( $J'=0.87$ ).

Among the three basins and two different depth ranges studied, the maximum number of molluscan species was determined at depths of 1000 m and 500 m in Çınarcık Basin (Fig. 4). The highest median value of the number of species and the maximum number of individuals were found at 1000 m and more depths in the Central Basin. Tekirdağ Basin was represented by the highest diversity index values, while Çınarcık Basin and the Central Basin, at 1000 m, had the highest median values. The five hundred meter depths of all basins showed the highest evenness index values (Fig. 4).

Pearson's correlation analysis between the mean scores of community parameters and environmental variables indicated that the number of individuals and the diversity index were significantly and positively correlated with depth and total organic carbon in sediment, respectively (Table 3). The other factor significantly and negatively correlated with the number of individuals in the area was salinity (Table 3).

Based on Bray-Curtis similarity values higher than 55%, three mollusc assemblages were determined in the area (Fig. 5). The stress value for the two-dimensional MDS plot was 0.16, showing a proper group separation. Stations Y18\_500, Y20\_500, Y30\_500 and Y33\_500 formed group A (average similarity: 56.28), all deeper stations (1000 m and deeper) formed group B (average similarity: 57.9) and stations Y24\_500, Y29\_500, Y39\_500 and Y40\_500 constituted group C (average similarity: 55.45). The deepest part of the bathyal zone seemed to be homogenous in terms of molluscan assemblages, aggregation in a large group called group B. However, the molluscan assemblages were heterogeneous at 500 m depth. There were two large groupings at 500 m depth, and 4 stations at 500 m depth were placed in distant locations on the MDS plot, which joined to other groups with low similarity index values (<40%). Vectors superimposed on the MDS plot indicated the relationships between environmental variables and patterns of similarity of the mollusc assemblages (Figure 5). According to the MDS plot and the correlation values between the environmental variables and MDS axes, the depth was found to be the main factor influencing the molluscan assemblages in the study area (Fig. 5). However, the granulometric features of sediment seemed to have impacted on the distribution of molluscan species at 500 m depth. As stations Y19\_500 and Y37\_500 had a different habitat structure (gravelly clay bottom), the molluscan assemblages found at these stations were very different from those with clayey habitats, resulting in the placement of these stations at distant locations on the MDS graph.

According to the SIMPER analysis, *Crenilabium exile*, *Parthenina flexuosa* and *Syrnola minuta* were the species most responsible for the similarity of groups A, B and C, respectively (Table 4). *Alvania cimicoides* was another species significantly contributing to the similarity in groups A and C.

Number of species and number of individuals varied significantly only between depths (ANOVA,  $P<0.05$ ). In addition, number of molluscan individuals changed significantly among basins (ANOVA,  $P<0.05$ ). The ANOSIM pairwise test detected a significant difference

**Table 3.** Pearson's correlation coefficients calculated between community parameters and environmental variables; numbers in bold represent statistically significant correlations ( $P<0.05$ ).


	Number of species	Number of individuals	Diversity index	Evenness index
Depth	-0.13	<b>0.56</b>	0.23	0.19
Temperature	-0.06	-0.22	-0.28	-0.28
Salinity	-0.12	<b>-0.41</b>	-0.04	0.00
pH	0.30	0.31	0.19	0.01
Oxygen	-0.37	0.29	-0.19	-0.01

**Table 4.** Species contributing significantly to the similarity within each assemblage (as shown in Fig. 5) and average similarity percentage.

Associations in Fig. 3	A		B		C	
<b>Average similarity</b>	<b>56.28</b>	<b>AA</b>	<b>57.90</b>	<b>AA</b>	<b>55.45</b>	<b>AA</b>
<i>Alvania cimicoides</i>	39.94	12.5			16.59	35.0
<i>Benthonella tenella</i>			8.12	45.4		
<i>Megastomia conoidea</i>					16.35	45.0
<i>Parthenina flexuosa</i>			<b>43.36</b>	447.7	8.52	80.0
<i>Syrnola minuta</i>					<b>17.75</b>	42.5
<i>Crenilabium exile</i>	<b>42.68</b>	22.5	35.40	173.1	11.89	115.0
<i>Roxania utriculus</i>					12.11	35.0
<i>Cylichna cylindracea</i>	17.38	10.0				
<i>Yoldiella philippiana</i>			4.01	23.1		
<i>Entalina tetragona</i>					9.81	17.5


Bold numbers indicate the highest score of species contribution in each assemblage.

AA: Average abundance (individuals  $m^{-2}$ ).



**Fig. 3:** Maximum number of species, faunal densities (number of individuals per  $m^{-2}$ ), diversity index and evenness index values determined at each station.





**Fig. 4:** Median values, 25-75 percentiles and minimum-maximum values of the number of species, faunal densities (number of individuals per m<sup>-2</sup>), diversity index and evenness index determined for each basin (n: number of samples; T: Tekirdağ Basin; C: Central Basin; ÇI: Çınarcık Basin).

in species assemblages between the Tekirdağ and the Central Basins (ANOSIM,  $P < 0.05$ ).

## Discussion

Analysis of the benthic samples taken from the bathyal zone of the Sea of Marmara showed that, compared to the western Mediterranean (Gofas *et al.*, 2014; Rueda *et al.*, 2015), the investigated area has a relatively poor molluscan fauna (a total of 28 species), like the other eastern parts of the Mediterranean (Bogi & Galil, 2004; Koutsoubas *et al.*, 2000; Mastroianni *et al.*, 2010). Among the species identified, *Akritogyra conspicua* and *Liostomia hansgei* (Fig. 6) are new records for the marine molluscan fauna of Turkey, while *Benthonella tenella*, *Odostomia silesui*, *Syrnola minuta* and *Crenilabium exile* (Fig. 6) are new ones for the molluscan fauna of the Sea of Marmara. *Akritogyra conspicua* is a deep-water species recorded from depths between 100-2400 m, and was previously reported as a rare species from the central and

western Mediterranean Sea (Warén, 1992). In this study, nine specimens of *Akritogyra conspicua* were found at depths between 500 and 1000 m. *Liostomia hansgei*, which was originally described from the Swedish west coast (Koster area) (Warén, 1991), is also a rarely distributed species in the Mediterranean Sea, known only from the Spanish coast (Peñas *et al.*, 1996). In our study, five specimens of this species were found at 1000 m depth.


Of the species that are new records for the Sea of Marmara, *Benthonella tenella* and *C. exile* were previously reported from the Levantine coasts of Turkey (Öztürk *et al.*, 2012; Öztürk, 2014), and *O. silesui* and *S. minuta* from the Aegean coast of Turkey (Öztürk *et al.*, 2013; Öztürk & Bitlis-Bakır, 2013).

In a previous study conducted by Ostroumoff (1896), a total of 49 mollusc species was reported in the bathyal zone of the region, of which 40 species were represented by their shells only. Out of the nine living molluscan species [*Aporrhais serresianus* (Michaud, 1828); *Odostomia unidentata* (Montagu, 1803); *Saccula commuta-*


*ta* (Philippi, 1844); *Yoldiella philippiana* (Nyst, 1845); *Delectopecten vitreus* (Gmelin, 1791); *Flexopecten glaber* (Linnaeus, 1758); *Kurtiella bidentata* (Montagu, 1803); *Abra longicallus* (Scacchi, 1835); *Kelliella miliaris* (Philippi, 1844)] reported by Ostroumoff (1896), only *O. unidentata* and *Y. philippiana* were encountered during our study. Besides, Ostroumoff (1896) reported live specimens of *K. miliaris* in the bathyal zones of the

Sea of Marmara, but no live individuals of this species were encountered during our study. However, many dead shells of this species were found in the area, at depths of 500-1000 m.

Another study dealing with the bathyal zone communities of the Sea of Marmara was carried out by Ritt *et al.* (2010) who reported 17 mollusc species from different microhabitats such as bioturbated sediment microhabi-



**Fig. 5:** MDS plot and dendrogram of the sampled stations, based on mollusc abundance data (sum of three replicates), and correlation of environmental variables with MDS axes, represented by superimposed vectors. Similarity among the stations was determined using the Bray-Curtis similarity index and was then superimposed on the MDS plot. The groups of stations (A, B and C) are highlighted in grey on the dendrogram.




**Fig. 6:** Species new to the marine molluscan fauna of Turkey (*Akritogyra conspicua* and *Liostomia hansgei*) and to the Sea of Marmara (*Benthonella tenella*, *Odostomia silesui*, *Syrrola minuta* and *Crenilabium exile*).

tat, reduced sediment microhabitat and carbonate crust microhabitat of cold seeps located in the north-east Central Basin. Although a limited number of samples were evaluated in that study, the number of molluscan species was high (23 species) as the samples were taken from the specific microhabitats of cold seeps, generally preferred by symbiont-bearing species (Salas & Woodside, 2002; Olu-Le Roy *et al.*, 2004; Brissac *et al.*, 2011).

*Lucinoma kazani* and *Myrtea amorpha*, which are symbiont-bearing lucinid species, were newly reported from the Sea of Marmara by Ritt *et al.* (2010). Among these two species, *L. kazani* was only found at one station (Y19\_500 m) with its congeneric *L. borealis* (Fig. 7), while *M. amorpha* was encountered at only two stations

(Y19\_500 m and Y36\_1000 m). *Falcidens guttuerosus*, *Putzeysia wiseri* and *Laeviphitus verduini*, which were identified in this study, were the other molluscan species reported by Ritt *et al.* (2010) in the cold seep microhabitats in the bathyal zones of the Sea of Marmara. According to these findings, some of our stations (i.e. Y19\_500 m and Y36\_1000 m) might have been close to the cold seeps favouring symbiont-bearing species.

During research carried out between 1994 and 1999 off the northern coast of Israel, at depths between 734 and 1558 m, 23 live molluscan species were reported (Bogi & Galil, 2004). In the northern part of Crete, Koutsoubas *et al.* (2000) recorded 31, 24, 22 and 10 molluscan species at depths of 500, 700, 1000 and 1570 m, respectively.



**Fig. 7:** External view of the left valve (A), internal view of the right valve (B), hinge line of the right (C) and the left (D) valves of *Lucinoma kazani*. External view of the left valve (E), internal view of the right valve (F), hinge line of the right (G) and the left (H) valves of *L. borealis*.

Mastrototaro *et al.* (2010) identified 35 molluscan species from the white coral bank off Cape Santa Maria di Leuca (Ionian Sea) at depths between 310 and 803 m. According to the results of all these studies, the molluscan assemblages in the bathyal zone of the Sea of Marmara were similar to those identified in the other parts of the eastern Mediterranean.

*Lucinoma borealis* is known to be distributed in shallower areas than those determined by this study (Salas & Woodside, 2002). This could be due to the distinctive ecological characteristics of the Sea of Marmara.

Gastropoda was the dominant group in terms of number of species and number of individuals. Gastropod dominance is due to the abundance of *Parthenina flexuosa*, which is a small pyramidellid mollusc known to be an ectoparasite, feeding on the body fluids of some invertebrates, such as polychaetes, molluscs and echinoderms (Robertson & Mau-Lastovicka, 1979). The high abundance of the echinoid echinoderm *Brissopsis lyrifera* (Forbes, 1841), with three specimens per square meter in the bathyal bottoms of the Sea of Marmara (unpublished data), could have stimulated the dense settlement of this ectoparasitic gastropod.

According to the MDS plot and the correlation values between the environmental variables and MDS axes, the depth was found to be the main factor affecting mol-

luscan assemblages in the study area. The granulometric features of the sediment also seem to have influence the distribution of molluscan species in the area. These factors are known to have strong impacts on the distribution patterns of zoobenthic organisms in the benthic ecosystem (Tselepidis *et al.*, 2000; Labrune *et al.*, 2007; Çınar *et al.*, 2012). Pancucci-Papadopoulou *et al.* (1999) note that sediment structure, depth and hydrological conditions could play a synergistic role in the discrimination of macrobenthic communities. According to Gray (2002), sediment structure and food availability could be considered as the main factors determining faunal biodiversity in the deeper zones of the seas.

As a result of this study, the number of molluscan species distributed in the 500 m and deeper zones of the Sea of Marmara increased to 42 (Table 5). Besides, some species on the list, such as *Myrtea spinifera* (Montagu, 1803) distributed in the shallower depths and quite similar to *Myrtea amorpha* (Sturany, 1896) might have been identified incorrectly.

This study presents the recent status of molluscan communities in the bathyal zone of the Sea of Marmara. In order to gain a better understanding of the deep-sea fauna of the Sea of Marmara, more detailed studies, using bottom-trawling and under-water equipment via ROV, are required.



**Table 5.** List of species distributed in the 500 m and deeper zones of the Sea of Marmara and references.

Species	Reference
<i>Falcidens gutturosus</i> (Kowalewsky, 1901)	Ritt <i>et al.</i> , 2010
<i>Putzeysia wiseri</i> (Calcara, 1842)	Ritt <i>et al.</i> , 2010
<i>Akritogyra conspicua</i> (Monterosato, 1880)	This study
<i>Bittium submamillatum</i> (de Rayneval & Ponzi, 1854)	This study
<i>Alvania cimicoides</i> (Forbes, 1844)	This study
<i>Benthonella tenella</i> (Jeffreys, 1869)	This study
<i>Psyllina inconspicua</i> (Alder, 1844)	This study
<i>Laeviphitus verduini</i> van Aartsen, Bogi & Giusti, 1989	Ritt <i>et al.</i> , 2010
<i>Ceratia proxima</i> (Forbes & Hanley, 1850)	This study
<i>Hyalia vitrea</i> (Montagu, 1803)	This study
<i>Liostomia hansgei</i> Warén, 1991	This study
<i>Mangelia nuperrima</i> (Tiberi, 1855)	This study
<i>Megastomia conoidea</i> (Brocchi, 1814)	This study
<i>Odostomia silesui</i> Nofroni, 1988	This study
<i>Odostomia unidentata</i> (Montagu, 1803)	This study
<i>Parthenina flexuosa</i> (Monterosato, 1874)	This study
<i>Parthenina interstincta</i> (Adams, J., 1797)	This study
<i>Syrnola minuta</i> H. Adams, 1869	This study
<i>Aporrhais serresianus</i> (Michaud, 1828)	Ostroumoff, 1896
<i>Euspira fusca</i> (Blainville, 1825)	Ostroumoff, 1896
<i>Pterotrachea coronata</i> Forsskål in Niebuhr, 1775	Ostroumoff, 1896
<i>Turbonilla micans</i> (Monterosato, 1875)	This study
<i>Crenilabium exile</i> (Jeffreys, 1870)	This study
<i>Roxania utriculus</i> (Brocchi, 1814)	This study
<i>Cylichna cylindracea</i> (Pennant, 1777)	This study
<i>Yoldiella philippiana</i> (Nyst, 1845)	Ostroumoff, 1896
<i>Yoldiella striolata</i> (Brugnone, 1876)	Ritt <i>et al.</i> , 2010
<i>Delectopecten vitreus</i> (Gmelin, 1791)	Ostroumoff, 1896
<i>Flexopecten glaber</i> (Linnaeus, 1758)	Ostroumoff, 1896
<i>Lucinoma borealis</i> (Linnaeus, 1767)	Ostroumoff, 1896
<i>Lucinoma kazani</i> Salas & Woodside, 2002	Ritt <i>et al.</i> , 2010
<i>Myrtea amorpha</i> (Sturany, 1896)	Ritt <i>et al.</i> , 2010
<i>Myrtea spinifera</i> (Montagu, 1803)	Ostroumoff, 1896
<i>Kurtiella bidentata</i> (Montagu, 1803)	Ostroumoff, 1896
<i>Abra longicallus</i> (Scacchi, 1835)	Ostroumoff, 1896
<i>Kelliella miliaris</i> (Philippi, 1844)	Ostroumoff, 1896
<i>Isorropodon perplexum</i> Sturany, 1896	Ritt <i>et al.</i> , 2010
<i>Timoclea ovata</i> (Pennant, 1777)	Ostroumoff, 1896
<i>Corbula gibba</i> (Olivi, 1792)	Ostroumoff, 1896
<i>Xylophaga dorsalis</i> (Turton, 1819)	Ostroumoff, 1896
<i>Cuspidaria cuspidata</i> (Olivi, 1792)	Sturany, 1895
<i>Entalina tetragona</i> (Brocchi, 1814)	This study

## Acknowledgements

We are much indebted to the crew of R/V Yunus-S for their help in collecting the benthic material. We also thank Melih Ertan Çınar for helping with the statistical analysis. This work has been supported financially by Scientific Research Projects of TUBITAK (Project number: 111Y268).

## References

- Albayrak, S., Balkis, N., 1996a. Benthic prosobranch gastropods of the Bosphorus. *IUFS Journal of Biology*, 59, 17-26.
- Albayrak, S., Balkis, N., 1996b. Bivalve fauna of the Bosphorus. *IUFS Journal of Biology*, 59, 1-15.
- Albayrak, S., Balkis, H., Balkis, N., 2004. Bivalvia (Mollusca) fauna of the Sea of Marmara. *Acta Adriatica*, 45, 9-26.

- Albayrak, S., Balkis, H., Zenetos, A., Kurun, A., Kubanc, C., 2006. Ecological quality status of coastal benthic ecosystems in the Sea of Marmara. *Marine Pollution Bulletin*, 52, 790-799.
- Balkis, H., 1992. A preliminary study on the macrobenthos of littoral of the Island of Marmara. *Bulletin of Institute of Marine Sciences and Geography, İstanbul University*, 9, 309-327.
- Beşiktepe, S.T., Sur, H.I., Özsoy, E., Latif, M.A., Oğuz, T. *et al.*, 1994. The circulation and hydrography of the Marmara Sea. *Progress in Oceanography*, 34, 285-334.
- Bogi, C., Galil, B.S., 2004. The bathy-benthic and pelagic molluscan fauna off the Levantine coast, eastern Mediterranean. *Bollettino Malacologico*, 39 (5-8), 79-90.
- Brissac, T., Rodrigues, C.F., Gros, O., Duperron, S., 2011. Characterization of bacterial symbioses in *Myrtea* sp. (Bivalvia: Lucinidae) and *Thyasira* sp. (Bivalvia: Thyasiridae) from a cold seep in the Eastern Mediterranean. *Marine Ecology*, 32, 198-210.
- Çinar, M.E., Katagan, T., Öztürk, B., Dagli, E., Açık, S. *et al.*, 2012. Spatio-temporal distributions of zoobenthos in Mersin Bay (Levantine Sea, eastern Mediterranean) and the importance of alien species in benthic communities. *Marine Biology Research*, 8 (10), 954-968.
- Clarke, K.R., Warwick, R.M., 2001. Change in marine communities: an approach to statistical analysis and interpretation, 2<sup>nd</sup> edition. PRIMER-E, Plymouth.
- Colombo, A., 1885. Raccolte Zoologiche Eseguite dal R. Piro-scafo Washington Nella Campagna Abissale Talassografica dell'Anno 1885. *Rivista Marittima*, 18, 22-53.
- Demir, M., 1952. *Benthic Invertebrates of Istanbul Strait and the Prince Islands*. İstanbul University, Faculty of Science, Hydrobiology Research Institute Publication, 3, 1-615.
- Erguvanli, K. 1995. *Jeology of Engineering*. Seç Publication and Distribution, 4<sup>th</sup> Press, İstanbul, 590 pp.
- Gaudette, H.E., Flight, W.R., Toner, L., Folger, W., 1974. An inexpensive titration method for the determination of organic carbon in recent sediments. *Journal of Sedimentary Petrology*, 44, 249-253.
- Gazioğlu, C., Gökaşan, E., Algan, O., Yücel, Z., Tok, B., *et al.*, 2002. Morphologic features of the Marmara Sea from multibeam data. *Marine Geology*, 190, 397-420.
- Gofas, S., Salas, C., Rueda, J.L., Canoura, J., Farias, C. *et al.*, 2014. Mollusca from a species-rich deep-water community in the Alboran Sea. *Scientia Marina*, 78, 537-553.
- Gray, J.S., 2002. Species richness of marine soft sediments. *Marine Ecology Progress Series*, 244, 285-297.
- Kaminski, M.A., Aksu, A., Box, M., Hiscott, R.N., Filipescu, S. *et al.*, 2002. Late Glacial to Holocene benthic foraminifera in the Marmara Sea: implications for Black Sea-Mediterranean Sea connections following the last deglaciation. *Marine Geology*, 190, 165-202.
- Koutsoubas, D., Tselepidis, A., Eleftheriou, A., 2000. Deep Sea Molluscan Fauna of the Cretan Sea (Eastern Mediterranean): Faunal, Ecological and Zoogeographical Remarks. *Senckenbergiana Maritima*, 30 (3/6), 85-98.
- Labrune, C., Gre'mare, A., Guizien, K., Amouroux, J.M., 2007. Long-term comparison of soft bottom macrobenthos in the Bay of Banyuls-sur-Mer (north-western Mediterranean Sea): A reappraisal. *Journal of Sea Research*, 58, 125-143.
- Marion, A.F., 1898. Notes sur la faune des Dardanelles et du Bosphore. *Annual Museum History Natural Marseille*, 1, 163-182.
- Mastrototaro, F., D'Onghia, G., Corriero, G., Matarrese, A., Maiorano, P. *et al.*, 2010. Biodiversity of the white coral bank off Cape Santa Maria di Leuca (Mediterranean Sea): An update. *Deep-Sea Research II*, 57, 412-430.
- McHugh, C.M.G., Grung, D., Giosan, L., Ryan, W.F.B., Mart, Y. *et al.*, 2008. The last reconnection of the Marmara Sea (Turkey) to the 237 World Ocean: A paleoceanographic and paleoclimatic perspective. *Marine Geology*, 255, 64-82.
- Oberling, J.J. 1969-1971. On the littoral Mollusca of Sea of Marmara. *Jahrbuch des Naturhistorischen Museums*, 4: 183-218.
- Olu-Le Roy, K., Sibuet, M., Fiala-Médioni, A., Gofas, S., Salas, C. *et al.*, 2004. Cold seep communities in the deep eastern Mediterranean Sea: composition, symbiosis and spatial distribution on mud volcanoes. *Deep-Sea Research I*, 51, 1915-1936.
- Ostroumoff, A., 1896. Otchet o dragirovka i planktonniyh ulovaht ekspeditsia "Selyanika". *Bulletin de l'Academie Imperiale des Sciences de St. Petersburg*, 5, 33-92.
- Öztürk, B., Topçu, E., Topaloğlu, B., 2012. The submarine canyons of the Rhodes basin and the Mediterranean coast of Turkey. p. 65-71. In: *Mediterranean Submarine Canyons: Ecology and Governance*, Würtz, M., (Ed). Gland, Switzerland and Malaga, Spain, IUCN.
- Öztürk, B., Bitlis-Bakır, B., 2013. Heterostropha species of the Turkish coasts: *Anisocycla*, *Eulimella*, *Puposyrnola*, *Syrnola*, and *Turbonilla* (Gastropoda, Heterobranchia). *Turkish Journal of Fisheries and Aquatic Sciences*, 13, 423-440.
- Öztürk, B., Bitlis-Bakır, B., Micali, P., 2013. Heterostropha species of the Turkish coasts: Odostomiinae Pelseneer, 1928 (Gastropoda, Heterobranchia, Pyramidellidae). *Turkish Journal of Fisheries and Aquatic Sciences*, 13, 139-157.
- Öztürk, B., 2014. *Shelled Molluscs of the Turkish coasts: informal group «Lower Heterobranchia»*. Ege University Press, Publication of the Faculty of Fisheries, No: 81, İzmir, 131 pp.
- Pancucci-Papadopoulou, A., Simbora, N., Zenetos, A., Thessalou-Legaki, M., Nicolaidou, A., 1999. Benthic invertebrate communities of NW Rodos (Rhodes) island (SE Aegean Sea) as related to hydrological regime and geographical location. *Israel Journal of Zoology*, 45, 371-393.
- Parsons, T.R., Matia, Y., Lalli, C.M., 1984. *A manual of chemical and biological methods for seawater analysis*. Pergamon Press, New York, 173 pp.
- Peñas, A., Templado, J., Martinez, J.L., 1996. Contribution to the knowledge of the Pyramidelloidea (Gastropoda, Heterostropha) in the Spanish Mediterranean coasts. *Iberus*, 14, 1-82.
- Ritt, B., Jozée, S., Caprais, J.C., Noël, P., Gauthier, O. *et al.*, 2010. First insights into the structure and environmental setting of cold-seep communities in the Marmara Sea. *Deep-Sea Research I*, 57, 1120-1136.
- Robertson, R., Mau-Lastovicka, T., 1979. The ectoparasitism of Boonea and Fargoa (Gastropoda: Pyramidellidae). *Biological Bulletin*, 157, 320-333.
- Rueda, J.L., Farias, C., Gallardo-Núñez, M., Gallardo-Roldán, H., Mateo, A. *et al.*, 2015. Molluscan assemblages from circalittoral and bathyal soft bottoms of the northern Albo-

- ran Sea. p.509-512. In: *Resúmenessobre el VIII Simposio MIA15*, 21-23 Septiembre 2015. Málaga.
- Salas, C., Woodside, J., 2002. *Lucinoma kazani* n. sp. (Mollusca: Bivalvia): evidence of a living benthic community associated with a cold seep in the Eastern Mediterranean Sea. *Deep Sea Research Part I: Oceanographic Research Papers*, 49 (6), 991-1005.
- Sturany, R., 1895. Bestimmungsliste der von Herrn Dr Konrad Natterer auf S. M. Schiff "Taurus" im Marmara-Meere Geddrechten mollusken. *Denkschriften der mathem-naturw*, 62, 119-121.
- Tortonese, E., 1959. Osservazioni sul Benthos del mar di Marmara e del Bosforo. *Rivista di Scienze Naturali*, 50, 18-26.
- Tselepides, A., Papadopoulou, N., Podaras, D., Plaiti, W., Koutsoubas, D., 2000. Macrobenthic community structure over the continental margin of Crete (South Aegean Sea, NE Mediterranean). *Progress in Oceanography*, 46, 401-428.
- Warén, A., 1991. New and little known Mollusca from Iceland and Scandinavia. *Sarsia*, 76, 53-124.
- Warén, A., 1992. New and little known "Skeneimorph" gastropods from the Mediterranean Sea and the adjacent Atlantic Ocean. *Bolletino Malacologico*, 27, 1-149.
- Yüksek, A., 1989. The investigations of littoral biota of southern coast of Marmara. *Bulletin of Institute of Marine Sciences and Geography, İstanbul University*, 6, 203-216.