

Mediterranean Marine Science

Vol 18, No 1 (2017)



An update on the feeding habits of fish in the Mediterranean Sea (2002-2015)

P. K. KARACHLE, K. I. STERGIUO

doi: [10.12681/mms.1968](https://doi.org/10.12681/mms.1968)

To cite this article:

KARACHLE, P. K., & STERGIUO, K. I. (2017). An update on the feeding habits of fish in the Mediterranean Sea (2002-2015). *Mediterranean Marine Science*, 18(1), 43–52. <https://doi.org/10.12681/mms.1968>

Table A. Feeding habits of Mediterranean marine fishes, based on studies published from 2000 to 2015. SM=sampling method (T=trawl; PT=pelagic trawl; H=hooks; N=nets; GN=gill nets; TN=trammel nets; PS=purse seiners; BS=beach seiners; SC=SCUBA; Sp=Speared; CL=commercial landings; LL=long line; Tr=traps; An=anaesthetic; FAD=fish aggregation devices; O=other); SF=sampling frequency (S=seasonal; M=monthly; BM=bimonthly; Da=daily; We=weekly; BWe=biweekly; 1=once, 2=twice, etc); LT=length type (TL=total length; FL=fork length; SL=standard length; LJFL=lower jaw standard length; DW=disk width); LR=length range (in cm); N=number of stomachs examined; Method: parameters estimated (F=frequency of occurrence; N=numerical percentage; W=percentage by weight; V=percentage by volume VC=percentage of empty stomachs; IRI=Index of Relative Importance; RA=relative abundance; MFI=modified food index); and Hb=habitat (as recorded in FishBase (Froese & Pauly, 2016); D=demersal; P=pelagic; BP=benthopelagic; RA=reef-associated; BaD=bathymedersal; BaP=bathypelagic). Asterisk denotes species/families/orders not included in Stergiou & Karpouzi (2002).

Species	Family	Area	SP	SM	SF	LM	LR	Method	N	Main prey	W (or N)	Hb	τ	SE	Reference
Elasmobranchii															
Carcharhiniformes															
1 <i>Galeus melastomus</i>	Scyliorhinidae	N-NW Aegean Sea	2001-2006	T	S	TL	23.3-54.0	W, VC	3	Cephalopoda	93.2	D	4.50	0.41	Karachle (2008); Karachle & Stergiou (2008, 2010b)
2 <i>Galeus melastomus</i>	Scyliorhinidae	Balearic Islands	2007-2009	T		TL	14.0-35.0	F, N, V, IRI, VC	170	Euphasiacea		D	3.34	0.46	Valls <i>et al.</i> (2011)
3 <i>Scyliorhinus canicula</i>	Scyliorhinidae	N-NW Aegean Sea	2001-2006	T	S	TL	10.0-63.0 24.1-45.1	W, VC	167 34	Euphasiacea, fish Cephalopoda, fish	50.2, 42.6	D	3.93 4.41	0.55 0.58	Karachle (2008); Karachle & Stergiou (2008, 2010b)
4 <i>Scyliorhinus canicula</i>	Scyliorhinidae	Sicilian Channel (Malta-Sicily)	6/2006	T	1	TL		N, W, VC	532	fish (T. trachurus, Macroramphosus scolopax), Crustacea (<i>Alpheus glaber</i> , <i>Chlorotocus crassicornis</i>), Cephalopoda	58.7, 19.3, 16.4	D	4.22	0.67	Gravino <i>et al.</i> (2010)
5 <i>Scyliorhinus canicula</i>	Scyliorhinidae	Balearic Islands	2007-2009	T		TL	11.0-53.0	F, N, V, IRI, VC	766	Macrura Reptantia, Polychaeta, fish		D	3.83	0.50	Valls <i>et al.</i> (2011)
6 <i>Scyliorhinus canicula</i>	Scyliorhinidae	Edremit Bay, Turkey		1998 T	M	TL	14.0-46.0 15.0-49.0	F, N, W, VC	66 60	Euphasiacea Euphasiacea <25.0: fish (<i>M. merluccius</i> , <i>S. pilchardus</i>) 25.0-45.0: fish (<i>S. pilchardus</i>) >45.0: fish (<i>S. pilchardus</i>)		D	3.37 3.48 4.50	0.47 0.50 0.80	
7 <i>Scyliorhinus canicula</i>	Scyliorhinidae	Foca, Izmir Bay, Turkey	9-11/2002	T		TL		F, N, W, IRI, VC	296	fish (<i>E. encrasicolus</i> , <i>G. niger</i>), Natantia	58.19, 18.05 (38.94, 23.89)	D	4.22	0.70	Filiz & Taskavak (2006)
8 <i>Scyliorhinus canicula</i>	Scyliorhinidae	Sardinia	2005-2008	T				F, N, W, IRI, VC	299	Crustacea (Decapoda, Mysidacea, Euphasiacea)	(67.0)	D	3.64	0.55	Mulas <i>et al.</i> (2010)
9 <i>Scyliorhinus canicula</i>	Scyliorhinidae	Aegean Sea, Greece	2005-2011	T,LL	S	TL	20.9-51.7	W, VC	432	fish, Cephalopoda, Crustacea	42.5, 30.8, 22.1	D	4.22	0.62	Kousteni (2015); Kousteni <i>et al.</i> (2015)
10 <i>Mustelus mustelus</i> *	Triakidae*	Gulf of Gabes, Tunisia	1/2002-12/2005	T,LL, GN		TL	34.0-158.5	F, N, W, IRI, VC	540	fish (Clupeidae, <i>D. annularis</i>), Cephalopoda (<i>Octopus vulgaris</i>)	53.8, 17.9 (34.82, 5.68)	D	4.26	0.68	Saidi <i>et al.</i> (2009a)
11 <i>Mustelus mustelus</i> *	Triakidae*	Aegean Sea, Turkey	2006-2007	T	S	TL	38.3-97.5	F, N, W, IRI, VC	158 84 146 81 71	<50: Crustacea 50-70: Crustacea 70-90: fish, Crustacea 90-110: fish >110: fish		D	3.50 3.70 4.21 4.41 4.41	0.56 0.57 0.72 0.75 0.75	
12 <i>Mustelus mustelus</i> *	Triakidae*	N Adriatic Sea	12/2005-4/2007	T		TL	50.5-152.5	F, N, W, IRI, VC	72	fish, Crustacea, Cephalopoda	40.15, 31.77, 27.29 (15.8, 76.47, 3.53)	D	4.17	0.61	Filiz (2009)
13 <i>Mustelus punctulatus</i> *	Triakidae*	Gulf of Gabes, Tunisia	1/2002-12/2005	T,LL, GN		TL	32.5-110.0	F, N, W, IRI, VC	133	fish (Sparidae, Carangidae), Cephalopoda (<i>Octopus vulgaris</i> , <i>Sepia officinalis</i>)	34.0, 29.7 (26.3, 12.23)	D	4.10	0.70	Saidi <i>et al.</i> (2009b)
14 <i>Mustelus punctulatus</i> *	Triakidae*	N Adriatic Sea	4/2005-4/2007	T				RA	9 30 59 35	<50: Crustacea 50-70: Crustacea, Sipuncula, Polychaeta 70-90: fish, Mollusca, Crustacea >90: Mollusca, fish		D	3.59 3.59 4.22 4.29	0.49 0.50 0.62 0.51	Gračan <i>et al.</i> (2013)
15 <i>Hexanchus griseus</i> *	Hexanchidae*	Sicily	2000-2003	TN,LL		TL	182-600		23	fish (<i>Mullus</i> sp., <i>Lepidopus caudatus</i> , <i>D. sargus sargus</i>), Cephalopoda (<i>Sepia officinalis</i> , <i>Octopus vulgaris</i>)		BaD	4.20	0.68	Celona <i>et al.</i> (2005)
16 <i>Hexanchus griseus</i> *	Hexanchidae*	Sea of Marmara, Black Sea, Turkey	11/1989-7/2000			TL	300.0-500.0	RA	9	fish (<i>P. saltatrix</i> , <i>M. merluccius</i> , <i>E. encrasicolus</i> , <i>Trachurus</i> spp.)		BaD	4.50	0.80	Kabasakal (2004)
Myliobatiformes*															
17 <i>Dasyatis pastinaca</i> *	Dasyatidae*	N-NW Aegean Sea	2001-2006	T	S	TL	50.1	W, VC	1	Natantia, Polychaeta	74.6, 25.4	D	3.46	0.53	Karachle (2008); Karachle & Stergiou (2008, 2010b)
18 <i>Dasyatis pastinaca</i> *	Dasyatidae*	Iskenderun Bay, Eastern Mediterranean Sea	4/1999-2/2000	T		TL	20.0-88.0	F, N, VC	251	Natantia, Stomatopoda	(81.3, 15.7)	D	3.70	0.59	Ismen (2003)
									20-29: Natantia 30-39: Natantia				3.70 3.60	0.58 0.57	

									40-49: Natantia, Stomatopoda		3.70	0.53		
									50-59: Natantia, Stomatopoda		3.80	0.55		
									60-69: Natantia, Stomatopoda, Brachyura		3.80	0.50		
									70-79: Natantia		3.70	0.57		
									>80: Natantia		3.60	0.59		
19	<i>Dasyatis pastinaca</i> *	Dasyatidae*	Cilician basin, northeastern Mediterranean	1999-2003	T	TL	14.6-100.9	F, N, IRI, VC	346	Natantia (<i>Metapenaeus stebbingi</i> , <i>Alpheus glaber</i> , <i>Charybdis longicollis</i> , <i>Penaeus semisulcatus</i>)	D	3.60	0.59	Yeldan <i>et al.</i> (2009)
20	<i>Pteroplatytrygon violacea</i> *	Dasyatidae*	Gulf of Trieste, Adriatic Sea	5-9/2004	T	DW	41.0-65.0	N	9	fish (<i>E. encrasicolus</i> , <i>C. macrophthalma</i>), Cephalopoda (<i>Sepia</i> sp.)	P	4.50	0.78	Mavric <i>et al.</i> (2004)
21	<i>Pteroplatytrygon violacea</i> *	Dasyatidae*	Gulf of Trieste, Adriatic Sea	4/2004-10/2005	T	DW		F, N, W, IRI, VC, †	84	fish (<i>E. encrasicolus</i> , <i>C. macrophthalma</i>), Cephalopoda (<i>Sepia</i> spp.)	P	4.50	0.73	Lipej <i>et al.</i> (2013)
22	<i>Myliobatis aquila</i> *	Myliobatidae*	Balearic Islands	2007-2009	T	TL	51.0-116.0	F, N, V, IRI, VC	23	Reptantia	BP	3.84	0.44	Valls <i>et al.</i> (2011)
23	<i>Myliobatis aquila</i> *	Myliobatidae*	north Adriatic, Croatia	1998-2002	T	DW	19.1-76.5	F, N, W, IRI, VC	165	Bivalvia (<i>Pterea hirundo</i> , <i>Cardium</i> sp.), Decapoda, Sipuncula (<i>Aspidosiphon mulleri</i>)	BP	3.37	0.43	Jardas <i>et al.</i> (2004)
Rajiformes														
24	<i>Dipturus nidarosiensis</i> *	Rajidae	Sardinia	2005-2011	T	TL	24.7-148.2	F, N, CV	23	Decapoda, fish	BaD	3.80	0.60	Follesa <i>et al.</i> (2012)
25	<i>Dipturus oxyrinchus</i> *	Rajidae	N-NW Aegean Sea	2001-2006	T	S TL	81.2	W, VC	1	Natantia	BaD	3.60	0.59	Karachle (2008); Karachle & Stergiou (2008, 2010b)
26	<i>Dipturus oxyrinchus</i> *	Rajidae	Saros Bay, North Aegean Sea, Turkey	3/2005-12/2007	T	TL	15.8-98.0	F, N, W, IRI, VC	124	Natantia (<i>Parapenaeus longirostris</i>), fish (<i>Trachurus</i> sp.)	BaD	3.82	0.63	Yigin & Ismen (2010a)
								F	8	10-19: Crustacea		3.48	0.54	
								F	6	20-29: Crustacea		3.67	0.56	
								F	9	30-39: Crustacea, Cephalopoda		4.04	0.49	
								F	10	40-49: Crustacea		3.71	0.62	
								F	6	50-59: Crustacea		3.60	0.59	
								F	12	60-69: Crustacea		3.60	0.59	
								F	19	70-79: Crustacea		3.97	0.61	
								F	18	80-89: Crustacea		3.70	0.62	
								F	9	90-99: Crustacea, fish		3.96	0.68	
27	<i>Dipturus oxyrinchus</i> *	Rajidae	Sardinia	2005-2010	T	TL	9.3-115.3	F, N, W, IRI, VC	255	Isopoda (<i>Lophogaster typicus</i>), Cephalopoda, fish	BaD	3.63	0.50	Mulas <i>et al.</i> (2015)
									140	<45.0: Isopoda (<i>Lophogaster typicus</i>)		3.41	0.44	
									56	45.0-75.0: Natantia (<i>Chlorothocous crassicornis</i>), Euphausiacea (<i>Meganctiphanes norvegica</i>)		3.75	0.54	
									59	>75.0: Cephalopoda (<i>Eledone</i> sp., <i>Octopus vulgaris</i>), fish		4.28	0.61	
28	<i>Leucoraja naevus</i> *	Rajidae	Balearic Islands	2007-2009	T	TL	22.0-52.0	F, N, V, IRI, VC	27	Natantia, fish	D	3.93	0.67	Valls <i>et al.</i> (2011)
29	<i>Raja asterias</i> *	Rajidae	Catalan Sea, W Mediterranean	2002-2003	T,CL	S TL		F, N, W, VC	102	Brachyura (<i>Liocarcinus depurator</i> , <i>Goneplax rhomboides</i>), fish (<i>M. barbatus</i>)	D	3.75	0.65	Navarro <i>et al.</i> (2013)
30	<i>Raja asterias</i> *	Rajidae	Tyrrhenian Sea, Italy	1999-2000	T	Da TL	21.0-46.5	N, CV	48	Brachyura (<i>Liocarcinus maculatus</i>), Natantia (<i>Alpheus glaber</i>)	D	3.60	0.61	Romanelli <i>et al.</i> (2007)
							46.0-56.0		43	Natantia (<i>Alpheus glaber</i>), Brachyura (<i>Liocarcinus depurator</i>)		3.85	0.65	
							55.0-62.5		38	Brachyura (<i>Liocarcinus depurator</i> , <i>Goneplax rhomboides</i>), fish		3.84	0.64	
31	<i>Raja asterias</i> *	Rajidae	N Tyrrhenian Sea, Italy	3/2001-8/2002	T	TL	12.0-64.0	F, N, W, IRI, VC	563	Brachyura (<i>Goneplax rhomboides</i>), fish	D	3.97	0.69	Serena <i>et al.</i> (2005)
32	<i>Raja asterias</i> *	Rajidae	S Ligurian Sea	3/2001-8/2002	T	TL	12.0-64.0	F, N, W, IRI, VC	563	Decapoda (<i>Goneplax rhomboides</i>)	D	3.60	0.60	Cuoco <i>et al.</i> (2005)
33	<i>Raja brachyura</i> *	Rajidae	Asinara Island, Sardinia	5-6/2002	TN	TL	22.5-95.5	F, N, W, IRI, VC	215	fish	D	4.48	0.79	Catalano <i>et al.</i> (2007)
									22.5-32.0: fish	95.42 (26.36)	4.45	0.79		
									32.1-42.0: fish	92.85 (53.49)	4.41	0.78		
									42.1-52.0: fish	99.45 (93.33)	4.50	0.80		
									>52.1: fish	99.99 (98.33)	4.50	0.80		
34	<i>Raja brachyura</i> *	Rajidae	Sardinia	6-7/2005, 1/2006	T	TL	18.8-95.5	F, N, IRI, VC	79	fish (<i>Gymnammodytes cicerellus</i>), Amphipoda (Gammaridae)	D	3.70	0.59	Follesa <i>et al.</i> (2010)
							18.8-48.5		51	Mysidacea, Amphipoda (Gammaridae)		3.40	0.48	
							51.0-80.0		14	fish (<i>Gymnammodytes cicerellus</i>)		4.50	0.77	
							84.6-95.5		9	fish (<i>Gymnammodytes cicerellus</i>)		4.50	0.78	

35	<i>Raja clavata*</i>	Rajidae	N-NW Aegean Sea	2001-2006	T	S	TL	25.6-46.5	W, VC	7	Natantia, fish, Brachyura	38.5, 36.3, 23.8	D	3.90	0.67	Karachle (2008); Karachle & Stergiou (2008, 2010b)		
36	<i>Raja clavata*</i>	Rajidae	Croatia, Adriatic Sea	1-12/2009	T	S	TL	14.0-75.1	F, N, W, IRI, VC	428	fish (<i>Gadiculus argenteus</i> , <i>T. capelanus</i> , <i>Maurolicus muelleri</i>), Natantia (<i>Alpheus glaber</i> , <i>Processa canaliculata</i>), Brachyura (<i>Liocarcinus corrugatus</i>)	28.7, 23.2, 17.4 (22.3, 29.7, 13.3)	D	3.80	0.62	Santic <i>et al.</i> (2012)		
										56	<25: Mysidacea, Amphipoda			3.35	0.49			
											25-35: Natantia			3.45	0.54			
											35-45: Natantia			3.60	0.59			
											45-55: Brachyura, Natantia			3.70	0.61			
											55-65: Natantia, fish			3.88	0.67			
										50	>65: fish, Cephalopoda			4.07	0.69			
37	<i>Raja clavata*</i>	Rajidae	Balearic Islands	2007-2009	T		TL	18.0-88.0	F, N, V, IRI, VC	266	fish, Macrura Reptantia, Natantia		D	4.00	0.60	Valls <i>et al.</i> (2011)		
											18.0-77.0			3.99	0.66			
											64.0-91.0			4.27	0.63			
38	<i>Raja clavata*</i>	Rajidae	Saros Bay, North Aegean Sea, Turkey	2/2005-12/2006	T		TL	10.7-88.0	F, N, W, IRI, VC	121	fish (<i>M. barbatus</i> , <i>E. encrasicolus</i> , <i>M. merluccius</i>), Brachyura (<i>Goneplax rhomboides</i>)	64.3, 22.9 (17.4, 50.8)	D	4.18	0.73	Yigin & Ismen (2010b)		
39	<i>Raja clavata*</i>	Rajidae	Sardinia	2005-2008	T				F, N, W, IRI, VC	242	Crustacea (Decapoda, Mysidacea)	(77.0)	D	3.57	0.53	Mulas <i>et al.</i> (2010)		
40	<i>Raja miraletus</i>	Rajidae	N-NW Aegean Sea	2001-2006	T	S	TL	22.6-33.9	W, VC	3	Brachyura, Cephalopoda	60.9, 31.3	D	3.82	0.54	Karachle (2008); Karachle & Stergiou (2008, 2010b)		
41	<i>Raja miraletus</i>	Rajidae	Sardinia	6-7/2005, 1/2006	T		TL	17.0-47.2	F, N, IRI, VC	100	Mysidacea, Amphipoda (Gammaridae), Brachyura (<i>Liocarcinus</i> spp.), Natantia (<i>Solenocera membranacea</i>)	(29.62, 25.32, 18.48, 16.21)	D	3.40	0.51	Follesa <i>et al.</i> (2010)		
											17.0-29.8			25	Amphipoda (Gammaridae), Mysidacea	(63.83, 40.57)	3.30	0.47
											30.5-40.0			47	Brachyura, Natantia	(54.52, 36.66)	3.50	0.58
											40.1-47.2			12	Natantia, Brachyura	(48.64, 26.88)	3.60	0.57
42	<i>Raja miraletus</i>	Rajidae	Balearic Islands	2007-2009	T		TL	24.0-43.0	F, N, V, IRI, VC	31	Macrura Reptantia, Natantia		D	3.60	0.57	Valls <i>et al.</i> (2011)		
43	<i>Raja miraletus</i>	Rajidae	Croatia, Adriatic Sea	1-12/2010	T	S	TL	13.5-40.0	F, N, W, IRI, VC	418	Natantia (<i>Processa canaliculata</i> , <i>Alpheus glaber</i>), Brachyura (<i>Macropipus tuberculatus</i> , <i>Galathea strigosa</i>), fish (<i>L. friessi</i>)	26.1, 24.9, 19.5 (35.8, 19.0, 6.1)	D	3.71	0.60	Santic <i>et al.</i> (2013)		
											<20: Mysidacea, Amphipoda			66		3.29	0.47	
											20-25: Brachyura, Mysidacea			101		3.42	0.55	
											25-30: Brachyura, Natantia			111		3.63	0.62	
											30-35: Natantia, Brachyura			91		3.73	0.64	
											>35: Natantia, Brachyura			49		3.73	0.64	
44	<i>Raja polystigma*</i>	Rajidae	Balearic Islands	2007-2009	T		TL	28.0-45.0	F, N, V, IRI, VC	15	Natantia		D	3.68	0.61	Valls <i>et al.</i> (2011)		
45	<i>Raja radula</i>	Rajidae	N-NW Aegean Sea	2001-2006	T	S	TL	21.8-32.0	W, VC	3	fish, Brachyura, Natantia	46.3, 28.1, 17.3	D	3.97	0.69	Karachle (2008); Karachle & Stergiou (2008, 2010b)		
46	<i>Raja radula</i>	Rajidae	Gulf of Gabes, Tunisia	1-12/2007	T,LL, GN	B	TL	13.4-80.0	F, N, W, IRI, VC	950	fish (Sparidae, Clupeidae), Natantia (<i>Metapenaeus monoceros</i> , <i>Sicyonia carinata</i>)	54.48, 19.99 (30.15, 24.69)	D	4.11	0.70	Kadri <i>et al.</i> (2013)		
											<40.0: Crustacea				3.50	0.50		
											40.0-55.0: Crustacea, fish				3.68	0.56		
											>55.0: fish				4.22	0.72		
47	<i>Rhinobatos rhinobatos</i>	Rhinobatidae	Gulf of Gabes, Tunisia		T		TL		N, W, IRI, VC		fish (Sparidae, Gobiidae)	71.83 (21.11)	D	4.28	0.73	Enajjar <i>et al.</i> (2007)		
48	<i>Rhinobatos rhinobatos</i>	Rhinobatidae	Iskenderun Bay, Eastern Mediterranean Sea	4/1999-2/2000	T		TL	22.2-120.0	F, N, VC	225	Stomatopoda, Natantia	(56.1, 23.3)	D	4.00	0.49	Ismen <i>et al.</i> (2007)		
											20-29: Natantia, Brachyura				3.50	0.52		
											30-39: Stomatopoda, Natantia				3.90	0.54		
											40-49: Natantia				3.70	0.54		
											50-59: Stomatopoda, Natantia				3.90	0.53		
											60-69: Natantia, Stomatopoda, fish				3.80	0.54		
											70-79: Natantia, Stomatopoda				3.80	0.52		
											80-89: fish, Stomatopoda, Natantia				4.20	0.63		
											90-99: fish, Stomatopoda, Natantia				4.10	0.66		
											>110-119: Brachyura, fish				4.00	0.71		
											>120: Brachyura, Stomatopoda				3.90	0.51		

Squaliformes

49	<i>Centrophorus granulosus*</i>	Centrophoridae*	Crete, Greece	7-8/2003	LL	TL	60.0-95.0	F, N, VC	43	fish (Scombridae), Cephalopoda (Histiotiethidae)	(63.89, 36.11)	BaD	4.50	0.65	Megalofonou & Chatzisprou (2006)	
50	<i>Dalatias licha</i>	Dalatiidae	N-NW Aegean Sea	2001-2006	T	S TL	38.0-40.2	W, VC	2	Cephalopoda	97.3	BaD	4.50	0.39	Karachle (2008); Karachle & Stergiou (2008, 2010b)	
51	<i>Dalatias licha</i>	Dalatiidae	Maghreb coast, Tunisia	1970-2007	T			RA	8	fish (<i>M. merluccius</i> , <i>G. melastomus</i>), Cephalopoda (<i>Sepiola</i> spp.)		BaD	4.50	0.80	Capape <i>et al.</i> (2008)	
52	<i>Etmopterus spinax</i>	Dalatiidae	Tuscany Archipelago	1985-2002	T			RA		9.5-19.0: Crustacea 19.5-29.0: Crustacea, fish 29.5-40.0: Crustacea, fish		BaD	3.69	0.56	Serena <i>et al.</i> (2006)	
53	<i>Etmopterus spinax</i>	Dalatiidae	Balearic Islands	2007-2009	T	TL	11.0-47.0	F, N, V, IRI, VC	46	Cephalopoda		BaD	4.39	0.55	Valls <i>et al.</i> (2011)	
54	<i>Squalus acanthias*</i>	Squalidae*	N Adriatic Sea	4/2005-4/2007	T			RA	157	fish (<i>E. encrasicolus</i> , <i>S. pilchardus</i>)		BP	4.23		Gračan <i>et al.</i> (2013)	
55	<i>Squalus acanthias*</i>	Squalidae*	Agios Efstratios Isl., N Aegean	26/6/2014	T	1 TL	22.5-40.0	F	60	Crustacea, Cephalopoda, fish		BP	3.61	0.54	Pori & Batzakas (2015)	
56	<i>Squalus acanthias*</i>	Squalidae*	SE Black Sea		T	TL	32-136	N	328	fish (<i>M. merlangus euxinus</i> , <i>Mullus barbatus</i> , <i>Gobius niger</i> , <i>S. sprattus phalericus</i>)	(73.37)	BP	4.40	0.78	Avsar (2001)	
57	<i>Squalus blainville*</i>	Squalidae*	Aegean Sea, Greece	2005-2011	T,LL	S TL	18.2-75.9	W, VC	211	Cephalopoda, fish	51.2, 41.7	D	4.42	0.60	Kousteni (2015); Kousteni <i>et al.</i> (2015)	
Torpediniformes																
58	<i>Torpedo marmorata</i>	Torpedinidae	N-NW Aegean Sea	2001-2006	T	S TL	8.8-37.3	W, VC	118	fish (<i>S. pilchardus</i> , <i>Gobius</i> spp.)	99.4	RA	4.39	0.67	Karachle (2008); Karachle & Stergiou (2008, 2010b)	
59	<i>Torpedo marmorata</i>	Torpedinidae	Gulf of Lions, NW Mediterranean	2001-2004	GN	TL	19.5-55.0	N, F, W, IRI, CV	102	fish (<i>Liza saliens</i>)	94.84 (86.0)	RA	4.50	0.78	Capape <i>et al.</i> (2007)	
60	<i>Torpedo torpedo</i>	Torpedinidae	Cullera coast, Spain	7/2004, 3-7/2005, 10/2005, 1/2006		TL	11.5-44.2	F, N, W, IRI, VC	63	fish	92.1 (50.8)	D	4.41	0.77	Jaramillo Londoño <i>et al.</i> (2011)	
61	<i>Torpedo torpedo</i>	Torpedinidae	Bizerte Lagoon, Tunisia	5/2010-4/2011	TN, GN	TL	10.0-43.5	F, N, W, IRI, VC	565	fish (<i>B. boops</i> , Gobiidae), Cephalopoda (<i>Sepia officinalis</i>)	49.21, 9.11 (39.31, 4.62)	D	4.04	0.66	El Kamel-Moutalibi <i>et al.</i> (2013)	
										10.0-26.9: fish (<i>M. barbatus</i> , Gobiidae)	56.43 (42.25)		4.02	0.69		
										27.0-43.5: fish (Gobiidae), Cephalopoda (<i>Sepia officinalis</i>)	47.02, 12.25 (37.25, 7.84)		4.05	0.65		
Actinopteri																
Anguilliformes																
62	<i>Conger conger</i>	Congridae	N-NW Aegean Sea	2001-2006	T	S TL	34.1-99.8	W, VC	31	fish (<i>C. macrophthalmus</i> , <i>E. encrasicolus</i>), Stomatopoda (<i>Squilla mantis</i>)	63.1, 25.1	D	4.18	0.58	Karachle (2008); Karachle & Stergiou (2008, 2012b)	
63	<i>Conger conger</i>	Congridae	Ionian Sea, Greece	6 & 10/2010	LL	TL	47.2-148.0	F, N, W, IRI, VC	44	fish (<i>Mora moro</i>)	96.37 (70.)	D	4.49	0.79	Anastasopoulou <i>et al.</i> (2012, 2013)	
64	<i>Conger conger</i>	Congridae	Adriatic Sea	7/2010-8/2011	LL	M TL	23.4-144.0	F, N, W, IRI, VC	317	fish (<i>H. dactylopterus</i> , <i>P. blennoides</i> , <i>B. boops</i>), Brachyura (<i>Eriphia verrucosa</i>)	72.47, 6.88 (46.37, 6.01)	D	4.30	0.73	Matic-Skoko <i>et al.</i> (2012)	
65	<i>Conger conger</i>	Congridae	central Adriatic Sea	2001-2002	T	TL	21.0-130.0	N, W, CV	170	fish (<i>Callionymus</i> sp., <i>G. niger</i>), Crustacea (<i>Goneplax</i> sp., <i>Munida</i> sp.)	49.0, 48.0	D	4.02	0.66	Vallisneri <i>et al.</i> (2007)	
66	<i>Conger conger</i>	Congridae	north Tunisian coasts	1/2008-12/2010	T	M TL	34.6-110.5	F, N, W, IRI, VC	288	fish (<i>S. aurita</i> , <i>S. pilchardus</i>), Cephalopoda (<i>Octopus defilippi</i>)	84.65, 10.30 (72.99, 4.21)	D	4.15	0.72	Sallami <i>et al.</i> (2015a)	
									61	<55.0: fish (<i>S. aurita</i>), Crustacea	66.78, 25.96 (44.23, 32.69)		3.94	0.66		
									67	55.0-67.5: fish (<i>S. aurita</i> , <i>S. pilchardus</i>)	92.05 (44.23)		4.20	0.72		
									86	67.5-80.0: fish (<i>S. aurita</i> , <i>P. pagrus</i> , <i>S. pilchardus</i>)	95.62 (86.15)		4.26	0.73		
									74	>80.0: fish (<i>S. aurita</i> , <i>S. pilchardus</i> , <i>D. vulgaris</i>), Cephalopoda (<i>Octopus defilippi</i>)	75.18, 22.37 (83.05, 1.69)		4.27	0.74		
67	<i>Gnathophis mystax</i>	Congridae	north Tunisian coasts	1/2010-12/2011	T	TL	17.0-36.6	F, N, W, IRI, VC	48	Crustacea, fish	92.64, 7.35 (89.47, 10.52)	D	3.51	0.60	Sallami <i>et al.</i> (2015b)	
68	<i>Muraena helena*</i>	Muraenidae*	Croatia, Adriatic Sea		LL	TL	27.5-100.0	F, N, W, IRI, VC	267	fish (<i>Phycis phycis</i>), Crustacea (<i>Pilumnus hirtellus</i> , <i>Galathea strigosa</i>)	66.27, 20.35 (69.23, 21.15)	RA	4.11	0.70	Matic-Skoko <i>et al.</i> (2010)	
69	<i>Muraena helena*</i>	Muraenidae*	NE Tunisia	1/2008-12/2010	CL	TL	47.2-109.2	F, N, W, IRI, VC	411	fish (<i>D. annularis</i> , <i>S. cabrilla</i>)	96.84 (78.83)	RA	4.27	0.74	Sallami <i>et al.</i> (2014)	
									223	47.2-69.7: fish (<i>D. annularis</i> , <i>S. cabrilla</i>)	94.16 (73.01)		4.19			
									188	70.0-109.2: fish (<i>D. annularis</i> , <i>S. cabrilla</i>)	98.01 (83.78)		4.21			
Aulopiformes																
70	<i>Saurida lessepsianus</i>	Synodontidae	Syria					RA		fish (<i>S. aurita</i> , <i>B. boops</i>)		RA	4.50	0.80	Ibrahim <i>et al.</i> (2010)	
71	<i>Synodus saurus</i>	Synodontidae	southern Tyrrhenian Sea, Sicily	6/2005-5/2006	T	4 TL		N, W, VC	224	fish (<i>P. acarne</i> , <i>S. aurita</i> , <i>B. boops</i>)	99.7 (94.2)	D	4.50	0.80	Esposito <i>et al.</i> (2009)	
Beloniformes*																
72	<i>Belone belone*</i>	Belonidae*	N-NW Aegean Sea	2001-2006	PS	S TL	27.2-53.5	W, VC	69	Crustacea, fish	40.4, 20.3	P	3.48	0.45	Karachle (2008); Karachle & Stergiou (2008, 2012b)	

73	<i>Belone belone*</i>	Belonidae*	Candarli & Izmir Bay, Aegean Sea, Turkey	1-12/2002	CL	M	TL	28.6-66.9	F, N, W, IRI, VC	597	Crustacea, fish		P	3.50	0.50	Sever <i>et al.</i> (2009)
74	<i>Belone belone*</i>	Belonidae*	Croatia, Adriatic Sea	2/2008-2/2009	PS		TL	23.1-72.0	F, N, VC	211	Euphasiacea, Copepoda	(59.5, 20.7)	P	3.16	0.34	Zorica & Cikes Kec (2012)
75	<i>Sargocentron rubrum</i>	Holocentridae	Syria						RA		Amphipoda, Isopoda, Decapoda		RA	3.36	0.52	Ibrahim <i>et al.</i> (2010)
76	<i>Alosa fallax*</i>	Clupeidae	N-NW Aegean Sea	2001-2006	T	S	TL	15.0-46.8	W, VC	27	fish (<i>E. encrasicolus</i> , <i>S. pilchardus</i>)	100.0	P	4.32	0.48	Karachle (2008); Karachle & Stergiou (2008)
77	<i>Alosa fallax*</i>	Clupeidae	Izmir Bay, Aegean Sea	11-12/2007	PS,T		FL	23.0-39.0	F, N, W, IRI, VC	208	fish (<i>E. encrasicolus</i>)	99.9(88.12)	P	4.50	0.80	Ceyhan <i>et al.</i> (2012)
78	<i>Sardina pilchardus</i>	Clupeidae	Thracian Sea, North Aegean	7/2007	PT	4	TL		N		Copepoda (<i>Acartia clausi</i> , <i>Euterpina acutifrons</i> , <i>Oncaea</i> spp.)	(86.45)	P	2.90	0.00	Nikolioudakis <i>et al.</i> (2012)
				7/2007							Diatoma (<i>Guinardia</i> spp.), Dinoflagellata (<i>Protoperidinium</i> spp.), Tintinnida (<i>Eutintinnus tubulosus</i>)	(40.36, 33.09, 9.28)		2.30	0.09	
				12/2007							Diatoma (<i>Pseudo-nitzschia</i> spp., <i>Chaetocerus</i> spp.), Dinoflagellata (<i>Neoceratium</i> spp.), Copepoda (<i>Centropages</i> spp.)	(60.76, 27.70, 8.55)		2.10	0.00	
				7/2008							Copepoda (<i>Oncaea</i> spp., <i>Euterpina acutifrons</i>), Appendicularia	(79.14, 7.07)		3.10	0.10	
				7/2008							Diatoma (<i>Coscinodiscus</i> spp., <i>Guinardia</i> spp.), Dinoflagellata (<i>Protoperidinium</i> spp., <i>Neoceratium</i> spp.)	(30.54, 30.51)		2.40	0.07	
				2/2009							Diatoma (<i>Rhizosolenia</i> spp., <i>Chaetocerus</i> spp.), Copepoda (<i>Oithona nana</i> , <i>Isias clavipes</i>), Crustacea larvae	(58.13, 26.20, 7.74, 7.21) (54.63, 16.80 (79.79, 8.17))		2.30 3.08	0.11 0.25	Sever <i>et al.</i> (2005)
79	<i>Sardina pilchardus</i>	Clupeidae	Ismir Bay, Aegean Sea	1-12/1997	PS,B	M	FL	9.6-14.9	F, N, W, IRI, VC	365	Copepoda (<i>Oithona nana</i> , <i>Isias clavipes</i>), Crustacea larvae	54.63, 16.80 (79.79, 8.17)	P	3.08	0.25	Sever <i>et al.</i> (2005)
80	<i>Sardina pilchardus</i>	Clupeidae	N-NW Aegean Sea	2001-2006	PS	S	TL	7.6-16.7	W, VC	752	Copepoda, Brachyura larvae	43.1, 38.9	P	3.14	0.29	Karachle (2008); Karachle & Stergiou (2008, 2014a)
81	<i>Sardinella aurita*</i>	Clupeidae	N-NW Aegean Sea	2001-2006	PS	S	TL	8.4-23.9	W, VC	230	Copepoda, Appendicularia	50.0, 22.1	P	3.20	0.32	Karachle (2008); Karachle & Stergiou (2008, 2014a)
82	<i>Sardinella aurita*</i>	Clupeidae	N Aegean Sea	10/2001	PS	S	TL	15.1-20.7	W, VC	30	Amphipoda, other Crustacea, Copepoda	36.23, 30.43, 18.84	P	3.08	0.29	Tsikliras <i>et al.</i> (2005)
				1/2002				14.5-20.0		30	Copepoda, other Crustacea, Ampipoda	58.92, 26.71, 10.96	P	3.04	0.20	
				4/2002				17.0-21.5		30	Copepoda, Decapoda larvae	33.75, 27.50	P	3.28	0.34	
				8/2002				16.0-19.4		30	Decapoda larvae, Amphipoda	39.56, 38.46	P	3.10	0.30	
				10/2001-8/2002						8	(15.0-16.0) Copepoda, Amphipoda	33.34, 33.33	P	3.08	0.32	
				10/2001-8/2002						24	(16.0-17.0) Decapoda larvae, Amphipoda	32.04, 24.57	P	3.20	0.36	
				10/2001-8/2002						10	(17.0-18.0) Amphipoda, Decapoda larvae	44.07, 28.44	P	3.13	0.35	
				10/2001-8/2002						40	(18.0-19.0) Siphonophora, Decapoda larvae	22.58, 20.32	P	3.31	0.32	
				10/2001-8/2002						22	(19.0-20.0) Copepoda	56.48	P	3.16	0.29	
				10/2001-8/2002						10	(20.0-21.0) Copepoda, Decapoda larvae	29.27, 11.71	P	3.21	0.34	
				10/2001-8/2002						6	(21.0-22.0) Copepoda	55.00	P	3.26	0.37	
83	<i>Sardinella aurita*</i>	Clupeidae	Sicily	4/2004-1/2005	PS	S	TL	9.3-31.5	F, N, W	254	Siphonophora (Diphyidae), fish larvae	41.73, 23.62 (1.70, 0.62)	P	3.54	0.52	Lomiri <i>et al.</i> (2008)
84	<i>Sardinella aurita*</i>	Clupeidae	Port Said, Egypt	10/2009-7/2010	PS	S	TL		F, N, VC	400	Phytoplankton, Copepoda	(49.9, 37.5)	P	2.60	0.11	Madkour (2012)
											(8.0-9.0) Phytoplankton, Detritus, Copepoda	(47.46, 25.42, 15.25)		2.50	0.16	
											(9.0-10.0) Phytoplankton, Copepoda, Detritus	(46.33, 25.42, 17.50)		2.60	0.17	
											(10.0-11.0) Phytoplankton, Copepoda, Detritus	(46.33, 23.73, 19.21)		2.40	0.10	
											(11.0-12.0) Phytoplankton, Copepoda, Detritus	(38.42, 36.16, 15.82)		2.50	0.09	
											(12.0-13.0) Copepoda, Phytoplankton	(47.46, 27.12)		2.80	0.17	
											(13.0-14.0) Copepoda, Phytoplankton	(40.11, 29.38)		2.50	0.09	
											(14.0-15.0) Copepoda, Phytoplankton	(64.97, 15.82)		2.90	0.10	
											(15.0-16.0) Copepoda	(86.44)		2.90	0.00	
85	<i>Sardinella aurita*</i>	Clupeidae	Syro-Lebanese waters						RA		Diatoma, Copepoda, Cladocera, fish larvae		P	3.20	0.36	Lakkis & Sabour (2010)
86	<i>Etrumeus golanii*</i>	Dussumieriidae*	El-Hammam to Rashid, Egypt	1-12/2008	CL	M	TL	9.0-25.0	F, N, W, MFI, VC	490	Natantia larvae, fish larvae (Engraulidae),	49.71, 30.65 (86.84, 6.51)	P	3.67	0.65	Osman <i>et al.</i> (2013)
											<15.0: Natantia larvae, fish larvae (Engraulidae)			3.65	0.63	
											16.0-20.0: Natantia larvae, fish larvae (Engraulidae)			3.65	0.62	
											>20.0: Natantia larvae, fish larvae (Engraulidae)			3.60	0.62	
87	<i>Engraulis encrasicolus</i>	Engraulidae	Ghazaouet, Algeria	9-12/2007	PS,T	M	TL		N, F,VC		Copepoda (<i>Oncaea mediterranea</i> , <i>Candacia longimana</i> , <i>Pleuromamma abdominalis</i>), Euphasiacea	(81.22, 8.71)	P	3.00	0.12	Bacha & Amara (2009)
88	<i>Engraulis encrasicolus</i>	Engraulidae	Benisaf, Algeria	9-12/2007	PS,T	M	TL		N, F,VC		Copepoda (<i>Candacia longimana</i> , <i>Temora stylifera</i> , <i>Oncaea mediterranea</i>)	(83.34)	P	3.00	0.10	Bacha & Amara (2009)

89	<i>Engraulis encrasicolus</i>	Engraulidae	Bejaia, Algeria	1-12/2007	PS, T	M	TL		N, F, VC	409	Copepoda (<i>Oncaea mediterranea</i>)	(93.04)	P	3.00	0.07	Bacha & Amara (2009)	
90	<i>Engraulis encrasicolus</i>	Engraulidae	Bay of Bénisaf, Algeria	7-12/2007	PS	M	TL	<11.8	N, F, VC		(G0) Copepoda (<i>Candacia longimana</i> , <i>Temora stylifera</i> , <i>Oncaea mediterranea</i>)	(95.94)	P	3.00	0.00	Bacha <i>et al.</i> (2009, 2010)	
91	<i>Engraulis encrasicolus</i>	Engraulidae	Bay of Bénisaf, Algeria	7-12/2007	PS	M	TL		N, F, VC		(G1) Copepoda (<i>Candacia longimana</i>)	(74.17)	P	3.00	0.15	Bacha <i>et al.</i> (2009, 2010)	
											(G2) Copepoda (<i>Candacia longimana</i> , <i>Oncaea mediterranea</i> , <i>Pleuromamma abdominalis</i>)	(76.62)		3.00	0.10		
92	<i>Engraulis encrasicolus</i>	Engraulidae	N-NW Aegean Sea	2001-2006	PS	S	TL	6.7-16.2	W, VC	759	Brachyura larvae, Euphasiacea larvae	28.8, 26.1	P	3.38	0.44	Karachle (2008); Karachle & Stergiou (2008, 2014a)	
93	<i>Engraulis encrasicolus</i>	Engraulidae	Izmir Bay, Turkey	12/1996-12/1997	PS		FL	6.9-14.0	F, N, W, IRI, VC	200	Copepoda, Thaliacea	41.91, 18.94 (77.74, 1.61)	P	3.29	0.38	Uğkun <i>et al.</i> (2003)	
										27	6.1-8.0: Copepoda	(82.69)		3.03	0.09		
										70	8.1-10.0: Copepoda	(71.16)		3.08	0.16		
										79	10.1-12.0: Copepoda	(88.25)		3.03	0.09		
										24	12.1-14.0: Copepoda	(75.37)		3.08	0.16		
94	<i>Engraulis encrasicolus</i>	Engraulidae	N Aegean Sea	7/2007	PT		TL		N		<10.0: Copepoda	(77.66)	P	3.04	0.13	Nikolioudakis <i>et al.</i> (2014)	
				7/2007	PT						>10.0: Copepoda	(84.96)		3.04	0.13		
				12/2007	PT						<10.0: Copepoda	(84.96)		3.01	0.07		
				12/2007	PT						>10.0: Copepoda	(78.04)		3.08	0.16		
				7/2008	PT						<10.0: Copepoda	(96.96)		3.00	0.00		
				7/2008	PT						>10.0: Copepoda	(97.72)		3.00	0.00		
				2/2000	PT						<10.0: Copepoda	(99.48)		3.00	0.00		
				2/2009	PT						>10.0: Copepoda	(99.32)		3.00	0.00		
Cyprinodontiformes*																	
95	<i>Aphanius fasciatus*</i>	Cyprinodontidae	Messolongi Lagoon, Messolongi, western Greece	3/1996-2/1997	N	M	TL		F, N, V, IRI, VC	859	Natantia (<i>Pandalus adspersus</i>), Plants, Isopoda	32.74, 13.3, 12.13 (10.70, 43.52, 3.38)	D	3.27	0.51	Leonardos (2008)	
96	<i>Aphanius fasciatus*</i>	Cyprinodontidae	Saltworks, Messolongi, western Greece		N	M	TL		F, N, V, IRI, VC	526	Isopoda (Sphaeromatidae), Insecta	32.42, 20.70 (7.66, 10.48)	D	3.25	0.48	Leonardos (2008)	
Gadiformes																	
97	<i>Merlangius merlangus*</i>	Gadidae	N-NW Aegean Sea	2001-2006	T	S	TL	14.1-29.1	W, VC	41	fish (<i>Gobius</i> spp., <i>E. encrasicolus</i>)	86.9	BP	4.38	0.73	Karachle (2008); Karachle & Stergiou (2008)	
98	<i>Merlangius merlangus*</i>	Gadidae	West Sea of Marmara	2000-2005	PS		FL		F, W, VC		Decapoda, fish	41.03, 38.21	BP	3.79	0.60	Artuz (2005a)	
99	<i>Merlangius merlangus*</i>	Gadidae	East Sea of Marmara	2000-2005	PS		FL		F, W, VC		Decapoda, fish	40.68, 36.57	BP	3.75	0.59	Artuz (2005a)	
100	<i>Micromesistius poutassou</i>	Gadidae	N-NW Aegean Sea	2001-2006	T	S	TL	9.2-24.0	W, VC	77	fish (<i>E. encrasicolus</i>), Amphipoda	75.6, 16.5	BaP	4.18	0.66	Karachle (2008); Karachle & Stergiou (2008)	
101	<i>Trisopterus capelanus</i>	Gadidae	Gulf of Valencia, Spain	10/1991-10/1994	CL	M	TL	8.5-23.0	F, N, W, VC	1276	Natantia (<i>Alpheus glaber</i>), fish (<i>Lesueurigobius friesii</i> , <i>Callionymus maculatus</i>)	47.6, 21.6 (33.72, 3.04)	BP	3.82	0.59	Morte <i>et al.</i> (2001)	
										558	<14.0: Natantia, Mysidacea			3.50	0.53		
										498	14.5-17.0: Natantia			3.60	0.56		
										220	>17.5: Natantia			3.70	0.61		
102	<i>Trisopterus capelanus</i>	Gadidae	N-NW Aegean Sea	2001-2006	T	S	TL	5.7-24.5	W, VC	174	fish (<i>L. suerii</i>), Natantia (Alpheidae)	55.2, 29.3	BP	4.13	0.64	Karachle (2008); Karachle & Stergiou (2008)	
103	<i>Gaidropsarus biscayensis</i>	Lotidae	N-NW Aegean Sea	2001-2006	T	S	TL	9.0-15.3	W, VC	65	Natantia, fish (Gobiidae)	50.8, 37.7	BP	3.93	0.67	Karachle (2008); Karachle & Stergiou (2008)	
104	<i>Gaidropsarus granti*</i>	Lotidae	Capo di Monte Santu, Sardinia	5/3/2007	TN		TL	31.9	RA	1	Natantia		D	3.60	0.59	Pais <i>et al.</i> (2008)	
105	<i>Gaidropsarus mediterraneus</i>	Lotidae	N-NW Aegean Sea	2001-2006	T	S	TL	8.5-14.5	W, VC	15	fish (Gobiidae), Brachyura	47.0, 36.4	D	3.95	0.61	Karachle (2008); Karachle & Stergiou (2008)	
106	<i>Coelorinchus caelorhincus</i>	Macrouridae	Sigacik Bay, Aegean Sea, Turkey	22/3/2003	T				F, N, W, IRI, VC	148	Copepoda (<i>Acartia</i> sp., <i>Aetideus armatus</i>), Decapoda	45.54, 30.95 (74.71, 18.87)	BP	3.20	0.34	Sever <i>et al.</i> (2008)	
107	<i>Coelorinchus labiatus*</i>	Macrouridae	Catalan Sea, W Mediterranean	27/6/1988-25/4/1991	T		TL		F, N, W, IRI, VC	15	Polychaeta, Amphipoda		BaD	3.12	0.33	Carrasson & Matallanas (2002)	
108	<i>Coryphaenoides guentheri*</i>	Macrouridae	Catalan Sea, W Mediterranean	27/6/1988-17/10/1988	T		TL		F, N, W, IRI, VC	14	Polychaeta, Foraminifera	31.9, 16.3	BaD	3.25	0.45	Carrasson & Matallanas (2002)	
										216	Amphipoda, Polychaeta	25.0, 15.7		3.28	0.46		
										264	Amphipoda, Polychaeta, Mysidacea	26.3, 24.4, 13.8		3.25	0.44		
109	<i>Coryphaenoides mediterraneus*</i>	Macrouridae	Catalan Sea, W Mediterranean	30/7/1987-29/7/1988	T		TL		F, N, W, IRI, VC	49	Natantia, Mysidacea, Polychaeta	28.5, 18.5, 13.3	BaP	3.36	0.49	Carrasson & Matallanas (2002)	
										114	Natantia, Mysidacea	43.1, 20.0		3.44	0.52		
110	<i>Hymenocephalus italicus</i>	Macrouridae	Cabrera Archipelago, Spain	8/2003-6/2004	T				F, N, W, IRI, VC	224	Mysidacea, Isopoda		BP	3.22	0.43	Fanelli & Cartes (2007)	
111	<i>Nezumia aequalis</i>	Macrouridae	Cabrera Archipelago, Spain	8/2003-6/2004	T				F, N, W, IRI, VC	258	Polychaeta, Amphipoda, Mysidacea		BP	3.18	0.41	Fanelli & Cartes (2007)	
112	<i>Trachyrincus scabrus</i>	Macrouridae	Catalan Sea, W Mediterranean	4/7/1988-25/4/1991	T		TL		F, N, W, IRI, VC	8	Copepoda, Mysidacea		BaD	3.14	0.28	Carrasson & Matallanas (2002)	
113	<i>Merluccius merluccius</i>	Merlucciidae	Alexandria, Egypt	3/2009-2/2010	T		TL	14.2-53.3	F, N, W, VC	549	fish (<i>C. rhonchus</i> , <i>S. aurita</i> , <i>E. gurnardus</i>)	93.74 (42.37)	D	4.36	0.77	Philips (2012)	
											<20.0: Crustacea	74.85 (25.0)		3.52	0.52		
											20.0-30.0: fish (<i>E. encrasicolus</i> , <i>S. aurita</i>)	64.63 (40.54)		4.09	0.72		

										>30.0: fish (<i>C. rhonchus</i> , <i>E. encrasicolus</i> , <i>S. aurita</i>)	94.04 (63.64)		4.44	0.79	
114	<i>Merluccius merluccius</i>	Merlucciidae	west Italy	1997-1998	T,GN M TL			F, N, W, IRI, VC	202	5.0-10.9: Euphasiacea (<i>Nictiphanes couchi</i>)		D	3.20	0.40	Carpentieri <i>et al.</i> (2005)
									430	11.0-15.9: Euphasiacea (<i>Nictiphanes couchi</i>), fish			3.66	0.56	
									564	16.0-20.9: fish (<i>S. pilchardus</i>)			4.41	0.78	
									454	21.0-25.9: fish (<i>S. pilchardus</i>)			4.41	0.78	
									555	26.0-30.9: fish (<i>S. pilchardus</i>)			4.41	0.78	
									224	31.0-35.9: fish (<i>S. pilchardus</i>)			4.41	0.78	
									139	36.0-40.9: fish (<i>Centracanthus cirrus</i>)			4.41	0.78	
									107	41.0-50.9: fish (<i>S. flexuosa</i> , <i>S. pilchardus</i>)			4.41	0.78	
									75	51.0-90.0: fish (Centracanthidae)			4.50	0.80	
115	<i>Merluccius merluccius</i>	Merlucciidae	N-NW Aegean Sea	2001-2006	T S TL	11.7-37.0	W, VC		21	fish (<i>E. encrasicolus</i>)	97.8	D	4.45	0.74	Karachle (2008); Karachle & Stergiou (2008)
116	<i>Merluccius merluccius</i>	Merlucciidae	Gulf of Castellammare	17/5-14/6/2005	T TL			N, IRI	30	6.0-12.0: Euphasiacea		D	3.36	0.46	Sinopoli <i>et al.</i> (2012)
									30	12.1-22.0: fish (<i>S. pilchardus</i>)			4.26	0.73	
									30	33.1-41.0: fish (<i>S. pilchardus</i>)			4.32	0.76	
			Termini Imerese, Sicily	17/5-14/6/2005	T TL			N, IRI	30	6.0-12.0: Euphasiacea			3.39	0.47	
									30	12.1-22.0: fish (<i>L. suerii</i>), Crustacea			3.85	0.59	
									12	33.1-41.0: fish (<i>S. pilchardus</i> , <i>E. encrasicolus</i>)			4.26	0.75	
			Sant'Agata, Sicily	17/5-14/6/2005	T TL			N, IRI	30	6.0-12.0: Euphasiacea			3.40	0.48	
									30	12.1-22.0: fish (<i>L. suerii</i> , <i>C. macrophthalma</i>), Crustacea			4.13	0.72	
									30	33.1-41.0: fish (<i>S. pilchardus</i> , <i>L. suerii</i>), Crustacea			4.26	0.75	
117	<i>Phycis blennoides</i>	Phycidae	Gulf of Valencia, Spain	10/1991-10/1994	CL M TL	9.5-33.5	F, N, W, VC		2631	Natantia (<i>Alpheus glaber</i>), fish	46.4, 24.2 (38.60, 3.77)	BP	3.89	0.62	Morte <i>et al.</i> (2002)
									647	<12.5: Natantia			3.60	0.55	
									1074	13-17: Natantia			3.70	0.54	
									659	17.5-21: Natantia			3.70	0.57	
									251	>21.5: Natantia			3.80	0.59	
118	<i>Phycis blennoides</i>	Phycidae	N-NW Aegean Sea	2001-2006	T S TL	8.1-37.4	W, VC		30	Anomura (Galatheididae), Natantia	64.4, 23.8	BP	3.55	0.59	Karachle (2008); Karachle & Stergiou (2008)
119	<i>Lophius budegassa</i>	Lophiidae	N-NW Aegean Sea	2001-2006	T S TL	5.0-38.4	W, VC		45	fish (<i>T. mediterraneus</i> , <i>S. pilchardus</i> , <i>S. hepatus</i>)	98.9	BaD	4.54	0.60	Karachle (2008); Karachle & Stergiou (2008)
120	<i>Lophius budegassa</i>	Lophiidae	North Adriatic	2005-2006	T TL			F, N, W, IRI, VC	283	fish (<i>M. merluccius</i>), Cephalopoda (<i>Illex coindetii</i>)	86.14, 8.98 (75.06, 4.59)	BaD	4.45	0.76	Stagnioni <i>et al.</i> (2013)
121	<i>Lophius budegassa</i>	Lophiidae	Tunisian coast	11/2004-1/2006	T M TL	15.0-70.0	F, N, W, IRI, VC		416	fish (<i>M. merluccius</i> , <i>M. barbatus</i> , <i>T. trachurus</i> , <i>T. mediterraneus</i>), Cephalopoda (<i>Todaropsis eblanae</i> , <i>Sepia orbignyana</i>)	87.91, 5.85 (78.05, 3.98)	BaD	4.47	0.52	Negzaoui-Garali <i>et al.</i> (2008)
									4	≤20.0: fish (Carangidae)			4.42	0.78	
									126	20.0-30.0: fish (Carangidae)			4.42	0.78	
									142	30.0-40.0: fish (Carangidae)			4.39	0.78	
									94	40.0-50.0: (Carangidae)			4.41	0.77	
									35	50.0-60.0: fish (Carangidae, Mullidae)			4.41	0.77	
									15	60.0-70.0: fish (Carangidae, Argentinidae)			4.37	0.76	
122	<i>Lophius piscatorius</i>	Lophiidae	N-NW Aegean Sea	2001-2006	T S TL	7.7-12.7	W, VC		6	fish (<i>T. capelanus</i> , <i>L. suerii</i>)	100.0	BaD	4.48	0.54	Karachle (2008); Karachle & Stergiou (2008)
123	<i>Diaphus metopoclampus</i> *	Myctophidae*	SW Sicily	21-24/10/2010, 5-9/5/2011	T SL	5.08-9.28	F, N, W, IRI, VC		296	Euphasiacea (<i>Nematoscelis megalops</i>), fish (Myctophidae)	31.0, 29.5 (12.11, 9.04)	BaP	3.66	0.58	Battaglia <i>et al.</i> (2014)
									110	<7.0: Euphasiacea, Copepoda (Calanoida)	48.8, 17.1 (10.64, 67.71)		3.44	0.48	
									173	>7.0: Euphasiacea, Natantia (Sergestidae), fish (Myctophidae)	33.6, 23.0, 21.2 (14.36, 4.83, 6.44)		3.59	0.54	
124	<i>Lampanyctus pusillus</i> *	Myctophidae*	Mallorca, Balearic Islands	12/2009	N 1 SL			F, N, W, IRI, VC		1.4-3.5: Copepoda (<i>Pleuromamma abdominalis</i> , <i>Pleuromamma</i> spp.)	(83.3)	BaP	3.10	0.16	Bernal <i>et al.</i> (2013)
										3.6-4.3: Copepoda (<i>Pleuromamma abdominalis</i> , <i>Pleuromamma</i> spp.), Euphasiacea	(68.4, 28.3)		3.05	0.18	
	Perciformes														
125	<i>Apogon imberbis</i>	Apogonidae	N-NW Aegean Sea	2001-2006	GN S TL	8.0-11.5	W, VC		37	Natantia, Mysidacea	69.9, 18.7	RA	3.54	0.56	Karachle (2008); Karachle & Stergiou (2008, 2010a)
126	<i>Parablennius gattorugine</i>	Blennidae	N-NW Aegean Sea	2001-2006	T S TL	13.4-17.9	W, VC		4	Macroalgae, Polychaeta	89.3, 10.7	D	2.11	0.09	Karachle (2008); Karachle & Stergiou (2008)
127	<i>Caranx crysos</i> *	Carangidae	Gulf of Gabes, Tunisia	6/2004-5/2006	PS,G FL	8.5-35.8	F, N, W, IRI, VC		1668	fish (<i>E. encrasicolus</i> , <i>S. aurita</i> , <i>S. pilchardus</i>)	89.6 (27.48)	RA	4.41	0.78	Sley <i>et al.</i> (2009)

										<18.0: fish, Crustacea	78.18, 21.80 (12.49, 87.42)	4.28	0.74		
										18.1-26.0: fish	90.74 (58.81)	4.41	0.78		
										>26.1: fish	98.23 (96.82)	4.48	0.80		
128 <i>Caranx rhonchus</i> *	Carangidae	N-NW Aegean Sea	2001-2006	PS	S	TL	18.0-19.8	W, VC	16	fish	100.0	BP	4.50	0.80	Karachle (2008); Karachle & Stergiou (2008)
129 <i>Caranx rhonchus</i> *	Carangidae	Gulf of Gabes, Tunisia	8/2004-7/2005	CL		TL	6.0-30.6	F, N, W, IRI, VC	777	fish (<i>E. encrasicolus</i>), Mollusca	81.4, 13.2 (31.7, 0.1)	BP	4.30	0.74	Sley <i>et al.</i> (2007)
130 <i>Caranx rhonchus</i> *	Carangidae	Gulf of Gabes, Tunisia	6/2004-5/2006	CL	M	TL	6.5-30.6	F, N, W, IRI, VC	1040	fish (<i>E. encrasicolus</i> , <i>S. aurita</i> , <i>S. pilchardus</i>)	80.7 (12.5)	BP	4.30	0.75	Sley <i>et al.</i> (2008)
										<16.0: Crustacea		3.68	0.56		
										16.1-22.0: Crustacea, fish		3.97	0.65		
										>22.0: fish, Crustacea		4.15	0.70		
131 <i>Naucrates ductor</i> *	Carangidae	Sicily	10/1994-1/1995	PS		SL	21.0-29.0	F, N, W, IRI, VC	80	Amphipoda (Hyperiidea), Mollusca (Atlantidae)	61.46, 28.21 (85.46, 6.27)	RA	3.37	0.56	Pipitone <i>et al.</i> (2000)
132 <i>Naucrates ductor</i> *	Carangidae	Aegean Sea, Greece			N	TL	12.0-33.0	RA		Crustacea larvae, Amphipoda (Hyperidae), Polychaeta (Alciopidae)		RA	3.19	0.35	Vassilopoulou <i>et al.</i> (2003)
133 <i>Trachurus mediterraneus</i>	Carangidae	Izmir and Candarli Bay, Aegean Sea	2008	PS,G		TL	9.7-25.3	F, N, W, IRI, VC	728	fish eggs and larvae, Copepoda	47.23, 27.29 (0.97, 79.49)	P	3.81	0.61	Bayhan <i>et al.</i> (2013)
134 <i>Trachurus mediterraneus</i>	Carangidae	off Varna and Bourgas, Bulgaria	5-9/2007	PT	M	TL	11.0-17.5	F, N, W, IRI, VC	1042	fish (<i>Sprattus sprattus ponticus</i> , <i>E. encrasicolus ponticus</i>), Amphipoda (<i>Gammarus subtypicus</i> , <i>Ampelisca diadema</i>), Cladocera (<i>Evadne tergestina</i> , <i>Podon polyphemoides</i>)	23.6, 15.5, 13.6	P	3.52	0.52	Yankova <i>et al.</i> (2008)
										11.0: Mysidacea		3.18	0.38		
										11.5: Mysidacea, Copepoda		3.17	0.36		
										12.0: Mysidacea, Copepoda		3.15	0.34		
										12.5: Mysidacea, Copepoda		3.15	0.34		
										13.0: Isopoda		3.29	0.49		
										13.5: Mysidacea		3.22	0.36		
										14.0: Mysidacea		3.18	0.36		
										14.5: fish, Crustacea		3.74	0.56		
										15.0: fish		4.01	0.66		
										15.5: fish, Mysidacea		3.87	0.61		
										16.0: fish, Mysidacea, Crustacea		3.59	0.52		
										16.5: fish, Mysidacea, Crustacea		3.59	0.52		
										17.0: fish, Mysidacea, Crustacea		3.59	0.50		
										17.5: fish, Polychaeta, Crustacea		3.67	0.53		
										18.0: fish, Crustacea		3.63	0.51		
135 <i>Trachurus mediterraneus</i>	Carangidae	N-NW Aegean Sea	2001-2006	PS,T	S	TL	7.0-25.8	W, VC	627	fish, Euphasiacea	58.3, 20.1	P	4.01	0.64	Karachle (2008); Karachle & Stergiou (2008, 2012a)
136 <i>Trachurus trachurus</i>	Carangidae	N-NW Aegean Sea	2001-2006	PS,T	S	TL	6.3-23.9	W, VC	133	Euphasiacea, fish, Amphipoda	28.4, 29.4, 20.1	P	3.58	0.50	Karachle (2008); Karachle & Stergiou (2008, 2012a)
137 <i>Trachurus trachurus</i>	Carangidae	Croatia, Adriatic Sea	1-12/1995	T	M	TL	12.9-37.6	F, N, W, IRI, VC	1200	fish (<i>Gadiculus argenteus</i> , <i>Maurolicus muelleri</i>), Euphasiacea (<i>Nyctiphanes couchii</i>)	65.7, 30.1 (3.4, 94.1)	P	4.08	0.69	Santic <i>et al.</i> (2005)
										<16.1: Euphasiacea		3.20	0.40		
										16.2-18.1: Euphasiacea		3.20	0.40		
										18.2-20.1: Euphasiacea		3.33	0.45		
										20.2-22.1: Euphasiacea, fish		3.46	0.49		
										22.2-24.1: Euphasiacea, fish		3.46	0.49		
										24.2-26.1: Euphasiacea, fish		3.33	0.45		
										26.2-28.1: Euphasiacea, fish		3.79	0.60		
										28.2-30.1: Euphasiacea, fish		3.86	0.62		
										30.2-32.1: fish, Euphasiacea		3.92	0.64		
										>32.2: fish, Euphasiacea		4.15	0.71		
138 <i>Trachurus trachurus</i>	Carangidae	Izmir and Candarli Bay, Aegean Sea	1-6, 9, 11/2003	PS		TL	10.2-28.4	F, N, W, IRI, VC	657	fish eggs and larvae	72.22 (15.92)	P	4.16	0.72	Bayhan & Sever (2009)
										10-12.9: fish		3.89	0.61		
										13-14.9: fish		3.97	0.67		
										15-16.9: fish		3.94	0.66		
										17-18.9: fish		4.17	0.72		
										19-20.9: fish		4.08	0.70		
										21-22.9: fish		4.13	0.71		
										23-24.9: fish		4.17	0.72		
										25-26.9: fish		4.18	0.73		
										27-28.9: fish		4.18	0.73		

139	<i>Spicara maena</i>	Centracanthidae	Algerian coasts	1999-2003		TL		F, N, VC	284	Copepods, chaetognaths, polychaets 8-10: Copepoda 10-12: Copepoda 12-14: Copepoda 14-16: Copepoda, Chaetognatha, Mysidacea 16-18: Copepoda, Amphipoda, Polychaeta 18-20: Copepoda, Polychaeta 20-22: Copepoda, Mysidacea, fish 22-24: fish, Mollusca	(71.25, 8.92, 7.83) (66.67) (96.62) (95.50) (68.5, 16.0, 7.5) (67.9, 14.9, 8.0) (65.8, 20.0) (32.7, 23.8, 7.2) (50.0, 50.0)	P	3.10 3.50 3.00 3.10 3.10 3.10 3.10 3.40 4.10	0.18 0.39 0.00 0.08 0.20 0.19 0.14 0.41 0.67	Harchouche <i>et al.</i> (2009)	
140	<i>Spicara maena</i>	Centracanthidae	N-NW Aegean Sea	2001-2006	PS,T, S GN	TL	9.0-20.2	W, VC	282	Copepoda, Mysidacea	54.3, 13.6	P	3.24	0.34	Karachle (2008); Karachle & Stergiou (2014b)	
141	<i>Spicara smaris</i>	Centracanthidae	N-NW Aegean Sea	2001-2006	PS,T, S GN	TL	7.0-18.5	W, VC	118	Copepoda (<i>Acartia</i> spp., <i>Oncaea</i> spp.), fish	63.5, 32.1	P	3.49	0.46	Karachle (2008); Karachle & Stergiou (2014b)	
142	<i>Cepala macrophthalma</i>	Cepolidae	N-NW Aegean Sea	2001-2006	T	S	TL	13.2-54.9	W, VC	195	Copepoda, Euphasiacea	45.7, 29.6	D	3.13	0.31	Karachle (2008); Karachle & Stergiou (2008)
143	<i>Cepala macrophthalma</i>	Cepolidae	Izmir Bay, Turkey	5/2005-6/2006	T			F, N, W, IRI, VC		11.0-13.9: Crustacea 14.0-16.9: Crustacea 17.0-19.9: Crustacea >20.0: Crustacea		D	3.00	0.00	Sever <i>et al.</i> (2010)	
144	<i>Clinitrachus argentatus*</i>	Clinidae*	northern Aegean and Marmara Seas, Turkey	1-12/2007	BS	M	TL	2.8-7.5	F, N, W, IRI, VC	59	Amphipoda, Brachyura	47.6, 15.2 (53.27, 6.25)	D	3.34	0.53	Ozen <i>et al.</i> (2010)
145	<i>Coryphaena hippurus*</i>	Coryphaenidae	Sicily, Ionian Sea	8/1994-1/1995, 7/1995-11/1995	FAD	SL		F, N, W, IRI, VC	132	fish (<i>E. encrasicholus</i> , <i>Glossanodon leioglossus</i>)	76.1 (48.77) 39.6, 15.2 (59.14, 7.53) 95.6 (75.0)	P	4.45	0.79	Castriota <i>et al.</i> (2007)	
			Sicily, Tyrrhenian Sea	8/1994-1/1995, 7/1995-11/1995	FAD	SL		F, N, W, IRI, VC	168	fish (<i>E. encrasicholus</i> , <i>P. bogaraveo</i>), Crustacea (Amphipoda Hyperidea, Euphaciacea), Cephalopoda	86.0, 8.2, 5.7 (31.8, 63.9, 1.2)	P	4.38	0.75	Castriota <i>et al.</i> (2007)	
								IRI		<30.0: fish 30.0-40.0: fish >40.0: fish			3.99 3.81 4.22	0.65 0.59 0.72		
146	<i>Ruvettus pretiosus*</i>	Gempylidae*	Cretan Sea, central Aegean Sea, Levantine Sea	10/2007-7/2008	LL	TL		F, N, CV	50	fish, Cephalopoda		BP	4.50	0.69	Vasilakopoulos <i>et al.</i> (2011)	
147	<i>Aphia minuta*</i>	Gobiidae	off Comacchio, Adriatic Sea	5/2003	T	TL	2.5-5.6	F, N, W, IRI, VC	338	Copepoda (<i>Temora longicornis</i> , <i>Acartia clausi</i>)	80.4 (88.6)	P	3.09	0.21	LaMesa <i>et al.</i> (2008)	
148	<i>Gobius niger</i>	Gobiidae	Gulf of Gabes, Tunisia	2/2009-1/2010	T	TL	6.4-15.7	F, N, W, IRI, VC, t	1055	Crustacea (Decapoda, Isopoda), Mollusca (Tellinidae)	60.0, 31.0	D	3.56	0.03	Hajji <i>et al.</i> (2013)	
									307	<10: Mollusca, Crustacea			3.54			
									342	10-11.4: Mollusca, Crustacea			3.57			
									406	>11.4: Crustacea, Mollusca			3.57			
149	<i>Gobius vittatus*</i>	Gobiidae	Croatia, Adriatic Sea	4/2001-3/2002	N,An	TL	1.92-5.4	F, N, W, IRI, VC	704	Polychaeta, Brachyura	27.3, 18.2	D	3.34	0.48	Kovacic (2007)	
										2.0-2.5: Mysidacea, Ostracoda 2.5-3.0: Bivalvia, Amphipoda 3.0-3.5: Polychaeta, Bivalvia, Gastropoda 3.5-4.0: Polychaeta, Gastropoda 4.0-4.5: Polychaeta, Bivalvia, Gastropoda 4.5-5.0: Polychaeta, Brachyura 5.0-5.5: Polychaeta, Brachyura	23.8, 21.4 23.8, 14.3 26.2, 19.1, 16.7 33.3, 16.7 16.7, 11.9, 11.9 26.2, 19.1 19.1, 16.7		3.33 3.25 3.25 3.27 3.46 3.36 3.33	0.44 0.40 0.41 0.42 0.50 0.47 0.45		
	<i>Lesueurigobius suerii*</i>	Gobiidae	N-NW Aegean Sea	2001-2006	T	S	TL	5.8-9.4	W, VC	141	Crustacea, Natantia, eggs	25.1, 20.0, 14.6	D	3.35	0.43	Karachle (2008); Karachle & Stergiou (2008, 2012b)
150																
151	<i>Neogobius melanostomus*</i>	Gobiidae	Black Sea, Romania		Tr			Ivlev		Amphipoda		D	3.44	0.54	Rosca <i>et al.</i> (2010)	
152	<i>Mesogobius batrachocephalus*</i>	Gobiidae	Black Sea, Romania		Tr			Ivlev		Amphipoda		D	3.45	0.54	Rosca <i>et al.</i> (2010)	
153	<i>Panticala platyrostris*</i>	Gobiidae	southeastern Black Sea	2005	Sp,N	TL		F, N, W, VC	241	fish, Brachyura, Amphipoda (Gammaridae)	45.9, 19.2, 13.6 (.2, 4.9, 65.3)	D	3.88	0.67	Engin & Seyhan (2010)	
154	<i>Tetrapturus belone*</i>	Istiophoridae *	Straits of Messina, Sicily	1995-2004	Sp	LJSL	89.0- 191.0	F, N, W, IRI, VC	69	fish (<i>B. belone</i> , <i>S. aurita</i> , <i>Scomberesox saurus</i>), Cephalopoda (<i>Tremoctopus violaceus</i>)	52.2, 47.8 (86.3, 12.1)	P	4.50	0.63	Castriota <i>et al.</i> (2008)	
155	<i>Coris julis</i>	Labridae	N-NW Aegean Sea	2001-2006	GN	S	TL	11.3-18.2	W, VC	78	Anomura, Crustacea	33.8, 29.7	RA	3.42	0.53	Karachle (2008); Karachle & Stergiou (2008, 2010a)

156	<i>Coris julis</i>	Labridae	Aegean Sea, Turkey	8/1999	N,Tr, O	SL	5.67-13.4	F, N, W, IRI, VC	28	Gastropoda, Crustacea		RA	3.41	0.45	Kabasakal (2001)
157	<i>Labrus bergylta</i> *	Labridae	Sea of Marmara	2004-2006	SC			F, W	158	Decapoda, Brachyura, Echinoidea	22.1, 15.9, 12.3	RA	3.24	0.41	Artuz (2004, 2005b)
158	<i>Labrus viridis</i>	Labridae	N-NW Aegean Sea	2001-2006	GN	S TL	19.5	W, VC	1	Gastropoda, Bivalvia	70.0, 30.0	RA	3.29	0.51	Karachle (2008); Karachle & Stergiou (2008, 2010a)
159	<i>Labrus viridis</i>	Labridae	Tunisia			TL	11.4-37.0	F, N, W, VC	583	Brachyura, Macrura Reptantia, fish	30.5, 28.6, 15.6 (14.3, 17.5, 9.3)	RA	3.82	0.49	Ouannes-Ghorbel & Bouain (2004)
160	<i>Symphodus ocellatus</i>	Labridae	Gabes, Tunisia			TL	7.4-14.4	F, N, W, VC	494	Isopoda, Macrura Reptantia	42.77, 26.59 (16.92, 8.76)	RA	3.57	0.49	Ouannes-Ghorbel <i>et al.</i> (2003)
161	<i>Symphodus ocellatus</i>	Labridae	Aegean Sea, Turkey	8/1999	N,Tr, O	SL	4.38-7.22	F, N, W, IRI, VC	20	Algae	60.99 (19.78)	RA	2.55	0.31	Kabasakal (2001)
162	<i>Symphodus roissali</i> *	Labridae	Aegean Sea, Turkey	8/1999	N,Tr, O	SL	3.81-8.27	F, N, W, IRI, VC	11	Algae, Gastropoda		RA	2.69	0.37	Kabasakal (2001)
163	<i>Symphodus tinca</i>	Labridae	Tunisia			CL TL	8.1-23.1	F, N, W, IRI, VC	1334	Macrura Reptantia, Isopoda	41.1, 20.6 (22.6, 17.8)	RA	3.71	0.48	Ouannes-Ghorbel & Bouain (2006)
										8.1-13.3: Crustacea, Mollusca 13.4-23.1: Crustacea, Mollusca	90.5, 9.5 (73.1, 26.9) 65.0, 8.12 (42.3, 33)		3.51 3.53	0.50 0.51	
164	<i>Symphodus tinca</i>	Labridae	N-NW Aegean Sea	2001-2006	GN	S TL	11.1-22.0	W, VC	212	Polychaeta	89.1	RA	2.95	0.25	Karachle (2008); Karachle & Stergiou (2008, 2010a)
165	<i>Thalassoma pavo</i> *	Labridae	Aegean Sea, Turkey	8/1999	N,Tr, O	SL	4.75- 10.25	F, N, W, IRI, VC	58	Crustacea		RA	3.42	0.43	Kabasakal (2001)
166	<i>Xyrichtys novacula</i> *	Labridae	N-NW Aegean Sea	2001-2006	GN	S TL	12.3-17.1	W, VC	12	Gastropoda, Bivalvia	37.7, 34.4	RA	3.37	0.51	Karachle (2008); Karachle & Stergiou (2008, 2010a)
167	<i>Xyrichtys novacula</i> *	Labridae	Selinunte, Sicily	10/2001	GN,T N	TL	12.2-18.1	N, F, V, IRI	64	Bivalvia (Tellinidae, <i>Nucula nucleus</i>), Anomoura	52.9, 17.9 (47.01, 7.49)	RA	3.24	0.43	Beltrano <i>et al.</i> (2006)
168	<i>Xyrichtys novacula</i> *	Labridae	Capo d'Orlando, Sicily	1994-2001	BS,H	TL		F, N, W, IRI, VC		Bivalvia, fish	34.1, 17.2	RA	3.49	0.49	Castriota <i>et al.</i> (2005)
169	<i>Mullus barbatus</i>	Mullidae	S Tyrrhenian Sea	6/2005-5/2006	T,BS	S TL	6.0-21.0	F, N, W, IRI, VC	667	Decapoda, Polychaeta, Bivalvia	26.7, 24.2, 21.4 (6.9, 9.0, 16.5)	D	3.33	0.45	Esposito <i>et al.</i> (2014)
									77	Mysidacea, fish	54.5, 19.9 (54.4, 2.0)		3.50	0.52	
									61	Polychaeta, Bivalvia, Decapoda	25.3, 19.1, 17.0 (2.9, 26.4, 7.8)		3.27	0.43	
									72	Polychaeta, Decapoda, Bivalvia	27.0, 17.3, 16.2 (4.3, 3.47, 9.7)		3.31	0.45	
									94	Decapoda, Bivalvia, Polychaeta	30.3, 24.7, 23.7 (5.1, 16.3, 12.5)		3.43	0.49	
170	<i>Mullus surmuletus</i>	Mullidae	N-NW Aegean Sea	2001-2006	GN	S TL	9.1-23.1	W, VC	55	Polychaeta, Natantia	46.3, 14.2	D	3.19	0.37	Karachle (2008); Karachle & Stergiou (2008)
171	<i>Mullus surmuletus</i>	Mullidae	M'diq, Marocco		2008	CL TL	8.2-38.1	F, N, W, IRI, VC	470	Macrura Reptantia, Amphipoda	27.9, 26.0	D	3.03	0.41	El Bakali <i>et al.</i> (2010)
										8-9: Amphipoda 9-10: Amphipoda 10-11: Amphipoda 11-12: Amphipoda 12-13: Amphipoda 13-14: Amphipoda 15-16: Amphipoda 16-17: Amphipoda 17-18: Decapoda, Amphipoda >18: Decapoda			3.30 3.20 3.20 3.30 3.30 3.30 3.40 3.40 3.40 3.50	0.46 0.43 0.46 0.44 0.46 0.46 0.48 0.47 0.49 0.50	
172	<i>Mullus surmuletus</i>	Mullidae	Cullera coast, Spain	7/2004, 3-7/2005, 10/2005, 1/2006		TL	14.7-28.5	F, N, W, IRI, VC	280	Brachyura, <i>Upogebia</i> sp.	49.2, 12.6 (20.0, 2.8)	D	3.55	0.51	Jaramillo Londoño <i>et al.</i> (2011)
173	<i>Nemipterus randalli</i> *	Nemipteridae*	Iskenderun Bay, Eastern Mediterranean Sea		2009	T TL	7.0-26.6	F, N, W, IRI, VC	117	Natantia (<i>Processa</i> sp.), Brachyura	50.1, 19.2 (62.92, 15.08)	D	3.70	0.62	Gurlek <i>et al.</i> (2010)
174	<i>Chromis chromis</i>	Pomacentridae	N-NW Aegean Sea	2001-2006	GN	S TL	8.6-13.3	W, VC	97	Copepoda, Appendicularia	43.5, 23.4	RA	3.25	0.37	Karachle (2008); Karachle & Stergiou (2008, 2010a)
175	<i>Pomatomus saltatrix</i> *	Pomatomidae*	N-NW Aegean Sea	2001-2006	PS	S TL	13.1-18.5	W, VC	6	fish	100.0	P	4.50	0.80	Karachle (2008); Karachle & Stergiou (2008)
176	<i>Pomatomus saltatrix</i> *	Pomatomidae*	Gulf of Gabes, Tunisia			TL	10.9-44.5	F, N, W, VC	654	fish (Clupeidae, Engraulidae), Cephalopoda	92.6, 5.0 (87.64, 3.37)	P	4.46	0.78	Dhieb <i>et al.</i> (2001)
177	<i>Sciaena umbra</i>	Sciaenidae	N-NW Aegean Sea	2001-2006	GN	S TL	12.2-16.0	W, VC	11	Natantia, Euphasiacea	40.0, 33.0	D	3.53	0.54	Karachle (2008); Karachle & Stergiou (2008)
178	<i>Sciaena umbra</i>	Sciaenidae	Gulf of Tunis, Tunisia	10/2008-11/2011	CL	M TL		IRI, VG	272	Crustacea		D	3.50	0.50	Charter <i>et al.</i> (2012)
179	<i>Euthynnus alletteratus</i>	Scombridae	north-eastern Sicily	8/2002-4/2004	N	TL	26.8-50.3	F, N, W, IRI, VC	187	fish (<i>Maurolicus muelleri</i> , <i>S. pilchardus</i>)	86.91 (92.01)	RA	4.38	0.76	Falautano <i>et al.</i> (2007)
										<30: fish 30-35: fish			4.50 4.50	0.80 0.80	

									35-40: fish	4.50	0.80					
									40-45: fish, Cephalopoda	4.50	0.77					
									>45: fish, Amphipoda	4.30	0.76					
180	<i>Euthynnus alletteratus</i>	Scombridae	Aegean & Ionian Seas, Greece	1995.1997-1999	PS	FL		F, N, W, VC	193 fish (<i>E. encrasicolus</i>), Amphipoda	94.0, 2.1 (75.4, 9.9)	RA	4.43	0.79	Zamboukas <i>et al.</i> (2001)		
181	<i>Sarda sarda</i>	Scombridae	NE Aegean Sea	5/2007-5/2009	PS,G	FL	26.6-65.5	F, N, W, IRI, VC	134 fish (<i>E. encrasicolus</i> , <i>Spicara</i> spp.)	94.7 (70.77)	P	4.46	0.79	Fletcher <i>et al.</i> (2013)		
182	<i>Sarda sarda</i>	Scombridae	Greek Seas (Aegean, Ionian Sea)	1997-2003	PS,G	TL, FL		F, N, W, IRI, VC	397 fish (<i>E. encrasicolus</i> , <i>S. pilchardus</i> , <i>S. aurita</i>)	99.9 (98.3)	P	4.48	0.79	Zamboukas <i>et al.</i> (2003), Zamboukas (2006)		
183	<i>Scomber colias</i> *	Scombridae	N-NW Aegean Sea	2001-2006	PS	S TL	8.8-26.8	W, VC	126 <45.0: fish (<i>E. encrasicolus</i> , <i>S. pilchardus</i>) 271 >45.0: fish (<i>E. encrasicolus</i> , <i>S. pilchardus</i>)	99.9 (98.6) 99.9 (98.0)	P	4.48	0.80	4.47	0.79	Karachle (2008); Karachle & Stergiou (2008, 2012a)
184	<i>Scomber scombrus</i>	Scombridae	N-NW Aegean Sea	2001-2006	PS	S TL	13.3-27.4	W, VC	204 fish (<i>E. encrasicolus</i>)	97.1	P	4.37	0.54	Karachle (2008); Karachle & Stergiou (2008, 2012a)		
185	<i>Anthias anthias</i> *	Serranidae	N-NW Aegean Sea	2001-2006	GN	S TL	12.7-16.6	W, VC	9 Crustacea	93.6	RA	3.54	0.52	Karachle (2008); Karachle & Stergiou (2008, 2010a)		
186	<i>Epinephelus costae</i> *	Serranidae	east coasts of Algeria	2001-2002		TL	18.9-90.0	F, N, W, VC	74 Ascidiacea, fish	52.27, 25.40 (47.87, 1.15)	D	3.39	0.46	Derbal & Kara (2004, 2007)		
187	<i>Epinephelus marginatus</i>	Serranidae	Balearic Islands	1998-2000	Sp,T	TL	13.4-105.6	F, N, W, VC	203 Cephalopoda (<i>Octopus vulgaris</i> , <i>Sepia officinalis</i>)	77.47 (16.73)	RA	4.38	0.46	Linde <i>et al.</i> (2004)		
188	<i>Epinephelus marginatus</i>	Serranidae	Syro-Lebanese waters					RA	fish, Cephalopoda, Crustacea		RA	4.10	0.63	Lakkis & Sabour (2010)		
189	<i>Epinephelus marginatus</i>	Serranidae	Balearic Islands		LL			F	fish, Brachyura, Cephalopoda		RA	4.13	0.62	Quetglas <i>et al.</i> (2001)		
190	<i>Serranus cabrilla</i>	Serranidae	N-NW Aegean Sea	2001-2006	GN,T	S TL	9.5-23.1	W, VC	34 fish, Brachyura	42.7, 20.5	D	3.90	0.67	Karachle (2008); Karachle & Stergiou (2008)		
191	<i>Serranus cabrilla</i>	Serranidae	Ligurian Sea	1996-1997	Sp			F, N, W, VC	49 Natantia (<i>Lysmata seticaudata</i>), Brachyura (<i>Xantho pilipes</i>)	34.5, 32.8 (50.1, 17.2)	D	3.84	0.66	Relini <i>et al.</i> (2002)		
192	<i>Serranus hepatus</i>	Serranidae	N-NW Aegean Sea	2001-2006	T	S TL	5.7-13.1	W, VC	99 Brachyura (<i>Goneplax</i> spp.), Crustacea, Natantia	29.2, 22.2, 17.2	D	3.77	0.63	Karachle (2008); Karachle & Stergiou (2008)		
193	<i>Serranus hepatus</i>	Serranidae	Ismir Bay, Aegean Sea	1,5,10/2002	T	TL	5.2-11.7	N, F, W, IRI, VC	603 Brachyura, Natantia	39.97, 35.42 (10.14, 17.87)	D	3.66	0.61	Bilecenoglu (2009)		
194	<i>Serranus hepatus</i>	Serranidae	Uzunada Island, Izmir Bay, Aegean Sea	1/12/2005	T	M		F, N, W, IRI, VC	278 Natantia, Brachyura	60.3, 29.34 (40.91, 27.27)	D	3.58	0.59	Yapici <i>et al.</i> (2012)		
195	<i>Serranus scriba</i>	Serranidae	N-NW Aegean Sea	2001-2006	GN	S TL	10.6-23.6	W, VC	81 fish (<i>S. tinca</i> , <i>Gobius</i> spp.), Brachyura, Natantia	50.9, 23.8, 18.6	D	3.94	0.66	Karachle (2008); Karachle & Stergiou (2008)		
196	<i>Siganus luridus</i>	Siganidae	Zakynthos Isl., Ionian Sea	5/2014	TN	1 TL	15.5-23.0	F, W, IRI	10 Algae (<i>Dictyota</i> sp., <i>Cystoseira</i> sp.)	88.52	RA	2.00	0.00	Athineou <i>et al.</i> (2015)		
197	<i>Siganus luridus</i>	Siganidae	Syria	2002		BM		RA	Algae (<i>Ulva</i> spp., <i>Cladophora</i> spp., <i>Enteromorpha</i> spp.)		RA	2.00	0.00	Sabour & Lakkis (2007)		
198	<i>Siganus rivulatus</i>	Siganidae	Syria	2002		BM		RA	Algae		RA	2.00	0.00	Sabour & Lakkis (2007)		
199	<i>Siganus rivulatus</i>	Siganidae	Syro-Lebanese waters					RA	algae		RA	2.00	0.00	Lakkis & Sabour (2010)		
200	<i>Boops boops</i>	Sparidae	N-NW Aegean Sea	2001-2006	T,PS,	S TL	11.2-19.9	W, VC	106 Euphysiacea, fish	55.2, 22.4	D	3.52	0.52	Karachle (2008, 2014); Karachle & Stergiou (2008)		
201	<i>Dentex dentex</i>	Sparidae	N-NW Aegean Sea	2001-2006	GN	S TL	11.7-15.3	W, VC	10 fish larvae, fish	52.4, 46.4	BP	4.49	0.80	Karachle (2008, 2014); Karachle & Stergiou (2008)		
202	<i>Diplodus annularis</i>	Sparidae	N-NW Aegean Sea	2001-2006	T,GN	S TL	6.1-17.5	W, VC	427 Crustacea, Polychaeta, eggs	47.0, 8.4, 7.6	BP	3.20	0.43	Karachle (2008, 2014); Karachle & Stergiou (2008)		
203	<i>Diplodus annularis</i>	Sparidae	Calich Lagoon, Sardinia	1998		TL	11.0-20.0	F, N, V, IRI, VC	130 Detritus, Mollusca	63.23, 21.42 (43.85, 23.36)	BP	2.59	0.29	Chessa <i>et al.</i> (2004)		
204	<i>Diplodus annularis</i>	Sparidae	Gulf of Annaba, Algeria	10/1999-9/2000		TL	10.4-18.8	F, N, W, VC	719 Algae, Ascidiacea (<i>Clavelina</i> sp., <i>Ciona</i> sp.)	25.76, 17.67 (4.16, 46.02)	BP	3.05	0.40	Nouacer <i>et al.</i> (2004)		
205	<i>Diplodus annularis</i>	Sparidae	Ligurian Sea	1996-1997	TN			F, N, VC	357 Hydrozoa (<i>Obelia dichotoma</i>), Polychaeta, Amphipoda (<i>Pariambus typicus</i>)	(30.8, 25.4, 26.3)	BP	3.26	0.43	Relini <i>et al.</i> (2002)		
206	<i>Diplodus puntazzo</i>	Sparidae	eastern central Adriatic, Croatia	12/1998-1/2000	BS	TL	15.2-41.5	F, N, W, IRI, VC	412 Bivalvia, Porifera	51.8, 23.1 (23.4, 8.8)	BP	3.12	0.36	Dulcic <i>et al.</i> (2006)		
207	<i>Diplodus sargus sargus</i>	Sparidae	Abu Qir Bay, Alexandria Egypt	8/2008-7/2009	CL	M TL		F, N, V, IRI, VC	403 Bivalvia, Polychaeta, Natantia	<24.0: Ophiuroidea, algae >24.0: Bivalvia (54.0, 12.0, 11.4)	D	3.40	0.45	Osman & Mahmoud (2009)		
208	<i>Diplodus sargus sargus</i>	Sparidae	Corfu, Ionian Sea, Greece	23/5/2004	N	TL		F, N, W, VC	24 Gasteropoda, Isopoda	10.0-15.0: Polychaeta, Natantia, algae 15.0-20.0: Natantia, Bivalvia 20.0-25.0: Bivalvia, fish (43.0, 31.6) (54.0, 14.0) (43.0, 33.0)	D	3.22	0.53	Primikiri <i>et al.</i> (2005)		
									5 15.5-20.5: Gastropoda, Isopoda	(47.0, 26.0)		3.21	0.53			
									12 20.5-25.5: Gastropoda	(62.0)		3.34	0.56			

209	<i>Diplodus vulgaris</i>	Sparidae	Abu Qir Bay, Alexandria Egypt	8/2008-7/2009	CL	M	TL		F, N, V, IRI, VC	7	25.5-33.0: Isopoda, Gastropoda, Amphipoda	(43.0, 26.0, 24.0)	3.25	0.54		
									W, VC	98	Polychaeta, Isopoda, Amphipoda	(61.2, 14.6, 11.4)	BP	3.20	0.37	Osman & Mahmoud (2009)
210	<i>Diplodus vulgaris</i>	Sparidae	N-NW Aegean Sea	2001-2006	T,GN	S	TL	9.0-16.7	W, VC	50	Polychaeta	90.2	BP	3.08	0.28	Karachle (2008, 2014); Karachle & Stergiou (2008)
211	<i>Lithognathus mormyrus</i>	Sparidae	Calich Lagoon, Sardinia	1998			TL	17.0-24.0	F, N, V, IRI, VC	128	Detritus, Amphipoda	42.19, 40.66 (30.56, 33.22)	D	2.78	0.38	Chessa <i>et al.</i> (2005)
212	<i>Lithognathus mormyrus</i>	Sparidae	Thracian Sea, North Aegean	1997-2000	CL		TL	15.6-31.0	F, N, V, IRI, VC	120	Bivalvia, Polychaeta, fish	29.4, 29.0, 16.7 (19.1, 53.8, 0.0)	D	3.42	0.46	Kallianiotis <i>et al.</i> (2005)
									IRI		(age <4) Polychaeta			3.20	0.35	
									IRI		(age 4-5) Polychaeta, Bivalvia			3.20	0.32	
									IRI		(age 6-7) Polychaeta, Bivalvia			3.10	0.34	
									IRI		(age 8-10) Polychaeta, Bivalvia, Echinoidea			3.20	0.33	
213	<i>Oblada melanura</i>	Sparidae	N-NW Aegean Sea	2001-2006	T,GN	S	TL	12.6-22.7	W, VC	56	Polychaeta, Crustacea	42.2, 13.3	BP	3.11	0.42	Karachle (2008, 2014); Karachle & Stergiou (2008)
214	<i>Oblada melanura</i>	Sparidae	Croatia, Adriatic Sea	1992-1993	BS		TL	11.3-29.3	F, N, W, IRI, VC	926	Copepoda (<i>Lucicutia flavirostris</i>), fish (<i>A. hepsetus</i>)	45.16, 24.9 (81.78, 10.21)	BP	3.43	0.46	Pallaoro <i>et al.</i> (2003)
										298	<17: Copepoda	76.23 (84.66)		3.23	0.31	
										360	17-22: Copepoda	75.31 (80.67)		3.26	0.34	
										268	>22: Copepoda, fish	47.95, 27.19 (75.98, 10.03)		3.53	0.49	
215	<i>Pagellus acarne</i>	Sparidae	N-NW Aegean Sea	2001-2006	T,GN	S	TL	10.5-19.2	W, VC	63	fish larvae, Cephalopoda, Polychaeta, Crustacea	23.5, 19.6, 12.7, 11.0	BP	3.84	0.55	Karachle (2008, 2014); Karachle & Stergiou (2008)
216	<i>Pagellus bogaraveo</i>	Sparidae	N-NW Aegean Sea	2001-2006	T,GN	S	TL	9.3-23.1	W, VC	72	fish	86.8	BP	4.43	0.76	Karachle (2008, 2014); Karachle & Stergiou (2008)
217	<i>Pagellus erythrinus</i>	Sparidae	N-NW Aegean Sea	2001-2006	GN	S	TL	8.4-16.4	W, VC	59	Posidonia oceanica, Natantia, Maroura reptantia	55.8, 17.5, 13.6	BP	3.30	0.39	Karachle (2008, 2014); Karachle & Stergiou (2008)
218	<i>Pagellus erythrinus</i>	Sparidae	Croatia, Adriatic Sea	2009	T		TL	10.0-26.1	F, N, W, IRI, VC	120	Natantia (<i>Alpheus dentipes</i> , <i>Alpheus</i> sp., <i>Processa canaliculata</i>), Bivalvia (<i>Cardium edule</i>)	35.6, 22.8 (33.2, 18.9)	BP	3.52	0.52	Santic <i>et al.</i> (2011)
219	<i>Pagrus pagrus</i>	Sparidae	N-NW Aegean Sea	2001-2006	GN	S	TL	10.2-15.5	W, VC	10	fish, Polychaeta	61.5, 32.1	BP	3.36	0.34	Karachle (2008, 2014); Karachle & Stergiou (2008)
220	<i>Sarpa salpa</i>	Sparidae	N-NW Aegean Sea	2001-2006	GN	S	TL	11.7-19.2	W, VC	25	Macroalgae, <i>Cymodocea</i> spp, <i>Posidonia oceanica</i>	70.3, 25.8, 2.7	BP	2.00	0.00	Karachle (2008, 2014); Karachle & Stergiou (2008)
221	<i>Spondyllosoma cantharus</i>	Sparidae	N-NW Aegean Sea	2001-2006	GN	S	TL	9.7-14.0	W, VC	82	Crustacea, Natantia, Ascidiacea	37.3, 15.2, 12.4	BP	3.39	0.46	Karachle (2008, 2014); Karachle & Stergiou (2008)
222	<i>Spondyllosoma cantharus</i>	Sparidae	eastern central Adriatic, Croatia	12/1998-1/2000	BS,T		TL	14.0-46.5	F, N, W, IRI, VC	432	Amphipoda, Polychaeta	53.0, 30.8 (27.6, 58.5)	BP	3.48	0.51	Dulcic <i>et al.</i> (2006)
											<22.0: Amphipoda, Ophiuroidea	71.1, 27.6 (22.6, 42.1)		3.44	0.51	
											>22.0: Amphipoda, Polychaeta	52.2, 44.9 (22.5, 65.6)		3.62	0.44	
223	<i>Trachinus draco</i> *	Trachinidae*	N-NW Aegean Sea	2001-2006	T	S	TL	15.0-30.5	W, VC	25	fish, Mysidacea	62.2, 16.5	D	4.19	0.66	Karachle (2008); Karachle & Stergiou (2008, 2012b)
224	<i>Uranoscopus scaber</i>	Uranoscopidae	N-NW Aegean Sea	2001-2006	T,GN	S	TL	8.7-26.9	W, VC	70	fish (<i>L. suerii</i> , <i>T. capelanus</i>)	91.2	D	4.43	0.75	Karachle (2008); Karachle & Stergiou (2008)
225	<i>Uranoscopus scaber</i>	Uranoscopidae	Alexandria, Egypt	7/2005-6/2006	T	M			F, N, W, VC	418	fish (<i>S. smarís</i> , <i>S. flexuosa</i>)	90.58 (34.79)	D	4.36	0.77	Rizkalla & Philips (2008)
											<15: fish			4.39	0.79	
											15-20: fish			4.40	0.78	
											>20: fish (<i>S. smarís</i> , <i>S. flexuosa</i>)			4.33	0.76	
226	<i>Uranoscopus scaber</i>	Uranoscopidae	Cullera coast, Spain	7/2004, 3-7/2005, 10/2005, 1/2006			TL	13.9-35.5	F, N, W, IRI, VC	139	fish	88.6 (68.2)	D	4.39	0.77	Jaramillo Londoño <i>et al.</i> (2011)
227	<i>Xiphias gladius</i>	Xiphiidae	Aegean Sea, Turkey	1999-2000	CL				F, N	108	fish (<i>S. pilchardus</i> , <i>E. encrasicolus</i>), Cephalopoda (<i>Sepietta oweniana</i> , <i>Todarodes sagittatus</i>)	(81.13, 17.86)	P	4.50	0.73	Salman (2004)
228	<i>Xiphias gladius</i>	Xiphiidae	S Aegean Sea, Greece	1/1994-9/1995	LL		LJFL	68.0-214.0	F, N, W, VC	96	fish (<i>S. scombrus</i>), Cephalopoda (<i>Todarodes sagittatus</i>)	49.38, 34.85 (74.89, 5.75)	P	4.33	0.63	Peristeraki & Tserpes (2001)
Pleuronectiformes																
229	<i>Arnoglossus laterna</i>	Bothidae	Gulf of Lions, NW Mediterranean	2000-2001	T	S	TL		W	76	depth 0-20m: Polychaeta, Amphipoda, Phoronida	30.8, 18.0, 16.1	D	3.30	0.44	Darnaude <i>et al.</i> (2004b)
										312	depth 30-50m: Natantia, Polychaeta	43.9, 28.8		3.39	0.50	
										208	depth 70-100m: Natantia, Mysidacea	41.7, 33.2		3.43	0.55	
230	<i>Arnoglossus laterna</i>	Bothidae	Gulf of Lions, NW Mediterranean	2000-2001	T	S	TL	2.0-13.2	W	741	Macrura Reptantia, Polychaeta	36.2, 24.3	D	3.56	0.42	Darnaude <i>et al.</i> (2004a)

231	<i>Arnoglossus laterna</i>	Bothidae	off Rhone River delta	2000-2001	T	S	TL	2.0-4.9	N, W	58	Cumacea, Polychaeta, Amphipoda	47.5, 21.5, 15.9	D	3.21	0.46	Darnaude (2005)
232	<i>Arnoglossus laterna</i>	Bothidae	Izmir Bay, Turkey, Aegean Sea	2002-2003	T		TL	8.1-13.2	F, N, W, IRI, VC	596	Natantia, Polychaeta	39.1, 26.0		3.38	0.50	
										540	Fish, Mysidacea	31.69, 21.85 (21.4, 76.86)	D	3.67	0.60	Bayhan <i>et al.</i> (2008)
233	<i>Arnoglossus laterna</i>	Bothidae	N-NW Aegean Sea	2001-2006	T	S	TL	4.5-16.9	W, VC	212	fish (<i>L. suerii</i> , Gobiidae)	85.4	D	4.35	0.74	Karachle (2008); Karachle & Stergiou (2008, 2011)
234	<i>Arnoglossus laterna</i>	Bothidae	Gulf of Castellammare	2005	T	1	TL		F, N, W, IRI	74	fish	62.3 (16.5)	D	4.15	0.72	Fanelli <i>et al.</i> (2009)
235	<i>Arnoglossus laterna</i>	Bothidae	Gulf of Termini Imerese	2005	T	1	TL		F, N, W, IRI	74	fish	80.2 (15.5)	D	4.32	0.76	Fanelli <i>et al.</i> (2009)
236	<i>Arnoglossus laterna</i>	Bothidae	Gulf of Sant' Agata	2005	T	1	TL		F, N, W, IRI	44	fish	77.7 (15.5)	D	4.29	0.76	Fanelli <i>et al.</i> (2009)
237	<i>Arnoglossus thori</i>	Bothidae	N-NW Aegean Sea	2001-2006	T	S	TL	9.1-11.2	W, VC	3	Crustacea, Natantia, Amphipoda	35.7, 28.5, 14.4	D	3.61	0.57	Karachle (2008); Karachle & Stergiou (2008, 2011)
238	<i>Bothus podas</i>	Bothidae	N-NW Aegean Sea	2001-2006	GN	S	TL	11.3-17.2	W, VC	22	Isopoda, Natantia, Polychaeta	38.4, 20.5, 14.1	D	3.39	0.53	Karachle (2008); Karachle & Stergiou (2008, 2011)
239	<i>Bothus podas</i>	Bothidae	Gulf of Gabes, Tunisia	5/2004-9/2005		We	TL	13.0-24.4	F,N,W,IRI, VC	749	Isopoda (<i>Cymodoce truncata</i>), Amphipoda (Gammaridae)	32.90, 27.79 (36.63, 32.47)	D	3.39	0.49	Abid <i>et al.</i> (2013)
										49	< 16: Amphipoda (Gammaridae), Isopoda (<i>Cymodoce truncata</i>)	32.75, 23.05 (39.96, 32.05)		3.47	0.54	
										577	16-21: Isopoda (<i>Cymodoce truncata</i>), Amphipoda (Gammaridae)	35.46, 28.29 (39.21, 32.26)		3.37	0.52	
										21	> 21: Isopoda (<i>Cymodoce truncata</i>), Amphipoda (Gammaridae)	32.93, 24.13 (45.24, 30.38)		3.45	0.53	
240	<i>Citharus linguatula</i>	Citharidae	Gulf of Lions, NW Mediterranean	2000-2001	T	S	TL	6.3-23.7	W	122	fish	75.7	D	4.33	0.72	Darnaude <i>et al.</i> (2004a)
241	<i>Citharus linguatula</i>	Citharidae	off Rhone River delta	2000-2001	T	S	TL	6.3-9.1	N, W	31	fish	64.8	D	4.06	0.69	Darnaude (2005)
242	<i>Citharus linguatula</i>	Citharidae	Ebro Delta, NW Mediterranean	2003	T	2	TL	16.1-23.7	N, W	51	fish	73.5		4.22	0.74	
											0-10 cm, fish (<i>L. suerii</i> , Gobidae), Mysidacea (<i>Leptomysis gracilis</i>)	66.79 (16.56)	D	4.27	0.75	de Juan <i>et al.</i> (2007)
											10-15 cm, fish (<i>L. suerii</i>)	96.20 (63.15)		4.46	0.79	
											15-25 cm, fish (<i>L. suerii</i> , <i>Pomatochistus</i> sp.)	98.9 (87.50)		4.49	0.80	
											0-10 cm, fish (<i>A. laterna</i> , <i>L. suerii</i>), Mysidacea (<i>Leptomysis gracilis</i>)	60.9, 21.08 (19.1, 67.89)		4.08	0.70	
											10-15 cm, fish (<i>L. suerii</i>), Mysidacea (<i>Leptomysis gracilis</i>)	45.7, 25.8 (3.79, 85.78)		3.91	0.66	
											15-25 cm, fish (<i>L. suerii</i>)	82.82 (22.47)		4.34	0.76	
243	<i>Citharus linguatula</i>	Citharidae	N-NW Aegean Sea	2001-2006	T	S	TL	3.9-24.3	W, VC	169	fish (<i>L. suerii</i> , <i>T. trachurus</i>)	76.6	D	4.34	0.69	Karachle (2008); Karachle & Stergiou (2008, 2011)
244	<i>Citharus linguatula</i>	Citharidae	Izmir Bay, Aegean Sea	1/2002-3/2003	T		TL		F, N, W, IRI, VC	432	Mysidacea, fish, Decapoda		D	3.47	0.50	Bayhan <i>et al.</i> (2009)
245	<i>Symphurus nigrescens</i>	Cynoglossidae	N-NW Aegean Sea	2001-2006	T	S	TL	6.4-11.9	W, VC	10	Brachyura, Polychaeta	66.6, 33.2	D	3.35	0.51	Karachle (2008); Karachle & Stergiou (2008, 2011)
246	<i>Buglossidium luteum</i>	Soleidae	Gulf of Lions, NW Mediterranean	2000-2001	T	S	TL	3.2-12.6	W	146	Polychaeta, Amphipods, Bivalvia	47.3, 12.2, 12.2	D	3.27	0.39	Darnaude <i>et al.</i> (2004a)
247	<i>Buglossidium luteum</i>	Soleidae	Gulf of Lions, NW Mediterranean	2000-2001	T	S	TL		W	30	depth 0-20m: Polychaeta, Bivalvia	46.0, 17.2	D	3.23	0.41	Darnaude <i>et al.</i> (2004b)
										31	depth 30-50m: Polychaeta, Bivalvia, Amphipoda	69.9, 8.3, 8.1		3.13	0.35	
										31	depth 70-100m: Natantia, Amphipoda, Mysidacea, Polychaeta	25.0, 23.5, 20.1, 18.1		3.31	0.50	
248	<i>Buglossidium luteum</i>	Soleidae	off Rhone River delta	2000-2001	T	S	TL	3.2-4.8	N, W	31	Polychaeta, Amphipoda, Cumacea	21.9, 17.4, 17.4	D	3.18	0.42	Darnaude (2005)
249	<i>Microchirus variegatus*</i>	Soleidae	N-NW Aegean Sea	2001-2006	GN	S	TL	8.3-12.6	W, VC	51	Polychaeta, Bivalvia	56.8, 14.8		3.16	0.37	
										3	Polychaeta	100.0	D	3.06	0.26	Karachle (2008); Karachle & Stergiou (2008, 2011)
250	<i>Monochirus hispidus*</i>	Soleidae	N-NW Aegean Sea	2001-2006	GN	S	TL	9.2-12.8	W, VC	24	Polychaeta, Sipuncula, Mollusca	69.2, 15.4, 14.4	D	3.19	0.32	Karachle (2008); Karachle & Stergiou (2008, 2011)
251	<i>Pegusa impar*</i>	Soleidae	Gulf of Lions, NW Mediterranean	2000-2001	T	S	TL	3.3-21.2	W	101	Bivalvia, Polychaeta	53.0, 16.3	D	3.20	0.37	Darnaude <i>et al.</i> (2004a)
252	<i>Pegusa lascaris*</i>	Soleidae	off Rhone River delta	2000-2001	T	S	TL	4.9-9.6	N, W	47	Polychaeta, Cumacea	65.6, 13.7	D	3.12	0.36	Darnaude (2005)
253	<i>Solea solea</i>	Soleidae	Gulf of Lions, NW Mediterranean	2000-2001	T	S	TL	16.0-21.2		32	Bivalvia	78.4		3.15	0.35	
										307	Polychaeta	71.0	D	3.15	0.32	Darnaude <i>et al.</i> (2004a)
254	<i>Solea solea</i>	Soleidae	Gulf of Lions, NW Mediterranean	2000-2001	T	S	TL		W	32	depth 0-20m: Polychaeta, Bivalvia	37.8, 23.6	D	3.34	0.43	Darnaude <i>et al.</i> (2004b)
										35	depth 30-50m: Polychaeta	88.3		3.08	0.28	
										30	depth 70-100m: Polychaeta, Holothuria	73.5, 9.2		3.11	0.30	
255	<i>Solea solea</i>	Soleidae	off Rhone River delta	2000-2001	T	S	TL	3.2-9.2	N, W	33	Polychaeta, Bivalvia, Amphipoda	34.5, 26.6, 17.7	D	3.21	0.40	Darnaude (2005)
										97	Polychaeta	75.9		3.14	0.31	
256	<i>Solea solea</i>	Soleidae	Syro-Lebanesse waters						RA		Mollusca, Crustacea, algae		D	3.20	0.39	Lakkis & Sabour (2010)
257	<i>Solea solea</i>	Soleidae	Port Said, Egypt	2006	T	M	TL		W, VC	550	Polychaeta, Copepoda	40.7, 18.8	D	2.77	0.22	El-mor & Ahamed (2008)

								9	11.5-12.4: Polychaeta, Copepoda	74.9, 25.2		3.05	0.22		
								13	12.5-13.4: Polychaeta, Copepoda	68.2, 16.4		3.10	0.30		
								26	13.5-14.4: Polychaeta, Copepoda	62.3, 20.9		3.07	0.27		
								38	14.5-15.4: Polychaeta, Copepoda	59.2, 20.2		3.04	0.27		
								52	15.5-16.4: Polychaeta, Copepoda	49.6, 19.9		3.04	0.27		
								97	16.5-17.4: Polychaeta, Mollusca	42.9, 20.5		3.06	0.29		
								103	17.5-18.4: Polychaeta, detritus	41.8, 21.7		2.79	0.22		
								63	18.5-19.4: Polychaeta, detritus	36.6, 20.9		2.68	0.20		
								60	19.5-20.4: algae, Polychaeta	40.3, 22.2		2.49	0.15		
								47	20.5-21.4: algae, detritus	48.8, 19.3		2.35	0.12		
								26	21.5-22.4: algae	64.1		2.28	0.08		
								16	22.5-23.4: algae	70.0		2.26	0.08		
258	<i>Synaptura lusitanica*</i>	Soleidae	Cullera coast, Spain	7/2004, 3-7/2005, 10/2005, 1/2006		TL	14.5-34.3	F, N, W, IRI, VC	476	Polychaeta	91.6 (98.8)	D	3.10	0.27	Jaramillo Londoño <i>et al.</i> (2011)
Scorpaeniformes															
259	<i>Scorpaena notata</i>	Scorpaenidae	Ligurian Sea	1996-1997		TN		F, N, W, VC	69	Brachyura (<i>Liocarcinus vernalis</i> , <i>Herbstia condyliata</i> , <i>Liocarcinus maculatus</i>)	82.4 (86.3)	D	3.58	0.57	Relini <i>et al.</i> (2002)
						Sp		F, N, W, VC	49	Brachyura (<i>Liocarcinus</i> sp., <i>Maja crispata</i>), Polychaeta	63.7, 22.9 (77.8, 3.7)		3.43	0.54	
260	<i>Scorpaena notata</i>	Scorpaenidae	N-NW Aegean Sea	2001-2006	T	S TL	8.3-17.8	W, VC	42	Brachyura (<i>Goneplax</i> spp., <i>Liocarcinus</i> spp.)	81.0	D	3.60	0.62	Karachle (2008); Karachle & Stergiou (2008)
261	<i>Scorpaena notata</i>	Scorpaenidae	Abruzzi, central Adriatic	7/2005-5/2006	T,GN	S TL	9.4-17.0	F, N, W, IRI, VC	67	Brachyura (<i>Goneplax rhomboides</i>)	89.2 (55.80)	D	3.53	0.60	Castriota <i>et al.</i> (2012)
262	<i>Scorpaena porcus</i>	Scorpaenidae	Abruzzi, central Adriatic	7/2005-5/2006	T,GN	S TL	9.0-24.6	F, N, W, IRI, VC	118	Brachyura (<i>Goneplax rhomboides</i>)	88.4 (75.6)	D	3.55	0.61	Castriota <i>et al.</i> (2012)
263	<i>Scorpaena porcus</i>	Scorpaenidae	N-NW Aegean Sea	2001-2006	T,GN	S TL	8.2-26.4	W, VC	96	Brachyura (<i>Ilia nucleus</i>), fish	47.3, 43.2	D	3.90	0.69	Karachle (2008); Karachle & Stergiou (2008)
264	<i>Scorpaena porcus</i>	Scorpaenidae	Ligurian Sea	1996-1997		TN		F, N, W, VC	88	Brachyura (<i>Liocarcinus vernalis</i>)	91.8 (43.0)	D	3.54	0.61	Relini <i>et al.</i> (2002)
265	<i>Scorpaena porcus</i>	Scorpaenidae	Black Sea, Romania			Tr		Ilev		Amphipoda		D	3.56	0.55	Rosca <i>et al.</i> (2010)
266	<i>Scorpaena porcus</i>	Scorpaenidae	southeastern Black Sea	1/2002-5/2003	T,TN	M TL		W	327	Brachyura (<i>Carcinus mediterraneus</i>), Natantia (<i>Crangon crangon</i>)	33.3, 16.6	D	3.58	0.55	Demirhan & Can (2009)
267	<i>Scorpaena porcus</i>	Scorpaenidae	Trabzon coast, SE Black Sea	11/2003-2/2004	T	TL	6.3-23.5	N, F, W, IRI, CV	262	fish (<i>M. barbatus ponticus</i> , <i>Hippocampus</i> sp.), Macrura Reptantia (<i>Upogebia pusilla</i>)	42.43, 31.28 (34.51, 21.96)	D	4.15	0.65	Bağcınar & Sağlam (2009)
								IRI	143	6-12: fish (<i>M. barbatus ponticus</i>), Brachyura (<i>Liocarcinus depurator</i>)			4.20	0.71	
								IRI	101	12-18: Macrura Reptantia (<i>Upogebia pusilla</i>), fish (<i>M. barbatus ponticus</i>), Brachyura (<i>Liocarcinus depurator</i>)			4.10	0.60	
								IRI	18	18-24: fish (<i>Hippocampus</i> sp., <i>M. barbatus ponticus</i>), Brachyura (<i>Liocarcinus depurator</i>)			4.20	0.67	
268	<i>Scorpaena scrofa</i>	Scorpaenidae	Cullera coast, Spain	7/2004, 3-7/2005, 10/2005, 1/2006		TL	8.9-37.0	F, N, W, IRI, VC	115	Stomatopoda (<i>Squilla</i> sp.)	99.5 (29.0)	D	4.08	0.64	Jaramillo Londoño <i>et al.</i> (2011)
269	<i>Scorpaena scrofa</i>	Scorpaenidae	Sardinia					N, F, W, IRI	64	fish (<i>C. macrophthalmus</i> , <i>M. surmuletus</i>), Cephalopoda (<i>Sepiella</i> sp.)	69.75, 17.10 (40.43, 2.13)	D	4.40	0.72	Cabiddu <i>et al.</i> (2010)
								IRI		<11: Crustacea, fish			3.90	0.62	
								IRI		11-21: fish			4.20	0.71	
								IRI		>21: fish			4.30	0.75	
270	<i>Chelidonichthys lucerna</i>	Triglidae	N-NW Aegean Sea	2001-2006	T	S TL	6.0-21.6	W, VC	15	Brachyura	80.3	D	3.64	0.63	Karachle (2008); Karachle & Stergiou (2008)
271	<i>Chelidonichthys lucerna</i>	Triglidae	Abruzzi, central Adriatic	7/2005-5/2006	T,GN	S TL	11.2-32.0	F, N, W, IRI, VC	387	Brachyura (<i>Goneplax rhomboides</i>)	78.8 (64.9)	D	3.66	0.63	Castriota <i>et al.</i> (2012)
272	<i>Chelidonichthys lucerna</i>	Triglidae	North Adriatic	5/2005-3/2007	T	TL	6.3-41.5	F, N, W, IRI, VC	1096	Brachyura (<i>Goneplax rhomboides</i> , <i>Liocarcinus depurator</i>), fish (<i>E. encrasicolus</i> , <i>G. niger</i>)	57.38, 39.74 (27.24, 6.72)	D	3.91	0.69	Stagnioni <i>et al.</i> (2012)
273	<i>Chelidonichthys lucerna</i>	Triglidae	Adriatic Sea	5/2005-3/2007	T	S TL	6.3-41.5	F, N, W, IRI, VC	1114	Crustacea, fish	58.01, 39.74 (89.69, 6.72)	D	3.90	0.68	Vallisneri <i>et al.</i> (2011)
274	<i>Chelidonichthys obscurus</i>	Triglidae	Gulf of Gabes, Tunisia	2/2000-7/2002	T	SL		N, F, W, CV	555	Mysidacea (<i>Gastrosaccus</i> sp., <i>Anchialina agilis</i>), Natantia (<i>Pontocaris lacazei</i>)	51.4, 20.1 (82.67, 2.99)	D	3.44	0.47	Boudaya <i>et al.</i> (2007)
								N, CV	50	<14.0: Mysidacea	(95.0)		3.20	0.40	
								N, CV	505	>14.0: Mysidacea	(80.0)		3.20	0.43	
275	<i>Eutrigla gurnardus</i>	Triglidae	N-NW Aegean Sea	2001-2006	T	S TL	6.3-14.8	W, VC	10	Natantia	93.9	D	3.58	0.58	Karachle (2008); Karachle & Stergiou (2008)
276	<i>Lepidotrigla cavillone</i>	Triglidae	N-NW Aegean Sea	2001-2006	T	S TL	8.8-11.7	W, VC	4	Crustacea	100.0	D	3.50	0.50	Karachle (2008); Karachle & Stergiou (2008)
277	<i>Trigloporus lastoviza</i>	Triglidae	N-NW Aegean Sea	2001-2006	T	S TL	16.9	W, VC	1	Euphysiacea, Brachyura	44.4, 41.7	D	3.32	0.49	Karachle (2008); Karachle & Stergiou (2008)
278	<i>Trigloporus lastoviza</i>	Triglidae	Gulf of Gabes, Tunisia	2/2000-7/2002	T	SL		N, F, W, CV	860	Natantia (<i>Sicyonia carinata</i> , <i>Pontocaris lacazei</i>), Mysidacea (<i>Gastrosaccus</i> sp.)	34.2, 27.6 (3.96, 91.83)	D	3.50	0.52	Boudaya <i>et al.</i> (2007)
								N, CV	146	<12.0: Mysidacea, other Crustacea	(40.0, 20.0)		3.40	0.47	

							N, CV	714	>12.0: Mysidacea	(70.0)		3.40	0.45			
Syngnathiformes																
279	<i>Fistularia commersonii</i> *	Fistulariidae*	N-NW Aegean Sea	2001-2006	GN	S	TL	92.0	W, VC	1	fish	100.0	RA	4.50	0.80	Karachle (2008); Karachle <i>et al.</i> (2004); Karachle & Stergiou (2008)
280	<i>Fistularia commersonii</i> *	Fistulariidae*	Lebanon	5/2003-11/2004	BS,G		TL		N, F, W, IRI	243	fish (<i>S. smar</i> , <i>B. boops</i>)	100.0	RA	4.50	0.80	Bariche <i>et al.</i> (2009)
											20-50: fish (Sparidae, Centracanthidae)	(100.0)		4.50	0.80	
											50-70: fish (Centracanthidae, Sparidae)	(100.0)		4.50	0.80	
											70-90: fish (Centracanthidae, Sparidae)	(100.0)		4.50	0.80	
											90-120: fish (Sparidae, Centracanthidae)	(100.0)		4.50	0.80	
281	<i>Fistularia commersonii</i> *	Fistulariidae*	Rhodes Island, Greece	9/2004-3/2005	T		SL	17.5-87.5	N, W, CV	245	fish (<i>B. boops</i> , <i>S. smar</i>)	100.0 (100.0)	RA	4.50	0.80	Kalogirou <i>et al.</i> (2006)
282	<i>Fistularia commersonii</i> *	Fistulariidae*	Syria	8/2004-9/2005	BS,T				RA	460	fish (<i>S. aurita</i> , <i>B. Boops</i> , <i>E. encrasicolus</i>)		RA	4.50	0.80	Saad & Sabour (2010)
283	<i>Fistularia commersonii</i> *	Fistulariidae*	Rhodes Island, Greece		2001	T	SL	14.1-73.4	N	37	fish (<i>S. smar</i> , <i>M. surmuletus</i> , <i>Thalassoma pavo</i>)		RA	4.50	0.80	Corsini <i>et al.</i> (2002)
284	<i>Hippocampus guttulatus</i> *	Syngnathidae	Rhodes Island, Greece	3/2004	T	1	SL	7.8-22.6	F, N, VC	279	Amphipoda, Anomura, algae	(34.31, 24.31, 20.83)	D	3.10	0.46	Kitsos <i>et al.</i> (2006, 2008)
285	<i>Hippocampus hippocampus</i>	Syngnathidae	Rhodes Island, Greece	3/2004	T	1	SL	6.9-10.4	F, N, VC	19	Amphipoda, Mysidacea, Anomura	(43.75, 15.63, 12.50)	D	3.20	0.47	Kitsos <i>et al.</i> (2006, 2008)
286	<i>Syngnathus abaster</i> *	Syngnathidae	Kleisova lagoon, Messolonghi	11/2011	N		TL	6.2-12.1	F, N, W, IRI, VC	50	Amphipoda, Mysidacea	60.0, 32.0 (14.0, 84.0)	D	3.28	0.50	Ramfos <i>et al.</i> (2013)
287	<i>Syngnathus typhle</i>	Syngnathidae	Kleisova lagoon, Messolonghi	11/2011	N		TL	6.2-20.8	F, N, W, IRI, VC	31	Natantia, Mysidacea	77.0, 22.0 (81.0, 12.0)	D	3.51	0.55	Ramfos <i>et al.</i> (2013)
Tetraodontiformes*																
288	<i>Lagocephalus sceleratus</i> *	Tetraodontidae*	Antalya Bay, Turkey	12/2008-1/2010	CL	M	TL	12.5-65.0	W	656	Natantia (Penaetidae), Brachyura (Portunidae)	54.0, 17.0	RA	3.73	0.60	Aydin (2011)
289	<i>Lagocephalus sceleratus</i> *	Tetraodontidae*	Rhodes Island, Greece	2008-2009	BS	S	TL	5.3-63.1	F, N, VC	290	Gastropoda, Cephalopoda	(46.84, 28.57)	RA	3.86	0.54	Kalogirou (2013)
Zeiformes																
290	<i>Zeus faber</i>	Zeidae	Saros Bay, North Aegean Sea, Turkey	9/2006-9/2008	T	M	TL	12.2-52.8	F, N, W, IRI, VC		fish (<i>E. encrasicolus</i>)	99.67 (95.49)	BP	4.50	0.80	Ismen <i>et al.</i> (2013)
											F	10-19: fish (<i>E. encrasicolus</i>)		4.40	0.78	
											F	20-29: fish (<i>E. encrasicolus</i>)		4.50	0.80	
											F	30-39: fish (<i>E. encrasicolus</i>)		4.50	0.80	
											F	40-49: fish (<i>E. encrasicolus</i>)		4.50	0.80	
											F	50-59: fish (<i>E. encrasicolus</i>)		4.50	0.80	

References

- Abid, S., Ouannes-Ghorbel, A., Jarboui, O., Bouain, A., 2013. Diet composition and feeding habits of the wide-eyed flounder, *Bothus podas*, in the Gulf of Gabes (Tunisia). *Marine Biodiversity*, 43, 149-161.
- Anastasopoulou, A., Mytilineou, Ch., Lefkaditou, E., Kavadas, S., Bekas, P. *et al.*, 2012. The diet of *Conger conger* (L. 1758) in the deep waters of the Eastern Ionian Sea. p. 190. In: *10th Panhellenic Symposium of Oceanography & Fisheries, Athens, 7-11 May 2012*. HCMR, Athens.
- Anastasopoulou, A., Mytilineou, Ch., Lefkaditou, E., Kavadas, S., Bekas, P. *et al.*, 2013. The diet and feeding ecology of *Conger conger* (L. 1758) in the deep waters of the Eastern Ionian Sea. *Mediterranean Marine Science*, 14(2), 365-368.
- Artüz, L. 2004, Feeding accustomation of the ballan wrasse in the Sea of Marmara *Labrus bergylta* Ascanius 1767. *Hidrobiologica, Publications Scientifiques*, 2004/10, 17-19.
- Artüz, M.L., 2005a. The diet and food consumption of whiting *Merlangius merlangus* merlangus (Linne) 1758 in the Sea of Marmara. *Hidrobiologica, Publications Scientifiques*, 2005/1, 23-26.
- Artüz, M.L., 2005b. Age and growth of the ballan wrasse in the Sea of Marmara *Labrus bergylta* Ascanius 1767. *Hidrobiologica, Publications Scientifiques*, 2005/3, 23-26.
- Athineou, C., Dimitriadis, C., Karris, G., Sourbès L., Koutsoubas, D., 2015. Preliminary results of feeding habits of the allochthonous herbivore species *Siganus luridus* in the National Marine Park of Zakynthos. p. 45-48. In: *11th Panhellenic Symposium of Oceanography & Fisheries, Mytilene, 13-17 May 2015*. HCMR, Athens.
- Avsar, D., 2001. Age, growth, reproduction and feeding of the spurdog (*Squalus acanthias* Linnaeus, 1758) in the South-eastern Black Sea. *Estuarine Coastal and Shelf Science*, 52, 269-278.
- Aydin, M., 2011. Growth, Reproduction and Diet of Pufferfish (*Lagocephalus sceleratus* Gmelin, 1789) from Turkey's Mediterranean Sea Coast. *Turkish Journal of Fisheries and Aquatic Sciences*, 11, 569-576.
- Bacha, M., Amara, R., 2009. Spatial, temporal and ontogenetic variation in diet of anchovy (*Engraulis encrasicolus*) on the Algerian coast (SW Mediterranean). *Estuarine, Coastal and Shelf Science*, 85, 257-264.
- Bacha, M., Amara, R., Benmansour, N., Brylinski, J.-M., Moali, A. *et al.*, 2009. Age, growth and feeding of anchovy (*Engraulis encrasicolus* L.) in the Bay of Bénisaf (SW Mediterranean, west Algerian coast). In: *4e Rencontres de l'Ichthyologie en France, Paris, 24-27 March 2009*.
- Bacha, M., Moali, A., Benmansour, N.-E., Brylinski, J.-M., Mahé, K. *et al.*, 2010. Relationships between age, growth, diet and environmental parameters for anchovy (*Engraulis encrasicolus* L.) in the Bay of Bénisaf (SW Mediterranean, west Algerian coast). *Cybium*, 34, 47-57.
- Bariche, M., Alwan, N., El-Assi, H. Zurayk, R., 2009. Diet composition of the Lessepsian bluespotted cornetfish *Fistularia commersonii* in the eastern Mediterranean. *Journal of Applied Ichthyology*, 25, 460-465.
- Başçınar, N.S., Sağlam, H., 2009. Feeding Habits of Black Scorpionfish *Scorpaena porcus*, in the South-Eastern Black Sea. *Turkish Journal of Fisheries and Aquatic Sciences*, 9, 99-103.
- Battaglia, P., Esposito, V., Malara, D., Falautano, M., Castriota, L. *et al.*, 2014. Diet of the spothhead lanternfish *Diaphus metopoclampus* (Cocco, 1829) (Pisces: Myctophidae) in the central Mediterranean Sea. *Italian Journal of Zoology*, 81(4), 530-543.

- Bayhan, B., Sever, T.M., 2009. Food and feeding habits of the Atlantic Horse Mackerel, *Trachurus trachurus*, from the Aegean Sea. *Zoology of the Middle East*, 46, 47-54.
- Bayhan, B., Sever, T.M., Kara A. 2013. Diet composition of the Mediterranean horse mackerel, *Trachurus mediterraneus* (Steindachner, 1868) (Osteichthyes: Carangidae), from the Aegean Sea. *Belgian Journal of Zoology*, 143 (1), 15-22.
- Bayhan, B., Sever, T.M., Taskavak, E., 2008. Age, length-weight relationships and diet composition of scaldfish *Arnoglossus laterna* (Walbaum, 1792) (Pisces: Bothidae) in Izmir Bay (Aegean Sea). *Journal of Animal and Veterinary Advances*, 7, 924-929.
- Bayhan, B., Sever, T.M., Taskavak, E., 2009. Age and feeding habits of Atlantic spotted flounder *Citharus linguatula* (Linnaeus, 1758) (Pisces: Pleuronectiformes) from central Aegean Sea of Turkey. *North-Western Journal of Zoology*, 5(2), 330-337.
- Beltrano, A.M., Cannizzaro, L., Vitale, S., Milazzo, A., 2006. Preliminary study on the feeding habits of cleaver wrasse, *Xyrichtys novacula* (Pisces: Labridae) in the Strait of Sicily (Mediterranean Sea). *e-Journal of Ichthyology*, 2, 50-54.
- Bernal, A., Olivar, M.P, Fernandez de Puellas, M.L., 2013. Feeding patterns of *Lampanyctus pusillus* (Pisces: Myctophidae) throughout its ontogenetic development. *Marine Biology*, 160, 81-95.
- Bilecenoglu, M., 2009. Growth and feeding habits of the brown comber, *Serranus hepatus* (Linnaeus, 1758) in Izmir Bay, Aegean Sea. *Acta Adriatica*, 50(1), 105- 110.
- Boudaya, L., Neifar, L., Taktak, A., Ghorbel, M., Bouain, A., 2007. Diet of *Chelidonichthys obscurus* and *Chelidonichthys lastoviza* (Pisces: Triglidae) from the Gulf of Gabes (Tunisia). *Journal of Applied Ichthyology*, 23, 646-653.
- Cabiddu, S., Pedoni, C., Pesci, P., Porcu, C., 2010. Trophic habits of red scorpionfish *Scorpaena scrofa* (Osteichthyes, Scorpaeniformes) in the Central Western Mediterranean. *Rapport de la Commission internationale de la Mer Méditerranée*, 39, 464.
- Capapé, C., Crouzet, S., Clement, C., Vergne, Y., Guelorget, O., 2007. Diet of marbled electric ray *Torpedo marmorata* (Chondichthyes: Torpedinidae) off the Languedocian coast (Southern France, Northern Mediterranean). *Annales, Series Historia Naturalis*, 17(1), 17-22.
- Capapé, C., Hemida, F., Quignard, J.-P., Ben Amor, M.M., Reynaud, C., 2008. Biological observations on a rare deep-sea shark, *Dalatias licha* (Chondrichthyes: Dalatiidae), off the Maghreb coast (south-western Mediterranean). *Pan-American Journal of Aquatic Sciences*, 3(3), 355-360.
- Carpentieri, P., Colloca, F., Cardinale, M., 2005. Feeding habits of European hake (*Merluccius merluccius*) in the central Mediterranean Sea. *Fishery Bulletin*, 103, 411-416.
- Carrasson, M., Matallanas, J., 2002. Diets of deep-sea macrourid fishes in the western Mediterranean. *Marine Ecology Progress Series*, 234, 215-228.
- Castriota, L., Falautano, M., Finoia, M.G., Consoli, P., Peda, C. *et al.*, 2012. Trophic relationships among scorpaeniform fishes associated with gas platforms. *Helgoland Marine Research*, 66, 401-411.
- Castriota, L., Finoia, M.G., Andaloro, F., 2005. Trophic interactions between *Xyrichtys novacula* (Labridae) and juvenile *Pagrus pagrus* (Sparidae) in the Central Mediterranean Sea. *e-Journal of Ichthyology*, 2, 54-60.
- Castriota, L., Finoia, M.G., Campagnuolo, S., Romeo, T., Potoschi, A. *et al.*, 2008. Diet of *Tetrapturus belone* (Istiophoridae) in the central Mediterranean Sea. *Journal of the Marine Biological Association of the U.K.*, 88(1), 183-187.

- Castriota, L., Pipitone, C., Campagnuolo, S., Romanelli, M., Poroschi, A. *et al.*, 2007. Diet of *Coryphaena hippurus* (Coryphaenidae) associated with FADs in the Ionian and Southern Tyrrhenian Seas. *Cybium*, 31 (4), 435-441.
- Catalano, B., Dalù, M., Scacco, U., Vacchi, M., 2007. New biological data on *Raja brachyura* (Chondrichthyes, Rajidae) from around Asinara Island (NW Sardinia, Western Mediterranean), *Italian Journal of Zoology*, 74(1), 55-61.
- Celona, A., De Maddalena, A., Romeo, T., 2005. Bluntnose sixgill shark, *Hexanchus griseus* (Bonnaterre, 1788), in the eastern north Sicilian waters. *Bollettino del Museo civico di Storia Naturale di Venezia*, 56, 137-151.
- Ceyhan, T., Akyol, O., Sever, T.M., Kara, A., 2012. Diet composition of adult twaite shad (*Alosa fallax*) in the Aegean Sea (Izmir Bay, Turkey). *Journal of the Marine Biological Association of the U.K.*, 92(3), 601-604.
- Chater, I., Romdhani, A., Ktari, M.H., Mahe, K., 2012. Diet composition and feeding intensity of brown meagre *Sciaena umbra* Linnaeus, 1758 (Teleostei, Sciaenidae) in the Gulf of Tunis. In: *14th European Congress of Ichthyology, Liège, 3-8 July 2012*.
- Chessa, L.A., Lanera, P., Pais, A., Plastina, N., Scardi, M. *et al.*, 2005. Observations on feeding habitus of *Lithognathus mormyrus* (Linnaeus, 1758) in the Calich lagoon (North Western Sardinia). *Biologia Marina Mediterranea*, 12(1), 492-495.
- Chessa, L.A., Pais, A., Scardi, M., Serra, S., Atzori, G., 2004. Observations on the feeding behaviour of *Diplodus annularis* (Linnaeus, 1758) in the Calich lagoon (North Western Sardinia). *Biologia Marina Mediterranea*, 11(2), 573-576.
- Corsini, M., Kondilatos, G., Economidis, P.S., 2002. Lessepsian migrant *Fistularia commersonii* from the Rhodes marine area. *Journal of Fish Biology*, 61, 1061-1062.
- Cuoco, C., Mancusi, C., Serena, F., 2005. Study of feeding habits of *Raja asterias* Delaroche, 1809 (Chondrichthyes, Rajidae). *Biologia Marina Mediterranea*, 12 (1), 504-508.
- Darnaude, A.M., 2005. Fish ecology and terrestrial carbon use in coastal areas: implications for marine fish production. *Journal of Animal Ecology*, 74, 864-876.
- Darnaude, A.M., Salen-Picard, C., Harmelin-Vivien, M.L., 2004a. Depth variation in terrestrial particulate organic matter exploitation by marine coastal benthic communities off the Rhone River delta (NW Mediterranean). *Marine Ecology Progress Series*, 275, 47-57.
- Darnaude, A.M., Salen-Picard, C., Polunin, N.V.C., Harmelin-Vivien, M.L., 2004b. Trophodynamic linkage between river runoff and coastal fishery yield elucidated by stable isotope data in the Gulf of Lions (NW Mediterranean). *Oecologia*, 138, 325-332.
- de Juan, S., Cartes, J.E., Demestre, M., 2007. Effects of commercial trawling activities in the diet of the flat fish *Citharus linguatula* (Osteichthyes: Pleuronectiformes) and the starfish *Astropecten irregularis* (Echinodermata: Asteroidea). *Journal of Experimental Marine Biology and Ecology*, 349, 152-169.
- Demirhan, S.A., Can, M.F., 2009. Age, growth and food composition of *Scorpaena porcus* (Linnaeus, 1758) in the southeastern Black Sea. *Journal of Applied Ichthyology*, 25, 215-218.
- Derbal, F., Kara, M.H., 2004. Alimentation de la badeche *Epinephelus costae* (Steindachner, 1878) des cotes de l'est Algerien. *Rapport de la Commission internationale de la Mer Méditerranée*, 37, 344.
- Derbal, F., Kara, M.H., 2007. Régime alimentaire de la badèche *Epinephelus costae* (Steindachner, 1875) (Serranidae) des côtes de l'Est Algérien. p. 67-69. In: *Second International Symposium on the Mediterranean Groupers, Nice, 13-17 May 2007*. Nice University, Nice.

- Dhieb, K., Ghorbel, M., Bouain, A., 2001. Regime alimentaire de *Pomatomus saltatrix* (Teleostei, Pomatomidae) dans le Golfe de Gabes, Tunisie. *Rapport de la Commission internationale de la Mer Méditerranée*, 36, 259.
- Dulcic, J., Lipej, L., Glamuzina, B., Bartulovic, V., 2006. Diet of *Spondyliosoma cantharus* and *Diplodus puntazzo* (Sparidae) in the Eastern Central Adriatic. *Cybium*, 30(2), 115-122.
- El Bakali, M., Talbaoui, M., Bendriss, A., 2010. Régime alimentaire du Rouget de roche (*Mullus surmuletus* L.) (Téléostéen, Mullidae) de la côte nord-ouest méditerranéenne du Maroc (région de M'diq). *Bulletin de l'Institut Scientifique, Rabat, section Sciences de la Vie*, 32(2), 87-93.
- El Kamel-Moutalibi, O., Mnasri, N., Boumaiza, M., Reynaud, C., Capape, C., 2013. Diet of common torpedo *Torpedo torpedo* (Chondrichthyes: Torpedinidae) from the Lagoon of Bizerte (northeastern Tunisia, central Mediterranean). *Cahier de Biologie Marine*, 54, 209-220.
- El-mor, M., Ahamed, A.I., 2008. Feeding habits of the common sole, *Solea vulgaris* (Quensel, 1806), from Mediterranean Sea, Port Said, Egypt. *Egyptian Journal of Aquatic Biology and Fisheries*, 12 (2), 51-61.
- Enajjar, S., Bradai, M.N., Bouain, A., 2007. Feeding habits of *Rhinobatos rhinobatos* in the Gulf of Gabes. *Rapport de la Commission internationale de la Mer Méditerranée*, 38, 468.
- Engin, S., Seyhan, K., 2010. Age, growth, reproduction and diet of the Flatsnout Goby, *Neogobius platyrostris* (Pallas, 1814), on the south-eastern Black Sea Coast of Turkey. *Zoology in the Middle East*, 50, 59-66.
- Esposito, V., Andaloro, F., Bianca, D., Natalotto, A., Romeo, T. *et al.*, 2014. Diet and prey selectivity of the red mullet, *Mullus barbatus* (Pisces: Mullidae), from the southern Tyrrhenian Sea: the role of the surf zone as a feeding ground. *Marine Biology Research*, 10 (2), 167-178.
- Esposito, V., Battaglia, P., Castriota, L., Finoia, M.G., Scotti, G. *et al.*, 2009. Diet of Atlantic lizardfish, *Synodus saurus* (Linnaeus, 1758) (Pisces: Synodontidae) in the central Mediterranean Sea. *Scientia Marina*, 73(2), 369-376.
- Falautano, M., Castriota, L., Finoia, M.G., Andaloro, F., 2007. Feeding ecology of little tunny *Euthynnus alletteratus* in the central Mediterranean Sea. *Journal of the Marine Biological Association of the U.K.*, 87, 999-1005.
- Fanelli, E., Cartes, J.E., 2007. Trophic relationships of two macrourids from the Balearic basin (Western Mediterranean): temporal changes in diets and in trophic levels. *Rapport de la Commission internationale de la Mer Méditerranée*, 38, 472.
- Fanelli, E., Badalamenti, F., D'Anna, G., Pipitone, C., 2009. Diet and trophic level of scaldfish *Arnoglossus laterna* in the southern Tyrrhenian Sea (western Mediterranean): contrasting trawled versus untrawled areas. *Journal of the Marine Biological Association of the U.K.*, 89, 817-828.
- Filiz, H., 2009. Diet composition of smooth-hound, *Mustelus mustelus* (Linnaeus, 1758), in Aegean Sea, Turkey. *Belgian Journal of Zoology*, 139 (1), 81-84.
- Filiz, H., Taşkavak, E., 2006. Food of lesser spotted dogfish, *Scyliorhinus canicula* (Linnaeus, 1758), in Foca (the Northeast Aegean Sea, Turkey) in autumn 2002. p. 60-68. In: *International Workshop on Mediterranean Cartilaginous Fish with Emphasis on the Southern and Eastern Mediterranean, Istanbul, 14-16 October 2006*. Turkish Marine Research Foundation, Istanbul.
- Fletcher, N., Batjakas, I.E., Pierce, G.J., 2013. Diet of the Atlantic bonito *Sarda sarda* (Bloch, 1793) in the Northeast Aegean Sea. *Journal of Applied Ichthyology*, 29, 1030-1035.

- Follesa, M.C., Cannas, R., Cabiddu, S., Alessandro, C.A.U., Mulas, A. *et al.*, 2012. Preliminary observations of the reproductive biology and diet for the Norwegian skate *Dipturus nidarosiensis* (Rajidae) from the Central Western Mediterranean Sea. *Cybium*, 36 (3), 473-477.
- Follesa, M.C., Mulas, A., Cabiddu, S., Porcu, C., Deiana, A.M. *et al.*, 2010. Diet and feeding habits of two skate species, *Raja brachyura* and *Raja miraletus* (Chondrichthyes, Rajidae) in Sardinian waters (central-western Mediterranean). *Italian Journal of Zoology*, 77 (1), 53-60.
- Gračan, R., Mladineo, I., Lazar, B., 2014. Insight into the diet composition and gastrointestinal parasite community of the common smooth-hound, *Mustelus mustelus* (Carcharhiniformes: Triakidae), in the northern Adriatic Sea. *Natura Croatica*, 23(1), 35-44.
- Gračan, R., Zavodnik, D., Krstinić, P., Dragičević, B., Lacković, G. *et al.*, 2013. Feeding ecology and trophic segregation of two sympatric shark species, *Mustelus punctulatus* and *Squalus acanthias*, in the North–Central Adriatic Sea. p. 6. In: *International Workshop on Conservation Biology, Koper, 14 May 2013*.
- Gravino, F., Dimech, M., Schembri, P.J., 2010. Feeding habits of the small-spotted catshark *Scyliorhinus canicula* (L., 1758) in the central Mediterranean. *Rapport de la Commission internationale de la Mer Méditerranée*, 39, 538.
- Gurlek, M., Erguden, S., Yaglioglu, D., Turan, F., Demirhan, S. *et al.*, 2010. Feeding habits of indo-pacific species *Nemipterus randalli* Russel, 1986 (Nemipteridae) in Iskenderun bay, Eastern Mediterranean Sea. *Rapport de la Commission internationale de la Mer Méditerranée*, 39, 539.
- Hajji, F., Ouannes-Ghorbel, A., Ghorbel, M., Jarboui, O., 2013. Feeding habits of the black goby, *Gobius niger* (Linnaeus, 1758), in the Gulf of Gabes (Southern Tunisia). *Cahier de Biologie Marine*, 54, 133-141.
- Harchouche, K., Maurin, C., Zerouali–Khodja, F., 2009. Régime alimentaire de *Spicara maena* (Centracanthidae) des eaux Algériennes. *Bulletin de la Société Zoologique de France*, 134 (1-2), 123-141.
- Ibrahim, A., Lahlah, M., Kassab, M.Y., Ghanem, W., Ogaily, S., 2010. *Signatus javus*, a new record from the Syrian waters, with reference to growth and feeding of two Lessepsian fish. *Rapport de la Commission internationale de la Mer Méditerranée*, 39, 544.
- Ismen, A., 2003. Age, growth, reproduction and food of common stingray (*Dasyatis pastinaca* L., 1758) in Iskenderun Bay, the eastern Mediterranean. *Fisheries Research*, 60, 169-176.
- Ismen, A., Arslan, M., Yigin, C.C., Bozbay, N.A., 2013. Age, growth, reproduction and feeding of John Dory, *Zeus faber* (Pisces: Zeidae), in the Saros Bay (North Aegean Sea). *Journal of Applied Ichthyology*, 29, 125-131.
- Ismen, A., Yigin, C., Ismen, P., 2007. Age, growth, reproductive biology and feed of the common guitarfish (*Rhinobatos rhinobatos* Linnaeus, 1758) in Iskenderun Bay, the eastern Mediterranean Sea. *Fisheries Research*, 84, 263-269.
- Jaramillo Londoño, A.M., Cantos, G., Porrás Castelló, R., Durà, V.B., 2011. *Composición de la dieta y estrategia alimentaria de cinco especies de peces bentónicos de la costa de Cullera (España)*. *Mediterránea*. Serie de Estudios Biológicos. Época II, n. especial, 41 pp.
- Jardas, I., Santic, M., Pallaoro, A., 2004. Diet composition of the eagle ray, *Myliobatis aquila* (Chondrichthyes: Myliobatidae), in the Eastern Adriatic Sea. *Cybium*, 28(4), 372-374.

- Kabasakal, H., 2001. Description of the feeding morphology and the food habits of four sympatric labrids (Perciformes, Labridae) from South-Eastern Aegean Sea, Turkey. *Netherlands Journal of Zoology*, 51(4), 439-455.
- Kabasakal, H., 2004. Preliminary observations on the reproductive biology and diet of the Bluntnose sixgill shark, *Hexacanthus griseus* (Bonnaterre, 1788) (Chondrichthyes: Hexanchidae), in Turkish Seas. *Acta Adriatica*, 45(2), 187-196.
- Kadri, H., Saïdi, B., Marouani, S., Bradai, M.N., Bouaïn, A., 2013. Food habits of the rough ray *Raja radula* (Chondrichthyes: Rajidae) from the Gulf of Gabès (central Mediterranean Sea). *Italian Journal of Zoology*, 80(1), 52-59.
- Kallianiotis, A., Torre, M., Argyri, A., 2005. Age, growth, mortality, reproduction, and feeding habits of the striped seabream, *Lithognathus mormyrus* (Pisces: Sparidae), in the coastal waters of the Thracian Sea, Greece. *Scientia Marina*, 69 (3), 391-404.
- Kalogirou, S., 2013. Ecological characteristics of the invasive pufferfish *Lagocephalus sceleratus* (Gmelin, 1789) in Rhodes, Eastern Mediterranean Sea. A case study. *Mediterranean Marine Science*, 14(2), 251-260.
- Kalogirou, S., Corsini Foka, M., Kondylatos, G., Wennhage, H., 2006. First results on the diet of the exotic fish *Fistularia commersonii* in SE Aegean Sea. p. 203. In: *8th Panhellenic Symposium of Oceanography and Fisheries, Thessaloniki, 6-8 June 2006*. HCMR, Athens.
- Karachle, P.K., 2008. *Feeding ecology of the most important fish stocks in North Aegean Sea, Greece*. PhD Thesis, Aristotle University of Thessaloniki, 277 pp.
- Karachle, P.K., 2014. Feeding habits for eleven sparids from the N-NW Aegean Sea. p. 218-220. In: *New Fisheries-related data from the Mediterranean Sea (April, 2014)*. Stergiou, K.I., Bobori, D.C., Ekmekçi, F.G., Gökoğlu, M., Karachle, P.K. *et al.* *Mediterranean Marine Science*, 15 (1), 213-224.
- Karachle, P.K., Stergiou, K.I., 2008. The effect of season and sex on trophic levels of marine fishes. *Journal of Fish Biology*, 72, 1463-1487.
- Karachle, P.K., Stergiou, K.I., 2010a. Food and feeding habits for eight fishes in the N Aegean Sea, Greece. p. 7-10. In: *14th Hellenic Conference of Ichthyologists, Piraeus, 6-9 May 2010*.
- Karachle, P.K., Stergiou, K.I., 2010b. Food and feeding habits of nine elasmobranch species in the N Aegean Sea. *Rapport de la Commission internationale de la Mer Méditerranée*, 39, 553.
- Karachle, P.K., Stergiou, K.I., 2011. Feeding and ecomorphology for seven flatfishes in the N-NW Aegean Sea (Greece). *African Journal of Marine Science*, 33(1), 67-78.
- Karachle, P.K., Stergiou, K.I., 2012a. Ecomorphology and feeding habits for four pelagic fishes from the North Aegean Sea. p. 76. In: *12th International Congress on the Zoogeography and Ecology of Greece and adjacent Regions, Athens, 18-22 June, 2012*.
- Karachle, P.K., Stergiou, K.I., 2012b. Diet and feeding habits for five fishes from the North Aegean Sea. p. 171. In: *10th Panhellenic Symposium of Oceanography & Fisheries, Athens, 7-11 May 2012*. HCMR, Athens.
- Karachle, P.K., Stergiou, K.I., 2014a. Feeding and ecomorphology of three clupeoids in the North Aegean Sea. *Mediterranean Marine Science*, 15(1), 9-26.
- Karachle, P.K., Stergiou, K.I., 2014b. Diet and feeding habits of *Spicara maena* and *S. smaris* in the N Aegean Sea. *Acta Adriatica*, 55(1), 75-84.
- Karachle, P.K., Triantaphyllidis, C., Stergiou, K.I., 2004. *Fistularia commersonii* Rüppell, 1838: A Lessepsian sprinter. *Acta Ichthyologica et Piscatoria*, 34 (1), 103-108.

- Kitsos, M.-S., Tzomos, Th., Anagnostopoulou, L., Koukouras, A., 2006. Diet composition of the seahorses, *Hippocampus guttulatus* Cuvier, 1829 and *Hippocampus hippocampus* (L., 1758) (Teleostei, Syngnathidae) in the Aegean Sea. p. 88. In: *8th Panhellenic Symposium of Oceanography and Fisheries, Thessaloniki, 6-8 June 2006*. HCMR, Athens.
- Kitsos, M.-S., Tzomos, Th., Anagnostopoulou, L., Koukouras, A., 2008. Diet composition of the seahorses, *Hippocampus guttulatus* Cuvier, 1829 and *Hippocampus hippocampus* (L., 1758) (Teleostei, Syngnathidae) in the Aegean Sea. *Journal of Fish Biology*, 72, 1259-1267.
- Kousteni, V., 2015. *Population characteristics and heavy metal concentrations in benthic sharks from the eastern Mediterranean Sea*. PhD Thesis, National and Kapodistrian University of Athens, 524 pp.
- Kousteni, V., Karachle, P.K., Megalofonou, P., 2015. Comparative feeding ecology of two demersal sharks from the self and slope off the Aegean Sea in the eastern Mediterranean. In: *Annual Symposium of the Fisheries Society of the British Isles, Plymouth, 27-31 July 2015*.
- Kovacic, M., 2007. Diet of *Gobius vittatus* (Gobiidae) in the Northern Adriatic Sea. *Vie et Millieu*, 57 (1/2), 29-35.
- La Mesa, M., Borne, D., Tirelli, V., Di Poi, E., Legovini, S. *et al.*, 2008. Feeding ecology of the transparent goby *Aphia minuta* (Pisces, Gobiidae) in the northwestern Adriatic Sea. *Scientia Marina*, 72(1), 99-108.
- Lakkis, S., Sabour, W., 2010. Marine biodiversity and fishery resources in Syro-lebanese coastal waters (Levantine basin). *Rapport de la Commission internationale de la Mer Méditerranée*, 39, 564.
- Leonardos, I., 2008. The feeding ecology of *Aphanius fasciatus* (Valenciennes, 1821) in the lagoonal system of Messolongi (western Greece). *Scientia Marina*, 72(2), 393-401.
- Linde, M., Grau, A.M., Riera, F., Massuti-Pascual, E., 2004. Analysis of trophic ontogeny in *Epinephelus marginatus* (Serranidae). *Cybium*, 28(1), 27-35.
- Lipej, L., Mavric, B., Paliska, D., Capapé, C., 2013. Feeding habits of the pelagic stingray *Pteroplatytrygon violacea* (Chondrichthyes: Dasyatidae) in the Adriatic Sea. *Journal of the Marine Biological Association of the U.K.*, 93(2), 285-290.
- Lomiri, S., Scacco, U., Mostarda, E., Andaloro, F., 2008. Size-related and temporal variation in the diet of the round sardinella, *Sardinella aurita* (Valenciennes, 1847), in the central Mediterranean Sea. *Journal of Applied Ichthyology*, 24, 539-545.
- Madkour, F.F., 2012. Feeding ecology of the round sardinella, *Sardinella aurita* (Family: Clupeidae) in the Egyptian Mediterranean waters. *International Journal of Environmental Science and Engineering*, 2, 83-92.
- Matić-Skoko, S., Ferri, J., Tutman, P., Skaramuca, D., Đikić, D. *et al.*, 2012. The age, growth and feeding habits of the European conger eel, *Conger conger* (L.) in the Adriatic Sea. *Marine Biology Research*, 8(10), 1012-1018.
- Matic-Skoko, S., Tutman, P., Marcelja, E., Skaramuca, D., Đikić, D. *et al.*, 2010. Feeding habits and trophic status of Mediterranean moray eel, *Muraena helena* L. 1758 in the Adriatic Sea – preliminary approach. *Rapport de la Commission internationale de la Mer Méditerranée*, 39, 583.
- Mavric, B., Jenko, R., Makovec, T., Lipej, L., 2004. On the occurrence of the pelagic stringray, *Dasyatis violacea* (Bonaparte, 1832), in the Gulf of Trieste (Norther Adriatic). *Annales, Series Historia Naturalis*, 14(2), 181-186.
- Megalofonou, P., Chatzistryou, A., 2006. Sexual maturity and feeding of the gulper shark, *Centrophorus granulosus*, from the eastern Mediterranean Sea. *Cybium*, 30(4) suppl., 67-74.

- Morte, M.S., Redon, M.J., Sanz-Brau, A., 2001. Feeding habits of *Trisopterus minutus capelanus* (Gadidae) off the Eastern coast of Spain (Western Mediterranean). *P.S.Z.N.: Marine Ecology*, 22(3), 215-229.
- Morte, M.S., Redon, M.J., Sanz-Brau, A., 2002. Diet of *Phycis blennoides* (Gadidae) in relation to fish size and season in the Western Mediterranean (Spain). *P.S.Z.N.: Marine Ecology*, 23(2), 141-155.
- Mulas, A., Bellodi, A., Cannas, R., Cau, A., Cuccu, D. *et al.*, 2015. Diet and feeding behaviour of longnosed skate *Dipturus oxyrinchus*. *Journal of Fish Biology*, 86, 121-138.
- Mulas, A., Bellodi, A., Pendugiu, A.A., Cabiddu, S., 2010. Preliminary data on the trophic interactions between *Scyliorhinus canicula* (Scyliorhinidae) and *Raja clavata* (Rajidae) in the central-western Mediterranean. *Rapport de la Commission internationale de la Mer Méditerranée*, 39, 596.
- Navarro, J., Coll, M., Preminger, M., Palomera, I., 2013. Feeding ecology and trophic position of a Mediterranean endemic ray: consistency between sexes, maturity stages and seasons. *Environmental Biology of Fish*, 96, 1315-1328.
- Nikolioudakis, N., Isari, S., Pitta, P., Somarakis, S., 2012. Diet of sardine *Sardina pilchardus*: an 'end-to-end' field study. *Marine Ecology Progress Series*, 453, 173-188.
- Nikolioudakis, N., Isari, S., Somarakis, S., 2014. Trophodynamics of anchovy in a non-upwelling system: direct comparison with sardine. *Marine Ecology Progress Series*, 500, 215-229.
- Nouacer, S., Derbal, F., Kara, M.H., 2004. Régime alimentaire du sparailon *Diplodus annularis* (L., 1758) du Golfe d'Annaba (Algerie Est). *Rapport de la Commission internationale de la Mer Méditerranée*, 37, 411.
- Osman, A.M., Mahmoud, H.H., 2009. Feeding Biology of *Diplodus sargus* and *Diplodus vulgaris* (Teleostei, Sparidae) in Egyptian Mediterranean Waters. *World Journal of Fish and Marine Sciences*, 1(4), 290-296.
- Osman, A.G.M., Farrag, M.M.S., Akel, E.S.H.Kh., Moustafa, M.A., 2013. Feeding behavior of lessepsian fish *Etrumeus teres* (Dekay, 1842) from the Mediterranean waters, Egypt. *Egyptian Journal of Aquatic Research*, 39, 275-282.
- Ouannes-Ghorbel, A., Bouain, A., 2004. Régime alimentaire de *Labrus viridis* (Pisces, Labridae) des côtes sud de la Tunisie. *Rapport de la Commission internationale de la Mer Méditerranée*, 37, 416.
- Ouannes-Ghorbel, A., Bouain, A., 2006. The diet of the peacock wrasse, *Symphodus (Crenilabrus) tinca* (Labridae), in the southern coast of Tunisia. *Acta Adriatica*, 47 (2), 175-182.
- Ouannes-Ghorbel, A., Bouain, A., Jarboui, O., Mrabet, R., 2003. Régime alimentaire de *Symphodus (Crenilabrus) ocellatus* des côtes sud de la Tunisie. *Bulletin de l'Institut National des Sciences et Technologies de la Mer de Salammbô*, 30, 33-37.
- Ozen, O., Altin, A., Ayyildiz, H., 2010. The diet of *Clinitrachus argentatus* (Blennioidei: Clinidae) in the northern Aegean and Marmara Seas. *Journal of Fish Biology*, 76, 1516-1519.
- Pais, A., Merella, P., Follesa, M.C., Garippa, G., Golani, D., 2008. New data on *Gaidropsarus granti* (Regan, 1903) (Gadiformes: Lotidae) from the Mediterranean Sea, with emphasis on its parasites. *Scientia Marina*, 72(3), 461-468.
- Pallaoro, A., Santic, M., Jardas, I., 2003. Feeding habits of the saddled bream, *Oblada melanura* (Sparidae), in the Adriatic Sea. *Cybium*, 27(4), 261-268.

- Peristeraki, P., Tserpes, G., 2001. Feeding habits of swordfish (*Xiphias gladius* L., 1758) in the S Aegean Sea. p. 25-28. In: *10th Hellenic Conference of Ichthyologists, Chania, 18-20 October 2001*.
- Philips, A.E., 2012. Feeding behavior of the European hake *Merluccius merluccius* Linnaeus, 1758 (Family: Gadidae) from Egyptian Mediterranean waters off Alexandria. *Egyptian Journal of Aquatic Research*, 38, 39-44.
- Pipitone, C., Campagnuolo, S., Andaloro, F., Potoschi, A., 2000. Diet of pilotfish, *Naucrates ductor* (Carangidae), associated with fads in Sicilian waters (Central Mediterranean). *Cybium*, 24(2), 415-418.
- Pori, A., Batzakas, J., 2015. Feeding habits of *Squalus acantias* from the North Aegean Sea. p. 41-44. In: *11th Panhellenic Symposium of Oceanography & Fisheries, Mytilene, 13-17 May 2015*. HCMR, Athens.
- Primikiri, A., Ramfos, A., Cladas, I., Koutsikopoulos, C., 2005. Stomach content analysis of the white seabream (*Diplodus sargus*) in Antinioti Lagoon (Corfu-western Greece). p. 19-22. In: *12th Hellenic Conference of Ichthyologists, Drama, 13-16 October 2005*.
- Quetglas, A., Renones, O., Goñi, R., 2001. Trophic interactions among grouper (*Epinephelus marginatus*), octopus (*Octopus vulgaris*) and red lobster (*Palinurus elephas*) in the Western Mediterranean. *Rapport de la Commission internationale de la Mer Méditerranée*, 36, 310.
- Ramfos, A., Kontochristos, N., Katselis, G., 2013. Feeding of *Syngnathus abaster* and *Syngnathus typhle* in Kleisova lagoon (Lagoon complex of Messolonghi-Aitoliko). p. 295-298. In: *15th Hellenic Conference of Ichthyologists, Thessaloniki, 10-13 October 2013*.
- Relini, G., Relini, M., Torchia, G., De Angelis, G., 2002. Trophic relationships between fishes and an artificial reef. *ICES Journal of Marine Science*, 59, S36–S42.
- Rizkalla, S.I., Philips, A.E., 2008. Feeding habits of the Atlantic stargazer fish *Uranoscopus scaber* Linnaeus, 1758 (Family: Uranoscopidae) in Egyptian Mediterranean waters. *Egyptian Journal of Aquatic Biology and Fisheries*, 12 (1), 1-11.
- Romanelli, M., Colasante, A., Scacco, U., Consalvo, I., Finioia, M.G. *et al.*, 2007. Commercial catches, reproduction and feeding habits of *Raja asterias* (Chondrichthyes: Rajidae) in a coastal area of the Tyrrhenian Sea (Italy, northern Mediterranean). *Acta Adriatica*, 48(1), 57-71.
- Rosca, I., Novac, A., Surugiu, V., 2010. Feeding selectivity of some benthic fish from the rocky bottom of the Romanian Black Sea coast (Agigea area). *Rapport de la Commission internationale de la Mer Méditerranée*, 39, 648.
- Saad, A.A., Sabour, W.G., 2010. Impact of invasive species on the biodiversity and fish stock; case study: *Fistularia commersonii* Ruppel, in the Syrian coast. *Rapport de la Commission internationale de la Mer Méditerranée*, 39, 652.
- Sabour, W., Lakkis, S., 2007. Diet and feeding habits of *Siganus rivulatus* and *S. luridus* two Red Sea migrants in the Syrian coastal waters (Eastern Mediterranean). *Rapport de la Commission internationale de la Mer Méditerranée*, 38, 584.
- Saidi, B., Bradai, M.N., Bouain, A., 2009b. Reproductive biology and diet of *Mustelus punctulatus* (Risso, 1826) (Chondrichthyes: Triakidae) from the Gulf of Gabès, central Mediterranean Sea. *Scientia Marina*, 73 (2), 249-258.
- Saidi, B., Enajjar, S., Bradai, M.N., Bouain, A. 2009a. Diet composition of smooth-hound shark, *Mustelus mustelus* (Linnaeus, 1758), in the Gulf of Gabes, southern Tunisia. *Journal of Applied Ichthyology*, 25 (Suppl.1), 113-118.
- Sallami, B., Ben Salem, M., Reynaud, C., Capapé, C., 2014. Diet of Mediterranean moray, *Muraena helena* (Actinopterygii: Anguilliformes: Muraenidae), from the north-

- eastern Tunisian coast (central Mediterranean). *Acta Ichthyologica et Piscatoria*, 44 (4), 273-283.
- Sallami, B., Ben Salem, M., Reynaud, C., Capapé, C., 2015a. Diet of European conger eel *Conger conger* (Osteichthyes: Congridae), from the northeastern coast of Tunisia (central Mediterranean). *Cahier de Biologie Marine*, 56, 253-262.
- Sallami, B., Ben Salem, M., Rafrafi-Nouira, S., El Kamel-Moutalibi, O., Reynaud, C. *et al.*, 2015b. Observations on the thinlip conger *Gnathophis mystax* (Osteichthyes: Congridae), from the Tunisian coast (central Mediterranean). *Annales, Series Historia Naturalis*, 25 (2), 131-140.
- Salman, A., 2004. The role of cephalopods in the diet of swordfish (*Xiphias gladius* Linnaeus, 1758) in the Aegean Sea. *Bulletin of Marine Science*, 74(1), 21-29.
- Šantić, M., Jardas, I., Pallaoro, A., 2005. Feeding habits of horse mackerel, *Trachurus trachurus* (Linnaeus, 1758), from the central Adriatic Sea. *Journal of Applied Ichthyology*, 21, 125-130.
- Šantić, M., Rađa, B., Paladin, A., Kovačević, A., 2011. Biometric properties and diet of common pandora, *Pagellus erythrinus* (Osteichthyes: Sparidae), from the Eastern Adriatic Sea. *Archives of Biological Sciences, Belgrade*, 63 (1), 217-224.
- Šantić, M., Rađa, B., Pallaoro, A., 2012. Diet and feeding strategy of thornback ray *Raja clavata*. *Journal of Fish Biology*, 81, 1070-1084.
- Šantić, M., Rađa, B., Pallaoro, A., 2013. Feeding habits of brown ray (*Raja miraletus* Linnaeus, 1758) from the eastern central Adriatic Sea. *Marine Biology Research*, 9, 316-323.
- Serena, F., Barone, M., Mancusi, C., Abella, A.J., 2005. *Reproductive biology, growth and feeding habits of Raja asterias Delaroché, 1809, from the north Tyrrhenian and south Ligurian Sea (Italy), with some notes on trends in landings*. Theme Session on Elasmobranch Fisheries Science CM2005/N:12. 2005 ICES Annual Science Conference 20–24 September 2005, 18 pp.
- Serena, F., Cecchi, E., Mancusi, C., Pajetta, R., 2006. Contribution to the knowledge of the biology of *Etmopterus spinax* (Linnaeus 1758) (Chondrichthyes, Etmopteridae). p. 388-394. In: *Deep Sea 2003: Conference on the Governance and Management of Deep-sea Fisheries. Part 2: Conference poster papers and workshop papers. Queenstown, New-Zealand, 1-5 December 2003*. Dunedin, New-Zealand, FAO Fisheries Proceedings 3/2.
- Sever, T.M., Bayhan, B., Bilge, G., Taskavak, E., 2009. Diet composition of *Belone belone* (Linnaeus, 1761) (Pisces: Belonidae) in the Aegean Sea. *Journal of Fish Biology*, 25, 702-706.
- Sever, T.M., Bayhan, B., Leblebici, S., 2010. Feeding habits of *Cepola macrophthalma* (Pisces: Cepolidae) in Izmir Bay, Aegean Sea. *Rapport de la Commission internationale de la Mer Méditerranée*, 39, 598.
- Sever, T.M., Bayhan, B., Taskavak, E., 2005. A preliminary study on the feeding regime of European pilchard (*Sardina pilchardus* Walbaum, 1792) in Izmir Bay, Turkey, Eastern Aegean Sea. *NAGA*, 28(3 & 4), 41-48.
- Sever, T.M., Filiz, H., Bayhan, B., Taskavak, E., Bilge, G., 2008. Food habits of the hollowsnout grenadier, *Caelorinchus caelorhincus* (Risso, 1810), in the Aegean Sea, Turkey. *Belgian Journal of Zoology*, 138 (1), 81-84.
- Sinopoli, M., Fanelli, E., D'Anna, G., Badalamenti, F., Pipitone, C., 2012. Assessing the effects of a trawling ban on diet and trophic level of hake, *Merluccius merluccius*, in the southern Tyrrhenian Sea. *Scientia Marina*, 76(4), 677-690.

- Sley, A., Jarboui, O., Bouain, A., 2007. Données préliminaires sur le régime alimentaire de *Caranx rhonchus* (Osteichthyens, Carangidae) dans le Golfe de Gabés (Tunisie). *Rapport de la Commission internationale de la Mer Méditerranée*, 38, 601.
- Sley, A., Jarboui, O., Ghorbel, M., Bouain, A., 2008. Diet composition and food habits of *Caranx rhonchus* (Carangidae) from the Gulf of Gabes (central Mediterranean). *Journal of the Marine Biological Association of the U.K.*, 88(4), 831-836.
- Sley, A., Jarboui, O., Ghorbel, M., Bouain, A., 2009. Food and feeding habits of *Caranx crysos* from the Gulf of Gabes (Tunisia). *Journal of the Marine Biological Association of the U.K.*, 89(7), 1375-1380.
- Stagioni, M., Montanini, S., Vallisneri, M., 2012. Feeding of tub gurnard *Chelidonichthys lucerna* (Scorpaeniformes: Triglidae) in the north-east Mediterranean. *Journal of the Marine Biological Association of the U.K.*, 92(3), 605-612.
- Stagioni, M., Montanini, S., Vallisneri, M., 2013. Feeding habits of anglerfish, *Lophius budegassa* (Spinola, 1807) in the Adriatic Sea, north-eastern Mediterranean. *Journal of Applied Ichthyology*, 29, 374-380.
- Stergiou, K.I., Karpouzi, V.S., 2002. Feeding habits and trophic levels of Mediterranean fish. *Reviews in Fish Biology and Fisheries*, 11, 217-254.
- Tsikliras, A.C., Torre, M., Stergiou, K.I., 2005. Feeding habits and trophic level of round sardinella (*Sardinella aurita*) in the northeastern Mediterranean (Aegean Sea, Greece). *Journal of Biological Research*, 3, 67-75.
- Türker Çakir, D., Torcu Koç, H., Erdoğan, Z., 2006. Some biological aspects of the lesser spotted dogfish *Scyliorhinus canicula* (Linnaeus, 1758) in Edremit Bay (the Northern Aegean Sea) Turkey. p. 17-27. In: *International Workshop on Mediterranean Cartilaginous Fish with Emphasis on the Southern and Eastern Mediterranean, Istanbul, 14-16 October 2006*. Turkish Marine Research Foundation, Istanbul.
- Uçkun, D., Sever, T.M., Toğulga, M., 2003. İzmir Körfezi'nde Hamsi (*Engraulis encrasicolus*, L., 1758)'nin Beslenme Özellikleri Üzerine Bir Ön Çalışma. *E.U. Journal of Fisheries & Aquatic Sciences*, 20 (1-2), 121-127.
- Vallisneri, M., Scapolatempo, M., Piccinetti, C., 2007. Preliminary biological data on the northeast Mediterranean conger eel *Conger conger* L., 1758. *Boletino del Instituto Espanol de Oceanografia*, 23(1-4), 111-114.
- Vallisneri, M., Stagioni, M., Montanini, S., Tommasini, S., 2011. Body size, sexual maturity and diet in *Chelidonichthys lucerna* (Osteichthyes: Triglidae) from the Adriatic Sea, north eastern Mediterranean. *Acta Adriatica*, 51(1), 141-148.
- Valls, M., Quetglas, A., Ordines, F., Moranta, J., 2011. Feeding ecology of demersal elasmobranchs from the shelf and slope off the Balearic Sea (western Mediterranean). *Scientia Marina*, 75(4), 633-639.
- Vasilakopoulos, P., Pavlidis, M., Tserpes, G., 2011. On the diet and reproduction of the oilfish *Ruvettus pretiosus* (Perciformes: Gempylidae) in the eastern Mediterranean. *Journal of the Marine Biological Association of the U.K.*, 91(4), 873-881.
- Vassilopoulou, V., Siapatis, A., Christides, G., Bekas, P., 2003. Life history aspects of the pilotfish (*Naucrates ductor*) in Geek waters. p. 182. In: *7th Panhellenic Symposium of Oceanography and Fisheries, Chersonissos, 6-9 May 2003*. HCMR, Athens.
- Yankova, M.H., Raykov, V.S., Frateva, P.B., 2008. Diet composition of Horse Mackerel, *Trachurus mediterraneus ponticus* Aleev, 1956 (Osteichthyes: Carangidae) in the Bulgarian Black Sea waters. *Turkish Journal of Fisheries and Aquatic Sciences*, 8, 321-327.
- Yapici, S., Filiz, H., Ozkan, O., 2012. Age, growth, reproduction and feeding habits of brown comber, *Serranus hepatus* (L., 1758) in eastern Aegean Sea. *Biharean Biologist*, 6(2), 99-107.

- Yeldan, H., Avsar, D., Manasırılı, M., 2009. Age, growth and feeding of the common stingray (*Dasyatis pastinaca*, L., 1758) in the Cilician coastal basin, northeastern Mediterranean Sea. *Journal of Applied Ichthyology*, 25 (Suppl. 1), 98-102.
- Yığın, C., Ismen, A., 2010a. Age, growth, reproduction and feed of longnosed skate, *Dipturus oxyrinchus* (Linnaeus, 1758) in Saros Bay, the north Aegean Sea. *Journal of Applied Ichthyology*, 26, 913-919.
- Yığın, C., Ismen, A., 2010b. Diet of Thornback ray (*Raja clavata* Linnaeus, 1758) in Saros Bay (the North Aegean Sea). *Rapport de la Commission internationale de la Mer Méditerranée*, 39, 700.
- Zamboukas, N., 2006. *Contribution to the study of biology and biochemical composition of the Atlantic bonito, *Sarda sarda* (Bloch, 1793), in Greek marine areas*. PhD Thesis, University of Athens, 302 pp.
- Zamboukas, N., Megalofonou, P., Apostolopoulou, M., 2003. Data on the diet of the Atlantic bonito (*Sarda sarda*) in the Aegean and Ionian Sea. p. 181. In: *7th Panhellenic Symposium of Oceanography and Fisheries, Chersonissos, 6-9 May 2003*. HCMR, Athens.
- Zamboukas, N., Megalofonou, P., Tomara, T., Yannopoulos, K., 2001. Preliminary results on the diet of little tunny (*E. alletteratus*) from the Aegean and Ionian Sea. p. 113-116. In: *10th Hellenic Conference of Ichthyologists, Chania, 18-20 October 2001*.
- Zorica, B., Cikes Kec, V., 2012. Preliminary observations on feeding habits of garfish *Belone belone* (L., 1761) in the Adriatic Sea. *Croatian Journal of Fisheries*, 70, 53-60.