

## Mediterranean Marine Science

Vol 19, No 2 (2018)



### Preliminary assessment of methanogenic microbial communities in marine caves of Zakynthos Island (Ionian Sea, Greece)

PARASKEVI POLYMENAKOU, MANOLIS MANDALAKIS, THANOS DAILIANIS, CHARALAMPOS DIMITRIADIS, MATEJ MEDVECKY, ANTONIOS MAGOULAS, VASILIS GEROVASILEIOU

doi: [10.12681/mms.14374](https://doi.org/10.12681/mms.14374)

#### To cite this article:

POLYMENAKOU, P., MANDALAKIS, M., DAILIANIS, T., DIMITRIADIS, C., MEDVECKY, M., MAGOULAS, A., & GEROVASILEIOU, V. (2018). Preliminary assessment of methanogenic microbial communities in marine caves of Zakynthos Island (Ionian Sea, Greece). *Mediterranean Marine Science*, 19(2), 284–289.  
<https://doi.org/10.12681/mms.14374>

## Preliminary assessment of methanogenic microbial communities in marine caves of Zakynthos Island (Ionian Sea, Greece)

PARASKEVI N. POLYMENAKOU<sup>1</sup>, MANOLIS MANDALAKIS<sup>1</sup>, THANOS DAILIANIS<sup>1</sup>,  
CHARALAMPOS DIMITRIADIS<sup>2</sup>, MATEJ MEDVECKY<sup>3,4</sup>, ANTONIOS MAGOULAS<sup>1</sup>  
and VASILIS GEROVASILEIOU<sup>1</sup>

<sup>1</sup>Hellenic Centre for Marine Research, Institute of Marine Biology, Biotechnology and Aquaculture,  
Gournes Pediados, Heraklion, Crete, Greece

<sup>2</sup>National Marine Park of Zakynthos, Zakynthos, Greece

<sup>3</sup>National Centre for Biomolecular Research, Faculty of Science, Masaryk University,  
Kotlářská 2, Brno 61137, Czech Republic

<sup>4</sup>Veterinary Research Institute, Brno, Czech Republic

Corresponding author: [polymen@hcmr.gr](mailto:polymen@hcmr.gr)

Handling Editor: Argyro Zenetos

Received: 31 August 2017; Accepted: 1 March 2018; Published on line: 18 June 2018

### Abstract

Mediterranean marine caves remain largely unexplored, while particularly limited information is available about microbial life in these unique environments. This study is a preliminary assessment of the composition of the active anaerobic microbial community colonizing the walls of newly explored systems of underwater caves and small cavities in Zakynthos Island. The interior of these caves is densely coated with egg-shaped, foam-shaped and filamentous biological structures that are characterised by a strong odour of hydrogen sulphide gas. A total of twelve structures scrapped from cave rocks were subjected to anaerobic cultivation for up to 208 days. Strong to moderate methanogenesis was observed in two different types of egg-shaped structures and one foam-like structure. Interestingly, this was observed in experiments that were performed at room temperature (i.e. 25°C), which is substantially lower than those typically considered optimum for methane production (e.g. 35°C). An analysis of the 16S rRNA genes revealed a clear dominance of archaea and bacteria closely related to known methane producers and sulphate reducers, including members of the families *Methanomicrobiaceae*, *Desulfobulbaceae*, *Desulfobacteraceae*, *Desulfuromonaceae*, *Campylobacteraceae*, *Marinifilaceae*, *Clostridiaceae*, Incertae Sedis – Family I & II. These results show that Mediterranean marine caves can host members of archaea and bacteria with potential biotechnological interest that deserves further investigation.

**Keywords:** 16S rRNA gene analysis; Anaerobic cultures; Mediterranean marine caves; Methanogens; Sulfate reducers.

### Introduction

Around the world, marine caves constitute biodiversity hotspots supporting unique diversity elements and species-rich communities of great ecological, scientific and conservation value (Gerovasileiou *et al.*, 2016). Recent biodiversity assessments of this habitat type revealed that Mediterranean marine caves harbour exceptionally high biodiversity (>2000 taxa belonging to 58 major taxonomic groups), including several rare and protected species (Gerovasileiou & Voultsiadou, 2012, 2014; Gerovasileiou *et al.*, 2015). However, insights into their microbiology are limited. Specifically, only 6 marine caves have been investigated regarding their microbial communities, namely, Grotta Azzurra and Grotta Sulfurea on the Tyrrhenian coasts of Italy (Mattison *et al.*, 1998; Canganella

*et al.*, 2002; 2007; Maugeri *et al.*, 2010); Gymnasium, Granchi and Mazzere caves on the Ionian coast of Sicily (Guido *et al.*, 2012, 2013, 2014, 2016; Sanfilippo *et al.*, 2015); and Kakoskali cave in Cyprus (Guido *et al.*, 2017). However, only the former two have been investigated for their microbiota, using molecular techniques, while studies on the other three caves involved the morphological identification of microbial structures.

Methane is a key component of the carbon cycle in many aquatic environments and it is an important end product in the anaerobic degradation of organic matter in marine and freshwater sediments, particularly in zones where more energy-yielding oxidants, such as sulphate, nitrate, or ferric iron, are depleted (Heyer, 1990). Cave ecosystems belong to this type of environments that can host methanogens, i.e. microorganisms that can gain en-

ergy from the conversion of a limited number of simple substrates into methane (Engel, 2010). During a recent investigation concerning marine cave biodiversity in the National Marine Park of Zakynthos – NMPZ (eastern Ionian Sea, Greece), cave and cavity systems densely colonized by egg-shaped, foam-shaped and filamentous biological structures were discovered. These systems were also characterized by a strong odour of hydrogen sulphide, which pointed to the presence of anaerobic activity. The latter is a fundamental prerequisite for the growth of methanogenic microorganisms, as they are all known to be obligate anaerobes. A number of biological samples, representing different morphological types of colonies, were collected from the caves and used for the preparation of anaerobic enrichment cultures. The latter were subsequently subjected to 16S ribosomal RNA gene analysis in order to obtain a preliminary assessment of the active sulphate reducers and methane producers, and evaluate their potential in the fields of bioenergy and bioremediation.

## Materials and Methods


### Sampling site

During a sampling campaign that took place in June 2015, microbial colonies were collected from marine

cave, cavity and fissure systems (semi- and entirely submerged) located at a depth range of 0-25 m, on the south-western coasts of Zakynthos Island (37.661° N, 20.860° E), within the National Marine Park of Zakynthos (Fig. 1a). Representative samples of six different types of biological colonies were collected from cave walls (two replicates) by SCUBA diving, placed in sterile caps at room temperature (total of 12 samples; Supplementary Fig. S1), transferred to the laboratory and immediately subjected to a series of cultivation experiments to enable enrichment of the anaerobic bacteria capable of producing methane (Supplementary Fig. S2). No particular pattern was identified with regard to the distribution of the different types of colonies in the studied caves.

### Enrichment and isolation of methanogenic microbes

Wolfe's anaerobic medium was selected for the enrichment and isolation of methanogenic microbial communities. Technical details about the preparation of this complex anaerobic medium, the transfer of medium aliquots into serum bottles under anaerobic conditions and the cultivation of methanogenic microbes are thoroughly described in Wolfe (2011) and are presented in Supplementary Materials & Methods. For the enrichment of methane-producing microorganisms, a total of 12 cultures were prepared in 100-mL serum bottles that



**Fig. 1:** a) Map showing the geographic location of the studied marine caves in the National Marine Park of Zakynthos, Greece; b-c) crevice colonized by methane producing egg-shaped and foam-like microbial structures; d) egg-shaped, e) foam-like and f-g) filamentous microbial structures found in the study area. Microbial structures are indicated by white arrows. (Photographs by T. Dailianis).



were inoculated with pieces of each biological structure (two replicates). The bottles were tightly sealed with Teflon-lined crimp caps to maintain anaerobic conditions and incubated inverted at room temperature (i.e. 25°C) in the dark (Supplementary Table S1). Production of methane was regularly assessed in all cultures by withdrawing headspace samples (100 µL) from the serum bottles using a gas-tight syringe and analyzing them on a Hewlett Packard 5890 gas chromatograph equipped with a flame ionization detector (GC-FID) and a Carboxen 1006 PLOT capillary column (30 m × 0.53 mm; Supelco, Inc.).

### 16S rRNA gene and phylogenetic analysis

The microbial methanogens enriched from the collected samples were characterized by cloning and sequencing of PCR-amplified 16S rRNA genes following the experimental procedure described by Polymenakou *et al.* (2005). Details about this procedure are provided in Supplementary Materials & Methods. The enrichment culture showing the highest methanogenic activity was selected for the extraction/isolation of genetic material and the amplification of the highly conserved ribosomal RNA gene (16S rDNA), which is typically used as a marker for the identification of microbial species. The sequencing reactions were undertaken using the 20-bp forward primer for bacterial clones Bac-27f 5'-AGAGTTTGATCCTGGCTCAG-3' (Lane, 1991) and the 16-bp forward primer for archaeal clones Arch-8f 5'-TCCGGTTGATCCTGCC-3' (Teske *et al.*, 2002). A total of 83 bacterial and 10 archaeal clones were successfully characterized during this investigation and were used for phylogenetic analysis. Approximately 600 bp long parts of 16S rDNA sequences were aligned by ClustalW (Larkin *et al.*, 2007). At this stage, identical sequences (i.e. sequences showing 100% similarity) were grouped into unique operational taxonomic units (OTUs) and were indicated as Bacterial OTU A-K and Archaeal OTU A-C (Supplementary Table S2). Phylogenetic analysis was performed by Bayesian statistics via MrBayes (Ronquist & Huelsenbeck, 2003) and tree topology was visualized via iTOL (Letunic & Bork, 2016). The 93 partial 16S rDNA sequences generated within the framework of this study were deposited in GenBank under accession numbers MF627323-MF627415.

### Results

The investigated marine cave systems of Zakynthos Island were characterized by the presence of different morphological biological structures forming small egg-shaped (2 types of colonies), foam-like (2 types of colonies) and filamentous mats (2 types of colonies), and pervaded by a strong odour of hydrogen sulphide gas, which is typically indicative of prevailing anaerobic conditions (Fig. 1; Supplementary Fig. S1). Out of 12 samples investigated over the course of a 208-days cultivation period, the two different types of egg-shaped structures (coded

as 1A/1B and 4A; Fig. 1; Supplementary Figs. S1, S2 and Table S1) and one type of foam-like structure (coded as 3A) (Figs. S1) indicated strong or moderate methanogenesis under an atmosphere of H<sub>2</sub>:CO<sub>2</sub> (up to 30% v:v methane in the headspace was measured).

Of the 93 bacterial and archaeal clones that were sequenced during this investigation, a total of 51 bacterial and 3 archaeal OTUs were identified and assigned to nine families (Fig. 2; Supplementary Table S2). More specifically, the microorganisms that dominated the enrichment anaerobic culture showing high methanogenesis were closely related to archaeobacterial strains of family *Methanomicrobiaceae*, and to bacterial strains of families *Desulfobulbaceae*, *Desulfobacteraceae*, *Desulfuromonaceae*, *Campylobacteraceae*, *Marinifilaceae*, *Clostridiaceae*, Incertae Sedis – Family I & II. Interestingly, two bacterial clones did not have any cultured representatives and formed a new cluster named “*ZakynCa*”.

### Discussion

Mediterranean marine caves are largely unexplored and only recently have been recognized as important biodiversity reservoirs that deserve further investigation (e.g. Gerovasileiou & Voultsiadou, 2012; Guido *et al.*, 2016). Regarding their microbiology, only few submarine caves on the Ionian coast of Sicily and Cyprus have been morphologically investigated. The submerged marine caves of Sicily are dominated by unique bioconstructions, named biostalactites, which are constituted of small serpulids and other metazoans associated with carbonate micrites (Guido *et al.*, 2012, 2013, 2014, 2016; Sanfilippo *et al.*, 2015). In these serpulid bioconstructions, micrite precipitates are probably induced by the presence of sulphate reducing bacteria (Guido *et al.*, 2012). Similar serpulid bioconstructions have been observed recently in submarine caves of Lesbos Island, North Aegean Sea (Sanfilippo *et al.*, 2017) and the Kakoskali submarine cave of Cyprus (Guido *et al.*, 2017). However, such bioconstructions were not found in the surveyed caves of Zakynthos. Extensive microbial biofilms/mats have also been found on the rocky surfaces and even on the bottom sediment of two submarine caves with internal sulphur springs, namely, Grotta Azzurra and Grotta Sufiurea in the Tyrrhenian Sea, Italy (Canganella *et al.*, 2002, 2007), which support a unique chemosynthetic ecosystem (Southward *et al.*, 1996).

The current study presents a preliminary assessment of the active microbial anaerobes residing in the biological structures covering the walls of newly discovered systems of submarine caves and cave-like formations in Zakynthos. Anaerobic enrichment cultures performed during this study indicated strong methanogenesis only in egg-shaped and one foam-like microbial structure.

Sequencing analysis of the 16S rRNA genes of collected microorganisms revealed the dominance of archaea and bacteria closely related to known methane producers of the genus *Methanoculleus* and sulphate reducers



palladium and gold) are some other important environmental remediation applications of SRBs (Barton & Fauque, 2009; Hussain *et al.*, 2016).

Overall, the newly discovered marine caves of Zakynthos host members of archaea and bacteria with potential biotechnological interest that deserve further investigation. In this preliminary study, we used enrichment cultures to characterize methanogenic microbial communities. Interestingly, strong to moderate methanogenesis was observed in experiments that were performed at room temperature (i.e. 25°C), which is substantially lower than those typically considered optimum for methane production (e.g. 35°C; Sánchez *et al.*, 2000). The ability of the isolated microbial communities to produce methane at the energetically less expensive ambient temperature implies potential large-scale exploitation. Finally, we would like to emphasize the necessity of an extensive survey on the chemistry/microbiology of Mediterranean marine caves and the application of modern molecular tools to unravel the metabolic capabilities of cave microbiomes and evaluate their potential applications in biotechnology.

## Acknowledgements

The authors acknowledge Christos Arvanitidis, Maria Sini, Georgios Chatzigeorgiou, Konstantinos Vatikiotis and the Management Agency of the National Marine Park of Zakynthos for providing assistance during fieldwork. This work was funded by the General Secretariat for Research and Technology-GSRT and Siemens A.G. under the “Programmatic agreements between Research Centres – GSRT 2015-2017” project.

## References

- Barton, L.L., Fauque, G.D., 2009. Biochemistry, physiology and biotechnology of sulfate-reducing bacteria. *Advances in Applied Microbiology*, 68, 41-98.
- Canganella, F., Bianconi, G., Gambacorta, A., Kato, C., Uematsu, K., 2002. Characterisation of Heterotrophic Microorganisms Isolated from the “Grotta Azzura” of Cape Palinuro (Salerno, Italy). *Marine Ecology*, 23, 1-10.
- Canganella, F., Bianconi, G., Kato, C., Gonzalez, J., 2007. Microbial ecology of submerged marine caves and holes characterised by high levels of hydrogen sulphide. *Reviews in Environmental Science and Biotechnology*, 6, 61-70.
- Dianou, D., Miyaki, T., Asakawa, S., Morii, H., Nagaoka, K. *et al.*, 2001. *Methanoculleus chikugoensis* sp. nov., a novel methanogenic archaeon isolated from paddy field soil in Japan, and DNA-DNA hybridization among *Methanoculleus* species. *International Journal of Systematic and Evolutionary Microbiology*, 51, 1663-1669.
- Engel, A.S., 2010. Microbial diversity of cave ecosystems. p. 219-238. In *Geomicrobiology: Molecular and Environmental Perspective*. Barton, L.L. *et al.* (Eds.), Springer Science+Business Media B.V.
- Friedrich, M., Springer, N., Ludwig, W., Schink, B., 1996. Phylogenetic positions of *Desulfobacterium glycolicum* gen. nov., sp. nov., and *Syntrophobacterium glycolicum* gen. nov., sp. nov., two new strict anaerobes growing with glycolic acid. *International Journal of Systematic and Evolutionary Microbiology*, 46, 1065-1069.
- Gerovasileiou, V., Voultsiadou, E., 2012. Marine caves of the Mediterranean Sea: a sponge biodiversity reservoir within a biodiversity hotspot. *Plos One*, 7, e39873. doi: 10.1371/journal.pone.0039873
- Gerovasileiou, V., Voultsiadou, E., 2014. Mediterranean marine caves as biodiversity reservoirs: a preliminary overview. p. 45-50. In: *Proceedings of the 1st Mediterranean Symposium on the conservation of Dark Habitats (Portorož, Slovenia, 31 October 2014)*. Bouafif, C., Langar, H., Ouerghi, A. (Eds). RAC/SPA, Tunis.
- Gerovasileiou, V., Chintiroglou, C., Vafidis, D., Koutsoubas, D., Sini, M. *et al.*, 2015. Census of biodiversity in marine caves of the Eastern Mediterranean Sea. *Mediterranean Marine Science*, 16, 245-265.
- Gerovasileiou, V., Martínez, A., Álvarez, F., Boxshall, G., Humphreys, W. *et al.*, 2016. World Register of marine Cave Species (WoRCS): a new Thematic Species Database for marine and anchialine cave biodiversity. *Research Ideas and Outcomes*, 2, e10451. doi: 10.3897/rio.2.e10451
- Guido, A., Mastandrea, A., Rosso, A., Sanfilippo, R., Russo, F., 2012. Micrite precipitation induced by sulphate reducing bacteria in serpulid bioconstructions from submarine caves (Syracuse, Sicily). *Rendiconti Online della Società Geologica Italiana*, 21, 933-934.
- Guido, A., Heindel, K., Birgel, D., Rosso, A., Mastandrea, A. *et al.*, 2013. Pendant bioconstructions cemented by microbial carbonate in submerged marine caves (Holocene, SE Sicily). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 388, 166-180.
- Guido, A., Mastandrea, A., Rosso, A., Sanfilippo, R., Tosti, F. *et al.*, 2014. Commensal symbiosis between agglutinated polychaetes and sulfate reducing bacteria. *Geobiology*, 12, 265-275.
- Guido, A., Rosso, A., Sanfilippo, R., Russo, F., Mastandrea, A., 2016. Submarine caves: a laboratory for geomicrobiology studies. *Rendiconti Online della Società Geologica Italiana*, 38, 62-64.
- Guido, A., Jimenez, C., Achilleos, K., Rosso, A., Sanfilippo, R. *et al.*, 2017. Cryptic serpulid-microbialite bioconstructions in the Kakoskali submarine cave (Cyprus, Eastern Mediterranean). *Facies*, 63, 21.
- Heyer, J., 1990. Der Kreislauf des Methans. Akademie-Verlag Berlin, 250 pp.
- Hussain, A., Hasan, A., Javid, A., Iqbal Qazi, J., 2016. Exploited application of sulfate-reducing bacteria for concomitant treatment of metallic and non-metallic wastes: a mini review. *3 Biotech*, 6, 119.
- Jaenicke, S., Ander, C., Bekel, T., Bisdorf, R., Dröge, M. *et al.*, 2011. Comparative and joint analysis of two metagenomic datasets from a biogas fermenter obtained by 454-pyrosequencing. *Plos One*, 6, e14519. doi: 10.1371/journal.pone.0014519
- Lane, D.J., 1991. 16S/23S rRNA sequencing. p. 115-175. In: *Nucleic acid techniques in bacterial systematics*. Stackebrandt, E., Goodfellow, M. (Eds). Wiley, New York.
- Larkin, M.A., Blackshields, G., Brown, N.P., Chenna, R., McGettigan, P.A. *et al.*, 2007. Clustal W and clustal X version 2.0. *Bioinformatics*, 23, 2947-2948.
- Letunic, I., Bork, P., 2016. Interactive tree of life (iTOL) v3: an online tool for the display and annotation of phylogenetic and other trees. *Nucleic Acids Research*, 44, W242-245.
- Liesack, W., Finster, K., 1994. Phylogenetic analysis of five strains of gram-negative, obligately anaerobic, sulfur-re-



- ducing bacteria and description of *Desulfuromusa* gen. nov., including *Desulfuromusa kysingii* sp. nov., *Desulfuromusa bakii* sp. nov., and *Desulfuromusa succinoxidans* sp. nov. *International Journal of Systematic Bacteriology*, 44, 753-758.
- Maestrojuan, G.M., Boone, D.R., Xun, L., Mah, R.A., Zhang, L., 1990. Transfer of *Methanogenium bourgense*, *Methanogenium marisnigri*, *Methanogenium olentangyi*, and *Methanogenium thermophilicum* to the genus *Methanoculleus* gen. nov., emendation of *Methanoculleus marisnigri* and *Methanogenium*, and description of new strains of *Methanoculleus bourgense* and *Methanoculleus marisnigri*. *International Journal of Systematic Bacteriology*, 40, 117-122.
- Mattison, R., Abbiati, M., Dando, P., Fitzsimons, M.F., Pratt, S.M. *et al.*, 1998. Chemoautotrophic microbial mats in submarine caves with hydrothermal sulphidic springs at Cape Palinuro, Italy. *Microbial Ecology*, 35, 58-71.
- Maugeri, T.L., Bianconi, G., Canganella, F., Danovaro, R., Gugliandolo, C. *et al.*, 2010. Shallow hydrothermal vents in the southern Tyrrhenian Sea. *Chemistry and Ecology*, 26, 285-298.
- Maus, I., Wibberg, D., Stantscheff, R., Eikmeyer, F.-G., Seffner, A. *et al.*, 2012. Complete genome sequence of the hydrogenotrophic, methanogenic archaeon *Methanoculleus bourgensis* strain MS2<sup>T</sup>, isolated from a sewage sludge digester. *Journal of Bacteriology*, 194, 5487-5488.
- Oren, A., 2014. The family Methanomicrobiaceae. p. 165-193. In: *The Prokaryotes – Other Major Lineages of Bacteria and the Archaea*. Rosenberg, E., DeLong, E.F., Lory, S., Stackebrandt, E., Thompson, F. (Eds), Springer-Verlag Berlin Heidelberg.
- Polymenakou, P.N., Bertilsson, S., Tselepidis, A., Stephanou, E.G., 2005. Bacterial community composition in different sediments from the Eastern Mediterranean Sea: a comparison of four 16S ribosomal DNA clone libraries. *Microbial Ecology*, 50, 447-462.
- Romesser, J.A., Wolfe, R.S., Mayer, F., Spiess, E., Walther-Mauruschat, A., 1979. *Methanogenium*, a new genus of marine methanogenic bacteria, and characterization of *Methanogenium cariaci* sp. nov. *Archives of Microbiology*, 121, 147-153.
- Ronquist, F., Huelsenbeck, J.P., 2003. MRBAYES 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics*, 19, 1572-1574.
- Sánchez, E., Borja, R., Weiland, P., Travieso, L., Martin, A., 2000. Effect of temperature and pH on the kinetics of methane production, organic nitrogen and phosphorus removal in the batch anaerobic digestion process of cattle manure. *Bioprocess Engineering*, 22, 247-252.
- Sanfilippo, R., Rosso, A., Guido, A., Mastrandrea, A., Russo, F. *et al.*, 2015. Metazoan/microbial biostalactites from present-day submarine caves in the Mediterranean Sea. *Marine Ecology*, 36, 1277-1293.
- Sanfilippo, R., Rosso, A., Guido, A., Gerovasileiou, V., 2017. Serpulid communities from two marine caves in the Aegean Sea, eastern Mediterranean. *Journal of the Marine Biological Association of the United Kingdom*, 97, 1059-1068.
- Southward A.J., Kennicutt M.C., Alcalá-Herrera J., Abbiati M., Airolidi L. *et al.*, 1996. On the biology of submarine caves with Sulphur springs: appraisal of <sup>13</sup>C/<sup>12</sup>C ratios as a guide to trophic relations. *Journal of the Marine Biological Association of the United Kingdom*, 76, 265-285.
- Teske, A., Hinrichs, K.-U., Edgcomb, V.P., de Vera Gomez, A., Kysela D. *et al.*, 2002. Microbial diversity in hydrothermal sediments in the Guaymas Basin: evidence for anaerobic methanotrophic communities. *Applied and Environmental Microbiology*, 68, 1994-2007.
- Wolfe, R., 2011. Techniques for cultivating methanogens. *Methods in Enzymology*, 494, 1-22.