

Mediterranean Marine Science

Vol 19, No 2 (2018)



Investigation on the genus *Squalus* in the Sardinian waters (Central-Western Mediterranean) with implications on its management

ANDREA BELLODI, CRISTINA PORCU, ALESSANDRO CAU, MARTINA FRANCESCA MARONGIU, RICCARDO MELIS, ANTONELLO MULAS, PAOLA PESCI, MARIA CRISTINA FOLLESA, RITA CANNAS

doi: [10.12681/mms.15426](https://doi.org/10.12681/mms.15426)

To cite this article:

BELLODI, A., PORCU, C., CAU, A., MARONGIU, M. F., MELIS, R., MULAS, A., PESCI, P., FOLLESA, M. C., & CANNAS, R. (2018). Investigation on the genus *Squalus* in the Sardinian waters (Central-Western Mediterranean) with implications on its management. *Mediterranean Marine Science*, 19(2), 256–272. <https://doi.org/10.12681/mms.15426>

Investigation on the genus *Squalus* in the Sardinian waters (Central-Western Mediterranean) with implications on its management

ANDREA BELLODI^{1,2}, CRISTINA PORCU^{1,2}, ALESSANDRO CAU^{1,2,3},
MARTINA F. MARONGIU^{1,2}, RICCARDO MELIS^{1,2}, ANTONELLO MULAS^{1,2},
PAOLA PESCI^{1,2}, MARIA C. FOLLESA^{1,2} and RITA CANNAS^{1,2}

¹Department of Life and Environmental Sciences, University of Cagliari via T. Fiorelli 1 - 09126 Cagliari, Italy

²CoNISMa Consorzio Nazionale Interuniversitario per le Scienze Mare,
Piazzale Flaminio, 9 - 00196 Rome, Italy

³Department of Architecture, Design and Urban Development, University of Sassari,
Palazzo Pou Salit, Piazza Duomo 6, 07041 Alghero, Italy

Corresponding author: abelloidi@unica.it

Handling Editor: Paraskevi Karachle

Received: 15 December 2017; Accepted: 23 March 2018; Published on line: 14 June 2018

Abstract

In the Mediterranean Sea, in addition to the two historically known species belonging to the *Squalus* genus, a third species, *Squalus megalops*, has been reported. Considering the high level of morphologic similarity of this species with the native species *S. blainville*, this study aims to evaluate the Central-Western Mediterranean spurdog population in order to test the hypothesis of the presence of two distinct species *S. blainville* and *S. megalops*. A total of 137 spurdogs, caught in the Sardinian waters, were analyzed morphologically and genetically after their subdivision into two groups depending on the number of the lateral processes in the chondrocranium basal plate. The CAP analysis, employing all body and chondrocranial measurements, revealed no clear segregation among the *a priori* assigned groups with a high misclassification percentage. Besides, no evident dissimilarities in teeth and dermal denticle morphology between the two groups were observed. All the 18 specimens which were genetically analyzed, by sequencing of the mtDNA marker COI, clustered together resulting to be *S. blainville*. All the obtained results indicated the presence, in the study area, of only one species, ascribable to *S. blainville*.

Keywords: *Squalus blainville*; *Squalus megalops*; Mediterranean Sea; taxonomy; mtDNA sequencing; morphology.

Introduction

The correct taxonomic identification of species provides a critical baseline that supports the rest of biological research (Last *et al.*, 2007). Generally, Elasmobranchs have suffered major taxonomic constraints that have led to misidentification issues related to by-catch and fisheries, which were usually solved by grouping data at higher taxonomic levels, such as genus or family (*e.g.* Zeeberg *et al.*, 2006; Coelho & Erzini, 2008).

Squalidae represent one of the most commercially targeted families among Elasmobranchs (Ebert *et al.*, 2013). Indeed, several species belonging to this family are landed by up to 50 countries in direct fisheries or as bycatch (Ebert *et al.* 2013). Their relatively high commercial value, in addition to *K*-selected life strategy that commonly characterizes Elasmobranchs, identifies this taxonomic group as exceptionally susceptible to fishing mortality. This particular situation, despite the considerable abundance and the wide habitat range of some species, could easily lead them to stock depletion (Ebert *et al.*, 2013).

Squalids belonging to the genus *Squalus* (Blainville, 1816), otherwise known as spurdogs, dogsharks and dogfishes, are among the most taxonomically problematic shark groups due to their strong morphological similarities. Until 2013, 25 species were known (Ebert *et al.*, 2013) including 14 species recognized as valid by Compagno *et al.* (2005) and 11 species added later from the Western Indo-Pacific Ocean by Last *et al.* (2007). In addition, considering the resurrection of *S. acutipinnis* (Regan 1908) by Viana & Carvalho (2016) from South Africa and the description of four new species (*S. albicaudatus*, *S. bahiensis*, *S. lobularis* and *S. quasimodo*) from the South-West Atlantic (Viana *et al.*, 2016), this number has recently been increased.

Squalus species have been divided into three main species groups, based on morphological features such as the relative position of the pectoral fins, the anterior nasal flap shape and skin colour (Bigelow & Schroeder 1957; Ebert *et al.*, 2010): 1) the '*acanthias* group'; 2) the '*mitsukurii* group' historically known as the '*blainville-fernandinus* group', and 3) the '*megalops* group', also

known as ‘the *brevirostris-cubensis* group’. However, a correct identification of several widespread species still remains doubtful. Besides, this particular condition has also been reinforced for the *Squalus* genus due to their high overlapping level of morphological features (Last *et al.*, 2007). Such classification uncertainties constituted an impediment to stakeholders, scientists and managers, somehow retarding the development of management measures because of the difficulties in evaluating the population status of several *Squalus* species.

In the Mediterranean Sea, two *Squalus* species commonly occur (Serena *et al.*, 2005; Serena *et al.*, 2009): the spiny spurdog *S. acanthias* (Linnaeus, 1758) belonging to the ‘*acanthias* group’ and the longnose spurdog *S. blainville* (Risso, 1827) belonging to the ‘*mitsukurii* group’. In this Basin, in the 1980s, Muñoz-Chápuli *et al.* (1984) and Muñoz-Chápuli & Ramos (1989) also recorded a third species, the piked spurdog *S. megalops* (Macleay, 1881), commonly distributed in the Eastern Atlantic and Indo-Pacific Oceans (Ebert *et al.*, 2013).

Despite the fact that *S. acanthias* shows diagnostic characters, such as the presence of white spots on the back or narrowly round to acutely angular rear tips and inner margins of the pectoral fins, which permit an easier identification and discrimination from the other two species (Bonello *et al.*, 2016), *S. blainville* and *S. megalops*, do show a very similar morphology. According to Muñoz-Chápuli *et al.* (1984) and Muñoz-Chápuli & Ramos (1989), *S. blainville* and *S. megalops* can be discriminated principally based on the number of chondrocranial lateral processes, in addition to other morphological features such as teeth and dermic denticles morphology. These findings have been confirmed by Marouani *et al.* (2012) in the Gulf of Gabès (southern Tunisia, central western Mediterranean Sea) through morphometric, meristic and genetic analyses, suggesting that *S. megalops* could be even more common than *S. blainville* in these waters. On the other hand, in a recent study, *S. blainville* was the only *Squalus* species identified in the Maltese waters (Bonello *et al.*, 2016). Indeed, the authors asserted that the species identification based only on morphological characteristics can easily lead to taxonomic misidentifications, especially when multiple anatomical characters (e.g. skull and teeth morphology) are used (Bonello *et al.*, 2016). Moreover, Veríssimo *et al.* (2017) reported that *S. blainville* and *S. megalops* are two names used almost interchangeably along the Eastern Atlantic and the Mediterranean Sea to identify the same species with the former mostly employed in the Mediterranean area while the latter in the Eastern Atlantic. Nevertheless, the results provided by those authors suggest that the ‘true’ *S. megalops* from Australia is not present in the eastern Atlantic and Mediterranean waters, but a different species that remains unidentified can occur (Veríssimo *et al.*, 2017).

Considering these last studies, the present paper aims to investigate the presence of the two species around Sardinian Sea through genetic and morphometric analyses, providing new evidences in order to solve the spurdogs taxonomic confusion in the investigated region.

Materials and Methods

A total of 137 spurdogs were sampled during experimental trawl surveys (MEDITS, Mediterranean International Trawl Survey, Bertrand *et al.*, 2000) and commercial hauls performed from 2010 to 2011 in Sardinian waters (Central Western Mediterranean Sea) at depths from 123 to 682 m (Fig. 1).

Once in the laboratory, specimens were measured (Total Length, TL) and weighed (Total Mass, TM). For the morphometric analysis, specimens were photographed with a digital camera (Nikon D90) in order to take 45 somatic measurements (expressed in millimetres). All measurements, including names and abbreviations, were defined according to Compagno (2001) and Last *et al.* (2007) and expressed in % of TL.

Each shark chondrocranium, after being extracted through a boiling process, was photographed in both dorsal and ventral view in order to obtain 16 measurements following Muñoz-Chápuli & Ramos (1989). Measurements were expressed in millimetres and in % of Total Length of Chondrocranium (TLC). The total number of vertebrae was counted after dissection. Teeth samples from both dental arches were extracted from each individual. Moreover, following Muñoz-Chápuli & Ramos (1989) and Marouani *et al.*, (2012), a skin portion was extracted from the lateral-dorsal area (anterior to the first dorsal spine) for the observation of dermal denticles.

According to Muñoz-Chápuli & Ramos (1989) the number of lateral processes of the chondrocranium basal plate allows the two spurdogs species *S. blainville* and *S. megalops* to be subdivided. For this reason, in the present study, specimens were subdivided into two groups: S1, hypothetically belonging to *S. blainville* (presenting a single lateral process) and S2, hypothetically belonging to *S. megalops* (presenting two lateral processes) (Fig. 2). This characteristic was preferred considering the uncertainty level typical of the other specific features suggested, subjected to corrosion, such as the teeth morphology, or characterized by a relatively high morphological variability degree due to the simultaneous presence of different development stages of the fast replacing rated structures, such as dermal denticles (Kemp, 1999).

Statistical analyses

Through a similarity matrix based on Euclidean distance, *a priori* multivariate differences in the morphological features of the species have been illustrated using the bi-plot produced after Canonical Analysis of Principal Coordinates CAP (Anderson & Willis, 2003) obtained through PRIMER (ver. PRIMER Permanova +) CAP routine. This analysis was chosen as a flexible method for constrained ordination on the basis of any distance or dissimilarity measure, which displays a cloud of multivariate points by reference to a specific *a priori* hypothesis; in our case the hypothesis was that different species of the genus *Squalus* are characterized by different morphological parameters. The routine was conducted on two




Fig. 1: Study area. The black dots represent the hauls in which specimens were caught.




Fig. 2: Dorsal (D) and ventral (V) view of dissected chondrocrania of specimens caught in the Sardinian waters, belonging to S1 group (male, TL= 594 mm) and S2 group (male, TL= 583 mm). White arrows indicate the processes of the chondrocranium basal plate.

data matrixes (and relative similarity matrixes) describing body parameters and chondrocranium. The cross-validation, given by the same routine, was used to further confirm (or reject) the *a priori* assignment of the species.

Moreover, a *t*-Student test (Zar, 1999) was conducted in order to test for differences in chondrocranial measurements between the two groups.

Genetic analysis

A subsample of 18 individuals were selected, based on the characteristics of their chondrocranium, and genetically analysed: 13 individuals (8 males and 5 females) presented two lateral processes and 5 individuals (all males) presented a single lateral process. Total genomic DNA was extracted from the tissues using a salting-out protocol (Miller *et al.*, 1988).

The primers (LCO1490: 5'-GGTCAACAAATCATAAAGATATTGG-3'; HCO2198: 5'-TAAACTTCAGGTGACCAAAAATCA-3') for the amplifications of mitochondrial COI gene were obtained from Folmer *et al.* (1994). The amplification was based on the following cycling parameters: 3 min at 94°C for the initial denaturation, followed by 37 cycles of 30 sec at 94°C, 45 sec at 50°C for the annealing of primers, and 60 sec at 72°C for extension, and then 4 min at 72°C for the final extension. The sequences were sequenced on both directions, aligned in MEGA v. 6 (Tamura *et al.*, 2013) and translated into aminoacidic sequences using the vertebrate genetic code to exclude the occurrence of codon stop and nuclear pseudogenes. Number of haplotypes, haplotype diversity [hd], and nucleotide diversity [π] were retrieved using DnaSP v. 5.1 (Librado & Rozas, 2009). Graphically, the haplotypes were arranged in a network with PopART (<http://popart.otago.ac.nz>) using the Median Joining method (Bandelt *et al.*, 1999).

The sequences obtained in this study were compared to COI sequences published for the three species of the genus *Squalus* reported to be present in the Mediterranean Sea (*S. acanthias*, *S. blainville*, and *S. megalops*) (Table S1). Moreover, the analyses also included sequences of the species included in Group I (*S. suckleyi*) and Group II (*S. cubensis*, *S. raoulensis*, *S. brevirostris*) in the *Squalus* phylogeny by Veríssimo *et al.* (2017). Sequences were retrieved from GenBank (<https://www.ncbi.nlm.nih.gov/>

genbank). *Cirrhigaleus australis* was used as outgroup (Veríssimo *et al.*, 2017). The list and details of the sequences used in the analyses are provided as supplementary table (Table S1).

The relationships among haplotypes were investigated with the Bayesian approach using MrBayes v. 3.1 (Huelsenbeck & Ronquist, 2001; Ronquist & Huelsenbeck, 2003). In MrBayes the analyses were performed using two parallel runs of 2 million generations each, using four chains, sampling every 100 generations, burnin 0.25, and saving branch lengths. The performance of the analyses was evaluated using the software Tracer v. 1.6 (Rambaut *et al.*, 2014). The tree was visualized with MEGA.

Results

According to the number of processes in the chondrocranium, out of the total 137 spurdogs, 19 were pooled in the S1 group (one process, 15 males and 4 females) and 118 were pooled in the S2 group (two processes, 55 males and 63 females) (Table 1).

Chondrocranium description

The chondrocranium measurements obtained are reported in Table 2 for S1 and Table 3 for S2. The distance between the posterior tip and the precerebral fenestra (PPF) was 62.93 and 63.33 in %TLC in S1 and S2 group respectively. In S1, the width across nasal capsules and the interorbital width were 54.74 and 28.25 in %TLC, while in S2 the same measurements were 55.34 and 28.34 in %TLC. Finally, the distance between the basal plate processes was equal to 31.32 in %TLC in S1 and 31.48 in %TLC in S2.

No significant differences in all chondrocranial measurements were found between the two groups (*t*-test $p > 0.05$) (Table 4).

Morphological description

Biometric data from S1 and S2 is reported in Tables 5 and 6, respectively. All studied specimens (S1 and S2) showed a fusiform and elongated body (Fig. 4). In both groups the head appeared slightly triangular from lateral view with a moderately long and sharp snout. The mouth,

Table 1. Number (N) and Total length (TL) range and mean (\pm SD, standard deviation) of the samples used in this study, for each sex and group (S1 and S2).

| Sex | S1 | | | S2 | | |
|---------|-----------|---------------|---------------------|------------|---------------|---------------------|
| | N | Range TL (mm) | Mean TL (\pm SD) | N | Range TL (mm) | Mean TL (\pm SD) |
| Males | 15 | 249-594 | 396.7 \pm 90.0 | 55 | 272-595 | 399.8 \pm 83.2 |
| Females | 4 | 361-792 | 523.3 \pm 234.35 | 63 | 207-834 | 433.3 \pm 140.9 |
| Total | 19 | 249-792 | 416.7 \pm 122.9 | 118 | 207-834 | 415.4 \pm 114.5 |

Table 2. Proportional cranial dimensions expressed as percentages of TLC (\pm SD) for specimens belonging to S1 group, compared with what reported for *S. blainville* by other authors in other world regions.

| References | | Muñoz-Chápuli & Ramos (1989) | | Marouani <i>et al.</i> (2012) | | Bonello <i>et al.</i> (2016) | | Present study | |
|---|-------|---------------------------------|------------------|---|------------------|--|------------------|-------------------------------|------------------|
| Study area | | Eastern Atlantic, Mediterranean | | Tunisian waters (Central Mediterranean) | | Maltese waters (Central Mediterranean) | | Central Western Mediterranean | |
| Measurements | Codex | <i>S. blainville</i> | | <i>S. blainville</i> | | One-lobed chondrocranial | | S1 | |
| Total length of chondrocranium range (mm) | TLC | 57.9-115.7 | | 48.5-88.5 | | | | 47.3-104.5 | |
| | | N | Mean \pm SD | N | Mean \pm SD | N | Mean \pm SD | N | Mean \pm SD |
| Posterior tip-precerebral fenestra | PPF | 9 | 63.85 \pm 1.36 | 23 | 65.46 \pm 3.02 | 23 | 61.58 \pm 3.34 | 16 | 62.93 \pm 2.84 |
| Length precerebral fenestra | LPF | 9 | 36.18 \pm 1.77 | 23 | 33.41 \pm 2.93 | 23 | 35.52 \pm 5.74 | 16 | 28.11 \pm 1.87 |
| Width precerebral fenestra | WPF | 9 | 14.95 \pm 1.45 | 23 | 19.12 \pm 3.05 | 23 | 23.08 \pm 5.49 | 16 | 19.62 \pm 1.79 |
| Width across nasal capsules | WNC | 9 | 54.39 \pm 1.73 | 23 | 54.35 \pm 2.36 | 23 | 54.47 \pm 4.83 | 16 | 54.74 \pm 1.93 |
| Interorbital width | IOW | 9 | 31.18 \pm 1.17 | 23 | 31.77 \pm 2.32 | 23 | 33.10 \pm 5.48 | 16 | 28.25 \pm 1.33 |
| Postorbital width | PsOW | 9 | 56.49 \pm 1.36 | 23 | 57.02 \pm 2.76 | 23 | 58.98 \pm 6.99 | 16 | 54.55 \pm 1.54 |
| Distance between orbital processes | OPD | 9 | 42.60 \pm 1.72 | 23 | 36.03 \pm 2.77 | 23 | 35.33 \pm 1.87 | 16 | 36.48 \pm 1.02 |
| Width between pterotic processes | PtPW | 9 | 37.52 \pm 1.53 | 22 | 39.52 \pm 2.33 | 23 | - | 16 | 37.65 \pm 0.96 |
| Width between hyomandibular facets | HFW | 9 | 45.61 \pm 1.16 | 22 | 45.73 \pm 3.39 | 23 | 43.85 \pm 2.56 | 16 | 44.69 \pm 0.99 |
| Posterior tip-rostral keel | PtRK | 9 | 64.79 \pm 1.35 | 22 | 68.10 \pm 2.76 | 23 | 70.65 \pm 6.35 | 16 | 63.83 \pm 1.97 |
| Length rostral keel | RKL | 9 | 20.05 \pm 2.94 | 22 | 19.96 \pm 2.13 | 23 | 14.32 \pm 2.16 | 16 | 22.58 \pm 1.64 |
| Subethmoidean width | SEtW | 9 | 17.22 \pm 1.12 | 22 | 14.37 \pm 2.12 | 23 | 17.01 \pm 2.16 | 16 | 14.70 \pm 1.11 |
| Width basal angle | BAW | 9 | 21.20 \pm 1.77 | 22 | 19.10 \pm 1.85 | 23 | 22.84 \pm 4.03 | 16 | 19.12 \pm 1.74 |
| Length basal plate | BpL | 9 | 39.53 \pm 1.62 | 22 | 46.61 \pm 2.53 | 23 | - | 16 | 40.12 \pm 1.36 |
| Width between processes of basal plate | BBpW | 9 | 30.39 \pm 1.01 | 22 | 31.37 \pm 2.25 | 23 | - | 16 | 31.32 \pm 0.68 |

Table 3. Proportional cranial dimensions expressed as percentages of TLC (\pm SD) for specimens belonging to S2 group, compared with what reported for *S. megalops* by other authors in other world regions.

| References | | Muñoz-Chápuli & Ramos (1989) | | Marouani <i>et al.</i> (2012) | | Bonello <i>et al.</i> (2016) | | Present study | |
|---|-------|---------------------------------|------------------|---|------------------|--|-------------------|-------------------------------|------------------|
| Study area | | Eastern Atlantic, Mediterranean | | Tunisian waters (Central Mediterranean) | | Maltese waters (Central Mediterranean) | | Central Western Mediterranean | |
| Measurements | Codex | <i>S. megalops</i> | | <i>S. megalops</i> | | Two-lobed chondrocranial | | S2 | |
| Total length of chondrocranium range (mm) | TLC | 32.0-83.8 | | 40.0-87.0 | | | | 34.3-109.2 | |
| | | N | Mean \pm SD | N | Mean \pm SD | N | Mean \pm SD | N | Mean \pm SD |
| Posterior tip-precerebral fenestra | PPF | 22 | 65.08 \pm 1.14 | 17 | 67.03 \pm 3.25 | 146 | 62.78 \pm 8.42 | 102 | 63.33 \pm 3.13 |
| Length precerebral fenestra | LPF | 22 | 35.7 \pm 1.02 | 17 | 31.95 \pm 1.61 | 146 | 35.75 \pm 4.25 | 102 | 28.13 \pm 1.72 |
| Width precerebral fenestra | WPF | 22 | 16.99 \pm 1.84 | 17 | 20.33 \pm 1.90 | 146 | 21.73 \pm 5.15 | 102 | 19.53 \pm 1.61 |
| Width across nasal capsules | WNC | 21 | 50.93 \pm 2.44 | 16 | 51.92 \pm 3.59 | 146 | 53.63 \pm 13.50 | 102 | 55.34 \pm 1.93 |
| Interorbital width | IOW | 22 | 28.57 \pm 1.26 | 16 | 28.72 \pm 1.79 | 146 | 31.85 \pm 5.75 | 102 | 28.34 \pm 1.19 |

(continued)

Table 3 continued

| References | Muñoz-Chápuli and Ramos (1989) | | Marouani <i>et al.</i> (2012) | | Bonello <i>et al.</i> (2016) | | Present study | |
|---|---------------------------------|--------------------|---|------------|--|-------------|-------------------------------|------------|
| Study area | Eastern Atlantic, Mediterranean | | Tunisian waters (Central Mediterranean) | | Maltese waters (Central Mediterranean) | | Central Western Mediterranean | |
| Measurements | Codex | <i>S. megalops</i> | <i>S. megalops</i> | | Two-lobed chondrocranial | | S2 | |
| Total length of chondrocranium range (mm) | TLC | 32.0-83.8 | 40.0-87.0 | | | | 34.3-109.2 | |
| Postorbital width | PsOW | 22 55.38±2.00 | 16 | 58.19±2.38 | 146 | 57.77±7.64 | 102 | 54.87±1.70 |
| Distance between orbital processes | OPD | 19 32.85±2.58 | 16 | 36.31±2.55 | 146 | 36.04±4.44 | 102 | 36.73±2.72 |
| Width between pterotic processes | PtPW | 22 37.3±1.24 | 16 | 39.80±2.16 | 146 | - | 102 | 38.06±1.58 |
| Width between hyomandibular facets | HFW | 22 45.62±1.16 | 16 | 47.06±2.04 | 146 | 43.74±3.65 | 102 | 44.96±1.27 |
| Posterior tip-rostral keel | PtRK | 22 63.6±3.12 | 16 | 68.43±3.21 | 146 | 68.79±14.58 | 102 | 64.21±2.03 |
| Length rostral keel | RKL | 22 22.82±2.69 | 16 | 21.02±3.16 | 146 | 14.97±5.89 | 102 | 21.89±1.66 |
| Subethmoidean width | SEtW | 22 15.57±1.31 | 16 | 13.60±1.54 | 146 | 16.33±3.53 | 102 | 15.08±1.20 |
| Width basal angle | BAW | 22 17.82±1.36 | 16 | 20.01±1.56 | 146 | 21.86±5.39 | 102 | 19.43±1.99 |
| Length basal plate | BpL | 22 40.56±1.09 | 16 | 46.24±1.75 | 146 | - | 102 | 40.41±1.78 |
| Width between processes of basal plate | BBpW | 22 31.08±0.85 | 17 | 33.39±3.61 | 146 | - | 102 | 31.48±1.12 |

Table 4. Comparison between chondrocranial measurements of spurdogs belonging to S1 and S2 groups from the Sardinian waters.

| Measurements | Codex | t-test | p-value |
|---|-------|--------|---------|
| Total length of chondrocranium range (mm) | TLC | | |
| Posterior tip-precerebral fenestra | PPF | -0.48 | 0.63 |
| Length precerebral fenestra | LPF | -0.03 | 0.97 |
| Width precerebral fenestra | WPF | 0.19 | 0.85 |
| Width across nasal capsules | WNC | -1.15 | 0.25 |
| Interorbital width | IOW | -0.30 | 0.76 |
| Postorbital width | PsOW | -0.69 | 0.49 |
| Distance between orbital processes | OPD | -0.36 | 0.72 |
| Width between pterotic processes | PtPW | -1.01 | 0.31 |
| Width between hyomandibular facets | HFW | -0.83 | 0.41 |
| Posterior tip-rostral keel | PtRK | -0.71 | 0.48 |
| Length rostral keel | RKL | 1.55 | 0.12 |
| Subethmoidean width | SEtW | -1.17 | 0.24 |
| Width basal angle | BAW | -0.58 | 0.56 |
| Length basal plate | BpL | -0.63 | 0.53 |
| Width between processes of basal plate | BBpW | -0.59 | 0.56 |

deeply convex, was situated on the ventral side and fitted 0.82 times in preoral length (POR) in S1 and 0.85 times in S2. The two groups shared the same teeth morphology (Fig. 3): teeth were similar in both jaws, looking small and compressed; the only sharp cuspid present seemed deeply turned towards the jaw termination, whereas the opposite margin appeared moderately rounded. Both

groups showed the same dental formula (12-13 / 12-13 in the upper jaw and 11-13 / 11-13 in the lower jaw).

Nostrils looked narrow, with well-developed nasal flaps. These structures, composed substantially by two lobes, were quite similar in the two groups with the external lobe considerably bigger than the internal one.

In both groups, the eye appeared relatively wide and




Fig. 3: Teeth of *Squalus sp.* from the Sardinian waters extracted from a S1 group male TL= 446 mm (teeth belonging to the higher and the lower jaw, A1 and A2 respectively) and a S2 group male TL= 470 mm (teeth belonging to the higher and the lower jaw, B1 and B2 respectively).




Fig. 4: Dermal denticles of *Squalus sp.* from the Sardinian waters. S1 group male TL= 552 mm (A) and a S2 group male TL= 634 mm (B). In both images an example of monocuspid (m) and tricuspid (t) typed denticle was highlighted.

more developed in length than in height; it fitted 4.70 and 4.54 times in head length (length at the 5th gill opening, PG5) for S1 and S2 group, respectively. The first dorsal fin was situated behind the pectoral fin and the pre-first dorsal length fitted 3.31 times in TL in S1 and 3.32 times in S2. In S1 and S2 groups, the first dorsal fin appeared more developed in length than in height; it fitted in length 1.79 times its height in both shark groups. Moreover, the first dorsal fin looked bigger than the second one, both

in length (1.26 times in S1 and 1.23 times in S2) and in height (1.87 times in S1 and 1.80 times in S2). The second dorsal fin length fitted 2.65 and 2.62 times its height in S1 and S2 respectively, looking mainly developed in length than in height, similarly to what was observed for the first dorsal fin. A strong spine with a triangular section was observed at the origin of each dorsal fin. The first dorsal spine length fitted 0.55 times in the fin base in both shark groups, while the second dorsal spine length

Table 5. Proportional dimensions expressed as percentages of TL (\pm SD) for specimens belonging to **S1** group, compared with what reported for *S. blainville* by other authors in other world regions.

| | | Muñoz-Chápuli and Ramos (1989) | | Marouani <i>et al.</i> (2012) | | Garrick (1960) | | Merrett (1973) | | Muñoz-Chápuli <i>et al.</i> (1984) | | Present study | |
|----------------------------|--------------|---------------------------------|-------------------------------|---|-------------------------------|----------------------|-------------------------------|---------------------------------|-------------------------------|------------------------------------|-------------------------------|-------------------------------|-------------------------------|
| | | Eastern Atlantic, Mediterranean | | Tunisian waters (Central Mediterranean) | | New Zealand | | Equatorial western Indian Ocean | | Mediterranean coasts of Spain | | Central Western Mediterranean | |
| | | <i>S. blainville</i> | | <i>S. blainville</i> | | <i>S. blainville</i> | | <i>S. blainville</i> | | <i>S. blainville</i> | | S1 | |
| N specimens | | 15 | | 9 | | 3 | | 4 | | 6 | | 19 | |
| Size range (mm, TL) | | 402-890 | | 630-960 | | 545-1008 | | 460-679 | | 560-730 | | 249-792 | |
| | Codex | N | Mean\pmSD | N | Mean\pmSD | N | Mean\pmSD | N | Mean\pmSD | N | Mean\pmSD | N | Mean\pmSD |
| Pre-inner nostril length | PNR | 15 | 3.41 \pm 0.65 | 9 | 4.30 \pm 0.15 | - | - | 4 | 4.22 \pm 0.45 | - | - | 16 | 2.31 \pm 0.35 |
| Preorbital length | POB | 15 | 5.55 \pm 0.78 | 9 | 6.19 \pm 0.53 | - | - | - | - | - | - | 16 | 4.44 \pm 0.44 |
| Preoral length | POR | 8 | 8.40 \pm 0.44 | 9 | 8.22 \pm 0.16 | - | - | 4 | 10.52 \pm 0.15 | - | - | 16 | 8.26 \pm 0.72 |
| Prebranchial length | PG1 | 15 | 16.68 \pm 0.78 | 9 | 17.02 \pm 0.59 | 3 | 17.16 \pm 0.25 | 4 | 19.5 \pm 0.4 | - | - | 16 | 15.47 \pm 1.01 |
| 5th gill opening | PG5 | 15 | 20.50 \pm 0.95 | - | - | - | - | - | - | - | - | 16 | 19.97 \pm 1.01 |
| Pre-first dorsal length | PD1 | 14 | 28.53 \pm 0.97 | 9 | 28.24 \pm 1.12 | 3 | 32.57 \pm 0.81 | 4 | 30.0 \pm 1.19 | - | - | 16 | 30.23 \pm 1.09 |
| Pre ventral length | SVL | 14 | 50.57 \pm 1.37 | 9 | 50.40 \pm 3.13 | 3 | 51 \pm 2.64 | 4 | 47.65 \pm 1.10 | - | - | 16 | 52.84 \pm 2.48 |
| Pre caudal length | PCL | 15 | 78.93 \pm 0.89 | 9 | 79.06 \pm 0.62 | 3 | 79.57 \pm 1.83 | 4 | 77.95 \pm 1.32 | - | - | 16 | 80.51 \pm 2.42 |
| Nostril-Labial furrow | NLF | - | - | - | - | - | - | - | - | - | - | 16 | 4.63 \pm 0.68 |
| Interdorsal space | IDS | 14 | 25.82 \pm 2.11 | 9 | 27.05 \pm 0.79 | 3 | 28.34 \pm 0.66 | 4 | 27.05 \pm 0.56 | - | - | 16 | 25.02 \pm 1.30 |
| Dorsal caudal space | DCS | 15 | 11.03 \pm 0.47 | 9 | 10.32 \pm 0.54 | 3 | 9.77 \pm 0.65 | 4 | 10.87 \pm 0.5 | - | - | 16 | 11.03 \pm 0.75 |
| Pectoral-pelvic space | PPS | 15 | 22.89 \pm 1.55 | 9 | 21.75 \pm 0.86 | - | - | - | - | - | - | 16 | 24.69 \pm 1.47 |
| Pelvic and caudal | PCA | 12 | 27.26 \pm 1.13 | - | - | - | - | - | - | - | - | 16 | 28.48 \pm 3.00 |
| Internarial space | INW | 15 | 4.53 \pm 0.48 | 9 | 4.16 \pm 0.26 | - | - | - | - | 6 | 4.48 \pm 0.29 | 16 | 4.59 \pm 0.43 |
| Between outer corners | ONW | 15 | 6.79 \pm 0.59 | - | - | - | - | - | - | - | - | 16 | 8.28 \pm 0.86 |
| Nostril length | NOW | 15 | 1.48 \pm 0.28 | - | - | - | - | - | - | - | - | 16 | 1.87 \pm 0.22 |
| Mouth width | MOW | 15 | 7.49 \pm 0.89 | 9 | 7.29 \pm 0.53 | - | 5.83 \pm 0.11 | 4 | 6.72 \pm 0.7 | - | - | 16 | 10.08 \pm 0.83 |
| Length of preoral cleft | MOL | 15 | 2.85 \pm 0.92 | - | - | - | - | - | - | - | - | 16 | 2.33 \pm 0.33 |
| Eye length | EYL | 15 | 4.03 \pm 0.39 | 9 | 3.86 \pm 0.23 | 3 | 4.37 \pm 0.21 | 4 | 5.22 \pm 0.12 | - | - | 16 | 4.24 \pm 0.42 |
| Spiracles | | | | | | | | | | | | | |

(continued)

Table 5 continued

| | | Muñoz-Chápuli and Ramos (1989) | | Marouani <i>et al.</i> (2012) | | Garrick (1960) | | Merrett (1973) | | Muñoz-Chápuli <i>et al.</i> (1984) | | Present study | |
|---|--------------|---------------------------------|----------------|---|----------------|----------------------|----------------|---------------------------------|----------------|------------------------------------|----------------|-------------------------------|----------------|
| | | Eastern Atlantic, Mediterranean | | Tunisian waters (Central Mediterranean) | | New Zealand | | Equatorial western Indian Ocean | | Mediterranean coasts of Spain | | Central Western Mediterranean | |
| | | <i>S. blainville</i> | | <i>S. blainville</i> | | <i>S. blainville</i> | | <i>S. blainville</i> | | <i>S. blainville</i> | | S1 | |
| N specimens | | 15 | | 9 | | 3 | | 4 | | 6 | | 19 | |
| Size range (mm, TL) | | 402-890 | | 630-960 | | 545-1008 | | 460-679 | | 560-730 | | 249-792 | |
| | Codex | N | Mean±SD | N | Mean±SD | N | Mean±SD | N | Mean±SD | N | Mean±SD | N | Mean±SD |
| Distance between tips | ISP | 15 | 8.11±0.90 | - | - | - | - | - | - | - | - | 16 | 8.45±0.32 |
| Gills | | | | | | | | | | | | | |
| First gill-slit height | GS1 | 15 | 1.95±0.23 | 9 | 1.85±0.26 | 3 | 1.9±0.42 | 4 | 1.77±0.27 | - | - | 16 | 2.01±0.25 |
| Third gill-slit height | GS3 | 15 | 2.23±0.26 | 9 | 2.20±0.17 | - | - | - | 1.85±0.31 | - | - | 16 | 2.11±0.30 |
| Fifth gill-slit height | GS5 | 15 | 2.49±0.42 | 9 | 2.07±0.27 | 3 | 2.33±0.21 | 4 | 2.04±0.12 | - | - | 16 | 2.26±0.21 |
| Intergill length (1 st and 5 th) | ING | 15 | 4.18±0.63 | 9 | 4.66±0.67 | - | - | - | - | - | - | 16 | 4.47±0.39 |
| First dorsal fin | | | | | | | | | | | | | |
| First dorsal length | D1L | - | - | 9 | 13.32±0.76 | - | - | - | - | - | - | 16 | 13.55±0.55 |
| First dorsal base length | D1B | 14 | 8.44±1.54 | 9 | 8.03±0.25 | 3 | 6.07±0.68 | 4 | 7.22±0.54 | - | - | 16 | 7.63±0.41 |
| First dorsal height | D1H | 15 | 8.09±0.61 | 9 | 7.07±0.7 | 3 | 8.03±0.15 | 4 | 8.6±0.91 | - | - | 16 | 7.56±0.66 |
| First dorsal inner margin | D1I | 15 | 6.10±0.53 | 9 | 5.40±0.28 | - | - | - | - | - | - | 16 | 5.87±0.44 |
| First dorsal spine length | D1ES | 14 | 4.32±0.71 | 9 | 5.06±0.3 | - | - | - | 4.15±1.30 | - | - | 16 | 4.19±0.44 |
| Second dorsal fin | | | | | | | | | | | | | |
| Second dorsal length | D2L | - | - | 9 | 9.45±0.31 | - | - | - | - | - | - | 16 | 10.73±1.27 |
| Second dorsal base length | D2B | 14 | 6.42±1.27 | 9 | 5.13±0.41 | 3 | 4.8±0.78 | 4 | 4.55±0.25 | - | - | 16 | 5.94±0.63 |
| Second dorsal height | D2H | 15 | 4.46±0.56 | - | - | - | - | - | - | - | - | 16 | 4.04±0.58 |
| Second dorsal inner margin | D2I | 15 | 4.79±0.46 | 9 | 4.29±0.21 | - | - | 4 | 4.2±1.01 | - | - | 16 | 4.57±0.89 |
| Second dorsal spine length | D2ES | 13 | 4.92±0.94 | 9 | 5.22±0.41 | - | - | - | - | 6 | 4.69±0.45 | 16 | 5.89±0.57 |
| Pectoral fin | | | | | | | | | | | | | |
| Pectoral length | P1L | - | - | 9 | 11.46±0.52 | - | - | - | - | - | - | 16 | 14.62±1.61 |
| Pectoral base length | P1B | 15 | 6.77±0.70 | 9 | 5.85±0.41 | - | - | 4 | 5.7±0.46 | - | - | 16 | 4.94±0.44 |
| Pectoral anterior margin | P1A | 15 | 13.99±1.02 | 9 | 13.31±0.95 | 3 | 14.43±0.91 | 4 | 15.05±0.91 | 6 | 13.63±0.85 | 16 | 13.20±0.51 |

(continued)

Table 5 continued

| | | Muñoz-Chápuli and Ramos (1989) | Marouani <i>et al.</i> (2012) | Garrick (1960) | Merrett (1973) | Muñoz-Chápuli <i>et al.</i> (1984) | Present study | | |
|---------------------------|-------|---------------------------------|---|----------------------|---------------------------------|------------------------------------|-------------------------------|----|------------|
| | | Eastern Atlantic, Mediterranean | Tunisian waters (Central Mediterranean) | New Zealand | Equatorial western Indian Ocean | Mediterranean coasts of Spain | Central Western Mediterranean | | |
| | | <i>S. blainville</i> | <i>S. blainville</i> | <i>S. blainville</i> | <i>S. blainville</i> | <i>S. blainville</i> | S1 | | |
| N specimens | | 15 | 9 | 3 | 4 | 6 | 19 | | |
| Size range (mm, TL) | | 402-890 | 630-960 | 545-1008 | 460-679 | 560-730 | 249-792 | | |
| | Codex | N | Mean±SD | N | Mean±SD | N | Mean±SD | N | Mean±SD |
| Pectoral posterior margin | P1P | 15 | 11.10±0.80 | 9 | 11.40±0.90 | | | 16 | 11.29±0.82 |
| Pectoral inner margin | P1I | 15 | 7.18±0.51 | 9 | 6.24±0.36 | | | 6 | 7.08±0.39 |
| Pelvic fin | | | | | | | | | |
| Pelvic anterior margin | P2A | 15 | 5.86±0.72 | 9 | 4.76±0.90 | 4 | 5.75±0.36 | 16 | 6.42±1.06 |
| Pelvic Length | P2L | 15 | 9.69±0.68 | 9 | 9.05±1.49 | | | 6 | 9.82±0.84 |
| Caudal fin | | | | | | | | | |
| Dorsal caudal margin | CDM | 15 | 21.10±0.54 | 9 | 20.74±0.90 | | | 16 | 19.77±1.55 |
| Preventral caudal margin | CPV | 14 | 11.08±0.70 | 9 | 10.15±0.99 | | | 16 | 9.78±0.67 |
| Trunk at pectoral origin: | | | | | | | | | |
| Trunk width | TRW | 8 | 11.72±0.94 | 9 | 10.00±0.94 | | | 16 | 12.88±0.71 |

appeared as long as the fin base, fitting 0.99 and 0.98 times the second dorsal base in S1 and S2, respectively. The interdorsal space fitted 1.21 times the pre-first dorsal length in S1 and 1.22 times in S2. The pectoral fins of both groups presented a large and almost straight anterior margin, culminating with a deeply rounded apex and their inner margin ends with a small rounded tip. The pectoral fin base fitted 2.67 and 2.68 times in the anterior margin length in S1 and S2, respectively. The pelvic fins instead were small and triangular with a rounded apex and almost straight anterior and posterior margin. The pelvic fin anterior margin fitted 1.82 times in fin length in S1 and 1.83 times in S2. The caudal peduncle appeared well developed with two solid lateral keels that origins behind the second dorsal base termination and ends below caudal fin insertion. The dorsal-caudal space fitted 2.27 times in the interdorsal space in S1 and 2.23 times in S2. The caudal fin presented an extended dorsal caudal margin (19.77 in %TL in S1; 20.41 in %TL in S2) without sub-terminal notch.

Both spurdog groups showed a uniform grey-brown coloration on the dorsal side while the ventral one and all the fins rear margins appeared paler. The eyes were bright

green when observed in live specimens. In Figure 4 dermal denticles obtained from S1 (Fig.4a) and S2 (Fig. 4b) specimens are showed. These structures appeared mostly monocuspid typed but, in every group it was possible to simultaneously discriminate some tricuspid denticles.

Analysis of Principal Coordinates CAP

The bi-plot produced after CAP analysis emphasized no clear segregation among the *a priori* assigned groups (S1 and S2), with a higher overlapping for the chondrocranium parameters compared to the somatic ones (Fig. 5b and Fig. 5a respectively). The cross-validation also showed an elevated percentage of misclassification (i.e., 41.03% for chondrocranium and 37.5 for somatic), further confirming that a considerable portion of samples did not follow the *a priori* grouping.

Genetic analysis

A 609 bp fragment of COI gene was obtained for the 18 individuals revealing a total of 7 haplotypes (Hd: 0.765), differing in 6 nucleotide positions (π : 0.00245).

Table 6. Proportional dimensions expressed as percentages of TL (\pm SD) for specimens belonging to **S2** group, compared with what reported for *S. megalops* by other authors in other world regions.

| N specimens | Marouani <i>et al.</i> (2012) | | Muñoz-Chápuli & Ramos (1989) | | Last <i>et al.</i> (2007) | | Last <i>et al.</i> (2007) | | Last <i>et al.</i> (2007) | | Chápuli <i>et al.</i> (1984) | | Present study | | |
|---------------------|---|---------|---------------------------------|------------------|---------------------------|-----------|---------------------------|-----------|---------------------------|-----------|----------------------------------|------------------|--------------------|---------------|-------------------------------|
| | Mean | Min-Max | Mean \pm SD | Mean | Min-Max | Mean | Min-Max | Mean | Min-Max | Mean | Min-Max | Mean \pm SD | Mean \pm SD | Mean \pm SD | S2 |
| Size range (mm, TL) | 330-695 | 9 | 318-742 | 34 | 373-527 | 6 | 328-384 | 3 | 414-541 | 4 | 485-680 | 24 | 118 | 207-834 | |
| | <i>S. megalops</i> | | <i>S. megalops</i> | | <i>S. megalops</i> | | <i>S. megalops</i> | | <i>S. megalops</i> | | <i>S. megalops</i> | | <i>S. megalops</i> | | |
| | Tunisian waters (Central Mediterranean) | | Eastern Atlantic, Mediterranean | | Southeastern Australia | | Queensland | | Western Australia | | Mediterranean, coasts of Morocco | | | | Central Western Mediterranean |
| Codex | PNR | 4.23 | 3.88-4.48 | 3.49 \pm 0.57 | 3.9 | 3.7-4.1 | 4.3 | 4.2-4.4 | 4.2 | 3.9-4.4 | - | 2.47 \pm 0.46 | | | |
| | POB | 6.83 | 6.18-7.27 | 6.01 \pm 0.51 | 7 | 6.4-7.5 | 7.2 | 7-7.4 | 7 | 6.4-7.4 | - | 4.53 \pm 0.73 | | | |
| | POR | 8.84 | 7.98-9.39 | 8.65 \pm 0.83 | 9.1 | 8.6-9.9 | 9.7 | 9.3-9.9 | 9.2 | 8.9-9.7 | - | 8.84 \pm 1.77 | | | |
| | PG1 | 17.88 | 16.76-19.79 | 16.62 \pm 0.88 | 18.5 | 17.7-19.8 | 18.9 | 18.6-19.2 | 18.3 | 17.8-19.1 | - | 15.89 \pm 1.68 | | | |
| | PG5 | - | - | 19.97 \pm 1.03 | - | - | - | - | - | - | - | 20.19 \pm 1.61 | | | |
| | PDI | 29.48 | 28.41-30.70 | 28.99 \pm 0.89 | 30.2 | 29.1-31.6 | 30.6 | 29.9-31.6 | 29.6 | 29.1-30.2 | - | 30.07 \pm 2.26 | | | |
| | SVL | 48.32 | 46.29-49.85 | 49.18 \pm 2.06 | 48.5 | 47.6-50.1 | 46.5 | 46.1-47.2 | 47.9 | 45.9-50.4 | - | 53.33 \pm 3.30 | | | |
| | PCL | 78.69 | 76.95-80.34 | 78.92 \pm 1.08 | 77.7 | 76.1-79.3 | 78.5 | 77.8-78.9 | 78.4 | 77.7-79.2 | - | 80.12 \pm 3.69 | | | |
| | NLF | | | | | | | | | | | | | | |
| | IDS | 25.42 | 22.77-27.59 | 24.49 \pm 1.9 | 24.8 | 24-25.3 | 24.6 | 23.2-25.8 | 25.3 | 23.7-26 | - | 24.73 \pm 1.89 | | | |
| | DCS | 10.93 | 9.49-11.81 | 11.16 \pm 0.60 | 10.4 | 9.5-10.9 | 12.2 | 11.5-12.7 | 10.7 | 9.9-12 | - | 11.09 \pm 1.08 | | | |
| | PPS | 21.72 | 20.29-23.46 | 23.10 \pm 2.64 | 22.3 | 20.9-26.1 | 19.1 | 18-20.3 | 22.6 | 20.5-24.6 | - | 24.42 \pm 2.29 | | | |
| | PCA | | | 27.73 \pm 1.44 | | | | | | | | 27.91 \pm 2.49 | | | |
| | INW | 3.98 | 3.71-4.19 | 3.83 \pm 0.26 | 4.5 | 4.3-4.7 | 4.7 | 4.6-4.9 | 4.5 | 4.2-4.8 | 3.82 \pm 0.20 | 4.66 \pm 0.44 | | | |
| | ONW | | | 6.68 \pm 0.57 | | | | | | | | 8.39 \pm 0.81 | | | |
| | NOW | | | 1.60 \pm 0.20 | | | | | | | | 1.89 \pm 0.26 | | | |

(continued)

Table 6 continued

| N specimens | Marouani <i>et al.</i> (2012) | | Muñoz-Chápuli & Ramos (1989) | | Last <i>et al.</i> (2007) | | Last <i>et al.</i> (2007) | | Last <i>et al.</i> (2007) | | Chápuli <i>et al.</i> (1984) | | Present study | |
|---|-------------------------------|-------------|-------------------------------|---------|-------------------------------|---------|-------------------------------|---------|-------------------------------|---------|-------------------------------|---------|-------------------------------|----|
| | Mean | Min-Max | Mean | Min-Max | Mean | Min-Max | Mean | Min-Max | Mean | Min-Max | Mean | Min-Max | Mean | SD |
| Size range (mm, TL) | <i>S. megalops</i> 330-695 | | <i>S. megalops</i> 318-742 | | <i>S. megalops</i> 373-527 | | <i>S. megalops</i> 328-384 | | <i>S. megalops</i> 414-541 | | <i>S. megalops</i> 485-680 | | <i>S. megalops</i> 207-834 | |
| Mouth width | 7.86 | 7.48-8.53 | 7.69-0.45 | 8.1 | 7.8-8.6 | 8.3 | 8-8.5 | 8.2 | 7.8-8.6 | - | - | - | 10.36±1.11 | |
| Length of preoral cleft | - | - | 3.30±1.10 | - | - | - | - | - | - | - | - | - | 2.50±0.49 | |
| Eye length | 4.19 | 3.59-4.84 | 4.27±0.53 | 4.8 | 4.4-5.4 | 5 | 4.9-5 | 4.8 | 4.3-5.3 | - | - | - | 4.44±0.68 | |
| Spiracles | | | | | | | | | | | | | | |
| Distance between tips | | | 8.10±0.60 | | | | | | | | | | 8.55±0.50 | |
| Gills | | | | | | | | | | | | | | |
| First gill-slit height | 1.99 | 1.51-2.60 | 2.00±0.30 | 2.3 | 2-2.4 | 1.9 | 1.8-1.9 | 2.2 | 1.9-2.4 | - | - | - | 2.10±0.32 | |
| Third gill-slit height | | | 2.09±0.24 | | | | | | | | | | 2.16±0.31 | |
| Fifth gill-slit height | 2.09 | 1.76-2.53 | 2.46±0.39 | 2.4 | 2.1-2.5 | 2.5 | 2.3-2.6 | 2.2 | 1.8-2.4 | - | - | - | 2.32±0.29 | |
| Intergill length (1 st and 5 th) | | | 4.48±0.92 | | | | | | | | | | 4.39±0.57 | |
| First dorsal fin | | | | | | | | | | | | | | |
| First dorsal length | 13.56 | 13.09-14.14 | - | 14.4 | 13.8-15.1 | 13.3 | 12.7-13.7 | 14 | 13.3-14.9 | - | - | - | 13.56±1.10 | |
| First dorsal base length | 7.63 | 7.27-8.02 | 8.06±0.75 | 8.2 | 7.9-8.9 | 7.6 | 7.2-8 | 8.3 | 7.7-8.9 | | | | 7.63±0.76 | |
| First dorsal height | 6.06 | 5.60-6.56 | 8.48±0.81 | 7 | 6.1-7.4 | 6.4 | 6.2-6.6 | 7.2 | 7-7.5 | | | | 7.57±0.88 | |
| First dorsal inner margin | 5.68 | 5.12-6.25 | 6.72±0.52 | 6.3 | 6.1-6.6 | 5.7 | 5.7-5.7 | 5.9 | 5.4-6.3 | | | | 5.85±0.74 | |
| First dorsal spine length | 4.58 | 4.09-5.08 | 4.63±0.49 | 3 | 2.4-3.3 | 3 | 2.9-3.2 | 3.3 | 3-3.4 | | | | 4.20±0.65 | |
| Second dorsal fin | | | | | | | | | | | | | | |
| Second dorsal length | 10.26 | 9.10-11.60 | - | 12 | 11-12.7 | 12.1 | 11.6-12.8 | 12.2 | 11.8-12.8 | | | | 11.01±0.97 | |
| Second dorsal base length | 5.28 | 4.64-6.13 | 6.98±1.40 | 7.1 | 6.4-7.5 | 7.2 | 6.9-7.6 | 7.5 | 7.1-8.2 | | | | 6.07±0.73 | |
| Second dorsal height | 3.49 | 3.03-4.23 | 5.20±1.1 | 4 | 3.6-4.6 | 3.7 | 3.2-4 | 3.9 | 3.7-4.3 | | | | 4.19±0.55 | |
| Second dorsal inner margin | 4.97 | 4.47-5.50 | 5.47±0.38 | 4.9 | 4.5-5.3 | 4.9 | 4.5-5.1 | 4.9 | 4.7-5 | | | | 4.70±0.70 | |
| Second dorsal spine length | 5.98 | 5.33-5.56 | 5.37±0.92 | 4.3 | 3.6-5 | 4.6 | 4-5 | 4.5 | 4.2-4.6 | | | | 6.00±0.67 | |

(continued)

Table 6 continued

| N specimens Size range (mm, TL) | Marouani <i>et al.</i> (2012) Tunisian waters (Central Mediterranean) | | Muñoz-Chápuli & Ramos (1989) Eastern Atlantic, Mediterranean | | Last <i>et al.</i> (2007) Southeastern Australia | | Last <i>et al.</i> (2007) Queensland | | Last <i>et al.</i> (2007) Western Australia | | Chápuli <i>et al.</i> (1984) Mediterranean, coasts of Morocco | | Present study Central Western Mediterranean | |
|------------------------------------|---|--------------------|---|--------------------|---|--------------------|---|--------------------|--|--------------------|--|--------------------|--|--------------------|
| | <i>S. megalops</i> | <i>S. megalops</i> | <i>S. megalops</i> | <i>S. megalops</i> | <i>S. megalops</i> | <i>S. megalops</i> | <i>S. megalops</i> | <i>S. megalops</i> | <i>S. megalops</i> | <i>S. megalops</i> | <i>S. megalops</i> | <i>S. megalops</i> | <i>S. megalops</i> | <i>S. megalops</i> |
| | 9 | 34 | 6 | 3 | 4 | 3 | 3 | 4 | 24 | 24 | 118 | 207-834 | | |
| | 330-695 | 318-742 | 373-527 | 328-384 | 414-541 | 485-680 | | | | | | | | |
| | Mean | Mean±SD | Mean | Mean | Mean | Mean | Mean | Mean | Mean | Mean | Mean±SD | Mean±SD | Mean±SD | Mean±SD |
| | Min-Max | | Min-Max | Min-Max | Min-Max | Min-Max | Min-Max | Min-Max | Min-Max | Min-Max | | | | |
| Pectoral fin | | | | | | | | | | | | | | |
| Pectoral length | 12.95 | 11.94-13.63 | | | | | | | | | | | | 14.59±1.23 |
| Pectoral base length | 5.61 | 5.15-6.22 | 5.3 | 4.4-5.7 | 5.3 | 4.9-5.8 | 4.9 | 4.4-5.3 | - | - | | | | 4.94±0.59 |
| Pectoral anterior margin | 14.12 | 13.03-15.25 | 14.3 | 13.6-14.9 | 12.5 | 12.3-12.6 | 14.3 | 13.7-15.1 | 15.26±0.96 | | | | | 13.28±1.10 |
| Pectoral posterior margin | 11.31 | 10.17-12.32 | 11.6 | 10.8-12.7 | 10.4 | 9.6-10.9 | 11.2 | 10.3-12.3 | | | | | | 11.21±0.94 |
| Pectoral inner margin | 7.40 | 6.61-8.26 | 8.2 | 7.4-9.2 | 8.4 | 7.7-8.8 | 9 | 8.4-9.7 | | | | | | 10.76±0.99 |
| Pelvic fin | | | | | | | | | | | | | | |
| Pelvic anterior margin | 5.24 | 5.02-5.72 | 6.36±0.65 | | | | | | | | | | | 6.34±1.39 |
| Pelvic Length | 10.38 | 9.56-11.32 | 11.21±1.15 | 10.5 | 9.9-11.5 | 10.6 | 9.9-11.2 | 10.4 | 9.9-10.8 | 11.55±0.95 | | | | 11.61±0.73 |
| Caudal fin | | | | | | | | | | | | | | |
| Dorsal caudal margin | 19.43 | 16.38-21.82 | 21.36±0.76 | 20.9 | 20-21.4 | 20.1 | 19.3-20.9 | 20.6 | 20.2-21.1 | | | | | 20.41±1.52 |
| Preventral caudal margin | 9.54 | 8.45-10.80 | 11.39±1.14 | 11 | 10.5-11.3 | 10.6 | 10.4-10.7 | 10.9 | 10.7-11 | | | | | 10.02±0.84 |
| Trunk at pectoral origin: | | | | | | | | | | | | | | |
| Trunk width | 10.36 | 8.55-11.84 | 11.29±0.59 | 12.1 | 11.2-13.2 | 10.8 | 10.3-11.7 | 12.2 | 11.3-14.5 | | | | | 12.85±1.09 |




Fig. 5: Biplot produced using CAP analysis for somatic (A) and chondrocranial (B) measurements.

All the newly generated COI sequences were deposited in GenBank (Accession numbers: MF828596-MF828613). The haplotype network (Fig. 6) showed the occurrence of two common haplotypes H1 and H3 (shared by 6 and 7 individuals, respectively), and five private haplotypes (H2, H4-H7). Both haplotypes H1 and H3 were shared by individuals with single and double lateral processes.

In the Bayesian tree (Fig. 7), the sequences of *Squalus* clustered in ten highly supported clades (c1-c10). All the newly obtained sequences of the Sardinian *S. blainville* were in the same clade (c2) with *S. blainville* individuals from the Mediterranean and the Eastern Atlantic. They were clearly different from the sequences of *S. megalops* (c3), *S. raoulensis* (c4), *S. brevirostris* (c5), and *S. cubensis* (c7). Clade c6 comprised a single divergent sequence of a specimen from Libya, originally identified as *S. blainville* (Kousteni *et al.*, 2016). Besides, sequences from Sardinia were strongly divergent from sequences in clades c1 (*S. acanthias* and *S. suckleyi*), and c8-c10 (individuals originally identified as *S. megalops* or *S. blainville* but belonging to distinct species still waiting formal description; Verissimo *et al.*, 2017).




Fig. 6: Median Joining network of the COI haplotypes. Each circle represents a haplotype and the area of the circle is proportional to the haplotype frequency. In white are shown the sequences of individuals with a single lateral process (S1), and in grey the sequences of individuals with double lateral processes (S2). All mutational steps are equal to 1 and represented with a vertical line. Haplotype codes correspond to sequences MF828596-MF828613 (Table S1).

Discussion

In the present study, although the observation of chondrocranial lateral processes initially allowed the investigated specimens to be subdivided into two groups, both morphological and genetic analysis revealed the presence of only one spurdog species in the Sardinian waters, the longnose spurdog (*S. blainville*). Indeed, the comparison of chondrocranial and body morphology of the spurdog specimens examined indicated that none of the considered measurements could discriminate the two squalid groups.

Comparing our results with the available data from literature, chondrocranial morphological measurements recorded in the present study were mostly coherent with others reported for *S. blainville* in other Mediterranean areas. The length of the precerebral fenestra (LPF) represented the only exception, which was smaller for both groups.

As far as the somatic data is concerned, in general, no major differences were found except for few measurements, regarding in particular the head and snout region. Indeed, for the S1 group pre-inner nostril length (PNR) and preorbital length (POB) appeared minor in terms of %TL than what was reported for *S. blainville* in the Mediterranean Sea by Muñoz-Chápuli *et al.* (1984); Muñoz-Chápuli & Ramos (1989) and Marouani *et al.* (2012), in the New Zealand waters (Garrick, 1960) and the




Fig. 7: Bayesian tree based on mitochondrial COI sequences. Bayesian posterior probabilities are next to the nodes. Clade c2, containing all the new Sardinian sequences is highlighted in black. Table S1 contains the complete list of sequences used.

Equatorial Western Indian Ocean (Merret, 1973). On the other hand, mouth width (MOW) of the Sardinian specimens looked larger compared to that reported in previous studies (Garrick, 1960; Merret, 1973; Muñoz-Chápuli & Ramos, 1989; Marouani *et al.*, 2012). The same situation occurred for the outer nostril corner width (ONW), even if it was possible to compare our results only with what was described by Marouani *et al.* (2012) because this measurement was not reported in the other cited papers. The only other relevant differences in the body morphology with respect to what is reported in literature for *S. blainville* were found in the pectoral fin measurements, more precisely in pectoral fin length (PIL) and pectoral inner margin (PII), both achieving higher values than those documented by other authors (Garrick, 1960; Merret, 1973; Muñoz-Chápuli & Ramos, 1989; Marouani *et al.*, 2012). Unfortunately, also in this case, it was possible to compare PIL values of Sardinian spurdogs only with Tunisian data (Marouani *et al.*, 2012).

Exactly the same situation occurred for the S2 group, in which the only relevant differences between the samples analysed in this study and what was described for *S. megalops* by other authors in Eastern Atlantic and Mediterranean Sea (Muñoz-Chápuli *et al.*, 1984; Muñoz-Chápuli & Ramos, 1989; Marouani *et al.*, 2012) and in Australian waters (Last *et al.*, 2007) coincided precisely with the same measurements previously reported for sharks belonging to the S1 group.

Besides the exact correspondence in the two shark groups of the morphological characters (both somatic and chondrocranial) that have reported disagreeing values from the literature could be a further indication of the presence of only one species.

Moreover, the observation of further characteristics, identified by other authors as different in the two spurdog species, such as teeth and dermal denticles, were not able to clearly discriminate the groups. In particular, S1 and S2 presented very similar teeth in both upper and lower dental arches. Furthermore, regarding the dermal denticles, every specimen analysed in this work presented, at the same time, both denticle shapes described as typical for *S. blainville* (tricuspid) and for *S. megalops* (monocuspid) (Muñoz-Chápuli *et al.*, 1984; Muñoz-Chápuli & Ramos, 1989; Marouani *et al.*, 2012). Considering the brief half-life and fast replacing rate of these structures (Kemp, 1999), this particular aspect could be due to a different development stage of denticles observed in the analysed skin portion (Kemp, 1999). Moreover, it is reported that some common diagnostic morphological features, such as dermal denticles, teeth and dorsal fin spines could vary in shape with the ontogenetic development (White *et al.*, 2013; Veríssimo *et al.*, 2014). Consequently, the dermal denticles morphology should be further investigated before it can be properly used as a suitable classification tool, as also suggested by Bonello *et al.* (2016) particularly for the genus *Squalus*.

All the specimens genetically analyzed in Sardinia, despite their morphological variability, clustered together, and resulted to be *S. blainville*. Both present and previous genetic data confirm that this taxon is widely distributed in the Mediterranean (Serena, 2005; Bat *et al.*, 2005; Serena *et al.*, 2009; Landi *et al.*, 2014; Bonello *et al.*, 2016; Kousteni *et al.*, 2016; Cariani *et al.*, 2017; Veríssimo *et al.*, 2017).

However, several taxonomic uncertainties still remain in this region with respect to the occurrence and distribution of additional *Squalus* species besides *S. blainville* and *S. acanthias*.

Recently, several studies highlighted the frequent misidentification of *Squalus* taxa in this area, and the inconsistent use of the names *S. blainville* and *S. megalops*, and even of *S. acanthias* (see Cariani *et al.*, 2017; Veríssimo *et al.*, 2017 and Table S1). For instance, the sequence available for a Mediterranean specimen originally identified as *S. megalops* (Marouani *et al.*, 2012) proved to be *S. blainville* (Veríssimo *et al.*, 2017). However, considering the finding of sporadic divergent sequences (Fig. 9 c6 and c8; Marouani *et al.*, 2012; Kousteni *et al.*, 2016; Veríssimo *et al.*, 2017) different from *S. blainville* (Fig. 9 c2), *S. acanthias* (Fig. 9 c1) but also *S. megalops* from Australia (Fig. 9 c3), the occurrence of a third species in the Mediterranean (apart from *S. acanthias* and *S. blainville*) cannot be ruled out.

In particular, the second sequence by Marouani *et al.* (2012) from a Mediterranean (Tunisian) specimen originally identified as *S. blainville*, clustered in c8 with indi-

viduals from Tropical West Africa, originally identified as *S. megalops* (Fig. 9 c8 or clade C *sensu* Veríssimo *et al.*, 2017). Nevertheless, as *S. megalops* is to be applied only to Australian spurdogs (Veríssimo *et al.*, 2017), which taxon name is to be used for the specimens with eastern Atlantic and Mediterranean origin remains uncertain (Veríssimo *et al.*, 2017).

The genetic and morphological analysis carried out in the present paper indicated the presence of only one spurdog species in Sardinian waters, ascribable to *S. blainville*. These results represent an important baseline for future assessment and management studies on Central-Western Mediterranean spurdog populations. However, considering the taxonomical confusion that characterizes the *Squalus* genus and the fact that a classification based only on morphological features can easily lead to misidentifications, as demonstrated in the present paper, additional studies combining genetics and morphology are welcomed and urgent.

References

- Anderson, M., Willis, T., 2003. Canonical analysis of principal coordinates: a useful method of constrained ordination for ecology. *Ecology*, 84, 511-525.
- Bandelt, H.J., Forster, P., Röhl, A., 1999. Median-joining networks for inferring intraspecific phylogenies. *Molecular Biology and Evolution*, 16, 37-48.
- Bat, L., Erdem, Y., Ustaoglu, S., Yardim, Ö., Satilmis, H.H., 2005. A Study on the Fishes of the Central Black Sea Coast of Turkey. *Journal of the Black Sea/Mediterranean Environment*, 11, 281-296.
- Bertrand, J. A., Gil de Sola, L., Papaconstantinou, C., Relini, G., Souplet, A., 2000. An international bottom trawl survey in the Mediterranean: the MEDITS programme. p. 76-93. In: *Demersal resources in the Mediterranean*. Bertrand J.A., Relini, G. (eds), *Proceedings of the Symposium, Pisa, 18-21 March 1998*. Actes de Colloques, 26. IFREMER, Plouzané. The general specifications of MEDITS surveys.
- Bigelow, H.B., Schroeder, W.C., 1957. A study of the sharks of the suborder Squaloidea. *Bulletin of the Museum of Comparative Zoology, Cambridge, Massachusetts*, 117, 1-150.
- Bonello, J., Bonnici, L., Ferrari, A., Cariani, A., Schembri, P., 2016. Not all that clear cut: Intraspecific morphological variability in *Squalus blainville* (Risso, 1827) and implications for identification of the species. *Journal of the Marine Biological Association of the United Kingdom*, 96, 1585-1596.
- Cariani, A., Messinetti, S., Ferrari, A., Arculeo, M., Bonello, J.J. *et al.*, 2017. Improving the Conservation of Mediterranean Chondrichthyan: The ELASMOMED DNA Barcode Reference Library. *PLoS One*, 12(1), e0170244. doi:10.1371/journal.pone.0170244
- Coelho, R., Erzini, K., 2008. Identification of deep water lantern sharks (Chondrichthyes: Etmopteridae) using morphometric data and multivariate analysis. *Journal of the Marine Biological Association of the United Kingdom*, 88, 199-204.
- Compagno, L.J.V., 2001. *Sharks of the world. An annotated and illustrated catalogue of shark species known to date. Vol. 2. Bullhead, mackerel and carpet sharks (Heterodontiformes, Lamniformes and Orectolobiformes)*. Report No. 1, vol. 2. Food and Agriculture Organisation of United Nations, Rome, 269 pp.
- Compagno, L.J.V., Dando, M., Fowler, S.L., 2005. *Sharks of the World*. Princeton University Press, 368 pp.
- Ebert, D.A., Fowler, S., Compagno, L.J.V., 2013. *Sharks of the world. A fully illustrated guide*. Wild Nature Press, Plymouth UK, 528 pp.
- Ebert, D.A., White, W.T., Goldman, K.J., Compagno, L.J.V., Daly-Engel, T.S. *et al.*, 2010. Resurrection and redescription of *Squalus suckleyi* (Girard, 1854) from the North Pacific, with comments on the *Squalus acanthias* subgroup (Squaliformes: Squalidae). *Zootaxa*, 2612, 22-40.
- Folmer, O., Black, M., Hoeh, W., Lutz, R., Vrijenhoek, R., 1994. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology*, 3, 294-299.
- Garrick, J.A.F., 1960. Studies on New Zealand Elasmobranchii. Part XII. The species of *Squalus* from New Zealand and Australia; and a general account and key to the New Zealand Squaloidea. *The Transactions and Proceedings of the Royal Society of New Zealand*, 88, 519-557.
- Huelsenbeck, J.P., Ronquist, F., 2001. MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics*, 17, 754-755.
- Kemp, N.E., 1999. Integumentary system and teeth. Chapter 2. p. 43-68. In: *Sharks, skates and rays: the biology of Elasmobranch fishes*. Hamlett, W.C. (Ed). The John Hopkins University Press.
- Kousteni, V., Kasapidis, P., Kotoulas, G., Megalofonou, P., 2016. Evidence of high genetic connectivity for the long-nose spurdog *Squalus blainville* in the Mediterranean Sea. *Mediterranean Marine Science*, 17(2), 371-383.
- Landi, M., Dimech, M., Arculeo, M., Biondo, G., Martins, R. *et al.*, 2014. DNA Barcoding for Species Assignment: The Case of Mediterranean Marine Fishes. *PLoS One*, 9, e106135. doi: 10.1371/journal.pone.0106135
- Last, P.R., White, W.T., Pogonoski, J.J., Gledhill, D.C., Yearsley, G.K. *et al.*, 2007. Application of a rapid taxonomic approach to genus *Squalus*. p. 1-10. In: *Description of new dogfishes of the genus Squalus (Squaloidea: Squalidae)*. Last, P.R., White, W.T., Pogonoski, J.J. (Eds) C.M.A.R.
- Librado, P., Rozas, J., 2009. DnaSP v5: a software for comprehensive analysis of DNA polymorphism data. *Bioinformatics*, 25, 1451-1452.
- Marouani, S., Chaâba, R., Kadri, H., Saidi, B., Bouain, A. *et al.*, 2012. Taxonomic research on *Squalus megalops* (Macleay, 1881) and *Squalus blainvillei* (Risso, 1827) (Chondrichthyes: Squalidae) in Tunisian waters (central Mediterranean Sea). *Scientia Marina*, 76, 97-109.
- Merrett, M.R., 1973. A new shark of the genus *Squalus* (Squalidae: Squaloidea) from equatorial western Indian Ocean; with notes on *Squalus blainvillei*. *Journal of Zoology*, 171, 93-110.
- Miller, S.A., Dykes, D.D., Polesky H.F., 1988. A simple salting out procedure for extracting DNA from human nucleated cells. *Nucleic Acids Research*, 16, 1215.
- Muñoz-Chápuli, R., Ramos, H., 1989. Morphological comparison of *Squalus blainvillei* and *S. megalops* in the eastern Atlantic, with notes on the genus. *Japanese Journal of Ichthyology*, 36, 6-21.
- Muñoz-Chápuli, R., Ramos, H., Garcia Garrido, L., 1984. *Squalus megalops*, McLeay, 1882, en el Mediterraneo. Notas sobre su diagnosis sistemática y distribución. *Bulletí de la Societat Catalana d'Ictiologia i Herpetologia*, 9, 16-21.
- Rambaut, A., Suchard, M., Xie, D., Drummond, A., 2014. Tracer v1.6. <http://beast.bio.ed.ac.uk/Tracer>.

- Ronquist, F., Huelsenbeck, J.P., 2003. MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics*, 19, 1572-1574.
- Serena, F., 2005. *Field identification guide to the sharks and rays of the Mediterranean and Black Sea*. FAO Species Identification Guide for Fisheries Purposes. Rome, 97 p.
- Serena, F., Papacostantinou, C., Relini, G., Gil De Sola, L., Bertrand, G.A., 2009. Distribution and abundance of spiny dogfish in the Mediterranean Sea based on the Mediterranean International Trawl Survey Program. p. 139-149. In: *Biology and Management of Dogfish Sharks*. Gallucci, V.F., McFarlane, G.A., Bargmann, G.G. (Eds). American Fisheries Society.
- Tamura, K., Stecher, G., Peterson, D., Filipiński, A., Kumar, S., 2013. MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Molecular Biology and Evolution*, 30, 2725-2729.
- Veríssimo, A., Cotton, C.F., Buch, R.H., Guallart, J., Burgess, H., 2014. Species diversity of the deep-water sharks (Squaliformes: Centrophoridae: *Centrophorus*) in North Atlantic waters - current status and taxonomic issues. *Zoological Journal of the Linnean Society*, 172, 803-830.
- Veríssimo, A., Zaera-Perez, D., Leslie, R., Iglésias, S.P., Séret, B. *et al.*, 2017. Molecular diversity and distribution of eastern Atlantic and Mediterranean dogfishes *Squalus* highlight taxonomic issues in the genus. *Zoologica Scripta*, 46, 414-428.
- Viana, S.T.D.F., de Carvalho, M.R., 2016. Redescription of *Squalus acutipinnis* Regan, 1908, a valid species of spiny dogfish from Southern Africa (Chondrichthyes: Squaliformes: Squalidae). *Copeia*, 2016, 539-553.
- Viana, S.T.D.F., de Carvalho, M.R., Gomes, U.L., 2016. Taxonomy and morphology of species of the genus *Squalus* Linnaeus 1758 from the southwestern Atlantic Ocean (Chondrichthyes: Squaliformes: Squalidae). *Zootaxa*, 4133, 1-89.
- White, W.T., Ebert, D.A., Naylor, G.J.P., Ho, H.C., Clerkin, P. *et al.*, 2013. Revision of the genus *Centrophorus* (Squaliformes: Centrophoridae): part 1 – Redescription of *Centrophorus granulatus* (Bloch & Schneider), a senior synonym of *C. acus* Garman and *C. niukang* Teng. *Zootaxa*, 3752, 35-72.
- Zar, J.H., 1999. *Biostatistical analysis*. 4th Edition. Prentice-Hall, Englewood Cliffs, 663 pp.
- Zeeberg, J., Corten, A., Graaf, E., 2006. Bycatch and release of pelagic megafauna in industrial trawler fisheries off Northwest Africa. *Fisheries Research*, 78, 186-195.