

Mediterranean Marine Science

Vol 19, No 3 (2018)



Redescription of the Mediterranean endemic parasitic isopod *Anuropodione ampliandra* (Codreanu, Codreanu & Pike, 1966) n. comb. (Crustacea: Isopoda: Bopyridae) from *Munida rutllanti* Zariquiey Alvarez, 1952, with new records from Turkey and a review of the genus *Anuropodione* Bourdon, 1967

CHRISTOPHER B. BOYKO, JASON D. WILLIAMS, AHMET ÖKTENER

doi: [10.12681/mms.17412](https://doi.org/10.12681/mms.17412)

To cite this article:

BOYKO, C. B., WILLIAMS, J. D., & ÖKTENER, A. (2018). Redescription of the Mediterranean endemic parasitic isopod *Anuropodione ampliandra* (Codreanu, Codreanu & Pike, 1966) n. comb. (Crustacea: Isopoda: Bopyridae) from *Munida rutllanti* Zariquiey Alvarez, 1952, with new records from Turkey and a review of the genus *Anuropodione* Bourdon, 1967. *Mediterranean Marine Science*, 19(3), 611–619. <https://doi.org/10.12681/mms.17412>

Redescription of the Mediterranean endemic parasitic isopod *Anuropodione ampliandra* (Codreanu, Codreanu & Pike, 1966) n. comb. (Crustacea: Isopoda: Bopyridae) from *Munida rutllanti* Zariquiey Alvarez, 1952, with new records from Turkey and a review of the genus *Anuropodione* Bourdon, 1967

CHRISTOPHER B. BOYKO^{1,2}, JASON D. WILLIAMS² and AHMET ÖKTENER³

¹Division of Invertebrate Zoology, American Museum of Natural History, Central Park West @ 79th St., New York, New York 10024, U.S.A.

²Department of Biology, Hofstra University, Hempstead, New York 11549, U.S.A.

³Department of Fisheries, Sheep Research Institute, Çanakale Street 7km., 10200, Bandırma, Balıkesir, Turkey

Corresponding author: cboyko@amnh.org

Handling Editor: Agnese Marchini

Received: 14 June 2018; Accepted: 10 August 2018; Published on line: 14 December 2018

Abstract

Seven species of bopyrid isopods were previously known from Turkish Mediterranean waters; a purported eighth species, *Orthione griffenis* Markham, 2004, is based on incorrect identification and likely represents *Epipenaeon ingens ingens* Nobili, 1906, a species already known from the area. In the present study, we report on specimens referable to *Pleurocrypta ampliandra* Codreanu, Codreanu & Pike, 1966, infesting *Munida rutllanti* Zariquiey Alvarez, 1952, collected from the southern Aegean Sea off Turkey; this species was formerly known only from the Mediterranean off Algeria and the Adriatic Sea. The species is redescribed and illustrated for the first time and transferred to *Anuropodione* Bourdon, 1967, which is reviewed and *A. australiensis* Bourdon, 1976, is transferred to *Allorbimorphus* Bourdon, 1976. A key to all species of *Anuropodione* and a list of all bopyrids found on squat lobsters and porcelain crabs in the Mediterranean are provided.

Keywords: Aegean Sea; bopyrid; endemic; Epicaridea; squat lobster; Turkey.

Introduction

The parasitic isopod family Bopyridae contains over 600 species (Boyko *et al.*, 2008 onwards), all of which are parasitic on calanoid copepods as larvae and decapod crustaceans as adults (Boyko & Williams 2009; Williams & Boyko 2012). Seven of these species have been reported from Turkish waters in the Mediterranean region: *Bopyrina ocellata* (Czerniavsky, 1868) from *Hippolyte inermis* Leach, 1816 (Artüz, 2017; Sea of Marmara); *Bopyrus squillarum* Latreille, 1802, from *Palaemon adspersus* Rathke, 1837, *P. elegans* Rathke, 1837, and *P. serratus* (Pennant, 1777) (Geldiay and Kocataş, 1972; Kirkim *et al.*, 2008); *Epipenaeon ingens ingens* Nobili, 1906, from *Penaeus aztecus* Ives, 1891 (a western Atlantic species introduced into the Mediterranean) (Korun *et al.*, 2013); *Pleurocrypta galataeae* Hesse, 1865, from *Galathea squamifera* Leach, 1814 (Geldiay and Kocataş, 1972); *P. longibranchiata* (Bate & Westwood, 1867) from *Galathea nexa* Embleton, 1834, and *G. squamifera* (Geldiay and Kocataş, 1972; Kirkim *et al.*, 2008); *P. microbranchiata* G. O. Sars, 1898, from *Galathea intermedia* Lilljeborg, 1851 (Geldiay and Kocataş, 1972; Kirkim *et al.*, 2008);

P. porcellanaelongicornis Hesse, 1876, from *Pisidia* cf. *bluteli* (Risso, 1816) and *Pisidia* sp. (Geldiay and Kocataş, 1972; Kirkim *et al.*, 2008).

Mediterranean records of a purported eighth species of bopyrid, *Orthione griffenis* Markham, 2004, found infesting *Penaeus kerathurus* (Forskål, 1775) (Uluköy *et al.*, 2008, cited by Ceyhan *et al.*, 2009) are obvious misidentifications. Although *Orthione griffenis* has been documented as an invasive species in mud shrimp (Decapoda: Gebiidea) along the west coast of the US (Williams & An 2009; Hong *et al.*, 2015; Boyko *et al.*, 2017), no species of *Orthione* infest penaeid shrimp (Markham, 2004). It is most likely that these bopyrids are actually *Epipenaeon ingens ingens* Nobili, 1906, originally described from the Red Sea but which entered the Mediterranean before 1968 and was first reported from Turkey on the host *P. semisulcatus* De Haan, 1844 (Bourdon, 1968; An *et al.*, 2014). This bopyrid is now also known in the Mediterranean from the introduced shrimp hosts *P. aztecus* and *P. stylirostris* (Stimpson, 1871) (Korun *et al.*, 2013; An *et al.*, 2014). Owens and Rothlisberg (1991) suggested that *E. ingens ingens* might eventually infest the Mediterranean endemic host *P. kerathurus* and this

prediction appears to have been born out; this is the first record of *E. ingens ingens* on any Mediterranean endemic shrimp host.

The squat lobster *Munida rutilanti* Zariquiey Alvarez, 1952, lives on muddy bottoms in depths between 130 and 540 m and is historically distributed in the eastern Atlantic from northwestern Spain south to Mauritania, including the Canary and Cape Verde Islands, as well as in the western Mediterranean and Aegean Sea (Fransen, 2014). However, the species has spread eastward into the eastern Mediterranean and the Adriatic Sea (Frogliola *et al.*, 2010; Petric *et al.*, 2010). *Munida rutilanti* has potential as a fishery due to its relatively large size (carapace length, excluding rostrum, up to 22 mm) and ability to be caught in large quantities (Kocak *et al.*, 2008; Frogliola *et al.*, 2010; Fransen, 2014) but does not currently have commercial value and is not a target species (Kocak *et al.*, 2008).

Four squat lobster species in the genus *Munida* Leach, 1820 (*M. intermedia* A. Milne Edwards & Bouvier, 1899, *M. rugosa* (Fabricius, 1775), *M. rutilanti*, and *M. tenuimana* G. O. Sars, 1872) have been reported from Turkish waters (Bakir *et al.*, 2014), all of which have been recorded bearing bopyrids elsewhere in the Mediterranean (Table 1) but no bopyrids have previously been reported on any Turkish specimens belonging to these species of *Munida*. Three species of *Pleurocrypta* have been reported from squat lobster hosts collected off Turkey, but

all were on hosts in the genus *Galathea* Fabricius, 1793 (Galatheidae) (Table 1; see also Boyko *et al.*, 2012; Bakir *et al.*, 2014).

The present paper reports for the first time the presence of *Anuropodione amphiandra* (Codreanu, Codreanu & Pike, 1966) n. comb., parasitic on *M. rutilanti* from Turkish waters. Examination of a large series of specimens allowed us to determine that this species does not belong to *Pleurocrypta* Hesse, 1865, the genus to which it was originally assigned, and to quantify the variability in the pleon morphology of males.

Material and Methods

Squat lobsters were collected at depths from 60 to 400 m with a commercially used bottom trawl (mesh size: 44 mm; time of haul period: 08:00-12:00 and 13:00-17:00; engine power of the commercial trawler: 700 hp; bottom type: muddy), off Babakale Port, North Aegean Sea, Turkey (39°25'2.32"N-39°28'11"N; 26°03'39.46"E-26°20'18"E) in March 2017 (93 specimens) and August 2018 (150 specimens).

Hosts and parasites were placed in 70% ethanol and parasites were subsequently removed from the host branchial chambers; male parasites were firmly attached to the ventral pleon region of females in all cases. Select parasites were dissected using a Wild M5 and an Olympus XZS12 stereomicroscope. Dissected parts were mounted

Table 1. List of bopyrid species known from squat lobsters and porcelain crabs (Anomura: Galatheaidea) in the Mediterranean Sea.

Bopyrid Species	Hosts	Mediterranean Localities	References
<i>Anuropodione amphiandra</i> (Codreanu, Codreanu & Pike, 1966) n. comb.	<i>Munida rutilanti</i> Zariquiey Álvarez, 1952	Algeria; Adriatic Sea	Codreanu, Codreanu & Pike, 1966; Petric <i>et al.</i> , 2010
<i>Pleurocrypta galathea</i> Hesse, 1865	<i>Galathea dispersa</i> Bate, 1859 <i>Galathea nexa</i> Embleton, 1834 <i>Galathea squamifera</i> Leach, 1814	France, Monaco, Turkey	Bourdon, 1968; Geldiay & Kocataş, 1972
<i>Pleurocrypta longibranchiata</i> (Bate & Westwood, 1867)	<i>Galathea nexa</i> Embleton, 1834; <i>Galathea squamifera</i> Leach, 1814	France, Turkey	Geldiay & Kocataş, 1972
<i>Pleurocrypta microbranchiata</i> G. O. Sars, 1898	<i>Galathea intermedia</i> Lilljeborg, 1851	Spain, Turkey	Bourdon, 1968; Geldiay & Kocataş, 1972
<i>Pleurocrypta piriformis</i> Bourdon, 1968	<i>Galathea strigosa</i> (Linnaeus, 1761)	Spain	Bourdon, 1968
<i>Pleurocrypta porcellanaelongicornis</i> Hesse, 1876	<i>Pisidia longicornis</i> (Linnaeus, 1767)	Turkey	Geldiay & Kocataş, 1972
<i>Pleurocrypta strigosa</i> Bourdon, 1968	<i>Galathea strigosa</i> (Linnaeus, 1761)	Spain, Italy	Bourdon, 1968
<i>Pseudione crenulata</i> G. O. Sars, 1898	<i>Munida intermedia</i> A. Milne Edwards & Bouvier, 1899 <i>Munida rugosa</i> (Fabricius, 1775) <i>Munida tenuimana</i> G. O. Sars, 1872	Spain, Italy	Zariquiey Alvarez, 1958; Castriota <i>et al.</i> , 2010

on slides in glycerine-gelatine mounting medium; pleopods were stained with methylene blue. Line drawings were made with a camera lucida drawing tube attached to Olympus compound or dissecting microscopes. Final images were created by tracing a scanned copy of the original sketch with a Wacom Cintiq pen display using Adobe Illustrator. In addition to conventional light micrographs, some specimens were imaged with a Macro pod Pro kit (MacroscopicSolutions) and resulting pictures were aligned and stacked with the focus stacking software Zerene Stacker (10–65 images from bottom to top of specimens).

Carapace length (CL, inclusive of rostrum) is provided as an indicator of size for the hosts. Isopod size is given as total body length (anterior margin of head to posterior margin of pleotelson). Measurements were made to 0.1 mm using an ocular micrometer, micro-scale tool (Electron Microscopy Services), or a micrometric programme (Pro-way).

Authors and dates of hosts and bopyrid parasites were verified using WoRMS (Boyko *et al.*, 2008 onwards). Bibliographic references are provided for authors and dates of all parasite taxa but not for those of host species.

Voucher specimens were deposited in the collections of the Muséum National d'Histoire Naturelle (MNHN) and the National Museum of Natural History, Smithsonian Institution, Washington, D.C. (USNM).

Results

Systematics

Family Bopyridae Rafinesque, 1815

Subfamily Pseudioninae Codreanu, 1967

Genus *Anuropodione* Bourdon, 1967

Type species: *Anuropodione senegalensis* Bourdon, 1967

Remarks

Prior to the present study, *Anuropodione* contained five species (Boyko *et al.*, 2008 onwards) but a review of these taxa shows that *A. australiensis* Bourdon, 1976, is not properly placed in this genus, *A. dubius* (Nierstrasz & Brender à Brandis, 1929) is questionably placed here, and *Pleurocrypta amphindra* should be transferred to *Anuropodione*. Examination of the description and illustrations of *A. australiensis* shows that this species is not correctly generically placed. The minimal development of the lateral plates on the pleon, expanded posterior lobe with acute ventral point on the first oostegite of the female, as well as the rounded, extended head and weakly projecting pleotelson and overall body proportions of the male indicate that this species is not congeneric with *A. senegalensis* Bourdon, 1967, the type species of *Anuropodione*. In many characters, *A. australiensis* is closer to *Allorbimorphus haigae* Bourdon, 1976, but it differs from that species in the females of *A. haigae* having only four pairs of biramous pleopods (four pairs biramous pleop-

ods, fifth pair uniramous in *A. australiensis*), uniramous uropods (no uropods in *A. australiensis*), and in males of *A. haigae* having all pleomeres distinct (1-3 pleomeres in *A. australiensis*). It is possible that *A. australiensis* should be placed in a new genus, but we refrain from such an action in the absence of material; however, we transfer the species to *Allorbimorphus* Bourdon, 1976, the genus to which it shares most characters, as *Allorbimorphus australiensis* (Bourdon, 1976) n. comb.

It is unclear if *A. dubius* is correctly placed in this genus. The female shows essentially no lateral plate development on the pleomeres compared to other species of *Anuropodione* and there are 5 uniramous pleopods, which contrasts with the typical 5 biramous pairs in species of *Anuropodione* (excepting *A. megacephalon*, which has 4 biramous and 1 uniramous pair). However, the pleon of the male greatly resembles that of one of the morphs found in *A. amphindra* and the morphology of the first oostegite and maxilliped (see Bourdon, 1972) are very similar to those of other *Anuropodione* species. Bourdon (1972) compared *Probopyrus dubius*, which was clearly not correctly placed in *Probopyrus* Giard & Bonnier, 1888, to all other species known where females lacked uropods: *Parioninella obovata* Shiino, 1958 (now *Ovoionella obovata* (Shiino, 1958); see Boyko, 2004)), *A. senegalensis*, and *Bopyrissa magellanica* (Nierstrasz & Brender à Brandis, 1929) and concluded that the male and female of *P. dubius* resembled those of *A. senegalensis* more than the other species. We concur with Bourdon's (1972) conclusion, inasmuch as the male of *O. obovata* is very different from that of *P. dubius* and the female of *B. magellanica* actually does have well-developed uropods, a fact not known to Bourdon (1972; see Williams *et al.*, submitted). Unfortunately, it will be difficult to recollect *A. dubius* and garner a larger sample size, as the host was only given by Nierstrasz & Brender à Brandis (1929) as "*Galathea spec.*"

Pleurocrypta amphindra Codreanu, Codreanu & Pike, 1966, belongs to *Anuropodione* on the basis of the females lacking well-developed uropods as well as maxilliped palps. Females of *Pleurocrypta* always have well-developed uropods (uniramous or biramous, depending on the species) and maxilliped palps. Codreanu *et al.* (1966) described the pleotelson of *Pleurocrypta amphindra* as bearing "exceptionnellement des uropodes rudimentaires"; in all specimens we examined, there are small ridges on the pleotelson but they are not true uropods. Bourdon (1972), however, did report very small, poorly developed uropods on some specimens of *A. senegalensis*.

Key to species of *Anuropodione* Bourdon, 1967, based on females

- 1a. Lateral plates of pleon weakly developed, closely adpressed ... *A. dubius* (Nierstrasz & Brender à Brandis, 1929) [Thailand, 27.4 m depth, infesting "*Galathea sp.*"]
- 1b. Lateral plates of pleon forming distinct lateral ex-

tensions, not closely adpressed ... 2

2a. Head length greater than that of first four pereomeres combined ... *A. megacephalon* Markham, 1974 [28°48'N; 85°02'W, Gulf of Mexico, 54-60 m depth, infesting *Munida pusilla* Benedict, 1902]

2b. Head length less than that of first four pereomeres combined ... 3

3a. Lateral plates of pleon thin, tapered distally ... *A. carolinensis* Markham, 1974 [35°05'N 74°33'W to 37°09'N 75°11.5'W, 83-662 m depth, infesting *Munida iris* A. Milne Edwards, 1880]

3b. Lateral plates of pleon expanded distally, ovate ... 4

4a. Lateral plates of pleon non-overlapping ... *A. senegalensis* Bourdon, 1967 [Senegal to Republic of the Congo, 200-300 m depth, infesting *Munida* sp. and *M. speciosa* von Martens, 1878]

4b. Lateral plates of pleon overlapping ... *A. amphandra* (Codreanu, Codreanu & Pike, 1966) n. comb. [Mediterranean Sea: Algeria to Turkey; Adriatic Sea, 60-400 m depth, infesting *Munida rutllanti* Zariquiey Alvarez, 1952].



Fig. 1: *Munida rutllanti* Zariquiey Alvarez, 1952, dorsal view of male specimen (USNM 1492241) with swollen right branchial chamber caused by *Anuropodione amphandra* (Codreanu, Codreanu & Pike, 1966) n. comb. Scale bar = 3 mm.

***Anuropodione amphandra* (Codreanu, Codreanu & Pike, 1966) n. comb.** (Figs. 1-4)

Pleurocrypta sp. Délye, 1955: 84 [Béni-Saf, west of Oran, Algeria, infesting *Munida iris* ssp. *rutllanti* [sic] = *M. rutllanti*]; Petric *et al.*, 2010: 1395-1403, fig. 7a [Adriatic Sea, infesting *M. rutllanti*].

“un bopírido sp.?” Zariquiey Alvarez, 1958: 101 [mention].

Pleurocrypta amphandra Codreanu, Codreanu & Pike, 1966: 1071-1072 [type locality: Béni-Saf, west of Oran, Algeria, 205-230 m; type host: *Munida iris rutllanti* Zariquiey Alvarez, 1952 = *M. rutllanti*]; Markham, 1974: 270 [mention], 271 [table], 265 [list]; Bourdon, 1976: 366 [mention]; Boyko & Williams, 2011: 285 [mention]; Boyko *et al.*, 2012: 9, 18 [list].

“le Bopyridae des *Munida iris rutllanti* [sic] Zariquiey Alvarez des côtes algériennes” Bourdon, 1968: 221 [mention].

“parasite (non encore décrit à notre connaissance) des *Munida iris rutllanti* [sic]” Bourdon, 1972: 833 [mention].

Material examined and used as basis for redescription

1 dextral ovigerous female (5.5 mm), 1 mature male (1.6 mm, pleon elongate with 3 pleomeres) from right branchial chamber of unknown sex and size *Munida rutllanti* Zariquiey Alvarez, 1952, off Babakale Port (39°25'2.32"N-39°28'11"N; 26°03'39.46"E-26°20'78"E, North Aegean Sea, Turkey)




Fig. 2: *Anuropodione amphandra* (Codreanu, Codreanu & Pike, 1966) n. comb., female specimen USNM 1492250 (A-I). A, dorsal view. B, ventral view. C, right antennule (top) and antenna (bottom). D, left barbula. E, left maxilliped, outer view. F, left oostegite 1, outer view. G, left oostegite 1, inner view. H, left pereopod 1. I, left pereopod 7. Scale bars = 2 mm (A, B), 250 µm (C, D, H), 500 µm (E-G, I).




Fig. 3: *Anuropodione amphandra* (Codreanu, Codreanu & Pike, 1966) n. comb., female specimen USNM 1492250 (A), personal collection (B-M). A, Pleon, dorsal view, B-F, pleopods 1-5, respectively, G-M, pereopods 1-7, respectively. Scale bars = 500 μ m (A), 1 mm (B-F), 250 μ m (G-M).



Fig. 4: *Anuropodione amphandra* (Codreanu, Codreanu & Pike, 1966) n. comb., male specimens USNM 1492250 (A-D), USNM 1492279 (E), USNM 1492241 (F). A, dorsal view. B, ventral view. C, mouthparts, left antennule (top), left antenna (bottom) and left pereopod 1. D, left pereopod 7. E, pleon with 5 segments. F, pleon with 3 segments. Scale bars = 250 μ m (A, B, E, F), 50 μ m (C, D).

in March 2017 or August 2018 (USNM 1492241; Fig. 1) (all other specimens cited below with same collection data); 1 dextral ovigerous female (8.6 mm), 1 mature male (4.8 mm, pleon elongate with 5 pleomeres), from right branchial chamber of unknown sex and size *M. rutllanti* (USNM 1492244); 1 dextral ovigerous female (7.5 mm), 1 mature male (1.7 mm, pleon short with all pleomeres coalesced), from right branchial chamber of unknown sex and size *M. rutllanti* (USNM 1492250); 1 sinistral ovigerous female (7.0 mm), 1 mature male (pleon elongate with 5 pleomeres), from left branchial chamber of unknown sex and size *M. rutllanti* (USNM 1492279); 1 dextral ovigerous female (5.5 mm), 1 mature male (pleon short with all pleomeres coalesced), from right branchial chamber of male *M. rutllanti* (15 mm CL) (USNM 1492233); 1 dextral ovigerous female (7.3 mm), infesting unknown size and sex *M. rutllanti* (MNHN-IU-2017-72).

Additional material examined

All specimens with same collection data as USNM 1492241: 1 sinistral ovigerous female (6.48 mm), 1 mature male (1.85 mm), infesting left branchial chamber of female *M. rutllanti* (36 mm CL); 1 sinistral ovigerous female (6.89 mm), 1 mature male (1.24 mm), infesting left branchial chamber of female *M. rutllanti* (37 mm CL); 1

dextral ovigerous female (6.94 mm), 1 mature male (1.57 mm), infesting right branchial chamber of female *M. rutllanti* (39 mm CL); 1 sinistral ovigerous female (7.09 mm), 1 mature male (1.28 mm), infesting left branchial chamber of female *M. rutllanti* (39 mm CL); 1 dextral ovigerous female (6.94 mm), 1 mature male (1.57), infesting right branchial chamber of female *M. rutllanti* (39 mm CL); 1 dextral ovigerous female (7.01 mm), 1 mature male (1.49 mm), infesting right branchial chamber of female *M. rutllanti* (39 mm CL); 1 dextral ovigerous female (7.65 mm), 1 mature male (1.76 mm), infesting right branchial chamber of female *M. rutllanti* (40 mm CL); 1 sinistral ovigerous female (7.09 mm), 1 mature male (1.72 mm), infesting left branchial chamber of female *M. rutllanti* (40 mm CL); 1 sinistral ovigerous female (7.47 mm), 1 mature male (1.56 mm), infesting left branchial chamber of female *M. rutllanti* (41 mm CL); 1 sinistral ovigerous female (7.22 mm), 1 mature male (2.47 mm), infesting left branchial chamber of female *M. rutllanti* (41 mm CL); 1 sinistral ovigerous female (7.29 mm), 1 mature male (2.65 mm), infesting left branchial chamber of female *M. rutllanti* (41 mm CL); 1 dextral ovigerous female (7.24 mm), 1 mature male (3.63 mm), infesting right branchial chamber of female *M. rutllanti*

(42 mm CL); 1 sinistral ovigerous female (7.74 mm), 1 mature male (1.82 mm), infesting left branchial chamber of female *M. rutllanti* (42 mm CL); 1 sinistral ovigerous female (7.87 mm), 1 mature male (1.74 mm), infesting left branchial chamber of female *M. rutllanti* (42 mm CL); 1 sinistral ovigerous female (7.96 mm), 1 mature male (2.10 mm), infesting left branchial chamber of female *M. rutllanti* (42 mm CL); 1 sinistral ovigerous female (6.65 mm), 1 mature male (2.31 mm), infesting left branchial chamber of female *M. rutllanti* (43 mm CL); 1 dextral ovigerous female (7.63 mm), 1 mature male (1.64 mm), infesting right branchial chamber of female *M. rutllanti* (43 mm CL); 1 sinistral ovigerous female (7.90 mm), 1 mature male (1.89 mm), infesting left branchial chamber of female *M. rutllanti* (44 mm CL); 1 sinistral ovigerous female (8.02 mm), 1 mature male (1.51 mm), infesting left branchial chamber of female *M. rutllanti* (45 mm CL); 1 sinistral ovigerous female (7.68 mm), 1 mature male (1.69 mm), infesting left branchial chamber of female *M. rutllanti* (45 mm CL); 1 dextral ovigerous female (7.67 mm), 1 mature male (2.99 mm), infesting right branchial chamber of female *M. rutllanti* (45 mm CL); 1 dextral ovigerous female (8.15 mm), 1 mature male (1.87 mm), infesting right branchial chamber of female *M. rutllanti* (46 mm CL); 1 sinistral ovigerous female (8.42 mm), 1 mature male (1.54 mm), infesting left branchial chamber of female *M. rutllanti* (47 mm CL); 1 sinistral ovigerous female (8.34 mm), 1 mature male (2.19 mm), infesting left branchial chamber of female *M. rutllanti* (50 mm CL); 1 sinistral ovigerous female (8.30 mm), 1 mature male (1.98 mm), infesting left branchial chamber of female *M. rutllanti* (52 mm CL).

Additional hosts examined

All specimens with same collection data as USNM 1492241: 1 female *M. rutllanti* (16 mm CL), with swollen left branchial chamber (parasite not present) (USNM 1492232); 1 male *M. rutllanti* (22.4 mm CL), with swollen right branchial chamber (parasite not present); host also infested with immature *Lernaeodiscus ingolfi* (1.0 x 2.5 cm) (see Remarks) (USNM 1492232).

Redescription

Female (based on USNM 1492250): body length 8.9 mm, maximal width 6.1 mm, head length 1.8 mm, head width 2.6 mm, pleon length, 2.7 mm. Head dextrally deflexed with ~10° distortion angle. Body broad anteriorly, tapered posteriorly, all body regions and segments distinct (Fig. 2A, B). Head large, subovate, moderately set into pereon; frontal lamina broad, posterior margin with small median projection (Fig. 2A); eyes absent. Antennules and antennae of 3 articles each, with setae at distal tip of distal segment of both; antennae with basal segment greatly expanded, approximately 3x width of second segment (Fig. 2C). Maxilliped with short, acute spur, palp absent or as a low non-articulated lobe with few marginal setae,

no setae on margins (Fig. 2E). Barbula with two short, smooth, falcate lateral projections with tapered tips; median region smooth (Fig. 2D). Oostegite surfaces smooth, completely enclosing brood chamber; setae on posterior margin of fifth oostegite (Fig. 2B); oostegite 1 with ovate anterior lobe, narrow triangular posterior lobe with short, rounded, posterolateral extension (Fig. 2F); inner ridge of oostegite 1 with numerous low digitate projections (Fig. 2G). Pereon of 7 distinct pereomeres, broadest across pereomere 3, tapering gently towards head, markedly towards pleon. Prominent coxal plates on pereomeres 1-7, those on 1-4 elongate, recurved ovals, those on 5-7 as posteriorly expanded flattened plates, flat on longer side of body, curled on shorter side of body; elongate, ovate dorsolateral bosses on pereomeres 1-4, smaller than corresponding coxal plates (Fig. 2A). Tergal projections absent. Pereopods subequal in size, each with large smooth-edged lobe projecting from dorsal surface of basis, size of basal lobe increasing on posterior pereopods, surface of lobe with minute scales; pereopods 1-3 spaced apart, 4-7 spaced closely together (Figs. 2H, I, 3G-M). Pleon of 6 distinct pleomeres, first 5 pleonal segments with lateral plates expanded posteriorly and overlapping from pleomeres 1-5 (Figs. 2A, 3A). Five pairs of biramous pleopods, visible in dorsal view, posteriorly extending well beyond margins of lateral plates; anterior pairs triangular, posterior pairs more elongate and tapered distally, all margins smooth (Figs. 2A, B, 3A-F); pleomere 6 lacking uropods (Figs. 2A, 3A).

Male (based on USNM 1492241, 1492250, 1492279): length 1.8 mm, maximal width 0.8 mm at pereomere 4, head length 0.3 mm, head width 0.5 mm, pleon length 0.5 mm. Head subovate, distinct from pereomere 1, all body segments distinct (Fig. 4A); prominent eyes on posterolateral border of head (Fig. 4A). Antennules of 3 articles each, setae on distal tip of article 3, setae subterminally on other articles (Fig. 4C); antennae of 4 articles each, tuft of setae on distal tip of terminal article (Fig. 4C). Pereon of 7 distinct pereomeres, broadest across pereomeres 3-5, tapering slightly anteriorly and posteriorly. Pereopods subequal in size, all articles distinct; dactyli of pereopods 1-3 proportionally longer than those of 4-7, propodi with ridges of scales on anterior portion underneath dactyl, tuft of setae on anterior tip of propodi opposed to tip of dactyli (Fig. 4C, D), meri and carpi slightly larger in posterior pairs. Pleon variable, ranging from one large, triangular, posteriorly tapered segment (Fig. 4A) with faint lateral undulations to 3 distinct segments (2 narrow pleomeres plus a globose pleotelson; Fig. 4F) or 5 distinct segments (4 narrow pleomeres plus elongate pleotelson showing lateral constriction; Fig. 4E). Midventral tubercles, pleopods and uropods absent; anal cone present but not projecting (Fig. 4B, E, F).

Variation

Females occur in both dextral and sinistral forms in approximately equal numbers (~40% dextral vs. ~60%

Table 2. Known variation in the number pleomeres for males of species in *Anuropodione* Bourdon, 1967.

Species	Number of Pleomeres (Number of Specimens Known)					Reference(s)
<i>A. amphandra</i> n. comb.	1 (29)	2 (2)	3 (1)	5 (4***)	6 (3)	Codreanu <i>et al.</i> , 1966; Present Study
<i>A. australiensis</i>	1 (1)		3 (1)			Bourdon, 1976
<i>A. carolinensis</i>	1 (147*)		3 (2)		6 (38)	Markham, 1974; Wenner & Windsor, 1979
<i>A. dubius</i>				5 (1)		Nierstrasz & Brender à Brandis, 1929; Bourdon, 1972
<i>A. megacephalon</i>	1 (1**)					Markham, 1974
<i>A. senegalensis</i>	1 (7***)	2 (2***)		5 (3***)		Bourdon, 1967; Bourdon, 1972

* 93 males with 1 pleomere but with lateral undulations suggesting incipient segmentation.

** Terminal segment with dorsal indications of 3 coalesced pleomeres.

*** Terminal pleomere with lateral undulations suggesting coalesced pleomeres.

sinistral). Male pleon morphology is highly variable (see redescription above); a similar level of variation was noted by Codreanu *et al.* (1966) for this species and is known across the genus as a whole (Table 2).

Ecology

The rate of parasitism by *A. amphandra* on *M. rutilanti* is relatively high compared to what is known for most bopyrid species (~1-5%; Cericola & Williams, 2015). At least 9 of the 93 squat lobsters (~9.7%) collected in March 2017 bore bopyrids and at least 16 of the 150 hosts (~10.7%) collected in August 2018 contained *A. amphandra* (these prevalences are conservative estimates because six additional parasitized hosts were collected but not assignable to a specific collecting date). All of the hosts were female, except for a single male that was co-infested with a rhizocephalan.

Remarks

Codreanu *et al.* (1966) reported that, of their nine males examined, four had all pleomeres fused into a single segment, two had two segments, and three had six segments and an overall elongate pleon. We found males with one, three, and five pleonal segments in the present material (Table 2). Codreanu *et al.* (1966) also described the pereomeres of the males as having “finement spinuleux” but these are merely small setae.

Codreanu *et al.* (1966) interpreted the variation in the number of pleomere segments as a response by males to incomplete inhibition of pleomere development by some kind of chemical secretion exuded by the female bopyrids. They labeled the males as being “néoténiques” but as that term specifically refers to the retention of larval fea-

tures in the adult, they applied it incorrectly in this situation. The assumption that male bopyrids are in some way prevented from transitioning from the male to female phase by the influence of females already present in or on the host has been mentioned in the literature (summarized in Trilles, 1999) but the mechanism is unknown and there are few studies which have experimentally demonstrated that it occurs (see Reverberi & Pitotti, 1943; Reverberi, 1947; Reinhard, 1949). Reinhard (1949) dealt with the transformation of cryptoniscus larvae, so the only studies that have shown males transforming into females are those on *Ione thoracica* Montagu, 1808 (Reverberi & Pitotti, 1943; Reverberi, 1947) and in this species the pleomeres of males are quite similar morphologically to those of the female and not variable in the number of segments, unlike in the present case with *Anuropodione* or with species of *Pleurocrypta* (Bourdon, 1965). In fact, in the case of the observed variability in the number of pleomeres of males in species of *Anuropodione*, it is unlikely that a male to female transition is occurring because if the male with one pleomere was transitioning to six pleomeres (the number possessed by the females), then that would mean the cryptoniscus larvae had all their pleomeres fused and subsequently separated again, a seemingly non-parsimonious developmental trajectory. The converse situation, that a male with six segments will transition to a male with one segment is also not probable, as females consistently present six pleomeres. It is more likely that in all species of *Anuropodione*, as with species of *Pleurocrypta*, the number of pleomeres of males is a polymorphic character lacking an evolutionary pressure for fixation at a particular number of segments. These genera provide an opportunity to test this hypothesis and examine the underlying genetic controls of the segmentation process.

One host (USNM 1492232) was doubly infested by *A.*

amphiandra n. comb. and an immature specimen of *Lernaeodiscus* cf. *ingolfi* Boschma, 1928. There is one prior record of a species of *Lernaeodiscus* on a *M. rutllanti* host, collected in the Ionian Sea (Øksnebjerg, 2000, who recorded the parasite as *L. ingolfi* and the host variously as *Munida iris*, *M. iris* ssp. *rutllanti* [sic], and *M. iris* ssp. *rutllanti* [sic]). However, because the host examined by Øksnebjerg (2000) bore a dried externa and the present specimen is immature, it is not certain that *M. rutllanti* hosts true *L. ingolfi*.

Petric *et al.* (2010) incorrectly stated that *M. rutllanti* was parasitized by two species of bopyrids: *Pleurocrypta amphiandra* (which they incorrectly credited Délye, 1955, with authorship of the species) and *P. intermedia* Bonnier, 1900 (which they incorrectly credited Codreanu *et al.*, 1966, with authorship and which is a synonym of *P. microbranchiata* G. O. Sars, 1898). *Pleurocrypta microbranchiata* has never been reported from *M. rutllanti*.

Acknowledgements

We gratefully acknowledge the assistance of the fisherman Mr. Şenol Demirtaş from the fishing boat *Sadik Huseyin* for making collections. Laure Corbari kindly made the MNHN specimens available for examination by the first author. We would also like to thank the reviewers for their constructive comments.

References

An, J., Boyko, C.B., Li, X., 2014. Review of the parasitic genus *Epipenaeon* Nobili, 1906 (Crustacea: Isopoda: Bopyridae), with new records and redescription of four species from China. *Journal of Natural History*, 48, 2027-2048.

Artüz, M. L., 2017. First record of *Bopyrina ocellata* (Isopoda, Bopyridae) from the Sea of Marmara, Turkey. *Crustaceana*, 90, 685-694.

Bakır, A.K., Katağan, T., Aker, H.V., Özcan, T., Sezgin, M. *et al.*, 2014. The marine arthropods of Turkey. *Turkish Journal of Zoology*, 38, 765-831.

Bate, C.S., Westwood, J.O., 1861-1868. *A history of the British sessile-eyed Crustacea*. Vol. 2. John Van Voorst: London.

Bonnier, J., 1900. Contribution à l'étude des épicarides. Les Bopyridae. *Travaux de la Station Zoologique de Wimereux*, 8, 1-476, pls. 1-41.

Boschma, H., 1928. Rhizocephala. *The Danish Ingolf-Expedition*, 3, 1-52, 1 map.

Bourdon, R., 1965. Remarques au sujet de la nouvelle espèce '*Pseudione convergens*' Stock, 1960 (epicaride de la famille des Bopyridae). *Cahiers de Biologie Marine*, 6, 173-179.

Bourdon, R., 1967. Sur trois nouveaux Bopyridae du Sénégal. *Bulletin de l'Institut Fondamental d'Afrique Noire*, Sér. A, 29, 107-122.

Bourdon, R., 1968. Les Bopyridae des Mers Européennes. *Mémoires du Muséum National d'Histoire Naturelle*, Série A, Zoologie 50, 77-424.

Bourdon, R., 1972. Sur quelques Bopyridae (Crustacea, Isopoda) parasites de galathéides. *Bulletin du Muséum National d'Histoire Naturelle*, 3^e Sér., 66, Zoologie 52, 817-838.

Bourdon, R., 1976. Les bopyres des porcellanes. *Bulletin du Muséum National d'Histoire Naturelle*, 3^e Sér., 359,

Zoologie, 252, 165-245.

Boyko, C.B., 2004. The Bopyridae (Crustacea: Isopoda) of Taiwan. *Zoological Studies*, 43, 677-703.

Boyko, C.B., Bruce, N.L., Hadfield, K.A., Merrin, K.L., Ota, Y. *et al.*, 2008 onwards. *World Marine, Freshwater and Terrestrial Isopod Crustaceans Database. Bopyridae Rafinesque, 1815. World Register of Marine Species* at: <http://www.marinespecies.org/aphia.php?p=taxdetails&id=1195> (Accessed 2 May 2018).

Boyko, C.B., Williams, J.D., 2009. Crustacean parasites as phylogenetic indicators in decapod evolution. p. 197-220. In: *Crustacean issues 18. Decapod crustacean phylogenetics*. Martin, J.W., Crandall, K.A., Felder, D.L. (Eds). CRC Press; Boca Raton.

Boyko, C.B., Williams, J.D., 2011. Chapter 9. Parasites and other symbionts of squat lobsters. p. 271-295. In: *The biology of squat lobsters*. Poore, G.C.B., Ah Yong, S.T., Taylor, J. (Eds). CSIRO Publishing, Melbourne.

Boyko, C.B., Williams, J.D., Markham, J.C., 2012. Recent and fossil Isopoda Bopyridae parasitic on squat lobsters and porcelain crabs (Crustacea: Anomura: Chirostyloidea and Galatheaidea), with notes on nomenclature and biogeography. *Zootaxa*, 3150, 1-35.

Boyko, C.B., Williams, J.D., Shields, J.D., 2017. Parasites (Isopoda: Epicaridea and Nematoda) from ghost and mud shrimp (Decapoda: Axiidea and Gebiidea) with descriptions of a new genus and a new species of bopyrid isopod and clarification of *Pseudione* Kossman, 1881. *Zootaxa*, 4365, 251-301.

Castriota, L., Falautano, M., Vivona, P., Andaloro, F., 2010. A new locality and host for *Pseudione crenulata* G. O. Sars, 1898 (Crustacea, Isopoda, Bopyridae) in the Mediterranean. *Acta Parasitologica*, 55, 291-295.

Cericola, M.J., Williams, J.D., 2015. Prevalence, reproduction and morphology of the parasitic isopod *Athelges takanoshimensis* Ishii, 1914 (Isopoda: Bopyridae) from Hong Kong hermit crabs. *Marine Biology Research*, 11, 236-252.

Ceyhan, T., Akyol, O., Erdem, M., 2009. Gökova körfezi (ege denizi) karides balıkçılığı. *E. U. Journal of Fisheries & Aquatic Sciences*, 26, 219-224.

Codreanu, R., 1967. Clasificarea evolutiva a bopirienilor, isopode parazite ale crustaceelor decapode si importanta lor biologica generala. *Studii si Cercetari de Biologie Seria Zoologie*, 19, 203-211.

Codreanu, R., Codreanu, M., Pike, R.B., 1966. *Pleurocrypta amphiandra* n. sp., bopyrien parasite branchial de *Munida iris rutllanti* [sic], galatheide d'Algérie et les effets d'interbiomorphose. In: Corradetti, A. (Ed.). *Proceedings of the First International Congress of Parasitology (Roma, 21-26 September 1964)*, 2, 1071-1072.

Czerniavsky, V., 1868. *Materialia ad zoographiam ponticam comparatam*. I. Studiosi Universitatis Charcoviensis.

Délye, G., 1955. Action d'un bopyrien sur les caractères sexuels de *Munida iris* ssp. *rutllanti* [sic] Zariquiey (décapode, anomoure). *Bulletin de la Société d'Histoire Naturelle de l'Afrique du Nord*, 46, 84-88.

Fransen, C.H.J.M., 2014. Anomurans. p. 236-248. In: *FAO species identification guide for fishery purposes. The living marine resources of the eastern central Atlantic. Vol. 1: Introduction, crustaceans, chitons and cephalopods*. Carpenter, K.E., De Angelis, N. (Eds). FAO: Rome.

Froggia, C., Gramitto, M.E., Morello, E.B., 2010. In less than 10 years the squat lobster *Munida rutllanti* has replaced *M. intermedia* in the western Pomo Pit (Central Adriatic). *Rapport du Commission Internationale Pour l'Exploration*

- Scientifique de la Mer Méditerranée*, 39, 519.
- Geldiay, R., Kocataş, A., 1972. Isopods collected in Izmir Bay, Aegean Sea. *Crustaceana Supplement*, 3, 19-30.
- Giard, A., Bonnier, J., 1888. Sur deux nouveaux genres d'épicarides (*Probopyrus* et *Palegyge*). *Comptes Rendus Hebdomadaires des Séances de l'Académie de Sciences*, 106, 304-306.
- Hesse, C.E., 1865. Recherches sur les crustacés rares ou nouveaux des côtes de France (complément du troisième article). *Annales des Sciences Naturelles, Zoologie*, Sér. 5, 3, 221-242, pl. 4.
- Hesse, C.E., 1876. Description des crustacés rares ou nouveaux des côtes de France (vingt-sixième article). *Annales des Sciences Naturelles, Zoologie et Paléontologies*, Sér. 6, 4, 1-48, pls. 7-9.
- Hong, J.-S., Lee, C.-L., Min, G.-S., 2015. Occurrence of *Orthione griffenis* Markham, 2004 (Isopoda: Bopyridae), parasite of the mud shrimp *Upogebia major* (De Haan, 1841) in South Korean waters, and its implications with respect to marine bioinvasion. *Journal of Crustacean Biology*, 35, 605-615.
- Kırkım, F., Kocataş, A., Katağan, T., Sezgin, M., 2008. A report on parasitic isopods (Crustacea) from marine fishes and decapods collected from the Aegean Sea (Turkey). *Türkiye Parazitoloji Dergisi*, 32, 382-385.
- Kocak, C., Leblebici, S., Ozaydin, O., Katagan, T., 2008. Some morphometric aspects of *Munida rutilanti* Zariquiey Alvarez, 1952 (Decapoda, Anomura, Galatheididae) in Sigacik Bay (southeastern Aegean Sea). *Crustaceana*, 81, 873-881.
- Korun, J., Gökoglu, M., Balci, B.A. Özvarol, Y., 2013. Infestation of brown shrimp, *Farfantepenaeus aztecus*, Ives (1891) (Penaeidae) by *Epipenaeon ingens*, Nobili (1906) (Isopoda, Bopyridae) from the Antalya Bay, Turkey. *Revue de Médecine Vétérinaire* 164, 559-563.
- Latreille, P.A., 1802. *Histoire naturelle generale et particulière, des crustacés et des insectes*. Vol. 7. F. Dufart: Paris.
- Markham, J.C., 1974. Six new species of bopyrid isopods parasitic on galatheid crabs of the genus *Munida* in the western Atlantic. *Bulletin of Marine Science*, 23, 613-648.
- Markham, J.C., 2004. New species and records of Bopyridae (Crustacea: Isopoda) infesting species of the genus *Upogebia* (Crustacea: Decapoda: Upogebiidae): The genera *Orthione* Markham, 1988, and *Gyge* Cornalia & Panceri, 1861. *Proceedings of the Biological Society of Washington*, 117, 186-198.
- Montagu, G., 1808. Description of several marine animals found on the south coast of Devonshire. *Transactions of the Linnean Society of London*, 9, 81-114, pls. 2-8.
- Nierstrasz, H.F., Brender à Brandis, G.A., 1929. Papers from Dr. Th. Mortensen's Pacific Expedition 1914-16. XLVIII. Epicaridea. I. *Videnskabelige Meddelelser Fra Den Dansk Naturhistoriske Forening i København*, 87, 1-44.
- Øksnebjerg, B., 2000. The Rhizocephala (Crustacea: Cirripedia) of the Mediterranean and Black Seas: taxonomy, biogeography, and ecology. *Israel Journal of Zoology*, 46, 1-102.
- Owens, L., Rothlisberg, P.C., 1991. Vertical migration and advection of bopyrid isopod cryptoniscid larvae in the Gulf of Carpentaria, Australia. *Journal of Plankton Research*, 13, 779-787.
- Petric, M., Ferri, J., Mladineo, I., 2010. Growth and reproduction of *Munida rutilanti* (Decapoda: Anomura: Galatheididae) and impact of parasitism by *Pleurocrypta* sp. (Isopoda: Bopyridae) in the Adriatic Sea. *Journal of the Marine Biological Association of the United Kingdom*, 90, 1395-1404.
- Rafinesque, C.S., 1815. *Analyse de la Nature ou Tableau de l'Univers et des Corps Organisés*. Palerme.
- Reinhard, E.G., 1949. Experiments on the determination and differentiation of sex in the bopyrid *Stegophryxus hyptius* Thompson. *Biological Bulletin*, 96, 17-31.
- Reverberi, G., 1947. Ancora sulla trasformazione sperimentale del sesso nei bopiridi. la trasformazione delle femmine giovanili in maschi. *Pubblicazione della Stazione Zoologica di Napoli*, 21, 81-91.
- Reverberi, G., Pittoti, M., 1943. Il ciclo biologico e la determinazione fenotipica del sesso di *Ione thoracica* Montagu, bopiride parassita di *Callianassa laticauda* Otto. *Pubblicazione della Stazione Zoologica di Napoli*, 19, 111-184.
- Sars, G.O., 1896-1899. *An account of the Crustacea of Norway with short descriptions and figures of all the species. Volume II. Isopoda*. Bergen Museum: Bergen.
- Shiino, S.M., 1958. Note on the bopyrid fauna of Japan. *Report of the Faculty of Fisheries, Prefectural University of Mie*, 3, 27-74, pl. 3.
- Trilles, J.-P., 1999. Ordre des isopodes sous-ordre des épicarides (Epicaridea Latreille, 1825). In: *Traité de Zoologie. Anatomie, Systématique, Biologie* (Pierre-P. Grassé). Tome VII, Fascicule III A, Crustacés Pécararides. Forest, J. (Ed.). *Mémoires de l'Institut Oceanographique, Monaco* 19, 279-352.
- Uluköy, G., Bulut, S., Özdemir, E., Erdem, M., 2008. An ectoparasite (*Orthione griffenis* Markham 2004) infestation, which being seen on caramote prawn (*Melicerthus kerathurus*, Forskål 1775) in Gökova Bay (in Turkish). *Ulusal Biyoloji Kongresi Özetler Kitabı*, 19, 306-307.
- Wenner, E.L., Windsor, N.T., 1979. Parasitism of galatheid crustaceans from the Norfolk Canyon and the Middle Atlantic Bight by bopyrid isopods. *Crustaceana*, 37, 293-303.
- Williams, J.D., An, J., 2009. The cryptogenic parasitic isopod *Orthione griffenis* Markham, 2004 from the eastern and western Pacific. *Integrative and Comparative Biology*, 49, 114-126.
- Williams, J.D., Boyko, C.B., 2012. The global diversity of parasitic isopods associated with crustacean hosts (Isopoda: Bopyroidea and Cryptoniscoidea). *PLoS ONE*, 7, e35350. doi:10.1371/journal.pone.0035350.
- Williams, J.D., Boyko, C.B., Madad, A.Z., submitted. Branchial parasitic isopods (Crustacea: Isopoda: Bopyridae: Pseudioninae) of hermit crabs (Crustacea: Decapoda: Paguroidea) from the western Pacific, with descriptions of a new genus and three new species. *Raffles Bulletin of Zoology*.
- Zariquiey Alvarez, R., 1958. Decápodos Españoles. XII. Acción de un bopirido sobre los caracteres sexuales de los especies del gén. *Munida* Leach. *Investigación Pesquera*, 11, 101-104.