

Mediterranean Marine Science

Vol 20, No 3 (2019)



Consequences of the experimental removal of *Sabella spallanzanii* (Gmelin, 1791) from the fouling assemblage of a Mediterranean harbour

CATALDO PIERRI, PAOLO COLANGELO, MICHELA DEL PASQUA, CATERINA LONGO, ADRIANA GIANGRANDE

doi: [10.12681/mms.19732](https://doi.org/10.12681/mms.19732)

To cite this article:

PIERRI, C., COLANGELO, P., DEL PASQUA, M., LONGO, C., & GIANGRANDE, A. (2019). Consequences of the experimental removal of *Sabella spallanzanii* (Gmelin, 1791) from the fouling assemblage of a Mediterranean harbour. *Mediterranean Marine Science*, 20(3), 476–486. <https://doi.org/10.12681/mms.19732>

Supplementary Data

Consequences of the experimental removal of *Sabella spallanzanii* (Gmelin, 1791) from the fouling assemblage of a Mediterranean harbour

Cataldo PIERRI, Paolo COLANGELO, Michela Del PASQUA, Caterina LONGO and Adriana GIANGRANDE

Mediterranean Marine Science, 2019, 20 (3)

Table S. List of benthic taxa detected during the manipulative experimental research, with the relative occurrence frequencies in the 4 sampling times (T1, T2, T3, T4) and in the 2 subsets referred to treatment S+ (with *Sabella spallanzanii*) and controls S- (without *S. spallanzanii*).

			S-				S+			
			T1	T2	T3	T4	T1	T2	T3	T4
Chlorophyta	<i>Bryopsis pennata</i>	Lamouroux, 1809	0.385	0.06	0	0	0.34	0.007	0	0
	<i>Chaetomorpha linum</i>	(Müller) Kützing, 1845	0	0	0	0	0	0	0	0.001
	<i>Ulva intestinalis</i>	Linnaeus, 1753	0.099	0.011	0	0	0.075	0	0	0
	<i>Ulva rotundata</i>	Bliding, 1968	0	0	0	0.001	0	0	0	0.002
Ochrophyta	<i>Dictyopteris polypodioides</i>	(De Candolle) Lamouroux, 1809	0	0	0	0	0.003	0	0	0
	<i>Dictyota dichotoma</i>	(Hudson) Lamouroux, 1809	0.007	0	0	0	0.014	0	0	0
	<i>Elachista fucicola</i>	(Velley) Areschoug, 1842	0	0	0.001	0.001	0	0	0.001	0
Rhodophyta	<i>Aglaothamnion feldmanniae</i>	Halos, 1965	0	0	0	0	0	0.002	0	0.001
	<i>Amphiroa rigida</i>	Lamouroux, 1816	0	0	0	0.003	0	0	0	0.002
	<i>Amphiroa</i> sp.	Lamouroux, 1812	0	0.001	0	0	0	0.001	0	0.001
	<i>Antithamnion cruciatum</i>	(Agardh) Nägeli, 1847	0.367	0.168	0.05	0.059	0.318	0.075	0.014	0.029
	<i>Ceramium tenerimum</i>	(Martens) Okamura, 1921	0	0.019	0.001	0.035	0	0.014	0	0.026
	<i>Chylocladia verticillata</i>	(Lightfoot) Bliding, 1928	0	0	0	0	0	0	0	0.001
	Encrusting Calcified Rhodophytes		0.003	0.009	0.018	0.01	0.003	0.004	0.005	0.008
	<i>Eupogodon planus</i>	(Agardh) Kützing, 1845	0	0	0	0	0	0	0.001	0
	<i>Nitophyllum albidum</i>	Ardissonne, 1875	0	0.003	0	0	0	0.002	0	0.002
	<i>Peyssonnelia squamaria</i>	(Gmelin) Decaisne ex Agardh, 1842	0	0	0	0.005	0	0	0	0.004
	<i>Spyridia filamentosa</i>	(Wulfen) Harvey, 1833	0	0.001	0.001	0	0	0.002	0	0
Porifera	<i>Leucetta</i> sp.	Haeckel, 1872	0	0.001	0	0.017	0	0.001	0	0.017
	<i>Paraleucilla magna</i>	Klautau, Monteiro & Borojevic, 2004	0	0.013	0.007	0.044	0.001	0.023	0.005	0.03
	<i>Sycon raphanus</i>	Schmidt, 1862	0	0	0	0.002	0	0.001	0	0.004
Cnidaria	<i>Aglaophenia picardi</i>	Svoboda, 1979	0	0.001	0.001	0	0	0.001	0	0
	<i>Clytia linearis</i>	(Thorneley, 1900)	0	0.003	0	0	0	0.001	0	0
	<i>Corydendrium parasiticum</i>	(Linnaeus, 1767)	0	0.001	0.001	0.02	0	0	0.007	0.023
	<i>Eudendrium racemosum</i>	(Cavolini, 1785)	0	0.014	0.006	0.016	0	0.013	0.006	0.008
	<i>Halecium</i> sp.	Oken, 1815	0	0	0.001	0	0	0	0	0
	<i>Hydrodendron mirabile</i>	(Hincks, 1866)	0	0	0	0.003	0	0	0	0.009
	<i>Obelia dichotoma</i>	(Linnaeus, 1758)	0	0.004	0.004	0.009	0	0.003	0.006	0.021
Mollusca	<i>Anomia ephippium</i>	Linnaeus, 1758	0	0.001	0.007	0.024	0	0.001	0.008	0.029
	<i>Hiatella rugosa</i>	(Linnaeus, 1767)	0	0	0	0.001	0	0	0	0
	<i>Limaria hians</i>	(Gmelin, 1791)	0	0	0	0.003	0	0	0	0.004
	<i>Mimachlamys varia</i>	(Linnaeus, 1758)	0	0.001	0.001	0.011	0	0.001	0.003	0.012
	<i>Musculus costulatus</i>	(Risso, 1826)	0	0.001	0.001	0	0	0	0.001	0

	<i>Musculus subpictus</i>	(Cantraine, 1835)	0	0	0.001	0.002	0	0	0.001	0.001
	<i>Mytilus galloprovincialis</i>	Lamarck, 1819	0.032	0.155	0.29	0.126	0.087	0.258	0.342	0.169
	<i>Ostrea edulis</i>	Linnaeus, 1758	0	0	0	0.006	0	0	0.002	0.006
	<i>Ruditapes decussatus</i>	(Linnaeus, 1758)	0	0	0	0	0	0	0	0.001
Annelida	<i>Branchiomma luctuosum</i>	(Grube, 1870)	0	0.011	0.03	0.12	0	0.026	0.038	0.114
	<i>Hydroides elegans</i>	(Haswell, 1883)	0.005	0.102	0.044	0.022	0.004	0.106	0.04	0.016
	<i>Polycirrus aurantiacus</i>	Grube, 1860	0	0.001	0.001	0.002	0	0	0	0.003
	<i>Sabella pavonina</i>	Savigny, 1822	0	0	0.002	0.004	0	0.001	0.016	0.005
	<i>Spirobranchus lamarcki</i>	(Quatrefages, 1866)	0	0.003	0.025	0.014	0	0.003	0.019	0.018
Bryozoa	<i>Amathia lendigera</i>	(Linnaeus, 1758)	0	0	0	0.003	0	0	0	0
	<i>Bugula neritina</i>	(Linnaeus, 1758)	0.033	0.003	0.008	0.001	0.088	0.002	0.003	0.002
	<i>Bugulina avicularia</i>	(Linnaeus, 1758)	0	0	0.001	0	0	0.001	0	0
	<i>Bugulina fulva</i>	(Ryland, 1960)	0.031	0.012	0.015	0.003	0.034	0.018	0.011	0.005
	<i>Bugulina stolonifera</i>	(Ryland, 1960)	0	0	0.005	0.001	0	0	0.01	0.001
	<i>Celleporaria brunnea</i>	(Hincks, 1884)	0.007	0	0	0	0.005	0	0	0
	<i>Cradoscrupocellaria bertholletii</i>	(Audouin, 1826)	0	0.003	0.001	0	0	0.002	0.003	0
	<i>Crisia eburnea</i>	(Linnaeus, 1758)	0	0.002	0.002	0.036	0	0.001	0.007	0.017
	<i>Cryptosula pallasiana</i>	(Moll, 1803)	0	0	0	0.011	0	0	0	0.025
	<i>Schizobrachiella sanguinea</i>	(Norman, 1868)	0.005	0.013	0.013	0.008	0.004	0.005	0.011	0.005
	<i>Schizoporella errata</i>	(Waters, 1878)	0	0.067	0.065	0.087	0	0.08	0.032	0.077
	<i>Scrupocellaria scruposa</i>	(Linnaeus, 1758)	0.013	0.001	0	0	0.018	0.001	0	0
Arthropoda	<i>Microeuraphia depressa</i>	(Poli, 1791)	0.005	0.007	0.006	0.003	0.004	0.004	0.006	0.006
Chordata	<i>Ascidia malaca</i>	(Traustedt, 1883)	0	0	0.001	0	0	0	0.001	0
	<i>Ascidiella aspersa</i>	(Müller, 1776)	0.001	0	0	0.012	0.002	0	0	0.031
	<i>Botryllus schlosseri</i>	(Pallas, 1766)	0	0.006	0	0.001	0	0.003	0.003	0
	<i>Ciona intestinalis</i>	(Linnaeus, 1767)	0	0	0.003	0	0	0	0.002	0
	<i>Clavelina lepadiformis</i>	(Müller, 1776)	0.001	0.065	0.058	0.073	0.001	0.104	0.051	0.028
	<i>Lissoclinum weigelei</i>	Lafargue, 1968	0	0	0	0	0	0	0.001	0
	<i>Phallusia mammillata</i>	(Cuvier, 1815)	0	0	0	0.023	0	0.001	0.008	0.001
	<i>Pyura microcosmus</i>	(Savigny, 1816)	0	0.005	0.072	0.236	0	0.012	0.052	0.172
	<i>Styela canopus</i>	(Savigny, 1816)	0	0	0	0	0	0	0.001	0
	<i>Styela plicata</i>	(Lesueur, 1823)	0.006	0.227	0.224	0.036	0.009	0.212	0.202	0.043
	<i>Trididemnum cereum</i>	(Giard, 1872)	0.001	0.01	0.034	0.008	0.001	0.013	0.083	0.019
	<i>Trididemnum inarmatum</i>	(Drasche, 1883)	0	0	0.002	0	0	0	0.001	0