

Mediterranean Marine Science

Vol 21, No 1 (2020)



Inventory and the biogeographical affinities of Annelida Polychaeta in the Algerian coastline (Western Mediterranean)

ALI BAKALEM, PATRICK GILLET, JEAN-PHILIPPE PEZY, JEAN-CLAUDE DAUVIN

doi: [10.12681/mms.20408](https://doi.org/10.12681/mms.20408)

To cite this article:

BAKALEM, A., GILLET, P., PEZY, J.-P., & DAUVIN, J.-C. (2020). Inventory and the biogeographical affinities of Annelida Polychaeta in the Algerian coastline (Western Mediterranean). *Mediterranean Marine Science*, 21(1), 157–182.
<https://doi.org/10.12681/mms.20408>

Inventory and the biogeographical affinities of Annelida Polychaeta in the Algerian coastline (Western Mediterranean)

Ali BAKALEM¹, Patrick GILLET², Jean-Philippe PEZY³and Jean-Claude DAUVIN³

¹ Ecole Nationale Supérieure Agronomique (ENSA), Avenue Hassan Badi, 16200 El Harrach, Algiers, Algeria

² Equipe de recherche Mer, Molécules, Santé EA 2160, Faculté des Sciences, UCO, 44 rue Rabelais, F-49008 Angers, France

³ Normandie Univ., UNICAEN, Laboratoire Morphodynamique Continentale et Côtière M2C,
UMR 6143 M2C, 24 rue des Tilleuls, F-14000 Caen, France

Corresponding author: jean-claude.dauvin@unicaen.fr

Handling Editor: Melih ÇINAR

Received: 10 May 2019; Accepted: 24 December 2019; Published on line: 27 April 2020

Abstract

From collected data analyzed to produce an inventory of all polychaetes in Algerian waters, the diversity of this group was estimated as 534 species. The largest families are Syllidae (66 species), Spionidae (37 species) and Terebellidae (27 species). The presence of those listed species along the Algerian coast was compared with their occurrence in nine other areas of the Mediterranean Sea, in the Black Sea and in four other more distant seas around the world. Comparisons are also made with respect to the biogeographical origin for each species. The polychaete fauna of the Algerian coast is among the richest of the Mediterranean Sea and comparable to the diversity reported for the French Mediterranean continental shelf and that in the Aegean Sea, but higher than that found in the nearby coastlines of Morocco and Tunisia. Most of the species found have an Atlantic origin and are present in the western Mediterranean Sea. This current inventory includes eight non-indigenous polychaete species found in the Algerian waters.

Keywords: Biogeography; inventory; non-native species; benthos.

Introduction

Establishing an inventory of marine species remains a challenging task despite the availability of numerous records (especially via the Internet), and the existing data already provided by taxonomists and ecologists. Coll *et al.* (2010) carried out the first complete inventory of marine species in the Mediterranean Sea at the end of the 2010s.

The entire Mediterranean region, with its smaller distinct areas, has long been recognized as rich in terms of marine biodiversity (Coll *et al.*, 2010, 2012; Altobelli *et al.*, 2017). In their review for the Mediterranean Sea, Coll *et al.* (2010) reported 16,848 different marine species, including 10,902 invertebrates. Moreover, these authors stressed that these numbers were probably still an underestimate and needed to be supplemented to include additional species found in some southern areas of the sea as well as those living in the deeper regions of the eastern basin. This gap in their records mostly concerns the coasts of North Africa, mainly relating to Libya and Algeria. Such an apparent lack of available information is probably due to the numerous data included in early

publications written in French and the large amount of grey literature including PhD theses that are often not accessible via the web.

The annelid polychaetes are amongst the most diversified of invertebrates and several publications report inventories for this group at the local, national and regional scale, including the coastlines along Morocco and Tunisia in the southern part of the western basin of the Mediterranean Sea.

Some inventories for polychaetes cover the entire Mediterranean Sea (Arvanitidis *et al.*, 2002; Castelli *et al.*, 2008; Coll *et al.*, 2010), while others are limited to certain areas: Moroccan waters (both the Atlantic and Mediterranean coasts) (Gillet, 2017), the Mediterranean coast of France (Laubier & Paris, 1962; Dauvin *et al.*, 2006), Italian waters (Castelli *et al.*, 1995, 2008), the Adriatic Sea (Musco, 2012; Mikac, 2015), the Tunisian coasts (Ayari *et al.*, 2009; Zaâbi *et al.*, 2012, Mosbahi *et al.*, 2017), the Aegean Sea (Ergen, 1976; Simboura & Nicolaidou, 1994, 2001; Arvanitidis, 2000; Ergen *et al.*, 2002, 2006; Faulwetter *et al.*, 2017), the Levantine Sea (Laubier, 1966; Ben-Eliahu, 1995; Çınar & Ergen, 2003; Çınar, 2005, 2007, 2009; Çınar *et al.*, 2003, 2014;

Mutlu *et al.*, 2010; Faulwetter *et al.*, 2017; Dorgham *et al.*, 2013, 2014), the Turkish coastline (Çinar *et al.*, 2014) and the Black Sea (Arvaniditis *et al.*, 2002; Şahin & Çinar, 2012). However, no complete inventory is available for the 1200km of Algerian coastline. The first studies of polychaetes in Algerian waters took place at the turn of the 20th century. Viguier (1886 to 1920) and Gravier & Dantan (1928) described several new species of polychaetes in the Bay of Algiers, including the genus *Alciope* Viguier, 1886, the genus *Frennia* Viguier, 1912 (*F. dubia* Viguier, 1912, *F. propinquua* Viguier, 1920), the genus *Iospilus* Viguier, 1886 (*I. phalacroides* Viguier, 1886), the genus *Maupasia* Viguier, 1886 (*M. coeca* Viguier, 1886), the genus *Paraiospilus* Viguier, 1911 (*P. affinis* Viguier, 1911), and the species *Nereis icosiensis* Gravier & Dantan, 1928. Most of the polychaetes found during this period were reported by Fauvel (1923, 1927a, 1927b) in the French publication ‘Faune de France, polychètes errantes et sédentaires’. Fauvel (1927b) wrote:

‘A l’exception des Mémoires de Viguier sur le plancton de la Baie d’Alger et de quelques mentions par Marenzeller (1888) on ne trouve que de bien rares indications disséminées dans d’autres travaux. Jamais à notre connaissance la Faune des Polychètes des Côtes d’Algérie n’a encore été l’objet de recherches suivies et cela est regrettable car cette étude fournirait sûrement bien des observations intéressantes’ translated as ‘Other than the memoirs of Viguier on the plankton in the Bay of Algiers and some references by Marenzeller (1888), it is very rare to find anything disseminated in other studies. To the best of our knowledge, polychaete fauna off the coast of Algeria has never been the subject of research, which is regrettable, as such a study would surely provide some interesting observations’.

Marcel (1962) studied the annual cycle of *Perinereis cultrifera*. In studies on the phanerogam meadows on the Algerian coastline near Bou Ismail, Le Gall (1969) identified 72 polychaetes species, whereas Falconetti (1970) listed 64 polychaete species from the maërl beds in the same bay. Separately, some data was acquired for the waters in the harbour of Algiers (Bakalem & Romano, 1983, 1988; Rebzani-Zahaf *et al.*, 1988; Rebzani-Zahaf, 1990, 1992). Populations of some nereids in shallow hard substrates from the eastern part of the Algerian coast were investigated by Rouabah & Scaps (2003) and also by Daas *et al.* (2010). In other work, Younsi *et al.* (2010) recorded the polychaetes of economic interest and Meghlaoui *et al.* (2015a,b) investigated the effects of local pollution on polychaetes. More recently Bakalem (2008) and Grimes (2010) studied the communities of soft bottom of all Algerian gulfs, bays and harbours, in reported work that includes inventories of polychaetes.

Despite this level of scientific interest, there has still been no complete inventory of all polychaetes found in Algerian waters. A total of 26 polychaete species were initially listed in the two monographs entitled ‘Faune de France’ (Fauvel, 1923, 1927a), whilst a further 69 species were later added by Gravier & Dantan (1928). Many further species have since been identified during the PhD research of Bakalem (2008) (444 species) and then by Grimes (2010) (506 species). All these previous documents from an essential basis for the current listing of polychaetes of the Algerian coast; other even more recent publications, (including as yet unpublished data and personal communications), have been used to supplement the complete inventory reported here.

The main objectives of this current study were:

- to present a current list of all polychaetes species




Fig. 1: Map of the Algerian coast with identification of the four study sectors with cited locations.

- recorded along the whole Algerian coastline;
- to attribute geographical affinities for each of the Algerian polychaete species;
- to compare the inventory of polychaete in the coastal waters off Algeria with similar studies compiled for other areas of the Mediterranean Sea.

Materials and Methods

General characteristics of the Algerian coastal zone

The Algerian coastline (Fig. 1) covers a variety of different habitats, ranging from rocky shorelines (sometimes with high cliffs), to sandy beaches and dunes found in many of the bays. The Algerian continental shelf is narrow, in contrast with those found in the north-western basin of the Mediterranean Sea, and especially the Gulf of Lion and the northern part of the Adriatic Sea. The distribution of soft sediments shows an increasing gradient (in terms of silt content) with sea depth; the seabed composition was characterized by a succession of fine sand, muddy sand, sandy mud and pure mud such as observed in the coastal seabed in the bay of Algiers and the bay of Bou Ismail (Bakalem, 2008; Bakalem *et al.*, 2020). Coarse sand and gravel make up the seabed near rocky shores, especially in the vicinity of the headlands bordering the bays (Grimes *et al.*, 2016). The shallow waters are also characterized by some specific habitats such as “seagrass zones” including *Posidonia oceanica* meadows and coralligenous formations both types having a high natural heritage value. The composition of the fauna in these habitats is characterized by a high diversity, and such areas are protected in many countries bordering the Mediterranean Sea against a range of environmental threats (Coll *et al.*, 2010, 2012; Altobelli *et al.*, 2017).

The circulation of the sea currents along the Algerian coast is mainly influenced by North Atlantic surface waters, which penetrate into the Mediterranean Sea through the Strait of Gibraltar. As they continue to progress eastward, the Atlantic waters gradually mix into the Mediterranean waters with its higher salinity. This movement is highlighted by the progressive impoverishment of Atlantic species travelling from the western to the eastern part of the Algerian coast (Dauvin *et al.*, 2013).

Data sources

The current study covers all the polychaetes that have been recorded along the Algerian coast, which runs 1200km from the Morocco to Tunisia. The study has been largely restricted to the continental shelf since there was little opportunity to access the continental slope or the offshore bathyal zones (Fig. 1). Zooplankton sampling in Algiers Bay (sector 2 in Fig. 1) enabled the collection of pelagic polychaetes and the subsequent description of several new species (Viguier, 1886, 1911, 1912, 1920; Gravier & Dantan, 1928). More recent studies have fo-

cused mainly on the soft-bottom communities found in the bays and gulfs of the Algerian continental shelf (Vaisière & Fredj, 1963; Le Gall, 1969; Falconetti, 1970; Bakalem, 2008; Grimes, 2010).

These results show a progressive change in the physico-chemical marine water characteristics along the the Algerian coast from the entrance of the Atlantic current in the West and its attenuation moving to the eastern part of the coast. Since the end of the 1970, the studies of Bakalem (2008), Grimes (2010) and Dauvin *et al.* (2013) on the soft-bottom macrobenthic communities of the Algerian shallow habitats from west to east have enabled the identification of four main sectors along the coast (see Dauvin *et al.*, 2013). This classification takes into account numerous characteristics based on: 1) the hydrology and local marine currents (Furnestin & Allain, 1962a,b,c; Benzohra & Millot, 1990; Millot, 1987, 1988; Millot *et al.*, 1990), 2) the hydrology, nutrients and phytoplankton (Bernard, 1952, 1956, 1958), and 3) biological data including ichthyology and macrobenthos (Le Danois, 1925; Maurin, 1962; Vaissière & Fredj, 1963; Bakalem, 2008; Grimes, 2010).

Sector 1 is located in the West before the Almeria-Oran front with hydrological characteristics similar than those of the Atlantic waters (Fig. 1). Sectors 3 and 4 correspond to Mediterranean hydrological water characteristics while the sector 2 corresponds to an intermediate situation between the western sector 1 and the eastern sectors.

Data reported here follows the model set out by Dauvin *et al.* (2013), where the presence of polychaetes is reported for the four sectors, which enables the identification of changes in diversity along the Algerian coast from west to east. Sector 1 comprises the western part of the Algerian coast from the Moroccan border to Cap Tenes, including the bays of Ghazaouet, Oran and Arzew. Sector 2 is the central part of the coastline from Cap Tenes to Azeffoun, including the bays of Bou Ismail and Algiers. Sector 3 is defined as the eastern part of Algerian coast from Azeffoun to the Cap de Garde including the Bejaia, Jijel and Skikda bays. Finally, Sector 4 corresponds to the most eastern Algerian coast from Cap de Garde to the Tunisian border, including the bays of Annaba and El Kala (Fig. 1). The coastlines of Sectors 1 and 2 are similar in length (300-350 km), whilst the coastline of Sector 3 is longer (400 km) and that of Sector 4 shorter (150 km).

Nomenclature and polychaete classification

The nomenclature and classification of polychaetes have been the subject of several successive studies (Fauchald & Rouse, 1997; Rouse & Fauchald, 1997; Hansson, 1998; Rouse & Gonzalo, 2016; Weigert & Bleidorn, 2016). The present work adopts the most recent form of the classification system for families based on polychaete phylogeny (Weigert & Bleidorn, 2016).

The individual species were classified by family, and these are listed alphabetically in Table 1. The species are validated following the nomenclature and classification given by the World Register of Marine Species (WoRMS)

Table 1. List of the Polychaetes recorded along the Algerian coast. The species are classified following the alphabetical order of the families. Year of collection: 1886, 1911, 1912, 1920: Viguier; 1923, 1927: Fauvel; 1928, Gravier & Dantan; 1963: Vaissière & Fredj; 1969: Le Gall; 1970: Falconetti; 2008: Bakalem; 2010: Grimes. S1: Sector 1... Status: Qs: Questionable species; Sc: species complex; NIS: Non-Indigenous Species. G: geographical origin, for legend see Figure 5; 1, 2, ... geographical areas, for legend see Figure 6.

	Species	Authority	Year of record	Status	G	S1	S2	S3	S4	1	2	3	4	5	6	7	8	9	10	11	12	13
Acoetidae	<i>Eupanthalis kinbergi</i>	McIntosh, 1876	1970	eamr	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Panhialis oerstedi</i>	Kinberg, 1856	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Polyodontes maxillosus</i>	(Ranzani, 1817)	2008	camrip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
Acroiridae	<i>Acrocirrus frontifilis</i>	(Grube, 1860)	2008	camip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Macrochaeta clavicornis</i>	(M. Sars, 1835)	2008	eam	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Naiades cantrainei</i>	Delle Chiaje, 1828	1886	camip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Rhynchoneereella capitata</i>	(Greeff, 1876)	1886	amip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Rhynchoneereella petersii</i>	(Langerhans, 1880)	1886	eam	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Torrea candida</i>	(Delle Chiaje, 1841)	1886	enip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Vanicadis longissima</i>	(Levinsen, 1885)	2008	enip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Alkmaria romijni</i>	Horst, 1919	2010	Qs	eam	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Amage adspersa</i>	(Grube, 1863)	2008	eam	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Amage gallasi</i>	Marion, 1875	2008	am	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
Ampharetidae	<i>Ampharete grubei</i>	Malmgren, 1865	2008	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Ampharete octocirrata</i>	(Sars, 1835)	2008	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Amphicteis gunneri</i>	(M. Sars, 1835)	2008	eam	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Anoplothrus gracilis</i>	(Malmgren, 1866)	2008	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Melitta palmata</i>	Grube, 1870	1970	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Arenticola marina</i>	Quarefages, 1866	2008	Qs	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Chloetia venusta</i>	(McIntosh, 1885)	1923	audouin & milne edwards,	eam	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Hermodice carunculata</i>	1830	2008	camr	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Hipponeoe gaudichaudi</i>	McIntosh, 1885	2008	enip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Notorygosa megalops</i>	Linnaeus, 1758	2008	Qs	eamr	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Aphrodita acutepennis</i>	(Savigny in Lamark, 1818)	2008	mip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
Aphroditidae	<i>Laemonice hystrix</i>	(Baird, 1865)	1963	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Pomponenia chrysocoma</i>	(Linnaeus, 1758)	1970	eam	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Arenicoloides branchialis</i>	(Audouin & Milne Edwards, 1833)	2008	eam	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
Arenicolidae	<i>Arenicoloides grubii</i>	Claparède, 1868	2008	e	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Branchiomaldane vincenti</i>	Langerhans, 1881 [as vincentii]	2008	cam	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
Bonellidae	<i>Maxmulleria gigas</i>	(Müller M., 1852)	2008	Qs&Sc	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Capitella capitata</i>	(Fabricius, 1780)	2008	(Mesnil, 1897)	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Capitella giardi</i>	Langerhans, 1880	2008	camrip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Capitella minima</i>	(Grube, 1846)	2008	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Dasybranchus caducus</i>	Eisig, 1887	1969	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Dasybranchus gjelae</i>	(Claparede, 1864)	2008	e	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Heteromastus filiformis</i>	Eisig, 1887	2010	NIS	mip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
Capitellidae	<i>Notomastus abernans</i>	Sars, 1851	1969	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Notomastus lineatus</i>	Claparède, 1869	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Notomastus profundus</i>	(Eisig, 1887)	2008	eam	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Pereziella clymenoides</i>	Harmelin, 1968	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Pseudocapitella incerta</i>	Fauvel, 1913	2008	e	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Pseudoleiocapitella fawveli</i>	Harmelin, 1964	2008																			

Continued

Table 1 continued

Continued

Table 1 continued

	Species	Authority	Year of record	Status	G	S1	S2	S3	S4	1	2	3	4	5	6	7	8	9	10	11	12	13
Euphrasinidae	<i>Euphrasina armatillo</i>	Sars, 1851	2008	earnr	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Euphrasina foliosa</i>	Audouin & H Milne Edwards, 1833	1928	earnr	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
Fabriciidae	<i>Euphrasina myrtosa</i>	Savigny in Lamarcq., 1818	2010	Qs	earnrip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Fabricia stellaris</i>	(Müller, 1774) (Bourne, 1883)	2008	Qs	earn	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
Flabelligeridae	<i>Pseudofabricia filamentosa</i>	(Malmgren, 1867) (Day, 1963)	2008	Sc	earnrip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Diplocirrus glaucus</i>	M. Sars, 1829 (Otto, 1820)	2010	em	earnrip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
Glyceraidae	<i>Flabelligera diplochaitus</i>	(Delle Chiaje, 1841) (Müller, 1776) (Claparède, 1869)	2008	Qs	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Pherusa monilifera</i>	(Sars in Sars, 1872)	2008	em	earn	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
Glyceridae	<i>Pherusa plumosa</i>	(Sars in Sars, 1872)	2008	em	earn	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Pironis erica</i>	(O.F. Müller, 1776)	1969	c	earn	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
Glycinidae	<i>Therochaeta flabellata</i>	(Örsted, 1843)	2008	earn	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Glycera alba</i>	Quatrefages, 1850	1970	em	earn	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
Goniadidae	<i>Glycera capitata</i>	Quatrefages, 1866	1928	earn	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Glycera fallax</i>	Grube, 1863	1928	c	earn	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
Hesionidae	<i>Glycera lapidum</i>	Schnarda, 1861	1928	c	earnrip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Glycera tessellata</i>	Lamarcq., 1818	2008	c	earn	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
Iospiidae	<i>Glycera tridactyla</i>	(Malmgren, 1866)	2008	earn	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Glycera unicornis</i>	Audouin & H Milne Edwards, 1866	1969	earnrip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
Lacydoniidae	<i>Glycinida nordmanni</i>	Örsted, 1843	2008	earn	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Gonioda emerita</i>	Gonioda maculata	1833	Örsted, 1843	2008	earnrip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
Lopadorrhynchidae	<i>Gonioda norvegica</i>	Gypsis propinqua	Marion & Bobretzky, 1875	2008	earn	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Hesione splendida</i>	Leocrates clapanedi	(Costa in Claparède, 1868)	2008	earnrip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
Lophiidae	<i>Nereimyra punctata</i>	Oxydromus agilis	(Müller, 1788)	2010	Qs	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Oxydromus flexuosus</i>	Oxydromus pallidus	(Ehlers, 1864)	1928	Qs	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
Lophiidae	<i>Psamnathë fusca</i>	Psamnathë fusca	(Delle Chiaje, 1827)	1928	earn	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Syllidia armata</i>	Johnston, 1836	2008	c	earnrip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
Lophiidae	<i>Iospius phalacroides</i>	Quatrefages, 1866	1928	earnrip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Paraiospius affinis</i>	Viguier, 1886	1886	am	earn	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
Lophiidae	<i>Phalacrophorus pictus</i>	Viguier, 1911	1920	e	earn	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Lacydoniidae Lacydonia miranda</i>	Greeff, 1879	1886	Qs	earn	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
Lopadorrhynchidae	<i>Lopadorrhynchus krohnii</i>	Marion & Bobretzky, 1875	2008	earnrip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Maupasia coeca</i>	(Claparède, 1870)	1886	em	earn	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
Pedinoecidae	<i>Pedinoecoma curtaum</i>	Viguier, 1886	1923	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Pelagobia longicirrata</i>	Reibisch, 1895	2010	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	

Continued

Table 1 continued

	Species	Authority	Year of record	Status	G	S1	S2	S3	S4	1	2	3	4	5	6	7	8	9	10	11	12	13
Lumbri- ridae	<i>Hilbigneris gracilis</i>	(Ehlers, 1868)	1969	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Lumbrierides acuta</i>	(Verrill, 1875)	2008	Qs	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Lumbrieriopsis paradoxa</i>	(Saint-Joseph, 1888)	1970	Qs	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Lumbrieris coccinea</i>	(Renier, 1804)	1970	Qs	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Lumbrieris inflata</i>	Moore, 1911	2008	Qs	camip	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Lumbrieris labrofimbriata</i>	Saint-Joseph, 1888	2008	Qs	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Lumbrieris laurrelli</i>	Audouin & Milne Edwards, 1834	2008	Qs	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Lumbrieris nonatoi</i>	Ramos, 1976	2008	Qs	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Lumbrieris pertensi</i>	Carrera-Parr, 2001	2008	NIS	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Scoleleoma emandibulata malbiti</i>	(Ramos, 1976)	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
Magelonidae	<i>Scoleleoma fragilis</i>	(O.F. Müller, 1776)	1923	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Scoleleoma funchalensis</i>	(Kinberg, 1865)	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Scoleleoma impatiens</i>	(Claparède, 1868)	1969	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Magelona allenii</i>	Wilson, 1958	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Magelona filiformis</i>	Wilson, 1959	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Magelona minuta</i>	Eliason, 1962	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Magelona mirabilis</i>	(Johnston, 1865)	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Magelona wilsoni</i>	Glémarec, 1966	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Axiothella constricta</i>	(Claparède, 1868)	2008	Sc	e	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Axiothella rubrocincta</i>	(Johnson, 1901)	2008	Sc	e	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
Maldanidae	<i>Chirrimia biceps</i>	(M. Sars, 1861)	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Euchymene collaris</i>	(Claparède, 1869)	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Euchymene lombrioides</i>	(Quatrefages, 1866)	1970	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Euchymene oerstedi</i>	(Claparède, 1863)	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Euchymene palmaritana</i>	(Grube, 1840)	1970	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Heterochymene robusta</i>	Arwidsson, 1906	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Johnstonia chymenoidea</i>	Quatrefages, 1866	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Leiochone leiopygos</i>	(Grube, 1860)	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Leiochone tricirrata</i>	Bellan & Reys, 1967	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Macrochymene santonensis</i>	(Rioja, 1917)	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
Nephytidae	<i>Maldane glebifex</i>	Grube, 1860	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Maldane sarsi</i>	Malmgren, 1865	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Metasyachis gotti</i>	(Izuka, 1902)	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Microtmaldane ornithocheaeta</i>	Mesnil, 1897	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Nicomache lumbicalis</i>	(Fabricius, 1780)	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Petaloprotus tericolus</i>	Quatrefages, 1866	1969	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Praxillella affinis</i>	(M. Sars in G.O. Sars, 1872)	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Praxillella gracilis</i>	(M. Sars, 1861)	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Praxillella lophoseta</i>	(Orlandi, 1898)	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Praxillella praetermissa</i>	(Malmgren, 1865)	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
Nephystidae	<i>Aglaophamus agilis</i>	(Langerhans, 1880)	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Inermonephytis inermis</i>	(Ehlers, 1887)	2008	Sc	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Nephtys caeca</i>	(Fabricius, 1780)	2008	Sc	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Nephtys ciliata</i>	(Müller, 1788)	2010	Sc	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Nephtys cirrosa</i>	Ehlers, 1868	2008	Sc	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Nephtys hombergii</i>	Savigny in Lamarck, 1818	1969	Sc	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Nephtys hyspéricis</i>	McIntosh, 1900	1923	Sc	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
Nephytidae	<i>Nephtys incisa</i>	Malmgren, 1865	2008	Sc	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Nephtys paradoxata</i>	Malin, 1874	2008	Sc	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	

Continued

Table 1 continued

Species	Authority	Year of re- cord	Status	G	S1	S2	S3	S4	1	2	3	4	5	6	7	8	9	10	11	12	13
<i>Alitta succinea</i>	(Leuckart, 1847)	2008	camip c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Ceratonereis (Compostetia) costae</i>	(Grube, 1840)	1970	amip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Ceratonereis (Compostetia) hircincola</i>	(Eisig, 1870)	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Eunereis longissima</i>	(Johnston, 1840)	1928	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Hedistie diversicolor</i>	(O.F. Müller, 1776)	2008	Sc	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Micronereis variegata</i>	Claparède, 1863	1969	Sc	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Neanthes acuminata</i>	(Ehlers, 1868)	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Neanthes fucata</i>	(Savigny, 1822)	1928	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Neanthes nubila</i>	(Savigny, 1822)	1928	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Nereis finchaleensis</i>	(Langerhans, 1880)	1928	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Nereis rava</i>	Ehlers, 1868	1928	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Nereis splendida</i>	Grube, 1840	1928	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Nereis zonata</i>	Malmgren, 1867	1928	Sc	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Perinereis cultirifera</i>	(Grube, 1840)	1928	Sc	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Perinereis macropus</i>	(Claparède, 1870)	1928	Sc	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Perinereis oliveriae</i>	(Horst, 1889)	2008	Sc	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Perinereis tenuisetsis</i>	(Faivel, 1915)	1928	Sc	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Platynereis coccinea</i>	(Delle Chiaje, 1822)	2008	Sc	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Platynereis dumetillii</i>	(Audouin & Milne Edwards, 1833)	1928	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Websterinereis glauca</i>	(Claparède, 1870)	1928	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Arabella geniculata</i>	(Claparède, 1868)	2008	Sc	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Arabella iricolor</i>	(Montagu, 1804)	1969	Sc	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Drilonereis filum</i>	(Claparède, 1868)	1969	Sc	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Halla parthenopeia</i>	(Delle Chiaje, 1822)	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Notocirrus scoticus</i>	McIntosh, 1869	1970	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Aponuphis bilineata</i>	(Baird, 1870)	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Aponuphis brementi</i>	(Faivel, 1916)	1969	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Diopatra neapolitana</i>	Delle Chiaje, 1841	1963	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Hyainoecta tubicola</i>	(O.F. Müller, 1776)	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Nothria conchylega</i>	(Sars, 1835)	1969	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Onuphis eremita</i>	Audouin & Milne Edwards, 1833	1969	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Paradiopatra quadricuspis</i>	(M. Sars in G.O. Sars, 1872)	1969	Qs	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Rhamphobrachium (Spinigerum) brevibrachiatum</i>	(Ehlers, 1875)	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Armandia cirrhosa</i>	Filippi, 1861	1928	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Armandia polyophthalma</i>	Kükenthal, 1887	1969	Sc	e	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Ophelia anomurexi</i>	Bellan & Costa, 1987	2008	Sc	e	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Ophelia barbata</i>	Faivel, 1927	2010	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Ophelia bicornis</i>	Savigny, 1822	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Ophelia neglecta</i>	Schneider, 1892	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Ophelia radiata</i>	(Delle Chiaje, 1828)	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Ophelia roscensis</i>	Augener, 1910	1970	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Ophelia acuminata</i>	Örsted, 1843	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
<i>Polyopthalmus pictus</i>	(Dujardin, 1839)	1928	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+

Continued

Table 1 continued

	Species	Authority	Year Of record	Status	G	S1	S2	S3	S4	1	2	3	4	5	6	7	8	9	10	11	12	13
	<i>Naineris laevigata</i>	(Grube, 1855) (Audouin & H Milne Edwards, 1833)	1969	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Orbinia latreillii</i>	(Savigny, 1822)	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Orbinia serulata</i>	(Claparède, 1868)	1969	Sc	emip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Phyllo foetida ligustica</i>	(Orlandi, 1896)	1927	Sc	emip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Phyllo grubei</i>	(McIntosh, 1910)	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Phyllo norvegicus</i>	(M. Sars in G.O. Sars, 1872)	2008	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Protoanicia oerstedii</i>	(Claparède, 1864)	2008	em	amr	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Scoloplos (Leodamas) chevalieri</i>	(Fauvel, 1902)	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Scoloplos armiger</i>	(Müller, 1776)	1969	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Scoloplos typicus</i>	(Eisig, 1914)	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Owenia oculata</i>	(Zachs, 1923)	2008	Qs	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Myriochela heeri</i>	Malmgren, 1867	2008	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Owenia fusiformis</i>	Delle Chiaje, 1844	1969	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Paracydonia paradoxa</i>	Fauvel, 1913	2008	emip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Aricidea (Acmira) assimilis</i>	Tebble, 1959	2008	Sc	amip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Aricidea (Acmira) catherinae</i>	Laubier, 1967	2008	Sc	emp	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Aricidea (Acmira) cerrutii</i>	Laubier, 1966	2008	Sc	am	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Aricidea (Acmira) lopezi</i>	Berkeley & Berkeley, 1956	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Aricidea (Acmira) simonae</i>	Laubier & Ramos, 1974	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Aricidea (Aricidea) capensis bansei</i>	Laubier & Ramos, 1974	2008	NIS	amip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Aricidea (Aricidea) fragilis</i>	Webster, 1879	2008	Sc	amip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Aricidea (Aricidea) pseudoarticulata</i>	Hobson, 1972	2008	Sc	emip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Aricidea (Strelzovia) claudiae</i>	Laubier, 1967	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Aricidea (Strelzovia) mediterranea</i>	(Laubier & Ramos, 1974)	2008	Sc	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Aricidea (Strelzovia) pseudanae</i>	Katzmann & Laubier, 1975	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Aricidea (Strelzovia) suecica meridionalis</i>	Laubier & Ramos, 1974	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Aricidea annae</i>	Laubier, 1967	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Cirrophorus branchiatus</i>	Hartman, 1965	2010	Sc	emip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Levinseria gracilis</i>	Ehlers, 1908	2008	Sc	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Paradoneis armata</i>	(Tauber, 1879)	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Paradoneis harpagonea</i>	Glémarec, 1966	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Paradoneis lyra</i>	(Storch, 1967)	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Paraorides neapolitana</i>	(Southern, 1914)	1969	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Paraorides fulgens</i>	(Cerruti, 1909)	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Paraorides paucibranchiata</i>	(Levinsen, 1884)	2008	Sc	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Paraorides tenera</i>	Cerruti, 1909	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Amphicteine auricomata</i>	Grube, 1873	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Lagus koreni</i>	(O.F. Müller, 1776)	1969	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Pectinaria belgica</i>	(Pallas, 1766)	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Petta pusilla</i>	Malmgren, 1866	1970	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Photoleia inornata</i>	Johnston, 1839	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+

Continued

Table 1 continued

Species	Authority	Year of record	Status	G	S1	S2	S3	S4	1	2	3	4	5	6	7	8	9	10	11	12	13	
<i>Eulalia bilineata</i>	(Johnston, 1840)	2008	Qs	ean	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Eulalia pusilla</i>	Ørstedt, 1843	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Eulalia viridis</i>	(Linnaeus, 1767)	2008	Sc	enr	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Eumida punctifera</i>	(Grube, 1860)	2008	Sc	en	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Eumida sanguinea</i>	(Ørstedt, 1843)	1928	Sc	enip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Hyperocone foliosa</i>	(Quatreffages, 1855)	2008	ean	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Lugia pierophora</i>	(Ehlers, 1864)	1970	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Mysta picta</i>	(Quatreffages, 1866)	1969	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Mysta siphodonta</i>	(Delle Chiaje, 1830)	1969	ean	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Mystides borealis</i>	Théel, 1879	2008	Qs	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Nerephylla paretii</i>	Blainville, 1828	2010	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Nerephylla pusilla</i>	(Claparède, 1870)	1970	ean	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Notophyllum foliosum</i>	(Sars, 1835)	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Paranaitis kosteriensis</i>	(Malmgren, 1867)	2008	Qs&Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Phyllocopte greenlandica</i>	Ørsted, 1842	2008	ean	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Phyllocopte lineata</i>	(Claparède, 1870)	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Phyllocopte maculata</i>	(Linnaeus, 1767)	1928	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Phyllocopte madeirensis</i>	Langerhans, 1880	1969	Sc	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Phyllocopte mucosa</i>	Ørsted, 1843	1970	ean	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Pseudomyctides limbata</i>	(Saint-Joseph, 1888)	1970	emip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Pterocirrus limbatus</i>	(Claparède, 1868)	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Pterocirrus macroceros</i>	(Grube, 1860)	1928	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Ancistrosyllis groenlandica</i>	McIntosh, 1878	2008	ean	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Ancistrosyllis hamata</i>	(Hartman, 1960)	2008	mip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Pilargis verrucosa</i>	Saint-Joseph, 1899	2008	ean	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Sigambra tentaculata</i>	(Treadwell, 1941)	2008	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Poecilochae-</i>																						
<i>Poecilochae-</i>	<i>Poecilochaeetus fulgoris</i>	Claparède in Ehlers, 1875	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Poecilochae-</i>	<i>Poecilochaeetus serpens</i>	Allen, 1904	1928	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Polygordii-</i>	<i>Polygordius appendiculatus</i>	Fraipont, 1887	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Polygordii-</i>	<i>Polygordius laetus</i>	Schneider, 1868	(Delle Chiaje, 1827)	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
			Michaelsen, 1892	2010	mip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
			(M. Sars, 1861)	2008	camip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
			Viguier, 1912	1912	e	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
			Viguier, 1920	1920	ean	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
			(Pallas, 1766)	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
			McIntosh, 1876	1970	ean	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
			(Grube, 1860)	2008	emip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
			(Grube, 1840)	1970	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
			(Johnston, 1839)	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
			(McIntosh, 1876)	2008	emip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
			(Grube, 1863)	1928	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
			(Ehlers, 1864)	1969	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
			(Grube, 1840)	2008	emip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
			Potts, 1910	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
			(Montagu, 1808)	1970	emip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
			(Linnaeus, 1758)	1969	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
			McIntosh, 1876	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
			(Malmgren, 1867)	1970	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
			(Delle Chiaje, 1830)	1969	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
			Savigny, 1822	1970	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
			(Ehlers, 1864)	1970	emip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	

Continued

Table 1 continued

	Species	Authority	Year of record	Status	G	S1	S2	S3	S4	1	2	3	4	5	6	7	8	9	10	11	12	13
Pontodoridae	<i>Pontodora pelagica</i>	Greeff, 1879	1886	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
Protodrilidae	<i>Megadrilus purpureus</i>	(Schneider, 1868)	2010	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Lygdamis muratus</i>	(Allen, 1904)	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
Sabellariidae	<i>Phalacrostemma cidariophilum</i>	Marenzeller, 1895	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Sabellaria alveolata</i>	(Linnaeus, 1767)	1969	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Sabellaria spinulosa</i>	(Leuckart, 1849)	2010	Qs	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Acromegalomma vesticulosum</i>	(Montagu, 1813)	2008	e	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Amphigiena mediterranea</i>	(Leydig, 1851)	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Bispira volutacornis</i>	(Montagu, 1804)	2008	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Branchiomma bombyx</i>	(Dalyell, 1853)	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Dialychine danerificta</i>	(Tovar-Hernández, Lucciano, Giangrande, 2007)	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Dialychine acutica</i>	(Claparède, 1869)	1969	e	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Dialychine collaris</i>	(Langerhans, 1881)	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Euchone rosea</i>	(Langerhans, 1884)	1970	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Eucione rubrocincta</i>	(Sars, 1862)	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Euratella salmacidis</i>	(Claparède, 1869)	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Hypsiconus stictophthalmos</i>	(Grube, 1863)	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Jasminoira candela</i>	(Grube, 1863)	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Jasminoira caudata</i>	Langerhans, 1880	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Jasminoira elegans</i>	Saint-Joseph, 1894	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Laonome kroyeri</i>	Malmgren, 1866	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Myxicola aesthetica</i>	(Claparède, 1870)	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Myxicola infundibulum</i>	(Montagu, 1808)	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Notaulax phaeotaenia</i>	(Schmarda, 1861)	2008	emrip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Amphicorina armstrongi</i>	(Claparède, 1864)	2008	emr	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Pseudopotamilla reniformis</i>	(Bruguière, 1789)	2008	eanip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Sabella discifera</i>	Graube, 1874	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Sabella pavonina</i>	Savigny, 1822	2008	eamip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Sabella spallanzanii</i>	(Gmelin, 1791)	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
Scalibregmatidae	<i>Ascerchocheilus intermedius</i>	(Saint-Joseph, 1894)	2010	Sc	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Scalibregma inflatum</i>	Rathke, 1843	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Sclerocheilus minutus</i>	Grube, 1863	1928	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	

Continued

Table 1 continued

	Species	Authority	Year Of record	Status	G	S1	S2	S3	S4	1	2	3	4	5	6	7	8	9	10	11	12	13
	<i>Apomatus similis</i>	Mariot & Bobretzky, 1875 (O. F. Müller, 1776)	2008	eam	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Ditrypa arietina</i>	(Faunel, 1923)	2008	NIS	eamip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Ficopomatus enigmaticus</i>	(Verrill, 1873)	2008	NIS	eamip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Hydroides dianthus</i>	Mörch, 1863	2008	NIS	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Hydroides dirampha</i>	(Haswell, 1883)	2008	NIS	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Hydroides elegans</i>	Gunnerus, 1768	2008	eam	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Hydroides norvegica</i>	Zibrowius, 1968 [original]	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Hydroides pseudouncinata</i>	(Philippi, 1844)	1927	2008	eam	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Metavernilia multicristata</i>	(Fabricius, 1779)	2008	emip	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Placostegus tridentatus</i>	(Lamarck, 1818)	2010	emr	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Protula intestinum</i>	(Montagu, 1803)	2010	emr	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Protula tubularia</i>	(Huxley, 1855)	2008	emr	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Serpulidae</i>	Claparède, 1870	2008	emr	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Salinacina incrassans</i>	(Marenzeller, 1893)	1927	2008	emr	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Seminervilia agglutinata</i>	(O. G. Costa, 1861)	2008	eam	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Serpula crenata</i>	Langerhans, 1880	2010	eamr	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Serpula concharium</i>	Rioja, 1917	2008	emr	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Serpula lobiancoi</i>	Linnaeus, 1767	1928	eam	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Serpula vermicularis</i>	(Zibrowius, 1968)	2008	eam	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Spiraserpula massiliensis</i>	(Quatrefages, 1866)	2008	eam	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Spirobranchus lamarchei</i>	(Linnaeus, 1758)	2008	eamr	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Spirobranchus triqueuer</i>	(Philippi, 1844)	2008	emr	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Vermilopsis infundibulum</i>	(O. G. Costa, 1861)	2008	emip	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Vermilopsis labata</i>	(Grube, 1862)	2008	eam	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Vermilionis striaticeps</i>	(Claparède, 1868)	1923	eam	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Euthaliensis oculata</i>	(Peters, 1854)	1970	em	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Fimbriosthenelais minor</i>	(Pruvot & Racovitza, 1895)	1970	em	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Labioleania yhenni</i>	(Malmgren, 1867)	2008	Qs	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Leanira hystericis</i>	Ehlers, 1874	2008	em	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Neoleaniira retrograna</i>	(Orsted, 1845)	1969	em	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Pelogenia arenosa</i>	(Delle Chiaje, 1830)	2008	em	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Pholidoides dorsipapillatus</i>	(Marenzeller, 1893)	1970	em	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Pistone remota</i>	(Southern, 1914)	Audouin & Milne Edwards in Cuvier, 1830	2008	emr	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Sigalion mathildae</i>	Delle Chiaje, 1830	1970	eam	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Sigalion squamosus</i>	(Johnston, 1833)	1969	emip	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Sthenelais boa</i>	(Ehlers, 1864)	1970	em	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Sthenelais limicola</i>	(Hansen, 1878)	2008	em	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Ephesiella abyssorum</i>	(Grefft, 1866)	2010	em	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Sphaerodoriidium claparedii</i>	(Webster & Benedict, 1887)	2008	em	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Sphaerodoriidium minutum</i>	(Rathke, 1843)	2008	Qs	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Sphaerodoriidium gracilis</i>	(M. Sars, 1851)	2008	em	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+
	<i>Sphaerodoridae</i>	<i>Sphincteridae</i>	<i>Sphaerodoridae</i>	Continued																		
	<i>Sphincter arcticus</i>																					

Table 1 continued

Species	Authority	Status	G	S1	S2	S3	S4	1	2	3	4	5	6	7	8	9	10	11	12	13
<i>Aonides oxycephala</i>	(Sars, 1862)	1969	enip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Aonides paucibranchiata</i>	Southern, 1914	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Aurospio banyulensis</i>	(Aubier, 1966)	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Boccardia polybranchia</i>	(Haswell, 1885)	2008	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Boccardella ligurica</i>	(Ferromière, 1898)	2010	emrip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Dipohydora armata</i>	(Langerhans, 1880)	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Dipohydora coeca</i>	(Örsted, 1843)	2008	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Dipohydora flava</i>	(Claparède, 1870)	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Dipohydora giardi</i>	(Mesnil, 1893)	2008	emip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Dipohydora quadrilobata</i>	(Jacobi, 1883)	2010	Qs&Sc	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Laonice cirrata</i>	(M. Sars, 1851)	1927	Sc	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Malacoceros fuliginosus</i>	(Claparède, 1868)	1969	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Malacoceros grandidi</i>	Quairefages, 1843	1969	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Malacoceros tetracerus</i>	(Schmarda, 1861)	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Microspio mecnikowianus</i>	(Claparède, 1869)	1969	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Paraprionospio pinnata</i>	(M. Sars, 1851)	2008	Sc	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Polydora ciliata</i>	(Johnston, 1838)	2008	Sc	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Polydora hoplura</i>	Claparède, 1868	2008	camip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Prionospio caspensi</i>	Laubier, 1962	2008	e	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Prionospio cirrifera</i>	Wirén, 1883	1928	Qs	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Prionospio decipiens</i>	Söderström, 1920	2010	Qs	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Prionospio ehlersi</i>	Fauvel, 1928	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Prionospio fallax</i>	Claparède, 1867	2008	Qs	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Prionospio steenstrupi</i>	Malmgren, 1867	1969	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Pseudopolydora antennata</i>	(Claparède, 1869)	2008	Sc	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Pygospio elegans</i>	Claparède, 1863	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Scolelepis (Parascolelepis) tridentata</i>	(Southern, 1914)	2008	(Rioja, 1918)	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Scolelepis (Scolelepis) cantabria</i>	(Audouin & Milne Edwards, 1833)	2008	(O.F. Müller, 1806)	1969	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Scolelepis (Scolelepis) foliosa</i>	(Mesnil, 1896)	2008	camip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Scolelepis (Scolelepis) squamata</i>	Bobretzky, 1870	1969	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Scolelepis bonieri</i>	(Rioja, 1918)	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Spio decorata</i>	(Claparède, 1870)	2008	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Spio multiculata</i>	Grube, 1860	2008	emr	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Spiophanes bombyx</i>	Laubier, 1964	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Spiophanes kroyeri</i>	(Buchanan, 1890)	2010	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
<i>Streblospio styrusoli</i>	(Ranzani, 1817)	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
Sternaspidae	<i>Sternaspis scutata</i>	(Claparède, 1863)	1886	Qs	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Amblyssyllis formosa</i>	Claparède, 1868	2010	Sc	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Anoplosyllis edentula</i>	(Gravier, 1900)	2008	emrip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Branchiosyllis exilis</i>	(Dujardin, 1851)	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Brania pusilla</i>	(Malmgren, 1867)	2008	Qs	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Epigamia alexandri</i>	Ehlers, 1864	1928	emr	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Euryssyllis tuberculata</i>	Marenzeller, 1875	2008	eamr	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Eusyllis assimilis</i>	Manjón & Bobretzky, 1875	2008	eam	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Eusyllis blomstrandii</i>	Castelli, Badalamenti & Lardici, 1987	2010	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Eusyllis lamelligera</i>	(Claparède, 1864)	2010	enip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Exogone (Parexogone) cognetti</i>																			
	<i>Exogone (Sylline) brevipes</i>																			

Continued

Table 1 continued

Continued

Table 1 continued

	Species	Authority	Year of record	Status	G	S1	S2	S3	S4	1	2	3	4	5	6	7	8	9	10	11	12	13
Syllidae	<i>Syllis vivipara</i>	Krohn, 1869	2008	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Symmerosyllis lamelligera</i>	(Saint-Joseph, 1886)	2010	eam	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Trypanosyllis (Trypanosyllis) coelitaca</i>	Claparède, 1868	1928	Sc	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Trypanosyllis zebra</i>	(Grube, 1860)	1928	Sc	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Virchowia clavata</i>	Langerhans, 1879	1886	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Amphithrite cirrata</i>	Müller, 1776	2008	eam	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Amphithrite rubra</i>	(Risso, 1826)	2008	em	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Amphithrite variabilis</i>	(Risso, 1826)	1970	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Eupolymlnia nebulosa</i>	(Grabe, 1860)	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Eupolymlnia nesidensis</i>	(Montagu, 1819)	2008	Sc	eamrip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Lanicus conchilega</i>	(Delle Chiaje, 1828)	1969	Sc	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Lisilla loveni</i>	(Pallas, 1766)	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Neoamphithrite affinis</i>	Malmgren, 1866	2010	Qs	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Neoamphithrite edwardsi</i>	(Quairefages, 1865)	2008	Qs	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Neoamphithrite figuilus</i>	(Dalyell, 1853)	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Nicolea venustula</i>	(Montagu, 1819)	1970	Sc	c	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Nicolea zostericola</i>	Ørsted, 1844	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Pista cretacea</i>	(Grube, 1860)	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Pista cristata</i>	(Müller, 1776)	1969	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Pista maculata</i>	(Dalyell, 1853)	2008	NIS	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Pista unibranchia</i>	Day, 1963	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Polycirrus aurantiacus</i>	Grube, 1860	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Polycirrus denticulatus</i>	Saint-Joseph, 1894	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Polycirrus haematodes</i>	(Claparède, 1864)	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Polycirrus medusa</i>	Grube, 1850	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Proctea graffi</i>	(Langerhans, 1884)	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Sireblosoma bairdi</i>	(Malmgren, 1866)	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Terebella lapidaria</i>	Linnaeus, 1767	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Thelepus cincinnatus</i>	(Fabricius, 1780)	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Thelepus serosus</i>	(Quairefages, 1866)	2008	Sc	emrip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	<i>Thelepus triserialis</i>	(Grube, 1855)	1927	Sc	emrip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	Tomopteri-dae	<i>Tomopteris (Johnstonella) levipes</i>	(Grefft, 1879)	1886	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
		<i>Tomopteris (Johnstonella) pacifica</i>	(Izuka, 1914)	1886	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	Traviidae	<i>Travisia forbesii</i>	Johnston, 1840	2010	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
	Tri-chobranchidae	<i>Octobranchus lingulatus</i>	(Grube, 1863)	2008	Sc&Sc	emrip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
		<i>Terebellides stroemii</i>	Sars, 1835	1927	Sc	emrip	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
		<i>Thrichobranchus glacialis</i>	Malmgren, 1866	2008	Sc	em	+	+	+	+	+	+	+	+	+	+	+	+	+	+	+	
		Total number of species			371	430	300	195	216	492	180	507	534	406	305	462	488	189	348	110	151	

(www.marinespecies.org; accessed on 25 January 2019). Comments are added for any questionable species present in either Algerian and Mediterranean waters. Uncertainty arose mainly for species complexes belonging to the Cirratulidae and Spionidae families.

In the current study, an abundant literature was consulted before attributing a status to each Algerian polychaete species (Çınar, 2003, 2005, 2009, 2013; Çınar & Ergen, 2003; Çınar *et al.*, 2003, 2014; Zenetos *et al.*, 2005, 2010, 2011, 2012, 2017; Gravili *et al.*, 2010; Gil, 2011; Occhipinti-Ambrogi *et al.*, 2011; Evangelopoulos *et al.*, 2015; Corriero *et al.*, 2016; Galil *et al.*, 2016; Ounifi-Ben Amor *et al.*, 2016; Faulwetter *et al.*, 2017; López & Richter, 2017).

To specify the status of a species found in Algerian waters, it was classified into one of three groups as proposed by Zenetos *et al.* (2011) and used by Faulwetter *et al.* (2017): Non-Indigenous Species (NIS), Questionable Species (Qs) and Species Complexes.

“Questionable species” are defined as either those reported as found in Algerian waters but whose presence still needs to be confirmed, or those, which have an unresolved taxonomic status. In this study, if records of any species are in doubt, they are not automatically classified as NIS if they have never been previously reported as NIS anywhere else in the Mediterranean. On the other hand, species already reported as NIS from other studies in the Mediterranean Sea but which still retain a questionable status, are listed as “NIS Questionable” to highlight this unresolved matter in the case of further assessments.

Species complexes correspond to those species known to form a complex of cryptic (morphologically indistinguishable but genetically different) or pseudo-cryptic (morphologically very similar) species (Nygren & Pleijel, 2011). The term ‘Species complex’ indicates that the name has been applied to a species as defined above.

Geographical patterns of polychaetes from the Algerian coasts

The distribution of the 534 polychaete species found in Algerian waters is categorised taking into account the distinct geographical areas of the Mediterranean and related maritime regions as proposed by Coll *et al.* (2010) and modified by Bakalem *et al.* (2020). These are, 1: African Atlantic Coast (from Cap Spartel to Cap Blanc); 2: North-Eastern Atlantic European waters; 3: Alboran Sea; 4: North-Western Mediterranean; 5: Algeria South-Western Mediterranean; 6: Ionian Sea; 7: Tunisian shelf and Libya; 8: Adriatic Sea; 9: Aegean Sea; 10: Black Sea; 11: Levantine Sea; 12: Red Sea and 13: Indo-Pacific oceans.

Each polychaete species found on the Algerian coast, is classified as one of fifteen groups reflecting their geographical origin (Bakalem *et al.*, 2020): eam (Mediterranean, European and African Atlantic species); em (Mediterranean and European Atlantic species); am (Mediterranean and African Atlantic coast species); e (endemic species); c (cosmopolitan species); emr (Mediterranean, European Atlantic and Red Sea species); emp (Mediterranean, European Atlantic and Indo-Pacific species); amr (African Atlantic, Mediterranean and Red Sea species); amip (African Atlantic, Mediterranean and Indo-Pacific species); mip (Mediterranean and Indo-Pacific species); emrip (European Atlantic, Mediterranean, Red Sea and Indo-Pacific species); eamrip (European and African Atlantic, Mediterranean, Red Sea and Indo-Pacific species); eamip (European and African Atlantic, Mediterranean and Indo-Pacific species); eamr (European and African Atlantic, Mediterranean and Red Sea species) and mrip (Mediterranean, Red Sea and Indo-Pacific species).

Hierarchical Cluster Analysis (HCA) based on Sørensen’s coefficient for the presence/absence of the species found in each of the four sectors of the Algerian coast and in the 13 geographical areas was carried out along with the construction of a dendrogram using the group average algorithm generated from the PRIMER V6 software (Clarke & Gorley, 2006).

Results

List of species

Studies on the presence of polychaetes along the Algerian coasts have been sporadic. Viguier (1886, 1911, 1912, 1920) and later, Gravier & Dantan (1928), all collected pelagic species from the Bay of Algiers, whilst Fauvel (1923, 1927a,b) focused on benthic species in the waters surrounding Algiers and Annaba. These focused studies only found a small number of polychaete families and species: between 8 and 18 families covering 25 to 69 species. More recently, Le Gall (1969) and Falconetti (1970) studied the Phanerogam meadows and maërl community respectively, (located in the Bay of Bou Ismail), and reported between them a total of 27 families and 72 species.

The present inventory of polychaete species is mainly compiled from data collected in three successive periods: the first was from 1886 to 1928, (leading to 28 families and 96 species), the second from 1963 to 1970, (leading to 44 families and 196 species), and finally from 1975 to 2010, (leading to the final list with 58 families and 534 species). During these three periods, the number of recorded species increased from 96 to 534. The PhD theses of Bakalem (2008) and Grimes (2010), which together, covered most of the soft-bottom habitats in the gulfs, bays and harbours along the Algerian coasts, listed the greatest numbers of polychaetes with 54 and 51 families and 444 and 506 species, respectively. It is noteworthy that there were no studies reporting new polychaetes between 1928 and 1963, and none have taken place since 2011.

The Syllidae (66 species representing 12% of the total diversity) is the most diversified family, followed by the Spionidae (37 species, 7%), the Terebellidae (27 species, 5%), the Serpulidae (25 species, 4.5%), the Polynoidae (24 species, 4.5%), the Sabellidae (23 species, 4%) and the Paraonidae (23 species, 4%).

Ten families (each with more than 20 species) account




Fig. 2: Number of polychaete species per family classified by decreasing number of species.




Fig. 3: Results of cluster analysis (Sørensen similarity) on the presence and absence of species, in the four defined sectors (S1 to S4 see Fig. 1) along the Algerian coast.




Fig. 4: Number of species present of the ten most diversified families in the four defined sectors (see Fig. 1) along the Algerian coast.

for 55% of polychaetes diversity (Fig. 2); seven families are contain between 10 and 20 species, whilst most of the remaining families (41) have fewer than 10 species; moreover, ten families are only represented by a single species (Fig. 2).

Out of the 534 species reported, 22 are planktonic, including five families with a strictly pelagic life style: Alciopidae (5 species), Lopadorhynchidae (4), Iosipilidae (3), Tomopteridae (2) and Pontodoridae (1). Five species, two of *Ancistrosyllis* (Pilargidae), two of Polynoidae and one Nereididae belonging to families of benthic polychaetes are all pelagic.

The previously defined geographical sector 2 was found to be the richest in terms of polychaete diversity with 56 families and 430 species (representing 81% of the total recorded species). Sector 1 with 48 families (371

species, 69%) supported the second largest diversity. The eastern sectors were relatively poorer in terms of diversity with 49 families (300 species, 56%) in sector 3 and 40 families (195 species, 37%) in sector 4. Figure 3 compares the similarities between the four sectors; sector 4 is very different from the three other sectors, whilst sectors 1 and 2 (in the western and central parts, which display the richest diversity), also show the greatest similarity. Families are represented by a similar number of species in all except sector 3 where the family of Syllidae is more diversified (Fig. 4).

Questionable species (Qs) and species complexes

Among the 534 species recorded along the Algerian coast, 45 correspond to the status “questionable species” and 38 are considered nowadays to be species complexes, these numbers respectively 8 and 6.5% of the total richness amongst the polychaete (Table 1). Further data will be required to confirm their presence in Algerian waters. It should be noted that most of these species have already been previously reported in other polychaete inventories (e.g., Faulwetter *et al.*, 2017).

The following taxa are classified as both questionable species and species complex: *Axiothella rubrocincta* (Johnson, 1901), *Capitella capitata* (Fabricius, 1780), *Chaetozone setosa* Malmgren, 1867, *Laonice cirrata* (M. Sars, 1851), *Lepidonotus squamatus* (Linnaeus, 1758), *Phyllodoce groenlandica* Örsted, 1842, *Protocirrineris chrysoderma*, (Claparède, 1868), and *Terebellides stroemii* Sars, 1835.

Ampharete acutifrons (Grube, 1860) may have been confused with *Ampharete lindstroemi* Hessle, 1917 in the Mediterranean Sea (Mikac, 2015) and in samples found in Algerian waters. It forms a species complex in its North Atlantic and Pacific distributions (Faulwetter *et al.*, 2017). *Ampharete grubei* Malmgren, 1865 is often reported as *A. acutifrons*, which is also present in Greek waters (Faulwetter *et al.*, 2017).

The specimens reported as *Chaetozone setosa* Malmgren, 1867 in the Mediterranean Sea and along the Algerian coast, corresponds to several possible species such as *Chaetozone gibber* Woodham & Chambers,

1994, *Chaetozone corona* Berkeley & Berkeley, 1941 or *Chaetozone carpenteri* McIntosh, 1911: it probably does not exist in the Mediterranean Sea (Çinar & Ergen, 2007; Faulwetter *et al.*, 2017).

Specimens of polychaetes from the Mediterranean previously named as *Terebellides stroemii* almost certainly belong to different species including *Terebellides gracilis* (Malm, 1874) and *Terebellides mediterranea* Parapar, Mikac and Fiege, 2013 (Faulwetter *et al.*, 2017).

Non-Indigenous Species (NIS)

Out of the 534 species reported along the Algerian coast, eight have the status NIS in the Mediterranean Sea: *Aricidea (Aricidea) fragilis* (Webster, 1879), *Ficopomatus enigmaticus* (Fauvel, 1923), *Hydrodoides dirampha* Mörch, 1863 and *H. elegans* (Haswell, 1883), *Lumbrineris perkinsi* Carrera-Parra, 2001, *Metasynchis gotoi* (Izuka, 1902), *Notomastus aberans* Day, 1957 and *Pista unibranchia* Day, 1963.

Seven species found along the Algerian coast were reported in the list compiled by Zenetos *et al.* (2017). They were *Ficopomatus enigmaticus* (Fauvel, 1923), *Hydrodoides dirampha* Mörch, 1863 and *H. elegans* (Haswell, 1883), *Lumbrineris perkinsi* Carrera-Parra, 2001, *Metasynchis gotoi* (Izuka, 1902), *Pista unibranchia* Day, 1963, and *Notomastus aberans* Day, 1957. *Aricidea (Aricidea) fragilis* (Webster, 1879) was a typically west-Atlantic species. It was present in the Adriatic and Tyrrhenian seas and in the Strait of Sicily (Langeneck *et al.*, 2018). The




Fig. 5: Number of species present in the different geographical groups. Legend: amr: African Atlantic, Mediterranean and Red Sea species, amip: African Atlantic, Mediterranean and Indo-Pacific species, mip: Mediterranean and Indo-Pacific species, mrip: Mediterranean, Red Sea and Indo-Pacific species, am: Mediterranean and African Atlantic species, emrip: European Atlantic, Mediterranean, Red Sea and Indo-Pacific species, eamr: European Atlantic, African Atlantic, Mediterranean, Red Sea and Indo-Pacific species, emr: European Atlantic, Mediterranean and Red Sea species, eamip: European Atlantic, African Atlantic, Mediterranean and Indo-Pacific species, emip: European Atlantic, Mediterranean and Indo-Pacific species, e: endemic species, c: cosmopolitan species, eam: European Atlantic, Mediterranean and African Atlantic species, em: European Atlantic and Mediterranean species.

status of NIS is also given for this species present along the Algerian coast.

Geographical patterns of Polychaetes of the Algerian coasts

Most of the Polychaetes found in Algerian waters have an Atlantic-Mediterranean origin (315 species, 50% of the total number recorded). 203 species (38%) are common to the Mediterranean Sea and the North-Eastern Atlantic coast, whereas 107 species (20%) are common to the Mediterranean Sea, the North-Eastern Atlantic and African Atlantic coast and only three species share a Mediterranean and African Atlantic distribution (Fig. 5). The cosmopolitan species (61) and endemic species (24) make up 11.5 and 5% of all species respectively. 122 species (23% of the total) show a wide distribution across the North-Eastern Atlantic Ocean, the Mediterranean and Red Seas as well as being found in the Indo-Pacific region. Only six species (1%) share a Mediterranean, Red Sea and Indo-Pacific distribution.

A total of 492 species from the European Atlantic Ocean and 507 from the North-Western Mediterranean Sea were reported as found along the Algerian coast; these species make up 92 and 95% respectively of the total inventory. The Aegean and Adriatic Seas have a high number of polychaete species in common with the Algerian coasts, respectively 488 (91%) and 462 (86%) species, whereas a significant number are in common with the Ionian Sea (406 species and 76%), the Levantine Sea (348 species, 66%) and with the Tunisian shelf extending to Libya (305 species, 57%). By contrast, the Alboran Sea (180 species, 34%) and the Black Sea (189 species, 35%) both share a relatively low number common of species with the Algerian coast. The number of species shared between the Algerian coast and the Red Sea (110 species, 21%) and the Indo-Pacific (151 species, 28%) was low.

The geographical distributions of the 534 species recorded from the Algerian coast (Table 1) are compared with their presence in 12 other geographical areas. Differences were analysed using the Sørensen coefficient (Fig. 6). The dendrogram showing the Sørensen similarity between the 13 geographical areas suggests the division of these areas into two main groups. The first group comprises only two areas, the Red Sea (labelled 12 in Fig. 6) and the Indo-Pacific region (13) whereas the second group encompasses all the 11 other areas. Within this second group, the Alboran Sea (3), the Black Sea (10) and the African Atlantic coast (1) show the lowest level of similarities with the other areas in the same group. However, the remaining eight areas display a high level of similarity and form an distinct pattern with the Tunisian shelf and Libya (7), Levantine Sea (11), Ionian Sea (6), Adriatic Sea (8), North-Eastern Atlantic European (2), Algeria (South-Western Mediterranean) (5), North-Western Mediterranean (4) and Aegean Sea (9). The highest similarities (>90%) are found between the North-Eastern Atlantic European waters (2) and the Algerian south-western Mediterranean (5) and also between the North-Western Mediterranean (4) and the Aegean Sea (9) (Fig. 6).

Discussion

The estimated total number of polychaete species varies depending on the author (Arvanitidis *et al.*, 2002; Coll *et al.*, 2010); the most recent census of polychaete species for the Mediterranean Sea is that of Faulwetter *et al.* (2017) who recorded 1,105 species. With 534 species, the polychaetes of the Algerian coasts represent 48% of the polychaete fauna found in the Mediterranean Sea and this diversity is appreciable in comparison with those noted in other areas of the Mediterranean Sea (Table 2).

However, the inventory of Faulwetter *et al.* (2017) do




Fig. 6: Results of cluster analysis (Sørensen similarity) on the species presence and absence in 13 geographical regions. 1: African Atlantic Coast (from Cap Spartel to Cap Blanc); 2: North-Eastern Atlantic European waters; 3: Alboran Sea; 4: North-Western Mediterranean; 5: Algerian South-Western Mediterranean; 6: Ionian Sea; 7: Tunisian Shelf and Libya coastline; 8: Adriatic Sea; 9: Aegean Sea; 10: Black Sea; 11: Levantine Sea; 12: Red Sea and 13: Indo-Pacific Oceans.

Table 2. Comparison of inventories of polychaetes species found in different areas of the Mediterranean Sea.

Author (s)	Areas	Species	Comments
Arvanitidis <i>et al.</i> , 2002	Mediterranean and Black seas	1,036	All species recorded including questionable species
Coll <i>et al.</i> , 2010	Mediterranean Sea	1,122	Few data in the southern part of the Mediterranean Sea
Faulwetter <i>et al.</i> , 2017	Mediterranean Sea	1,105	Number standardised to currently accepted names in 2017
Gillet, 2017	Morocco (Atlantic and Mediterranean)	321	Continental shelf species
Dauvin <i>et al.</i> , 2006	Mediterranean French continental shelf	623	Continental shelf species
	Gulf of Lions	641	Continental and bathyal species
	Gulf of Lions	487	Continental shelf species
	Provence-Alpes-Côte d'Azur coast	516	Continental shelf species
	Corsica	256	Continental shelf species, few data around Corsica
Castelli <i>et al.</i> , 2008	Italian waters	876	Including questionable species
Mikac, 2015	Adriatic Sea	764	Including questionable species
Ayari <i>et al.</i> , 2009	Tunisia	289	Few data for the southern part of the Tunisia
Zaâbi <i>et al.</i> , 2012	Cap Bon Peninsula, Tunisia	212	North-east coast of Tunisia
Zaâbi <i>et al.</i> , 2012	Tunisia	375	New cumulative number
Fersi <i>et al.</i> , in preparation	Tunisia	443	Indicative total number
Arvanitidis, 2000	Aegean Sea	592	0-1000 m, except Turkish coast
Simboura & Nicolaïdou, 2001	Greek waters	753	Census from 1891 to 2000
Arvanitidis <i>et al.</i> , 2002	Western Basin of the Mediterranean	884	All species recorded including questionable species
	Central Basin of the Mediterranean	528	All species recorded including questionable species
	Levantine Basin of the Mediterranean	451	All species recorded including questionable species
	Aegean Sea	493	All species recorded including questionable species
Faulwetter <i>et al.</i> , 2017	Greek waters	836	Including 142 questionable species
Çinar <i>et al.</i> , 2014	Mediterranean coasts of Turkey	705	Including questionable species
Çinar, 2005	Cyprus waters	456	All species recorded including questionable species
Şahin & Çinar, 2012	Black Sea	238	New cumulative number
Dorgham <i>et al.</i> , 2013	Egyptian Mediterranean waters	283	Including questionable species
This study	Algerian waters	534	Species mainly found on the continental shelf

not include 45 of the 534 Algerian species. Moreover, this value of 534 species may also be under-estimated due to: 1) the low level of study and sampling in the eastern part of the Algerian coast (sectors 3 and 4); 2) the few studies of exceptional habitats (including the phanerogams, shallow and deep hard substratum as well as coralligenous formations and the bathyal benthic zone), and 3) the absence of molecular analyses applied to those questionable species and species complexes, those Non-Native species and those species classified as “cosmopolitan”.

The richness of polychaete species along the coast of Algeria is comparable to that reported for the continental shelf of the French Mediterranean coast (Dauvin *et al.*, 2006) and also to that for the Greek part of the Aegean Sea (Arvanitidis, 2000) (Table 2). This richness is higher

than that found in bordering countries such as Morocco, with 321 species listed including the Atlantic coast of Morocco (Gillet, 2017), and Tunisia with 443 species (Ayari *et al.*, 2009; Zaâbi *et al.*, 2012). It is also higher than that observed for the Black Sea (Şahin & Çinar, 2012). Nevertheless, an even greater diversity of polychaetes have been found for Greek waters (Faulwetter *et al.*, 2017), along the Italian coast (Castelli *et al.*, 2008), in Turkish waters (Çinar *et al.*, 2014), and in the Adriatic Sea (Mikac, 2015).

In the coasts of Algeria, the families of Syllidae, Spionidae, Polynoidae, Phyllodocidae, Terebellidae, Serpulidae and Sabellidae, account for most of the diversity, as is the case in along the French Continental shelf (Dauvin *et al.*, 2006), the Greek coasts (Faulwetter *et al.*, 2017), the

Turkish coasts (Çinar *et al.*, 2014), and Adriatic coastline (Mikac, 2015); however, in these areas, there is also the significant presence of the family Maldanidae.

The percentage of questionable species and species complexes remains high in the case of studies along the Algerian coastline (15%), but this number is lower than that given by Faulwetter *et al.* (2017) for such types of species present in Greek waters (23%).

Arvanitidis *et al.* (2002) reviewed 1,036 species of polychaetes for the Mediterranean Sea, including 41% that are considered as cosmopolitan, 29% as Atlantic-Mediterranean and 19% as endemic polychaetes. By comparison, the Mediterranean, European and African Atlantic species are dominant along the Algerian coastal waters accounting for 60%. The European Atlantic-Mediterranean species (38%) and the cosmopolitan species (12%) are the dominant groups on the Algerian coasts. The number of endemic species for Algeria is low (24 species, making up just 5% of the recorded species and indeed, lower than the values reported for other areas of the Mediterranean Sea such as the French Mediterranean coast (Dauvin *et al.*, 2006), the Greek coasts (Simboura & Nicolaïdou, 2001) and the Aegean Sea (Arvanitidis *et al.*, 2002).

The close similarity between the Algerian polychaetes and those found in both the European and African coasts of the Atlantic Ocean reflects the influence of the Atlantic waters along the Moroccan coasts extending towards Tunisia in the south of the Mediterranean Sea, which is inevitably related to the connexion via the Strait of Gibraltar (Fig. 6). This same observation has previously been highlighted for all the macrobenthic species of the Algerian coasts (Dauvin *et al.*, 2013). Moreover, the polychaete fauna in Algerian waters is similar to those of the North-Western basin of the Mediterranean Sea off Spain, France and Italy (Fig. 6). Furthermore, Algerian polychaete fauna displays a great similarity to that of the Aegean Sea (both Greek and Turkish coasts), suggesting a high level of similarity of Polychaete fauna at the scale of the Mediterranean Sea. The Black Sea, due its particular physico-chemical characteristics shows little similarity in terms of polychaete fauna with the different areas of the Mediterranean Sea.

Faulwetter *et al.* (2017) pointed out that the regional lists of species inevitably reflect the research and sampling effort. He added that their reliability depends on the presence of zoological experts who are competent in the identification of polychaetes, noting that some groups or species of polychaetes can be very difficult to identify (Dauvin, 2005). The richness and geographic origin of a species in a given area can be estimated by making comparisons. Faulwetter *et al.* (2017) stressed that each inventory carried out on Polychaetes has given rise to a relatively large number of species reported in the same region. Faulwetter *et al.* (2017) go on to explain that this may be partly due by the actual geographic distribution of species and partly due to the incorrect identification of some species.

Comprehensive revisions and updated identification keys are lacking for many of the taxa studied in the Medi-

terranean yet protocols established a century ago, such as those of Fauvel (1923, 1927a), are still commonly used in many laboratories as the main tools for species identification. It is unfortunate that the level of research on Mediterranean Polychaete Fauna performed in the 1990s was not maintained to the present day. The taxonomic effort remains to be implemented and will be a challenge for polychaete experts around the world. On the other hand, British Synopses, Fauna Iberica and other publications on the taxonomy and phylogeny of polychaetes remain very useful tools for the identification of the Mediterranean polychaetes. Furthermore, molecular analysis needs to be developed to resolve the problems of cryptic and sibling polychaete species, not only for the Mediterranean Sea but also for the wider area of the North-Eastern Atlantic Ocean. The World Register of Marine Species (WoRMS) has become an indispensable tool, not only for naming existing species and problems of synonymy, but also for describing their actual geographical distribution.

The Mediterranean Sea is recognised as a rich area for biodiversity. It is also impacted on by a range of damaging factors such as habitat loss and degradation, climate change, pollution and the invasion of NIS (Coll *et al.*, 2010, 2012; Altobelli *et al.*, 2017). A regular evaluation of the marine biodiversity of target groups (such as the polychaetes) in certain target areas (such as the wide range of soft-bottom habitats in the Bay of Bou Ismail) is strongly recommended. Such monitoring activity can reveal the long-term changes in biodiversity under the increased influence of climatic and human pressures in Algerian coastal environments.

The analyses by Arvanitidis *et al.* (2002) show differences in the Mediterranean Sea between the western basin, where the polychaete studies are satisfactory, and the eastern basin where the knowledge of polychaetes remains insufficient. Furthermore, Spanish, French and Italian inventories cover the northern coasts of the western basin, whereas there is relatively sparse data for the southern coast (North Africa) of this same basin. For the central part of the Mediterranean Sea, these inventories take into account the Ionian Sea (Italy and Greece). Finally, for the eastern basin, the inventories carried out concern mostly the Aegean Sea (Ergen, 1976; Simboura, 1987; Arvanitidis, 1994, 2000), the Levantine Sea, mainly the coasts of Cyprus and Israel (Ben-Eliahu, 1972, 1991, 1995; Ben-Eliahu & Fiege, 1995, 1996; Russo, 1997), the Egyptian coasts (Abd-Elnaby, 1999; Selim, 1978, 1996) and Turkish waters (Ergen & Çinar, 1997). The data relating to these areas remain sporadic.

Since the beginning of new millennium there has been a considerable research effort on polychaetes with a great deal of effort targeted at those areas with a low level of data: Algeria (Bakalem, 2008; Grimes, 2010), Tunisia (Ayari *et al.*, 2009; Zaâbi *et al.*, 2012; Ayari *et al.*, in preparation), Greece (see Faulwetter, 2017), Turkey (see Çinar *et al.*, 2014), Cyprus (Çinar, 2005), and Egypt (Abd-Elnaby, 2008, 2009; Dorgham *et al.*, 2013, 2014). These very recent results have shown that the diversity of polychaete fauna is relatively homogenous at the scale of the whole Mediterranean Sea and that the differences between the

West and East (as noted by Arvanitidis *et al.*, 2002) may only be due to the lack of information and thus not correctly reflect the real situation. Future research using molecular analysis will be crucial to assess the representability of questionable species and species complex, which are numerous amongst the polychaete species reported.

Acknowledgements

The authors wish to thank Dr Colin Burton for the English language revision and Pr. Melih Ertan Çinar, (Ege University, Izmir, Turkey), for his very useful remarks and suggestions with respect to the successive versions of this paper.

References

- Abd-Elnaby, F.A., 1999. Composition and distribution of some bottom fauna associations along the Alexandria coast, Mediterranean Sea. MSc Thesis, Alexandria University, Egypt.
- Abd-Elnaby, F.A., 2008. New recorded species of Magelonidae and Maldanidae (Annelida: Polychaeta) from El-Tina Bay, Mediterranean coast of Egypt. *Egyptian Journal of Aquatic Biology & Fisheries*, 12, 119-141.
- Abd-Elnaby, F.A., 2009. Polychaete Study in Northeastern Mediterranean Coast of Egypt. *World Journal of Fish and Marine Sciences*, 1, 85-93.
- Altobelli, C., Perzia, P., Falautano, M., Consoli, P., Canese, S et al., 2017. Mediterranean banks in EBSA area: Hotspots of biodiversity under threat. *Marine Environmental Research*, 131, 57-68.
- Arvanitidis, C., 1994. Systematic and bionomic study of the macrobenthic Polychaeta (Annelida) of the northern Aegean. PhD thesis, Aristotelian University of Thessaloniki (in Greek).
- Arvanitidis, C., 2000. Polychaete fauna of the Aegean Sea: inventory and new information. *Bulletin of Marine Science*, 60, 73-96.
- Arvanitidis, C., Bellan, G., Drakopoulos, P., Valavanis, V., Dounas, C. et al., 2002. Seascape biodiversity patterns along the Mediterranean and the Black Sea: lessons from the biogeography of benthic polychaetes. *Marine Ecology Progress Series*, 244, 139-152.
- Ayari, R., Muir, A., Aissa, P., Paterson, G., Afli, A., 2009. An updated list of Polychaetous annelids from Tunisian coasts (Western Mediterranean Sea). *Cahiers de Biologie Marine*, 50, 1-13.
- Ayari, R., Fersi, A., Afli, A., Bakalem, A., Baffreau, A., Boudaya, L., Khedri, I., Mosbahi, N., Neifar, L., Dauvin, J.C. (in preparation). New data of Tunisian polychaetous annelids.
- Bakalem A., Romano, J.C. 1983. Pollution et peuplements benthiques dans la région algéroise (baie d'Alger et baie de Bou Ismail). *VI^e Journées Etudes de Pollution Marine, Cannes*, CIESM, 687-697.
- Bakalem, A., Romano, J.C. 1988. Les peuplements benthiques du port d'Alger : 3. Les polychètes. *Rapport de la Commission de la Mer Méditerranée*, 31, 2, 24.
- Bakalem, A., 2008. Les peuplements des sables fins de la côte algérienne. Thèse de Doctorat d'Etat, Université des Sciences et de la Technologie, Algérie.
- Bakalem, A., Hassam, N., Oulmi, Y., Martinez, M., Dauvin, J.C., 2020. Diversity and biogeographical patterns of soft-bottom macrobenthos from the Bay of Bou Ismail (central Algerian coast). *Regional Studies of Marine Science*, 33, 100938.
- Ben-Eliahu, M.N., 1972. Littoral Polychaeta from Cyprus. *Tethys*, 4, 85-94.
- Ben-Eliahu, M.N., 1991. Red Sea serpulids (Polychaeta) in the eastern Mediterranean. *Ophelia Supplement*, 5, 515-528.
- Ben-Eliahu, M.N., 1995. A list of Polychaeta along the Levant coast. *Haasiana*, 1, 78-93.
- Ben-Eliahu, M.N., Fiege, D., 1995. Polychaeta from the continental shelf and slope of Israel collected by the "Meteor" 5, Expedition (1987). *Senckenbergiana Maritima*, 25, 85-105.
- Ben-Eliahu, M.N., Fiege, D., 1996. Serpulid tube worms (Annelida: Polychaeta) of the central and eastern, Mediterranean with particular attention to the Levant basin. *Senckenbergiana Maritima*, 28, 1-52.
- Benzohra, M., Millot, Cl., 1990. Analyse des caractéristiques hydrologiques et circulation des masses d'eau dans la zone côtière algérienne. Variabilité à moyenne échelle du bassin algérien. *Campagne Médiproduction*, 1986, 88-89.
- Bernard, F., 1952. Eaux Atlantiques et Méditerranéennes au large de l'Algérie. I. Hydrographie, sels nutritifs et phytoplancton en 1950. *Annales de l'Institut Océanographique*, Paris, 27, 1-8.
- Bernard, F., 1956. Eaux Atlantiques et Méditerranéennes au large de l'Algérie : II. Courants et nannoplancton de 1951 à 1953. *Annales de l'Institut Océanographique*, Paris., 31, 231-334.
- Bernard, F., 1958 ; Le courant atlantique en Méditerranée. *Commission Internationale d'Exploration Scientifique de la Mer Méditerranée*, 14, 97-100.
- Castelli, A., Abbiati, M., Badalamenti, F., Bianchi, C.N., Cantone, G. et al., 1995. Annelida Polychaeta, Pogonophora, Echiura, Sipuncula. In: Minelli A, Ruffo S, La Posta S (eds), *Checklist delle specie della fauna italiana*. Calderini, Bologna, 19, 1-45.
- Castelli, A., Bianchi, C.N., Cantone, G., Çinar, M.E., Gambi, M.C. et al., 2008. Annelida Polychaeta. *Biologia Marina Mediterranea*, 15, 323-373.
- Çinar, M.E., 2003. Ecology of Syllidae (Annelida: Polychaeta) from Northern Cyprus (Eastern Mediterranean Sea). *Bulletin of Marine Science*, 72, 795-811.
- Çinar, M.E., 2005. Polychaetes from the coast of northern Cyprus (eastern Mediterranean Sea), with two new records for the Mediterranean Sea. *Cahiers de Biologie Marine*, 46, 143-159.
- Çinar, M.E., 2007. Re-description of *Timarete punctata* (Polychaeta: Cirratulidae) and its occurrence in the Mediterranean Sea. *Scientia Marina*, 71, 755-764.
- Çinar, M.E., 2009. Alien polychaete species (Annelida: Polychaeta) on the southern coast of Turkey (Levantine Sea, eastern Mediterranean), with 13 new records for the Mediterranean Sea. *Journal of Natural History*, 43, 2283-2328.
- Çinar, M.E., 2013. Polychaetes (Annelida: Polychaeta) associated with *Posidonia oceanica* along the coasts of Turkey and northern Cyprus. In "First National Workshop on *Posidonia oceanica* (L.) Delile on the coasts of Turkey". 19-20

- September 2013, Gökçeada, Turkey”, 77-95.
- Çinar, M.E., Ergen, Z., 2003. Eusyllinae and Syllinae (Annelida: Polychaeta) from Northern Cyprus (Eastern Mediterranean Sea) with a checklist of species reported from the Levant Sea. *Bulletin of Marine Science*, 72, 769-793.
- Çinar, M.E., Ergen, Z., 2007. The presence of *Chaetozone corona* (Polychaeta: Cirratulidae) in the Mediterranean Sea: an alien or a native species? *Cahiers de Biologie Marine*, 48, 339-346.
- Çinar, M.E., Ergen, Z., Benli, H.A., 2003. Autolytinae and Exogoninae (Annelida: Polychaeta) from Northern Cyprus (Eastern Mediterranean Sea) with a checklist of species reported from the Levant Sea. *Bulletin of Marine Science*, 72, 741-767.
- Çinar, M.E., Dağlı, E., Kurt Şahin, G., 2014. Checklist of Annelida from the Coasts of Turkey. *Turkish Journal of Zoology*, 38, 1-31.
- Clarke, K.R., Gorley, R.N., 2006. PRIMER V6: User Manual/Tutorial. PRIMER-E. Plymouth.
- Coll, M., Piroddi, C., Steenbeek, J., Kaschner, K., Ben Rais Lasram, F. et al., 2010. The biodiversity of the Mediterranean Sea: estimates, patterns, and threats. *PLoS ONE*, 5, e11842.
- Coll, M., Piroddi, C., Albouy, C., Ben Rais Lasram, F., Cheung, W.W.L. et al., 2012. The Mediterranean Sea under siege: spatial overlap between marine biodiversity, cumulative threats and marine reserves. *Global Ecology and Biogeography*, 21, 465-480.
- Corriero, G., Pierri, C., Accoroni, S., Alabiso, G., Bavestrello et al., 2016. Ecosystem vulnerability to alien and invasive species: a case study on marine habitats along the Italian coast. *Aquatic Conservation: Marine and Freshwater Eco-system*, 26, 392-09.
- Daas, T., Younsi, M., Daas-Maamcha, O., Gillet, P., Scaps, P., 2010. Reproduction, population dynamics and production of *Nereis falsa* (Nereididae: Polychaeta) from the rocky coast of El-Kala National Park, Algeria. *Helgoland Marine Research*, 65, 165-173.
- Dauvin, J.C., 2005. Expertise in coastal zone environmental impact assessments. *Marine Pollution Bulletin*, 50, 107-100.
- Dauvin, J.C., Bacheler, G., Bellan, G., 2006. Biodiversity and biogeographic relationships of the polychaete fauna in French Atlantic and Mediterranean waters. *Scientia Marina*, 70, 259-267.
- Dauvin, J.C., Grimes, S., Bakalem, A., 2013. Marine Biodiversity on the Algerian Continental Shelf (Mediterranean Sea). *Journal of Natural History*, 47, 1745-1765.
- Dorgham, M.M., Hamdy, H., El-Rashidy, H.H., Atta, M.M., 2013. First records of polychaetes new to Egyptian Mediterranean waters. *Oceanologica*, 55, 235-267.
- Dorgham, M.M., Hamdy, H., El-Rashidy, H.H., Atta, M.M., Musco, L., 2014. Distribution patterns of shallow water polychaetes (Annelida) along the coast of Alexandria, Egypt (eastern Mediterranean). *Mediterranean Marine Science*, 15, 635-649.
- Ergen, Z., 1976. Investigations on the taxonomy and ecology of Polychaeta from Izmir Bay and its adjacent areas. Scientific Report of the Faculty of Science of the Ege University, Izmir, Turkey, 209, 1-73.
- Ergen, Z., Çinar, M.E., 1997. Polychaeta of Antalya Bay (Mediterranean Sea coast of Turkey). *Israel Journal of Zoology*, 43, 229-241.
- Ergen, Z., Dora, C., Çinar, M.E., 2002. Seasonal analysis of polychaetes from the Gediz River Delta (Izmir Bay, Aegean Sea). *Acta Adriatica*, 43, 29-42.
- Ergen, Z., Çinar, M.E., Dagli, E., Kurt, G., 2006. Seasonal dynamics of soft-bottom polychaetes in Izmir Bay (Aegean Sea, eastern Mediterranean). *Scientia Marina*, 70, 197-207.
- Evagelopoulos, A., Poursanidis, D., Papazisi, E., Gerovasiliou, V., Katsiaras, N. et al., 2015. Records of alien marine species of Indo-Pacific origin at Sigri Bay (Lesvos Island, north-eastern Aegean Sea). *Marine Biodiversity Records*, 8, 1-10.
- Falconetti, C., 1970. Etude faunistique d'un faciès : la “Gravellote” ou maërl de Castiglione (Algérie). *Tethys*, 1, 1057-1096.
- Fauchald, K., Rouse, G.W., 1997. Polychaete systematics: past and present. *Zoologica Scripta*, 26, 71-138.
- Faulwetter, S., Simboura, N., Katsiaras, N., Chatzigeorgiou, G., Arvanitidis, C., 2017. Polychaetes of Greece: an updated and annotated checklist. *Biodiversity Data Journal*, 5, e20997.
- Fauvel, P., 1923. Faune de France. 5. *Polychètes errantes*. Paris, Paul Lechevalier, 448 pp.
- Fauvel, P., 1927a. Faune de France. 16. *Polychètes sédentaires*. Paris, Paul Lechevalier, 494 pp.
- Fauvel, P., 1927b. Sur quelques Polychètes d'Algérie et de Tunisie. *Comptes Rendus de l'Association Française de l'Avancement des sciences, Congrès de Constantine*, 270-271.
- Furnestin, J., Allain, C., 1962a. Hydrologie de la Méditerranée occidentale au nord du 42e parallèle en automne 1958 (campagne du navire «Président Théodore Tissier» du 27 septembre au 18 octobre 1958). *Revue des Travaux de l'Institut des Pêches Maritimes*, 26, 133-161.
- Furnestin, J., Allain, C., 1962b. L'hydrologie algérienne en hiver (campagne du «Président Théodore Tissier», février 1960). *Revue des Travaux de l'Institut des Pêches Maritimes*, 26, 277-308.
- Furnestin, J., Allain, C., 1962c. Nouvelles observations sur l'hydrologie de la Méditerranée occidentale (entre Alger et le 40e parallèle). Campagne de la «Thalassa», hiver 1961. *Revue des Travaux de l'Institut des Pêches Maritimes*, 26, 309-317.
- Galil, B.S., Marchini, A., Occhipinti-Ambrogi, A., 2016. East is east and West is west? Management of marine bioinvasions in the Mediterranean Sea. *Estuarine, Coastal and Shelf Science*, 201, 7-16.
- Gil, J., 2011. The European fauna of Annelida Polychaeta. Ph.D. Thesis, Universida de Lisboa, Portugal.
- Gillet, P., 2017. Liste des Annélides Polychètes du Maroc. *Bulletin Institut Scientifique, Rabat, Maroc*, 39, 17-24.
- Gravier, C., Dantan, J.L., 1928. Pêches nocturnes à la lumière dans la baie d'Alger. *Annales de l'Institut Océanographique, Paris*, 5, 1-185.
- Gravili, C., Belmonte, G., Cecere, E., Denitto, F., Giangrande, A. et al., 2010. Nonindigenous species along the Apulian coast, Italy. *Chemistry and Ecology*, 26, Supplement, 121-142.
- Grimes S., 2010. Peuplements benthiques des substrats meu-

- bles de la côte algérienne : taxonomie, structure et statut écologique. Thèse de Doctorat d'Etat ès Sciences, Université Es Séria, Oran.
- Grimes, S., Dauvin, J.C., Bakalem, A., 2016. Annotated checklist of marine Algerian Crustacean Decapods. *Mediterranean Marine Science*, 17, 384-395.
- Hansson, H.G., 1998. NEAT (North East Atlantic Taxa): South Scandinavian marine Annelida check-list. Internet pdf [<http://www.tmbi.gu.se>].
- Laubier, L., Paris, J., 1962. Faune marine des Pyrénées-Orientales. 4. Annelides Polychètes. *Vie et Milieu*, 13, 1-82.
- Laubier, L., 1966. Sur quelques Annélides Polychètes de la région de Beyrouth. *Miscellaneous papers in the natural sciences, American University of Beirut*, 5, 9-23.
- Langeneck, J., Mazzotti, C., Mikac, B., Scirocco, T., Castelli, A., 2018. *Aricidea fragilis* (Annelida: Paraonidae) in the Mediterranean Sea: overlooked native or alien species? *The European Zoological Journal*, 85, 267-272.
- Le Danois, E., 1925. Recherches sur les fonds chalutables des côtes de Tunisie et d'Algérie en 1924. *Mémoires de l'Office Scientifique et Techniques des Pêches maritimes*, 3, 1-55.
- Le Gall, J.Y., 1969. Etude de l'endofaune des pelouses de Zostéracées superficielles de la baie de Castiglione. *Téthys*, 1, 395-420.
- López, E., Richter, A., 2017. Non-indigenous species (NIS) of polychaetes (Annelida: Polychaeta) from the Atlantic and Mediterranean coasts of the Iberian Peninsula: an annotated checklist. *Helgoland Marine Research*, 71, 1-17.
- Marcel, R. 1962. Cycle annuel de *Perinereis cultrifera* Grube (Annélide Polychète) à Alger. *Mémoires Société Nationale des Sciences Naturelles et Mathématiques de Cherbourg*, 49, 39-54.
- Maurin, C., 1962. Etude des fonds chalutables de la Méditerranée Occidentale (écologie et pêche). *Revue des Travaux de l'Institut des Pêches Maritimes*, 26, 163-218.
- Meghlaoui, Z., Daas, T., Snani, M., Daas-Maamcha, O., Scaps, P., 2015a. Annual variations of the presence of Nereididae (Annelida: Polychaeta) from intertidal rocky shores along the east coast of Algeria. *Check List the Journal of Biodiversity Data*, 11, 1-9.
- Meghlaoui, Z., Guemouda, M., Snani, M., Daas, T., Maamcha, O., Scaps, P., 2015b. Effects of oil pollution on polychaete annelids in Algerian East coast. *Journal of Entomology and Zoology Studies*, 3, 339-343.
- Mikac, B., 2015. A sea of worms: polychaete checklist of the Adriatic Sea. *Zootaxa*, 3943, 1-172.
- Millot, C., 1987. Circulation in the Western Mediterranean Sea. *Oceanologica Acta*, 10, 143-149.
- Millot, C., 1988. Surface circulation in the Algerian Basin during 1984. *Oceanologica Acta* 9, 119-131.
- Millot, C., Taupier - Letage, I., Benzohra, M., 1990. The Algerian eddies. *Earth Science Reviews*, 27, 203-219.
- Mosbahi, N., Dauvin, J.C., Neifar, L., 2017. Polychaete fauna from the intertidal zone of the Kneiss Islands. *Mediterranean Marine Science*, 18, 215-228.
- Musco, L., 2012. Ecology and diversity of Mediterranean hard-bottom Syllidae (Annelida): a community-level approach. *Marine Ecology Progress Series*, 461, 107-119.
- Mutlu, E., Çınar, M.E., Ergev, M.B., 2010. Distribution of soft-bottom polychaetes of the Levantine coast of Turkey, eastern Mediterranean Sea. *Journal of Marine Systems*, 79, 23-35.
- Nygren, A., Pleijel, F., 2011. From one to ten in a single stroke – resolving the European *Eumida sanguinea* (Phyllodocidae, Annelida) species complex. *Molecular phylogenetics and evolution*, 58, 132-141.
- Occipinti-Ambrogi, A., Marchini, A., Cantone, G., Castelli, A., Chimenz, C. et al., 2011. Alien species along the Italian coasts: an overview. *Biological Invasions*, 13, 215-37.
- Ounifi-Ben Amor, K., Rifi, M., Ghanem, R., Draief, I., Zaouali, J. et al., 2016. Update of alien fauna and new records from Tunisian marine waters. *Mediterranean Marine Science*, 17, 124-143.
- Rebzani-Zahaf C. 1990. Les peuplements macrobenthiques du port d'Alger. Evolution spatio-temporelle. Impact de la pollution. Thèse Magistère, USTHB. Alger.
- Rebzani-Zahaf C., Bakalem A., Romano, J.C. Sebane, N. Bel-lan, G., 1988. Le peuplement benthique du port d'Alger. Evolution spatio-temporelle. *Rapport de la Commission internationale de la Mer Méditerranée*, 31, 147.
- Rebzani-Zahaf, C., 1992. Le peuplement macrobenthique du port d'Alger : impact de la pollution. *Hydroécologie Appliquée*, 4, 91-103.
- Rouabah, A., Scaps, P., 2003. Life cycle and population dynamics of *Perinereis cultrifera* from the Algerian Mediterranean Coast. *Marine Ecology*, 24, 85-99.
- Rouse, G.W., Fauchald, K., 1997. Cladistics and polychaetes. *Zoologica Scripta*, 26, 139-204.
- Rouse, G.W., Gonzalo, G., 2016. Phylum Annelida: the Segmented (and Some Unsegmented) Worms. Chapt 14. p. 531-602. In: *Invertebrates*. Third Edition R.C. Brusca, W. Moore, S.M. Shuster. Sunderland, Massachusetts, Sinauer Associates, available online at <https://global.oup.com/ushe/product/invertebrates-9781605353753>.
- Russo, A.R., 1997. Epifauna living on sublittoral seaweeds around Cyprus. *Hydrobiologia*, 344, 169-179.
- Şahin G. K., Çınar M. E., 2012. A check-list of polychaete species (Annelida: Polychaeta) from the Black Sea. *Journal of the Black Sea Mediterranean Environment*, 18, 10-48.
- Selim, S.A., 1978. Systematic and distributional studies of polychaetes in the Eastern Harbour, Alexandria. MSc. thesis, Alexandria University, Egypt.
- Selim, S.A., 1996. Notes on the distribution of polychaetes along Alexandria coast, Egypt. *Bulletin of High Institute of Public Health*, 26, 341-350.
- Simboura, N., 1987. Study of the distribution and variations of the benthic polychaete fauna in three regions of the Aegean Sea and especially in the Island of Rhodes. MSc Thesis. National and Kapodistrian University of Athens, Athens.
- Simboura, N., Nicolaidou, A., 1994. Checklist and bibliography of polychaetes from Greece with some recent additions. *Mémoires du Muséum National d'Histoire Naturelle*, 162, 640-648.
- Simboura, N., Nicolaidou, A., 2001. The Polychaetes (Annelida, Polychaeta) of Greece: checklist, distribution and ecological characteristics. *Monographs on Marine Sciences*, 4, 1-115.
- Vaissière, R., Fredj, G., 1963. Contribution à l'étude de la faune benthique du plateau continental de l'Algérie. *Bulletin de l'Institut Océanographique de Monaco*, 60, 5-83.

- Viguier, C., 1886. Etudes sur les animaux inférieurs de la baie d'Alger. Recherches sur les Annélides pélagiques. *Archives de zoologie expérimentale et générale*, 14, 347-442.
- Viguier, C., 1911. Nouvelles études sur le plancton de la baie d'Alger. *Annales des Sciences Naturelles, Paris, série zoologie*, 9, 187-267.
- Viguier, C., 1912. Nouvelles études sur le plancton de la baie d'Alger. Trois Aphrotidiens pélagiques. *Annales des Sciences Naturelles, Paris, série zoologie*, 9, 84-99.
- Viguier, C., 1920. Nouvelles études sur le plancton de la baie d'Alger. Annélides pélagiques. *Annales des Sciences Naturelles, Paris, série zoologie*, 10, 256-271.
- Weigert, A., Bleidorn, C., 2016. Current status of annelid phylogeny. *Organisms Diversity and Evolution*, 16, 345-362.
- Younsi, M., Daas, T., Daas, O., Scaps, P., 2010. Polychaetes of commercial interest from the Mediterranean East Coast of Algeria. *Mediterranean Marine Science*, 11, 185-189.
- Zaâbi, S., Gillet, P., Chambers, S., 2012. Inventory of Polychaete species from the Cap Bon peninsula, western Mediterranean Sea, with new records. *Mediterranean Marine Science*, 13, 36-48.
- Zenetos, A., Cinar, M., Pancucci-Papadopoulou, M., Harmelin, J., Furnari, G. *et al.*, 2005. Annotated list of marine alien species in the Mediterranean with records of the worst invasive species. *Mediterranean Marine Science*, 6, 63-118.
- Zenetos, A., Gofas, S., Verlaque, M., Çinar, M.E., García Raso, J.E. *et al.*, 2010. Alien species in the Mediterranean Sea by 2010. A contribution to the application of European Union's Marine strategy framework directive (MSFD). Part I. Spatial distribution. *Mediterranean Marine Science*, 11, 381-493.
- Zenetos, A., Katsanevakis, S., Poursanidis, D., Crocetta, F., Damalas, D. *et al.*, 2011. Marine alien species in Greek Seas: Additions and amendments by 2010. *Mediterranean Marine Science*, 12, 95-120.
- Zenetos, A., Gofas, S., Morri, C., Rosso, A., Violanti, D. *et al.*, 2012. Alien species in the Mediterranean Sea by 2012. A contribution to the application of European Union's Marine Strategy Framework Directive (MSFD). Part 2. Introduction trends and pathways. *Mediterranean Marine Science*, 13, 328-352.
- Zenetos, A., Çinar, M.E., Crocetta, F., Golani, D., Rosso, A., 2017. Uncertainties and validation of alien species catalogues: The Mediterranean as an example. *Estuarine, Coastal and Shelf Science*, 191, 171-187.