



## **Mediterranean Marine Science**

Vol 21, No 3 (2020)

Vol 21, n3



# Pinna nobilis in the Greek seas (NE Mediterranean): on the brink of extinction?

MARIA ZOTOU, PAVLOS GKRANTOUNIS, ELPIDA KARADIMOU, KONSTANTINOS TSIRINTANIS, MARIA SINI, DIMITRIS POURSANIDIS, MARTA AZZOLIN, THANOS DAILIANIS, ELENI KYTINOU, YIANNIS ISSARIS, VASILIS GERAKARIS, MARIA SALOMIDI, POLY LARDI, ALEXIS RAMFOS, VASILIS AKRIVOS, EFTHIMIOS SPINOS, CHARALAMPOS DIMITRIADIS, DIMITRIS PAPAGEORGIOU, ATHANASIOS LATTOS, IOANNIS A. GIANTSIS, BASILE MICHAELIDIS, VASILIKI VASSILOPOULOU, ANASTASIA MILIOU, STELIOS KATSANEVAKIS

doi: 10.12681/mms.23777

## To cite this article:

ZOTOU, M., GKRANTOUNIS, P., KARADIMOU, E., TSIRINTANIS, K., SINI, M., POURSANIDIS, D., AZZOLIN, M., DAILIANIS, T., KYTINOU, E., ISSARIS, Y., GERAKARIS, V., SALOMIDI, M., LARDI, P., RAMFOS, A., AKRIVOS, V., SPINOS, E., DIMITRIADIS, C., PAPAGEORGIOU, D., LATTOS, A., GIANTSIS, I. A., MICHAELIDIS, B., VASSILOPOULOU, V., MILIOU, A., & KATSANEVAKIS, S. (2020). Pinna nobilis in the Greek seas (NE Mediterranean): on the brink of extinction?. *Mediterranean Marine Science*, *21*(3), 575–591. https://doi.org/10.12681/mms.23777

## Pinna nobilis in the Greek seas (NE Mediterranean): on the brink of extinction?

## Maria ZOTOU<sup>1</sup>, Pavlos GKRANTOUNIS<sup>1</sup>, Elpida KARADIMOU<sup>1</sup>, Konstantinos TSIRINTANIS<sup>1</sup>, Maria SINI<sup>1</sup>, Dimitris POURSANIDIS<sup>2,3</sup>, Marta AZZOLIN<sup>4,5</sup>, Thanos DAILIANIS<sup>6</sup>, Eleni KYTINOU<sup>1,7</sup>, Yiannis ISSARIS<sup>7</sup>, Vasilis GERAKARIS<sup>7</sup>, Maria SALOMIDI<sup>7</sup>, Poly LARDI<sup>7</sup>, Alexis RAMFOS<sup>8</sup>, Vassilis AKRIVOS<sup>8</sup>, Efthimios SPINOS<sup>9</sup>, Charalampos DIMITRIADIS<sup>10</sup>, Dimitris PAPAGEORGIOU<sup>1</sup>, Athanasios LATTOS<sup>11</sup>, Ioannis A. GIANTSIS<sup>12</sup>, Basile MICHAELIDIS<sup>11</sup>, Vasiliki VASSILOPOULOU<sup>13</sup>, Anastasia MILIOU<sup>14</sup> and Stelios KATSANEVAKIS<sup>1</sup>

<sup>1</sup>Department of Marine Sciences, University of the Aegean, 81100 Mytilene, Greece

<sup>2</sup> Foundation for Research and Technology - Hellas (FORTH), Institute of Applied and Computational Mathematics,

N. Plastira 100, Vassilika Vouton, 70013 Heraklion, Greece

<sup>3</sup> TerraSolutions Marine Environment Research, Heraklion, Crete, Greece, 71601

<sup>4</sup>Gaia Research Institute Onlus, Milokopi, 20300 Loutraki, Greece

<sup>5</sup> Life and System Biology Department, University of Torino, 10100 Torino, Italy

<sup>6</sup> Institute of Marine Biology, Biotechnology & Aquaculture (IMBBC),

Hellenic Centre for Marine Research (HCMR), 71500 Heraklion Crete, Greece

<sup>7</sup>Institute of Oceanography, Hellenic Centre for Marine Research (HCMR), 19013, Anavyssos, Greece

<sup>8</sup> Department of Animal Production, Fisheries & Aquaculture, University of Patras, 302 00 Messolonghi, Greece

<sup>9</sup> Directorate of Rural Economy and Veterinary Science, Regional Unit of Cephalonia, 281 00 Argostoli, Greece

<sup>10</sup> National Marine Park of Zakynthos, 1 Eleftheriou Venizelou str, Z.C, 29100 Zakynthos, Greece

<sup>11</sup>Laboratory of Animal Physiology, Department of Zoology, School of Biology, Faculty of Science,

Aristotle University of Thessaloniki, 54124, Thessaloniki, Greece

<sup>12</sup>Department of Animal Science, Faculty of Agricultural Sciences, University of Western Macedonia, 53100, Florina, Greece

<sup>13</sup>Institute of Marine Biological Resources and Internal Waters,

Hellenic Centre for Marine Research (HCMR), 16452 Argyroupoli, Greece

<sup>14</sup>Archipelagos Institute of Marine Conservation, Marine Research Base, Pythagorio, 83103, Samos, Greece

Corresponding author: stelios@katsanevakis.com

Handling Editor: Stelios SOMARAKIS

Received: 8 July 2020; Accepted: 24 July 2020; Published on line: 15 September 2020

#### Abstract

The Mediterranean endemic fan mussel *Pinna nobilis* is suffering an ongoing basin-scale mass mortality event (MME) since 2016. As most Mediterranean populations have collapsed, the species has been declared as Critically Endangered in the IUCN Red List of threatened species. In an effort to track the progress of the MME and provide updated information on the status of the species in the Greek seas, data collected through dedicated surveys and opportunistic assessments during 2019 and 2020 have been compiled. During surveys conducted at 258 sites, a total of 14,589 fan mussels were recorded, of which 81.1% were dead. Of the remaining 2,762 live individuals, 256 were juveniles. Two marine areas that still sustain living populations were identified, namely Kalloni Gulf (Lesvos Island), and Laganas Bay (Zakynthos Island). The inner part of Kalloni Gulf appears to maintain the largest surviving population of the species in the eastern Mediterranean, with an abundance estimate of 684,000 individuals (95% confidence interval: 322,000-1,453,000). Solitary, potentially resistant, scattered individuals were recorded at several sites. Other previously abundant populations that had been assessed in the past, specifically those of Lake Vouliagmeni (Korinthiakos Gulf), Souda Bay (Crete) and Gera Gulf (Lesvos Island) with a total of ~350,000 individuals, have now been wiped out. Our results document the collapse of most *P. nobilis* populations throughout the Greek seas. The MME has progressed substantially between early 2019 and mid-2020, as indicated by the increase in mortality at sites consecutively monitored multiple times. This work highlights the urgent need for continuous monitoring of surviving populations and calls for immediate implementation of an effective protection and management strategy that will ensure the persistence of surviving individuals and the production of resistant offspring.

**Keywords**: Pinnidae; pen shell; population assessment; abundance; critically endangered; *Haplosporidium pinnae*; conservation status; mass mortality events.

## Introduction

The Mediterranean endemic fan mussel *Pinna nobilis* Linnaeus, 1758 is the largest bivalve in the Mediterranean Sea and one of the largest worldwide. Its elongated triangular shell can reach a length of up to 120 cm (Schultz & Huber, 2013), and its lifespan can exceed 45 years (Rouanet *et al.*, 2015). Until recently, the species was well-distributed at a variety of habitats throughout all Mediterranean ecoregions, at depths between 0.5 m and 60 m (Zavodnik *et al.*, 1991; Butler *et al.*, 1993).

Part of the shell is buried and anchored by byssus threads in soft sediments usually covered with seagrass meadows, mainly Posidonia oceanica (Rabaoui et al., 2010; Vázquez -Luis et al., 2014), but also Cymodocea nodosa (Russo, 2012; Kersting & García-March, 2017), Zostera spp. (Russo, 2012; Prado et al., 2014; Kersting & Garcia-March, 2017) and the long-established alien Halophila stipulacea (Katsanevakis & Thessalou-Legaki, 2009). The species may also occur on unvegetated sandy bottoms (Richardson et al., 1999; Katsanevakis, 2006), estuaries (Addis et al., 2009), pebbly bottoms (Zavodnik et al., 1991; Richardson et al., 1999) or sandy patches among hard substrates (Kersting & Garcia-March, 2017; Tsatiris et al., 2018). It is a habitat-forming species, as the surface of its shell provides a hard substrate for many epibionts on a soft bottom surrounding (Giacobbe, 2002; Addis et al., 2009; Rabaoui et al., 2009). It can thus function as an ecosystem engineer, and very dense populations can create biogenic reefs (Katsanevakis, 2016). Being a filter feeding mollusc, P. nobilis plays a key ecological role, retaining large amounts of organic matter and reducing turbidity (Trigos et al., 2014), while enhancing benthic-pelagic coupling (Basso et al., 2015).

Over the years, human stressors have caused the decline of fan mussel populations across the Mediterranean basin (De Gaulejac, 1995; Richardson et al., 1999; Garcia-March, 2005; Katsanevakis, 2007; Deudero et al., 2015; Basso et al., 2015). Exploitation of the species can be traced back to antiquity; in addition to its exploitation for human consumption, its byssus threads were used to produce a precious textile, "sea silk" (Maeder, 2008). Furthermore, the degradation of P. oceanica meadows a vital fan mussel habitat - as a result of several human activities (e.g. trawling, illegal fishing methods, uncontrolled anchoring) is increasingly threatening the survival of the species (Katsanevakis et al., 2008; Vázquez-Luis et al., 2015; Basso et al., 2015). Additional stressors affecting fan mussel populations include climate change, marine pollution, and invasive species (Basso et al., 2015; Kersting et al., 2019).

Consequently, *P. nobilis* has been listed as an endangered species and is protected under the EU Habitats Directive (92/43/EEC, Annex IV), the Protocol for Specially Protected Areas and Biological Diversity in the Mediterranean of the Barcelona Convention (Annex II), and the national legislation of most Mediterranean countries. However, despite its protection status, commercial exploitation of the species continues illegally in some countries, with collected individuals being destined either for consumption as a delicacy (Katsanevakis *et al.*, 2011), or to fulfil decorative purposes (Kersting *et al.*, 2019).

Until the autumn of 2016, no parasitic or bacterial infection had been reported as constituting a threat for the survival of fan mussel populations (Darriba, 2017). A mass mortality event (MME) of P. nobilis along the SE coasts of the Iberian Peninsula, with mortality rates exceeding 90% within a few weeks, alarmed the scientific community. By June 2017, the MME had expanded northwards (Balearic Islands), affecting all age classes of fan mussel populations at all depths and habitat types, with mortalities reaching up to 100% at some locations (Vázquez-Luis et al., 2017). Histological examination of affected individuals showed heavy inflammatory host response and severe general dysfunction (Darriba, 2017). Thereafter, severe mortalities of P. nobilis populations were reported from additional locations along the Spanish coastline, as well as France and Italy (Catanese et al., 2018: Carella et al., 2019: Panarese et al., 2019). Soon after the first reports, new MME records were reported in Greece, Cyprus, Turkey, Algeria, Tunisia, Morocco and Croatia, with mortality rates often reaching 100% (Katsanevakis et al., 2019; Kersting et al., 2019; IUCN, 2019; Čižmek et al., 2020). Due to the severe decline of its global population and its high risk of extinction, in 2019, P. nobilis was classified as Critically Endangered in the IUCN Red List of Threatened Species (Kersting et al., 2019).

The search for the cause of this large-scale MME that has affected most Mediterranean populations has puzzled the scientific community. Two pathogens have been identified and associated with the disease affecting P. nobilis populations; a newly described, highly species-specific pathogenic protozoon of unknown origin, named Haplosporidium pinnae (Catanese et al., 2018), and a Mycobacterium sp. (Carella et al., 2019). The coexistence of both pathogens in some affected populations has been reported (Garcia-March et al., 2020, Lattos et al., 2020; Carella et al., 2020; Čižmek et al., 2020). Furthermore, in some cases, increased sea temperature seems to favour the proliferation of another opportunistic pathogen in P. nobilis, namely Vibrio mediterranei (Rodriguez et al., 2018), which was recently found to intensify pathogenicity and cause increased mortality of stabled fan mussels at temperatures of about 25-26°C (Prado et al., 2020).

Despite the dramatic population declines in impacted areas, there are still some surviving populations across the Mediterranean [e.g. Fangar Bay (Spain), Mar Menor (Spain), Rhone Delta (France), Etang de Thau (France), Diana and Urbino (Corsica), the inner part of Kalloni Gulf, Lesvos (Greece) and the Venice lagoon (Italy)], possibly due to local environmental conditions that delay the expansion of the disease or are unfavourable for the parasites (Catanese *et al.*, 2018; Katsanevakis *et al.*, 2019; Kersting *et al.*, 2019; Cabanellas-Reboredo *et al.*, 2019).

In Greece, no unusual mortality of the species had been recorded before the summer of 2018, when anecdotal information about an ongoing MME in the Aegean Sea triggered the first mortality assessments. In early autumn

2018, a survey conducted along the coasts of Lesvos Island provided a mean mortality estimate of 93%, thus confirming the spread of the disease in the Aegean Sea. The P. nobilis population in Gera Gulf had an estimated mortality between 91% and 100%, while in Kalloni Gulf mortalities varied from 36% (northern, inner part of the gulf) to 99% (southern, outer part of the gulf). Respective molecular analyses confirmed the presence of H. pinnae in infected individuals from Gera Gulf (Katsanevakis et al., 2019). Later on, Lattos et al. (2020) confirmed the presence of both H. pinnae and Mycobacterium sp. in specimens collected from Thermaikos Gulf and the north Aegean (Kalloni Gulf and Limnos island) during 2018-2019. Currently, it is clear that the Mediterranean MME has affected previously thriving P. nobilis populations in Greek waters. However, the lack of past information on the population status of the species in the Aegean and Ionian seas, with the exception of a few localities (Katsanevakis, 2006; Katsanevakis & Thessalou-Legaki, 2009; Tsatiris et al., 2018), constitutes a challenge for the assessment of its status at national level (Katsanevakis et al., 2019).

Approximately one year after the onset of the MME in Greece, several research teams across the country initiated visual surveys to assess fan mussel mortalities and investigate the expansion of the MME. By compiling information collected through such dedicated surveys, conducted in 2019 and 2020, as well as other opportunistic records, this work aims to provide the most comprehensive and updated information on the health status of *P. nobilis* populations in the Greek seas.

## **Materials and Methods**

#### General approach for data collection

This study presents the state of P. nobilis populations as a mosaic of individual case studies, conducted in different marine areas, in order to assess the mortality rates of impacted populations in the Greek seas, and estimate the current extent of the MME. A number of independent surveys were conducted between February 2019 and June 2020. In exceptional cases, certain populations have been monitored occasionally over the past years (Katsanevakis, 2009; Katsanevakis & Thessalou-Legaki, 2009; Tsatiris et al., 2018), thus allowing a 'Before & After MME' comparison of their population status. However, since long term monitoring data of marine benthic communities is particularly scarce in Greece, most of the information provided herein comes from one-off assessments, carried out after the onset of the MME. Aiming to cover a large geographical scale, the dataset consists of eight dedicated case studies focusing on P. nobilis MMEs and opportunistic assessments from 70 additional sites carried out within the framework of various projects (Table 1).

In order to assess the mortality of *P. nobilis*, both dead and live individuals were counted during visual surveys, lasting 20-120 minutes. Only recently dead individuals were counted, excluding individuals presumed to be dead for longer periods (i.e. according to the type and degree of fouling), or having died due to mechanical damage (e.g. due to anchoring) (Katsanevakis *et al.*, 2019).

**Table 1.** Sampling methodology followed by the research groups in each marine area. "45 min" refers to 45 minutes respectively of visual observations applying the protocol described in Case Study 1, "30 min" refers to the same protocol but less survey time, "Opportunistic" refers to opportunistic records without applying a strict protocol, "Transect" refers to observations along strip transects and "Distance" when Distance sampling methodology was applied.

Geographic Area	Sampling methodology	
	2019	2020
Saronikos Gulf	45 min	-
Petalion Gulf	45 min; Transect	-
South Evoikos Gulf	45 min; Opportunistic	Opportunistic
North Evoikos Gulf	45 min	Opportunistic
Maliakos Gulf	-	Distance
Kalloni Gulf (Lesvos Island)	45 min; Distance	45 min
Gera Gulf (Lesvos Island)	45 min	45 min
inner Thermaikos Gulf	Opportunistic	Distance
outer Thermaikos Gulf	45 min	Opportunistic
Toronean Gulf	-	Opportunistic
Korinthiakos Gulf	30 min; Opportunistic	-
Elounda Bay (Crete)	45 min; Opportunistic	-
N.M.P.A.N.S.	45 min; Opportunistic	-
N.M.P.Z.	Transects	-
Other Areas	Opportunistic; 45 min	Opportunistic

## Case study 1: Saronikos Gulf, South and North Evoikos Gulfs, Petalion Gulf, Maliakos Gulf

In September 2019, four marine areas in Greece (Saronikos Gulf, Petalion Gulf, South and North Evoikos Gulfs) were investigated by conducting a dedicated survey targeting the mortality of P. nobilis. These gulfs are adjacent and interconnected. The narrow Evripus Strait links the South and North Evoikos Gulfs. Several of the surveyed locations are included in the Natura 2000 European network of protected areas (Site Codes: GR3000005, GR3000004, GR2420016, GR2440002). Published data from previous surveys (e.g. Sini et al., 2017), anecdotal information (including communication with local dive centres, divers, fishermen, and research scientists), combined with Google Earth images, were used to select the 32 sampling sites, targeting shallow waters covered with seagrass (Fig. 1 CS 1). The surveys were conducted over hard, soft or mixed substrates, with the dominant vegetation being P. oceanica, C. nodosa and Cystoseira spp., at a depth range of 0.5 and 20 m (Supplementary file). Each visual survey lasted 45 minutes and was conducted using either SCUBA equipment or free diving, depending on the depth and the visibility of each location. Live individuals were characterized as "presumed healthy" (clearly unaffected populations) or "presumed moribund", based on the delayed valve-closing reflex, as described in Vázquez-Luis et al. (2017). Juvenile individuals were recorded separately.

Additional records from two independent research groups that investigated parts of the aforementioned ma-

rine areas were also included, since the data recorded in March 2019 and between March and June 2020 allow comparison of the species status in three different periods. In March 2019, 9 sites were investigated at a depth range between 0 and 14 m, while in early 2020, 7 sites were visited at a depth range between 0 and 27 m.

#### Case study 2: Kalloni Gulf – Lesvos Island

Kalloni Gulf is the larger of the two shallow, semi-enclosed embayments of Lesvos Island (NE Aegean Sea), with an approximate surface area of 130 km<sup>2</sup> (Kefalas *et al.*, 2016). Kalloni Gulf is a productive marine ecosystem that supports a variety of ecologically valuable habitats and communities (Evagelopoulos & Koutsoubas, 2008; Kefalas *et al.*, 2016; Sini *et al.*, 2019), and is included in the Natura 2000 European network of protected areas (Site Code: SCI GR4110004). Before the spread of the MMEs in the eastern Mediterranean, Kalloni Gulf hosted a large *P. nobilis* population that, however, had never been quantitatively assessed (Katsanevakis *et al.*, 2019).

In a dedicated study conducted in June 2019, a total of 20 sites were surveyed in Kalloni Gulf, following the same protocol as the one described in Case Study 1. The sampling sites were systematically positioned every  $\sim$ 3 km along the coastline (Fig. 1 CS 2&3), at a depth range of 0-9 m, and were investigated by free diving. Furthermore, additional surveys, following the same protocol as the one described in Case Study 1, provided data for different periods (4 sites visited in August-September 2019)



*Fig. 1:* Map of Greece depicting the sampling sites selected for the assessment of the status of *Pinna nobilis* populations. Inset maps indicate the sampling sites used during the dedicated case studies (CS 1-8), whereas the main map indicates other scattered sampling sites.

and 3 sites in May-June 2020), at a depth range between 0.5 and 3 m.

The June 2019 assessment indicated a low mortality population in the northern, inner part of Kalloni. To estimate the abundance of the presumably unaffected (or less affected) population in the inner part of Kalloni Gulf, 10 sampling sites were chosen in a systematically random way along the coastline (i.e. one site was randomly chosen, and the others were sequentially positioned at a constant distance of 1 km) (Fig. 2). At each site, a 200-m transect, perpendicular to the coastline, was set and surveyed following the line transect distance sampling method (Buckland et al., 2001) using SCUBA diving. Distance sampling has been widely used for the estimation of P. nobilis abundance in the Aegean Sea (Katsanevakis, 2006; Katsanevakis & Thessalou-Legaki, 2009; Tsatiris et al., 2018). This method properly accounts for imperfect detection (Buckland et al. 2001), given that fan mussel detectability can be very low, especially for small individuals (Katsanevakis & Thessalou-Legaki, 2009), and if ignored can lead to substantially biased estimates. For each detected fan mussel, one observer recorded its size (shell width in cm) and perpendicular distance, y (cm), from the line transect. A hazard-rate function,

 $g(y) = 1 - exp\left[-\left(\frac{y}{\sigma}\right)^{-\nu}\right]$ , was used to model detectability (Buckland *et al.*, 2001). The total area sampled (Fig. 2) was 1.7 km<sup>2</sup>, estimated using ArcMap 10.1.

#### Case study 3: Gera Gulf – Lesvos Island

Gera Gulf is a semi-enclosed mesotrophic embayment in the south-eastern part of Lesvos Island (Fig. 1 CS 2&3) with a total surface of 40.099 km<sup>2</sup> (Tsatiris et al., 2018). Close to the coastline there is a narrow zone (with a total area of 0.10 km<sup>2</sup>) of mixed bottoms, while further offshore there are extensive areas of sandy (13.7 km<sup>2</sup>) and muddy (24.8 km<sup>2</sup>) sediments, the latter covering the central part of the gulf. Scattered P. oceanica meadows (1.21 km<sup>2</sup>) can be found along the coastline of the Gulf, with the largest detected meadow located in the south-western part (Tsatiris et al., 2018). Gera Gulf is included in the Natura 2000 European network of protected areas (Site Code: GR4110005). According to the assessment carried out by Tsatiris et al. (2018) in 2016 (i.e. two years before the beginning of the MME in the area), the estimated total fan mussel population was ~213,000 individuals (95% CI: 97,600-466,000), with a population density peak at a depth range of 1.5-8 m (practically zero at depths >15 m), and the highest densities in *P. oceanica* meadows, followed by mixed bottoms. Dedicated follow-up monitoring surveys at three sites of high population density during August-October 2018, provided mortality estimates of between 91% and 100%, while the total collapse of the Gera fan mussel population was confirmed in December 2018 when no live individual was recorded (Katsanevakis et al., 2019).

During July-August 2019, 7 sites in Gera Gulf (Fig. 1 CS 2&3) were investigated through 45-min standardized free-diving surveys, by applying the same protocol



*Fig. 2:* Map of the inner part of Kalloni Gulf (Lesvos Island) showing the ten transects positioned perpendicular to the coastline used to estimate the abundance of live fan mussels.

as the one described in Case Study 1. The depths in the surveyed sites varied between 0.5 and 5 m. An additional site in Gera Gulf was also visited in June 2020, at a depth range of between 0.5 and 4 m, using the same protocol.

## Case study 4: Thermaikos Gulf

Thermaikos Gulf is located at the north-western Aegean Sea, and its inner part is considered a heavily polluted area due to urban, industrial, and agricultural wastes (Arditsoglou & Voutsa, 2012). It receives the outflow of five rivers (Aliakmon, Axios, Loudias, Gallikos, Pineios), with part of the extensive wetland and estuarine system of the former four designated as a Natura 2000 site (Site Code: GR1220010). It is characterized by high nutrient content, particularly favouring a high biodiversity of filter feeders, such as bivalves. Among the various bivalve species of the Gulf, a large *P. nobilis* population had been reported in the past (Katsanevakis *et al.*, 2008), with densities locally exceeding one individual per m<sup>2</sup> (Galinou-Mitsoudi *et al.*, 2006).

Fan mussel population mortality in Thermaikos Gulf was assessed by two independent research groups. The first group focused on the inner part of the Gulf, while the second one on the outer part. In the inner Thermaikos, a total of 8 sites were investigated (Fig. 1 CS 4). Initially, 2 sites were sampled during May 2019. However, following the reports of local fishermen relating to unusual mortalities of the species, these 2 sites were revisited and assessed during July 2019, along with 6 additional sites. The depth range surveyed was between 3 and 9 meters, and both live and dead individuals were recorded. The dedicated survey in the outer Thermaikos Gulf, conducted between July and October 2019, included 13 sampling sites that were investigated following the methodology described in Case Study 1 (Fig. 1 CS 4). The depth range investigated was between 0 and 18 m, on soft and mixed substrates, covered with Cymodocea nodosa, Zostera noltei, Cystoseira spp. or P. oceanica. Furthermore, opportunistic records obtained from different research groups that investigated the area during different periods were also included. A total of 8 additional sites were visited in March and June 2020, at a depth range between 0 and 10 m.

## Case study 5: Korinthiakos Gulf

Korinthiakos Gulf is a narrow embayment providing a link between the Aegean and Ionian seas through the Corinth Canal and the Rion-Antirion Strait, respectively (Perissoratis *et al.*, 2000). It is 115 km long and its width ranges from 5 to 30 km, with a maximum depth of more than 900 m in the centre of the basin (Lykousis *et al.*, 2007; Papageorgiou *et al.*, 2017). This semi-enclosed basin is of significant geologic interest due to its peculiar morphological characteristics that resemble a small-scale ocean basin (with shelf, slope and deep basin) (Perissoratis *et al.*, 2000) and is included in the Natura 2000 European network of protected areas (Site Code: GR2530007). On the northern side, the substrate is mainly rocky, in contrast to the sandy and muddy southern coasts (Papageorgiou *et al.*, 2017), with mud deposits being dominant at the deeper parts of the basin (Poulos *et al.*, 1996). Korinthiakos Gulf supports a rich biodiversity of marine flora and fauna (Issaris *et al.*, 2012), with significant populations of marine mammals (Papageorgiou *et al.*, 2017; Frantzis & Herzing, 2002).

In Korinthiakos Gulf, a total of 12 sites were investigated, between May and August 2019, at a depth range of 0 to 30 m (Fig. 1 CS 5). Visual surveys lasting 30 minutes were carried out either with SCUBA or free diving, on hard or mixed substrates. Both live and dead individuals were counted. Juvenile individuals were considered separately. Additionally, opportunistic records from two other research groups were obtained from 5 sites that were surveyed in October 2019, at a depth range of 1 to 35 m (Fig. 1 CS 5).

## Case study 6: Elounda – Crete

Elounda Bay is a semi-enclosed embayment located on the NE coast of the island of Crete (Fig. 1 CS 6). It covers a surface area of 6.5 km<sup>2</sup>, and at its northern part is connected to the open sea via an elongated canal, 1400 m long and 700 m wide. At its southern side, the bay is also connected to the open sea through an artificially created narrow strait, 10 m wide and 140 m long. The western part of Elounda Bay is generally characterized by a sandy coast, while the eastern part is characterized by a rocky coastline. The bay's depth reaches a maximum of 8.8 m and the substrate consists of fine sand and silt, with rare small rocky reefs. The vegetation consists mainly of the seagrass C. nodosa at shallow depths and the green macroalga Caulerpa prolifera at depths greater than 5 m. P. nobilis has been one of the dominant sessile invertebrates found in Elounda Bay, while the area is also considered an important habitat for the growth of juvenile fish species, contributing to the conservation and maintenance of the local marine biological resources (Koulouri et al., 2016).

The 10 sampling sites for the assessment of *P. nobilis* population covered a geographically representative area in the bay as well as deeper locations outside the bay (Fig. 1 CS 6). The surveys were conducted in June, August and October 2019. Sampling was performed with SCUBA diving, at a depth range of 1 to 7 m within the bay, and 4 to 20 m outside the bay, according to the protocol described in Case Study 1.

## Case study 7: National Marine Park of Alonissos -Northern Sporades - N.M.P.A.N.S.

Established in 1992, the National Marine Park of Alonissos and Northern Sporades (NMPANS), is Greece's first marine park and one of Mediterranean's largest marine protected areas covering an area of approximately 2,200 km<sup>2</sup> (Natura 2000 Site Code: GR1430004). The shallow coastal seascape is predominantly covered by rocky reefs and seagrass meadows. The NMPANS is important for marine mammals, such as cetaceans and the Mediterranean Monk Seal.

Between June and October 2019, sampling was carried out at 11 sites (Fig. 1 CS 7). At each site, free roaming SCUBA diving surveys were performed for 45 minutes at depths ranging from 2 to 45 m. Furthermore, an additional site was surveyed as an opportunistic record by another research group, in April 2019, at a depth range of 26 to 30 m.

#### Case study 8: Laganas Bay – NMPZ – Zakynthos Island

This case study was conducted in the National Marine Park of Zakynthos (NMPZ), which is located in the southernmost part of Zakynthos Island, eastern Ionian Sea (Fig. 1 CS 8), and hosts one of the most important rookeries of the threatened marine turtle *Caretta caretta* in the Mediterranean (Margaritoulis, 2005). Acknowledging the prime ecological importance of this area, the NMPZ was formally established in 1999 (Presidential Decree 906D/1999) and covers an area of 83.3 km<sup>2</sup> in Laganas Bay.

In August 2019, 35 sampling sites were selected in a systematic way along the rocky part of the NMPZ coastline, to estimate the population of *P. nobilis* along the shallow habitats of the NMPZ (with consecutive sites having a 200 m distance between them) (Fig. 1 CS 8). At each sampling site, a 200-m transect was set parallel to the coastline using a diving reel. Fan mussel individuals were recorded (number of live and dead individuals, number of juveniles) along each transect by snorkelling, during 20-min long surveys at depths between 0 and 10 m.

#### Data from other areas

Apart from case studies focusing on *P. nobilis* MME, additional opportunistic records of fan mussel populations were also included herein, in order to cover as much of the Greek coastline as possible, and to depict the status of the species in different marine areas (Fig. 1).

Other data originates from different sites located along the coasts of mainland Greece, 28 Aegean Sea islands, 6 Ionian Sea islands, and the island of Crete (Supplementary file). Among these opportunistic records of *P. nobilis* populations, Souda Bay in Crete and Lake Vouliagmeni have been assessed in the past, since both locations used to host large *P. nobilis* populations (Katsanevakis & Thessalou-Legaki, 2009; Katsanevakis, 2009).

## Results

#### General description of the dataset

Overall, 258 visual surveys were carried out between February 2019 and June 2020, by 9 research groups. In total, 14,589 (dead or live) *P. nobilis* individuals were recorded at a depth range of 0-55 m. Among the recorded individuals, 18.9% were alive and 81.1% dead. Of the 2,762 live individuals, 9.3% were juveniles, most of which were found in Kalloni Gulf, indicating successful recruitment in specific areas despite the ongoing MME.

## Case study 1: Saronikos Gulf, Evoikos Gulfs, Petalion Gulf, Maliakos Gulf

From the data collected in March 2019, a total of 1,949 individuals were recorded, 101 of which were found alive, resulting in a mortality rate estimate of 94.8%. However, these records refer to two marine areas; Petalion Gulf where no live individuals were found, thus the recorded mortality rate was 100%, and South Evoi-kos Gulf where out of 133 individuals recorded, only 32 were found dead, resulting in a mortality rate estimate of 24.1% (Fig. 3a).

During the dedicated Case Study carried out in September 2019, 895 *P. nobilis* individuals were recorded, 670 of which were found dead, resulting in a total mortality rate estimate of 74.9% in the study area. By September 2019, *P. nobilis* populations had totally collapsed in Saronikos Gulf, Petalion Gulf, and S. Evoikos Gulf, where no *P. nobilis* individuals were found alive (Fig. 3b). Live individuals were only found along the coasts of N. Evoikos Gulf, where the population appeared to be in relatively good state, with an estimated mortality rate of 31.4%. Among the 225 live individuals, 2.7% were noted as moribund and 6.7% were juveniles. Dead juveniles were also present in most marine areas visited.

In many locations around Saronikos Gulf, where no individuals of P. nobilis were recorded, old and broken shells of individuals presumed to be dead for a long time, indicated the past presence of the species in the area. Similarly, at the sampling sites of Petalion Gulf, numerous 'old' shells were noted but not included in the assessment, as they did not match the assessment criteria of the current survey. Overall, 73.3% of all dead individuals originated from the S. Evoikos Gulf where not a single individual was found alive at the two locations visited that hosted dense populations of large fan mussels. The N. Evoikos Gulf was the only Case Study 1 area with recorded live fan mussels. However, the mortality rates at different locations in the Gulf varied. In September 2019, most of the sites located along the western coastline of the N. Evoikos Gulf sustained surviving individuals, and populations displayed a relatively low mortality rate (22.5%).

However, the situation deteriorated in 2020. Two hundred and eighty four (284) dead individuals were reported from the Evoikos and Maliakos gulfs and not a single live



*Fig. 3:* Mortality assessment of *Pinna nobilis* populations at the sites of Case Study 1 (Saronikos Gulf, Evoikos Gulfs, Petalion Gulf, Maliakos Gulf), in March 2019 (a), September 2019 (b), and March-June 2020 (c). The size of the pie charts represents the number of recorded individuals. The red and green colour in the pie charts show the proportion of dead and live individuals, respectively.

fan mussel (mortality rate: 100%). This highlighted the collapse of the last surviving populations in the area (Fig. 3c).

#### Case study 2: Kalloni Gulf

During the dedicated visual surveys conducted in Kalloni Gulf (June 2019), a total of 2,675 *P. nobilis* individuals were recorded, of which 1,021 live and 1,654 freshly dead, resulting in an average mortality of 61.8%. Based on the spatial pattern of mortalities, the area of Kalloni Gulf can be divided into three distinct subareas (Fig. 4a), namely, i) the southern seaward part of the Gulf where no fan mussel was recorded, ii) the central part of the Gulf, where the assessment revealed a higher than 70% mortality, and an increased number of old dead shells that were severely damaged or stacked in the upper swash zone, and iii) the north-eastern inner part of the Gulf (Fig. 4a), where mortality rates were relatively low ranging between 14.1% and 23.4%, apart from one site where dead individuals reached 50.8%. Throughout the Gulf, 94.4% of the total recorded individuals were adults, of which 1592 (63%) were dead. Juveniles (149 individuals or 5.6% of the total records), were present at only half of the assessment sites, and displayed a mortality rate of 41.6%.

Distance sampling was applied in June 2019 in the north-eastern part of Kalloni Gulf, where a total of 815 live individuals were recorded along 10 line transects (Fig. 2). The shell width of recorded individuals ranged between 4 and 18 cm, with a mean value ( $\pm$  SD) of 11.2  $\pm$  2.3 cm. Mean visibility across all transects was estimated at 2.1  $\pm$  1.5 m. The probability of *P. nobilis* detection was estimated at 0.19  $\pm$  0.08 for 5-m wide transects. The total abundance of the species in the surveyed zone was estimated at 684,000 individuals (95% CI: 322,000-1,453,000).

Subsequently, four sites in the inner part of the Gulf



*Fig. 4:* Mortality assessment of *Pinna nobilis* populations at the sites of Case Study 2 (Kalloni Gulf), in June 2019 (a), August-September 2019 (b), and May-June 2020 (c). The size of the pie charts represents the number of recorded individuals. The red and green colour in the pie charts show the proportion of dead and live individuals, respectively.

were revisited in August-September 2019. A total of 526 individuals were recorded, of which only 33 were alive; the estimated mortality rate was 93.7% (Fig. 4b). Finally, three sites were surveyed during late May-June 2020. A total of 1,233 individuals were recorded, of which only 133 were found alive (estimated mortality rate: 89.2%) (Fig. 4c); 30.8% of the surviving individuals were juveniles.

## Case study 3: Gera Gulf

In 2019, 469 dead individuals and only one live were recorded in Gera Gulf (Fig. 5 CS 3). The live individual was a juvenile found at 0.8 m on mixed bottom. Hence, the estimated average mortality rate in Gera Gulf was 99.8%. In June 2020, no live individual was recorded during a revisit to one of the sites.

## Case study 4: Thermaikos Gulf

During the first investigation in the inner Thermaikos Gulf (May 2019), two sites were visited and a total of 150 live individuals were counted (0% mortality), of which 23.3% were juveniles. Two months later (July 2019), the same two sites were revisited, along with six additional sites. A total of 1,092 individuals were recorded, of which 227 were dead (mortality rate estimate: 20.1%) (Fig. 6a). Among the 865 live individuals, 42 were juveniles (4.9%). However, opportunistic records obtained from visual surveys conducted in June 2020 in the inner Thermaikos Gulf indicated the collapse of the aforementioned surviving populations, since 254 individuals were counted, and all of them were found dead (Fig. 6c).

During the dedicated survey in the outer Thermaikos Gulf (late July-October 2019), a total of 2,221 *P. nobilis* individuals were recorded, of which only 2 were alive (mortality rate estimate: 99.9%) (Fig. 6b). No *P. nobilis* individuals were detected at several locations around the Thermaikos Gulf. However, in some cases broken shells washed ashore indicated the past presence of the species in the area.

In 2020, five more sites were investigated in the outer Thermaikos Gulf and the Toronean Gulf (Chalkidiki peninsula). The survey revealed 543 dead and no live ones thus indicating 100% mortality in the area (Fig. 6c).



*Fig. 5:* Mortality assessment of *Pinna nobilis* populations at the sites of Case Studies 3 – Gera Gulf (July-August 2019, June 2020), 7 – NMPANS (June-October 2019), and 8 – Laganas Bay (August 2019), as well as other sites not included in a specific case study. The size of the pie charts represents the number of recorded individuals. The red and green colour in the pie charts show the proportion of dead and live individuals, respectively.



*Fig. 6:* Mortality assessment of *Pinna nobilis* populations at the sites of Case Study 4 (Thermaikos Gulf), in May-July 2019 (a), July-October 2019 (b), and June 2020 (c). The size of the pie charts represents the number of recorded individuals. The red and green colour in the pie charts show the proportion of dead and live individuals, respectively.

## Case study 5: Korinthiakos Gulf

During May-August 2019, 155 dead and only 21 live individuals were recorded in the Korinthiakos Gulf, resulting in an overall mortality rate estimate of 88% (Fig. 7a). Of the remaining live individuals, 5 were juveniles. Resampling of certain sites in October 2019 indicated 100% mortality at the investigated sites, as all 50 recorded individuals were dead (Fig. 7b).

## 385 (77.3%) were dead (Fig. 8a). At two sites, all individuals were dead (Fig. 8a), whereas at the other two sites, both live and dead individuals were recorded. Among the counted individuals, 13 were juveniles, of which 4 were dead. Moribund individuals (n = 20) were encountered among the live ones, indicating that the MME was in progress at the time of the survey.

In August and October 2019, six additional sites were surveyed both within Elounda Bay and in the marine area just outside the bay. During these revisits, all 125 recorded individuals were dead, indicating that between June and August/October all fan mussels were infected and died (Fig. 8b).

#### Case study 6: Elounda Bay

Overall, 498 *P. nobilis* individuals were counted at the four sites surveyed in Elounda Bay in June 2019, of which



*Fig.* 7: Mortality assessment of *Pinna nobilis* populations at the sites of Case Study 5 (Korinthiakos Gulf), in May-August 2019 (a) and October 2019 (b). The size of the pie charts represents the number of recorded individuals. The red and green colour in the pie charts show the proportion of dead and live individuals, respectively.



*Fig. 8:* Mortality assessment of *Pinna nobilis* populations at the sites of Case Study 6 (Elounda), in June 2019 (a) and October 2019 (b). The size of the pie charts represents the number of recorded individuals. The red and green colour in the pie charts show the proportion of dead and live individuals, respectively.

#### Case study 7: N.M.P.A.N.S.

Overall, 87 *P. nobilis* individuals were counted at the 11 sites surveyed in N.M.P.A.N.S. between June and October 2019, and all were found dead (100% mortality estimate) (Fig. 5 CS 7). More than half of the counted individuals were observed at Tsoukalia Bay (G8 in the Supplementary file), where 40 freshly dead individuals were found uprooted, lying on the seabed. No live individuals were recorded during a previous opportunistic survey conducted in April 2019 thus indicating that the collapse of the fan mussel population in the area, might have occurred before summer 2019.

#### Case study 8: Laganas Bay

During August 2019, a total of 55 *P. nobilis* individuals were recorded along transects of 7 km total length, of which 33 were alive and 22 dead (mortality rate estimate: 40%) (Fig. 5 CS 8). The maximum density was estimated at 11 individuals per 200 m transect length. The presence of dead individuals was restricted mainly to 6 sites. Healthy populations of *P. nobilis* with no dead individuals were recorded at 11 sites, whereas the presence of juveniles (27.3% of the live individuals) was evident at 4 sites.

#### Data from other sites

Overall, 1,134 *P. nobilis* individuals were recorded at 70 sites in several marine areas of the Greek seas (see Supplementary File), of which only 64 (5.6 %) were found alive. At most sites, no live specimens were found (Fig. 5 'Other Sites'). The Gulf of Argostoli (Kefalonia

Island, Ionian Sea) was surveyed by two research groups, during two different periods (July and November 2019). During the first visit (July 2019), all 56 recorded individuals were alive. Five months later (November 2019) only one live juvenile was found among 310 dead individuals, indicating that the specific population was severely affected during the summer of 2019. Similarly, only one surviving fan mussel was reported from Limnos Island (NE Aegean Sea), among a dense population of 299 dead individuals. In Kalamaki Viotias (Korinthiakos Gulf), two live individuals were recorded in April 2019; however, in July 2019 all specimens recorded in the area were dead. In the marine Lake Vouliagmeni (Korinthiakos Gulf), where P. nobilis populations had been thriving before the onset of the MME (Katsanevakis, 2006), the estimated mortality rate reached 89.7%, while the few live individuals recorded were considered moribund. In Souda Bay (NW Crete), only dead individuals were encountered in July 2019, indicating that a previously recorded dense population (Katsanevakis & Thessalou-Legaki, 2009) had been wiped off.

#### Integrated mapping of P. nobilis status in Greece

All data collected within the framework of this study were combined to depict the status of *P. nobilis* in the Greek seas by June 2020 (Fig. 9). Healthy or less affected populations are restricted to the inner Kalloni Gulf and Laganas Bay. Pooling together all records, most live individuals were recorded before June 2019, but mortality rates gradually increased, reaching 95% in 2020, at the surveyed sites (Fig. 10). Despite the inherent representativeness bias of our dataset (as different sites were surveyed during each time interval), this result indicates that the *P. nobilis* MME has been progressing rapidly during



*Fig. 9:* Integrated map of *P. nobilis* status in Greece. Red coastlines represent the sites where fan mussel populations have collapsed. Inner Kalloni Gulf, and Laganas Bay (right panels) are the only known areas with surviving populations. Coastlines with no colour represent fan mussel populations that have not been assessed.



*Fig. 10:* Temporal progress of the MME in the Aegean Sea in 2019, compiling data obtained from all case studies and opportunistic records. The stacked bars were created by including all dead and live individuals recorded during each period, while the line refers to the average mortality rate among surveyed sites during each period (treating all sites equally in the estimation of the mean, ignoring differences in abundance).

2019 and 2020 and devastating many fan mussel populations in several coastal areas of the Greek seas (in some cases only within a couple of months).

## Discussion

This study documents the large-scale progression of the *P. nobilis* MME throughout the Greek seas during 2019 and 2020 and the collapse of fan mussel populations in most of the marine areas investigated. Infection by pathogens seems to have affected populations in both protected and unprotected areas, in the Aegean and Ionian seas, at all longitudes and latitudes. The spread of the disease in Greek waters, similarly to the ongoing spatial expansion of mortality events in other Mediterranean ecoregions (Čižmek *et al.*, 2020), indicates that the fan mussel MME that started in 2016 is still in progress, and no local population in the Mediterranean may be considered safe.

Few past assessments of *P. nobilis* populations in Greece allowed the estimation of local loss due to the on-

going MME. In Vouliagmeni Lake (Korinthiakos Gulf) the fan mussel population was monitored between 2004 and 2007, and its abundance varied between ~4,200 and 8,300 individuals (Katsanevakis, 2009). During May 2019, the survey of a single site in Vouliagmeni Lake indicated a mortality rate of more than 89%, whereas live individuals were clearly affected. This translates into at least 3,700-7,400 dead individuals (assuming the abundance of the species in 2019 remained within the 2004-2007 range) by May 2019. Similarly, in Souda Bay (NW Crete) in 2007, the estimated total abundance of the species was approximately 130,900 individuals (with a 95% CI of 100,600-170,400) (Katsanevakis & Thessalou-Legaki, 2009). According to our results, this dense population has greatly been affected, since at the sites surveyed during June and July 2019 only dead individuals were encountered. A local professional diver (Nikos Giannoulakis interviewed by D. Poursanidis) also reported full mortality of the species up to 60 m depth. Although further surveys would be needed to confirm that all fan mussels in the entire gulf are dead, it is highly likely that the entire population of ~130,900 individuals is lost. Finally, a study conducted in Gera Gulf (Lesvos Island) in 2016, revealed the greatest (until then) fan mussel population recorded in Greece, with an estimated abundance of 213,300 (CI: 97,600-466,000) individuals (Tsatiris et al., 2018). Visual surveys conducted during August-October and December 2018 provided mortality rate estimates greater than 90% and 100% respectively, indicating the collapse of the previously thriving population (Katsanevakis et al., 2019). Revisiting the area during the summers of 2019 and 2020 (Case Study 3) confirmed the great loss of this population, since the estimated mortality of fan mussels in the gulf reached 99.8%, with only a single live individual recorded. Hence, in these three areas alone, it is estimated that ~350,000 P. nobilis individuals were lost due to the MME. Obviously, this is only a small fraction of the total number of lost individuals in the Greek seas.

It is promising for the survival of the species that surviving populations were found in two areas (Kalloni Gulf, Laganas Bay). Laganas Bay is a protected area, and although not dense, the fan mussel population there appeared to maintain a relatively healthy state in 2019, also benefiting from management measures that are in place, such as prohibition of anchoring and trawling, regulation of maritime traffic, and seasonal fishing closures.

The fan mussel population in Kalloni Gulf should be considered as the largest currently living population in the eastern Mediterranean. The abundance of live individuals estimated by the present study (~684,000 individuals) is the highest ever recorded in the eastern Mediterranean. Moreover, as this survey (Case Study 2) covered only part of the Gulf, it is quite likely that the number of live individuals in the Gulf exceeded one million by June 2019. Despite the high mortality rates estimated after June 2019 from opportunistic records (August-September 2019: 93.7%, May-June 2020: 89.2%), translated into hundreds of thousands of dead individuals, juvenile *P. nobilis* were also present in the studied area (30.8% of the live individuals recorded in May-June 2020), indicating successful reproduction in 2019.

In North Evoikos Gulf, consistently strong currents prevailing in the Evripus Strait presumably delayed the arrival of the pathogen from the South Evoikos Gulf, and as a result, the populations surveyed in September 2019 appeared to be unaffected. However, 2020 records have confirmed the expansion of the MME, and consequently the collapse of these populations.

The two areas with important live fan mussel populations in 2019-2020 (Kalloni Gulf and Laganas Bay) should be regularly monitored, also adopting targeted research activities aiming to shed light on the role of environmental conditions for the resilience of the species. Proper management aimed at minimizing accidental mortality due to human interactions (e.g. anchoring and fishing activities) or poaching should be pursued, as local anthropogenic stressors can further threaten the viability of the surviving populations. Experience gained since 2016 from the conservation of the remaining fan mussel populations (García-March et al., 2020), will allow effective management of the remaining Greek populations. These populations may potentially act as sources of recolonization for other areas where fan mussels are now locally extinct. For example, the remaining population in Kalloni Gulf could be the source of the identified single juvenile in the neighbouring Gera Gulf where the population was entirely lost in 2018. Recolonization of Gera Gulf and other coastal areas around Lesvos Island will benefit from safeguarding the offspring production of Kalloni Gulf. The fact that juveniles have been observed in Kalloni Gulf (and in other areas) indicates successful reproduction.

Areas with surviving populations are very important for the conservation of the species, and systematic regular monitoring is highly recommended. Field studies in different areas and depths might reveal more locations with live populations. Deep sites are under-sampled (the vast majority of surveys reported herein are at depths <30 m), and thus the status of *P. nobilis* populations in the deeper part of their depth range (i.e. 30-60 m) needs further investigation. The resistance of surviving populations might be due to local abiotic conditions (e.g. salinity or temperature) that either control or delay the possible expansion of the pathogen, resulting in natural refugia against the pathogen(s), or improving the host's health and immune system (Cabanellas-Reboredo et al. 2019). Assessment of genetic variation of surviving populations is therefore considered essential to track potential recolonization events of affected areas in the near future and support sound management decisions.

Even in heavily affected populations, a small number of surviving individuals, potentially resistant to the disease, were found. Such likely resistant strains may produce resistant offspring, but being isolated decreases the probability of successful reproduction, as reproductive success depends on the proximity of other individuals spawning synchronously. Hence, a proper strategy is needed to assist viability and increase the chances of reproduction of these resistant strains, e.g. by transplanting them in groups in well-protected areas (Katsanevakis, 2016). Nevertheless, it should be confirmed that these individuals did not survive due to reasons other than being resistant (e.g. due to local hydrodynamics, physicochemical conditions or just chance), otherwise transplantation might have a negative outcome.

Excluding the eight case studies, data from other areas used for this study consisted of reports and observations made by scientists, assessing the ongoing fan mussel MME or gathering this information as a side project. As a result, the geographical range of available data does not fully cover the entire Greek coastline and is not representative of the species bathymetric distribution. Anecdotal information provided by citizens (divers, fishermen etc.) has been published in social media, reporting mortalities at other locations (not included herein). The participation of citizen scientists through a dedicated structured online platform and a proper verification strategy may prove useful for filling the gap of knowledge and contribute valuable information, helping scientists to understand the spatiotemporal progress of the disease (Vázquez-Luis et al. 2017; Cabanellas-Reboredo et al., 2019).

Apart from Haplosporidium pinnae (Vázquez-Luis et al., 2017; Catanese et al., 2018; Cabanellas-Reboredo et al., 2019), and Mycobacterium sp. (Carella et al., 2019, 2020), which have been identified as the primary suspects of this Mediterranean pandemic, the actual spread of the pathogenic organism/s may be reinforced by the wider deterioration of the coastal marine environment and the effects of climate change. For example, the increase in frequency and extent of mucilaginous events (Goffredo & Dubinsky, 2014) has been linked to sea warming, and these aggregates can act as carriers of microorganisms, facilitating the spread of pathogenic bacteria (Danovaro et al., 2009). Increased temperature can facilitate pathogen outbreaks and intensify the pathogenicity of specific species, such as in the case of Vibrio mediterranei infecting P. nobilis (Prado et al., 2020). The observed multitude of pathogens associated with the P. nobilis MMEs suggests that the immune status of the animals could be involved (Carella et al., 2020), and it is likely that the spreading of the pathogens associated with the disease is favoured by a, yet unknown, common driver (Carella et al., 2019). Whether the latter may be an increase in sea water temperature, linked per se to episodes of mass mortality of sessile organisms both at Mediterranean (Garrabou et al., 2019) and at global scale (Smale et al., 2019), or its positive impact on other causes/events, such as the formation of marine mucilage aggregates or the transformation of a previous symbiont into a pathogen due to changes in the inner conditions of the host (Darriba, 2017), it seems that potential cascading impacts of climate change may play an important role in driving biological changes, thus altering the status of vulnerable marine ecosystem components. More research is needed to understand the real cause of the P. nobilis MME, but time is limited as the fan mussel populations are collapsing and the species is on the brink of extinction.

#### Acknowledgements

Case studies 1, 2, 3 and 4 were conducted within the framework of the PinnaSOS project, supported by the MSc program "Integrated Coastal Management" of the Department of Marine Sciences, University of the Aegean, and the "Development of the scientific diving lab" project (code: 70091) funded by the Research Unit of the University of the Aegean. Case study 7 was financed by the Interreg MED programme AMAre (Actions for Marine Protected Areas). Case study 8 was conducted within the framework of the BLUECOAST ("Climate-Smart Coastal Practices for Blue governance") (Interreg IPA CBC Programme "Greece-Albania 2014-2020", Ref. No. 4763/12-10-2016) project, co-funded by the European Union and the Greek government. EK, YI, MSi, and SK were also supported by the "Development of innovative methods for the study of marine food webs" project (MIS 5004302) under the call for proposals "Supporting researchers with an emphasis on new researchers" (EDULLL 34); the project is co-financed by Greece and the European Union (European Social Fund - ESF) under the Operational Programme Human Resources Development, Education and Lifelong Learning 2014-2020. Finally, opportunistic records across Greece made by YI, VG, MSa, and PL were supported by the ongoing monitoring projects "Water Framework Directive (WFD) Monitoring Network for Coastal and Transitional Waters in Greece, 2018-2023" and "MSFD implementation in Greece, 2018-2023".

### References

- Addis, P., Secci, M., Brundu, G., Manunza, A., Corrias, S. et al., 2009. Density, size structure, shell orientation and epibiontic colonization of the fan mussel *Pinna nobilis* L. 1758 (Mollusca: Bivalvia) in three contrasting habitats in an estuarine area of Sardinia (W Mediterranean). *Scientia Marina*, 73 (1), 143-152.
- Arditsoglou, A., Voutsa, D., 2012. Occurrence and partitioning of endocrine-disrupting compounds in the marine environment of Thermaikos Gulf, Northern Aegean Sea, Greece. *Marine Pollution Bulletin*, 64 (11), 2443- 2452.
- Basso, L., Vázquez-Luis, M., García-March, J., Deudero, S., Alvarez, E. *et al.*, 2015. The pen shell, *Pinna nobilis*: A review of population status and recommended research priorities in the Mediterranean Sea. *Advances in Marine Biolo*gy, 71, 109-160.
- Buckland, S.T., Anderson, D.R., Burnham, K.P., Laake, J.L., Borchers, D.L. et al., 2001. Introduction to Distance Sampling. Oxford University Press, Oxford, 432 pp.
- Butler, A., Vicente, N., de Gaulejac, B., 1993. Ecology of the pterioid bivalves *Pinna bicolor* Gmelin and *Pinna nobilis* L. *Marine Life*, 3 (1-2), 37-45.
- Cabanellas-Reboredo, M., Vázquez -Luis, M., Mourre, B., Alvarez, E., Deudero, S. *et al.*, 2019. Tracking a mass mortality outbreak of pen shell *Pinna nobilis* populations: A collaborative effort of scientists and citizens. *Nature Scientific Reports*, 9, 13355.

- Carella, F., Aceto, S., Pollaro, F., Miccio, A., Iaria, C. *et al.*, 2019. A mycobacterial disease is associated with the silent mass mortality of the pen shell *Pinna nobilis* along the Tyrrhenian coastline of Italy. *Scientific Reports*, 9, 2725.
- Carella, F., Elisabetta, A., Simone, F., Fulvio, S., Daniela, M. et al., 2020. In the Wake of the Ongoing Mass Mortality Events: Co-occurrence of Mycobacterium, Haplosporidium and Other Pathogens in Pinna nobilis Collected in Italy and Spain (Mediterranean Sea). Frontiers in Marine Science, 7, 48.
- Catanese, G., Grau, A., Valencia, J.M., García-March, J.M., Álvarez, E. et al., 2018. Haplosporidium pinnae sp. nov., a haplosporidan parasite associated with massive mortalities of the fan mussel, Pinna nobilis, in the Western Mediterranean Sea. Journal of Invertebrate Pathology, 157, 9-24.
- Čižmek, H., Čolić, B., Gračan, R., Grau, A., Catanese, G., 2020. An emergency situation for pen shells in the Mediterranean: the Adriatic Sea, one of the last *Pinna nobilis* shelters, is now affected by a mass mortality event. *Journal* of Invertebrate Pathology, 173, 107388
- Danovaro, R., Fonda Umani, S., Pusceddu, A., 2009. Climate Change and the Potential Spreading of Marine Mucilage and Microbial Pathogens in the Mediterranean Sea. *PLoS ONE*, 4 (9), e7006.
- Darriba, S., 2017. First haplosporidan parasite reported infecting a member of the Superfamily Pinnoidea (*Pinna nobilis*) during a mortality event in Alicante (Spain, Western Mediterranean). Journal of Invertebrate Pathology, 148, 14-19.
- De Gaulejac, B., 1995. Successive hermaphrodism with asynchronous maturation of *Pinna nobilis* (L.) (Bivalvia: Pterioidea). Comptes Rendus de l'Academie des Sciences Paris-Series III. Sciences de la vie. *Biologie et Pathologie Animal*, 318, 99-103.
- Deudero, S., Vázquez-Luis, M., Alvarez, E., 2015. Human Stressors Are Driving Coastal Benthic Long-Lived Sessile Fan Mussel *Pinna nobilis* Population Structure More than Environmental Stressors. *PLoS ONE*, 10 (7), e0134530.
- Evagelopoulos, A., Koutsoubas, D., 2008. Seasonal community structure of the molluscan macrofauna at the marine-lagoonal environmental transition at Kalloni solar saltworks (Lesvos Island, NE Aegean Sea, Greece). *Journal of Natural History*, 42, 597-618.
- Frantzis, A., Herzing, D.L., 2002. Mixed-species associations of striped dolphins (*Stenella coeruleoalba*), short-beaked common dolphins (*Delphinus delphis*), and Risso's dolphins (*Grampus griseus*) in the Gulf of Corinth (Greece, Mediterranean Sea). Aquatic Mammals, 28 (2), 188-197.
- Galinou-Mitsoudi, S., Vlahavas, G., Papoutsi, O., 2006. Population study of the protected bivalve *Pinna nobilis* (Linaeus, 1758) in Thermaikos Gulf (North Aegean Sea). *Journal of Biological Research*, 5, 47-53.
- García-March, J.R., 2005. Aportaciones al conocimiento de la Biologia de Pinna nobilis Linneo, 1758 (Mollusca: Bivalvia) en el litoral mediterraneo Iberico. Publication Service of the University of Valencia, Valencia, 332 pp.
- García-March, J.R., Tena-Medialdea, J., Henandis, S., Vázquez-Luis, M., López, D. *et al.*, 2020. Can we save a marine species affected by a highly infective, highly lethal, waterborne disease from extinction? *Biological Conservation*, 243, 108498.
- Garrabou, J., Gómez-Gras, D., Ledoux, J.-B., Linares, C., Ben-

auna at the marme-la- tion, abundance and

- Katsanevakis, S., Leftaditou, E., Galinou-Mitsoudi, S., Koutsoubas, D., Zenetos, A., 2008. Molluscan species of minor commercial interest in Hellenic seas: Distribution, exploitation and conservation status. *Mediterranean Marine Science*, 9 (1), 77-118.
- Katsanevakis, S., Poursanidis, D., Issaris, Y., Panou, A., Petza, D. *et al.*, 2011. "Protected" marine shelled molluscs: thriving in Greek seafood restaurants. *Mediterranean Marine Science*, 11(2), 429-438.
- Katsanevakis, S., Tsirintanis, K., Tsaparis, D., Doukas, D., Sini, M. et al., 2019. The cryptogenic parasite Haplosporidium pinnae invades the Aegean Sea and causes the collapse of Pinna nobilis populations. Aquatic Invasions, 14 (2), 150-164.
- Kefalas, E., Castritsi-Catharios, J., Zouganelis, G.D., 2016. A study of the benthic community structure of Demospongiae (Porifera) in Kalloni gulf at Lesvos Island (NE Aegean Sea, Greece). *Regional Studies in Marine Science*, 5, 12-18.
- Kersting, D.K. & García-March, J.R., 2017. Long-term assessment of recruitment, early stages and population dynamics of the endangered Mediterranean fan mussel *Pinna nobilis* in the Columbretes Islands (NW Mediterranean). *Marine*

soussan, N. *et al.*, 2019. Collaborative database to track mass mortality events in the Mediterranean Sea. *Frontiers in Marine Science*, 6, 707.

- Giacobbe, S., 2002. Epibiontic mollusc communities on *Pinna* nobilis L. (Bivalvia, Mollusca). Journal of Natural History, 36 (12), 1385-1396.
- Goffredo, S., Dubinsky, Z. (Eds), 2014. *The Mediterranean Sea: its History and Present Challenges.* Springer, Dordrecht, 678 pp.
- Issaris, Y., Katsanevakis, S., Pantazi, M., Vassilopoulou, V., Panayotidis, P. *et al.*, 2012. Ecological mapping and data quality assessment for the needs of ecosystem-based marine spatial management: case study Greek Ionian Sea and the adjacent gulfs. *Mediterranean Marine Science*, 13 (2), 297-311.
- IUCN, 2019. Mediterranean Noble Pen Shell Crisis (*Pinna no-bilis*) January 2020 Update. https://www.iucn.org/news/ mediterranean/202001/mediterranean-noble-pen-shell-crisis-pinna-nobilis-january-2020-update (Assessed 15 January 2020).
- Katsanevakis, S., 2006. Population ecology of the endangered fan mussel *Pinna nobilis* in a marine lake. *Endangered Species Research*, 1, 51-59.
- Katsanevakis, S., 2007. Growth and mortality rates of the fan mussel *Pinna nobilis* in Lake Vouliagmeni (Korinthiakos Gulf, Greece): a generalized additive modelling approach. *Marine Biology*, 152 (6), 1319-1331.
- Katsanevakis, S, 2009. Estimating abundance of endangered marine benthic species using Distance Sampling through SCUBA diving: the *Pinna nobilis* (Mollusca: Bivalvia) example. p. 81-115. In: *Endangered Species: New Research*. Columbus, A.M., Kuznetsov, L. (Eds). Nova Science Publishers, New York.
- Katsanevakis, S., 2016. Transplantation as a conservation action to protect the Mediterranean fan mussel *Pinna nobilis*. *Marine Ecology Progress Series*, 546, 113-122.
- Katsanevakis, S., Thessalou-Legaki, M., 2009. Spatial distribution, abundance and habitat use of the protected fan mussel *Pinna nobilis* in Souda Bay, Crete. *Aquatic Biology*, 8, 45-54.

Environmental Research, 130, 282-292.

- Kersting, D., Benabdi, M., Čižmek, H., Grau, A., Jimenez, C. *et al.*, 2019. *Pinna nobilis*. The IUCN Red List of Threatened Species 2019, e.T160075998A160081499, 24 pp.
- Koulouri, P., Kalogirou, S., Maidanou, M., Koutsoubas, D., Dounas, C., 2016. Fish and cephalopod assemblage structure of green alga *Caulerpa prolifera* (Chlorophyta) meadow in the eastern Mediterranean Sea (Elounda Bay, Crete Island). *Regional Studies in Marine Science*, 3, 33-41.
- Lattos, A., Giantsis, I.A., Karagiannis, D., Michaelidis, B., 2020. First detection of the invasive Haplosporidian and Mycobacteria parasites hosting the endangered bivalve *Pinna nobilis* in Thermaikos Gulf, North Greece. *Marine Environmental Research*, 155, 104889.
- Lykousis, V., Sakellariou, D., Moretti, I., Kaberi, H., 2007. Late Quaternary basin evolution of the Gulf of Corinth: Sequence stratigraphy, sedimentation, fault-slip and subsidence rates. *Tectophysics*, 440 (1-4), 29-51.
- Maeder, F., 2008. Sea-silk in aquincum: first production in antiquity. 109-118. proof p. In: Purpureae Vestes: Textiles and Dyes in Antiquity. Vestidos, textiles y tintes : estudios sobre la producción de bienes de consumo en la antigüedad : actas del II Symposium Internacional sobre Textiles y Tintes del Mediterráneo en el Mundo Antiguo (Atenas, 24 al 26 de noviembre, 2005). II. Giner, C.A., Karali, L. (Eds). Universitat de València, València.
- Margaritoulis, D., 2005. Nesting Activity and Reproductive Output of Loggerhead Sea Turtles, *Caretta caretta*, Over 19 seasons (1984-2002) at Laganas Bay, Zakynthos, Greece: The Largest Rookery in the Mediterranean. *Chelonian Conservation and Biology*, 4(4), 916-929.
- Panarese, R., Tedesco, P., Chimienti, G., Latrofa, M.S., Quaglio, F. *et al.*, 2019. *Haplosporidium pinnae* associated with mass mortality in endangered *Pinna nobilis* (Linnaeus 1758) fan mussels. *Journal of Invertebrate Pathology*, 164, 32-37.
- Papageorgiou, M., Christopoulou, O., Kostopoulou, T., 2017. Urban pressure in coastal areas of Greece: the case of Corinthiakos Gulf. p. 515-525. In: Proceedings of the International Conference on Changing Cities III: Spatial, Design, Landscape & Socio-economic Dimensions, Syros Island, 26-30 June 2017. Grafima Publications, Thessaloniki.
- Perissoratis, C., Piper, D.J.W., Lykousis, V., 2000. Alternating marine and lacustrine sedimentation during late Quaternary in the Gulf of Corinth rift basin, central Greece. *Marine Geology*, 167 (3-4), 391-411.
- Poulos, S.E., Collins, M.B., Pattiaratchi, C., Cramp, A., Gull, W. et al., 1996. Oceanography and sedimentation in the semi-enclosed, deep-water Gulf of Corinth (Greece). *Marine Geology*, 134 (3-4), 213-235.
- Prado, P., Caiola, N., Ibáñez, C., 2014. Habitat use by a large population of *Pinna nobilis* in shallow waters. *Scientia Marina*, 78 (4), 555-565.
- Prado, P., Carrasco, N., Catanese, G., Grau, A., Cabanes, P. et al., 2020. Presence of Vibrio mediterranei associated to major mortality in stabled individuals of Pinna nobilis L. Aquaculture, 519, 734899.
- Rabaoui, L., Tlig-Zouari, S., Cosentino, A., Ben Hassine, O.K., 2009. Associated fauna of the fan shell *Pinna nobilis* (Mollusca: Bivalvia) in the northern and eastern Tunisian coasts.

Scientia Marina, 73, 129-141.

- Rabaoui, L., Tlig-Zouari, S., Katsanevakis, S., Ben Hassine, O.K., 2010. Modelling population density of *Pinna nobilis* (Bivalvia) on the eastern and southeastern coast of Tunisia. *Journal of Molluscan Studies*, 76 (4), 340-347.
- Richardson, C.A., Kennedy, H.A., Duarte, C.M., Proud, S.V.P., 1999. Age and growth of the fan mussel *Pinna nobilis* from S.E. Spanish Mediterranean seagrass *Posidonia oceanica* meadows. *Marine Biology*, 133, 205-212.
- Rodríguez, S., Balboa, S., Olveira, G., Montes, J., Moreno, D. et al., 2018. First report of mass mortalities in natural population of *Pinna nobilis*. A microbial perspective. FEMS7-3233. In: *The 7th Congress of European Microbiologists* (*FEMS 2018*), Valencia, Spain, 9-13 July 2017. FEMS, Valencia.
- Rouanet, E., Trigos, S., Vicente, N., 2015. From youth to death of old age: the 50-year story of a *Pinna nobilis* fan mussel population at Port - Cros Island (Port - Cros National Park, Provence, Mediterranean Sea). *Scientific Reports of Port -Cros National Park*, 29, 209-222.
- Russo, P., 2012. Segnalazione di una grande colonia di *Pinna* nobilis (Linnaeus, 1758) nella Laguna di Venezia. Notiziario della Società Italiana di Malacologia, 30 (1), 31-34.
- Schultz, P.W.W., Huber, M., 2013. Revision of the worldwide recent Pinnidae and some remarks on fossil European Pinnidae. Acta Conchyliorum, 13, 1-164.
- Sini, M., Katsanevakis, S., Koukourouvli, N., Gerovasileiou, V., Dailianis, T. *et al.*, 2017. Assembling Ecological Pieces to Reconstruct the Conservation Puzzle of the Aegean Sea. *Frontiers in Marine Science*, 4, 347.
- Sini, M., Garrabou, J., Trygonis, V., Koutsoubas, D., 2019. Coralligenous formations dominated by *Eunicella cavoli*ni (Koch, 1887) in the NE Mediterranean: biodiversity and structure. *Mediterranean Marine Science*, 20(1), 174-188.
- Smale, D.A., Wernberg, T., Oliver, E.C.J., Thomsen, M., Harvey, B.P., et al., 2019. Marine heatwaves threaten global biodiversity and the provision of ecosystem services. Nature Climate Change, 9, 306-312.
- Trigos, S., García-March, J.R., Vicente, N., Tena, J., Torres, J., 2014. Utilization of muddy detritus as organic matter source by the fan mussel *Pinna nobilis*. *Mediterranean Marine Science*, 13(3), 667-674.
- Tsatiris, A., Papadopoulos, V., Makri, D., Topouzelis, K., Manoutsoglou, E. *et al.*, 2018. Spatial distribution, abundance and habitat use of the endemic Mediterranean fan mussel *Pinna nobilis* in Gera Gulf, Lesvos (Greece): comparison of design-based and model-based approaches. *Mediterranean Marine Science*, 19 (3), 642-655.
- Vázquez-Luis, M., Alvarez, E., Barrajon, A., García-March, J.R., Grau, A. *et al.*, 2017. S.O.S. *Pinna nobilis*: a mass mortality event in western Mediterranean Sea. *Frontiers in Marine Science*, 4, 109.
- Vázquez-Luis, M., Borg, J.A., Morell, C., Banach-Esteve, G., Deudero, S., 2015. Influence of boat anchoring on *Pinna nobilis*: a field experiment using mimic units. *Marine and Freshwater Research*, 66 (9), 786-794.
- Vázquez-Luis, M., March, D., Alvarez, E., Alvarez-Berastegui, D., Deudero, S., 2014. Spatial distribution modelling of the endangered bivalve *Pinna nobilis* in a Marine Protected Area. *Mediterranean Marine Science*, 15 (3), 626-634.

Zavodnik, D., Hrs-Brenko, M., Legac, M., 1991. Synopsis on the fan shell *Pinna nobilis* L. In the eastern Adriatic Sea. p. 169-178. In : *Les Espèces Marines à protéger en Méditerra*- née. Boudouresque, C.F., Avon, M., Gravez, V. (Eds). Gis posidonie Parc scientifique et technologique de Luminy, Marseille.

## Supplementary data

The following supplementary information is available online for the article:

Detailed information on all 258 surveyed sites (date, location, coordinates, depth, predominant habitats, number of live and dead fan mussels).