

Mediterranean Marine Science

Vol 22, No 1 (2021)

Vol 22, No 1 (2021)



The dual impact of *Ostreopsis cf. ovata* on *Mytilus galloprovincialis* and *Paracentrotus lividus*: Toxin accumulation and pathological aspects

ANGELA SARDO, RACHELE ROSSI, VITTORIO SOPRANO, PATRIZIA CIMINIELLO, ERNESTO FATTORUSSO, PAOLA CIRINO, ADRIANA ZINGONE

doi: [10.12681/mms.24160](https://doi.org/10.12681/mms.24160)

To cite this article:

SARDO, A., ROSSI, R., SOPRANO, V., CIMINIELLO, P., FATTORUSSO, E., CIRINO, P., & ZINGONE, A. (2021). The dual impact of *Ostreopsis cf. ovata* on *Mytilus galloprovincialis* and *Paracentrotus lividus*: Toxin accumulation and pathological aspects. *Mediterranean Marine Science*, 22(1), 59–72. <https://doi.org/10.12681/mms.24160>

The dual impact of *Ostreopsis cf. ovata* on *Mytilus galloprovincialis* and *Paracentrotus lividus*: Toxin accumulation and pathological aspects

Angela SARDO, Rachele ROSSI, Vittorio SOPRANO, Patrizia Ciminiello, Ernesto FATTORUSSO,
Paola CIRINO and Adriana ZINGONE

Mediterranean Marine Science, 2021, 22 (1)

Table S1. Cell ingestion and mussel toxicity in the 24 h experiments in which *Mytilus galloprovincialis* were fed with *Ostreopsis cf. ovata* cultures. Toxins were always below the detection limit at the chemical analyses.

Experiment number	Replicate	Ingested <i>O. cf. ovata</i> (cells · 10 ⁴ g ⁻¹ mussel WW)	N° dead mice (death times)
1	1	3.24	3 (<2h)
	2	3.15	3 (2 <5 h; 1 6-22h)
	3	3.43	3 (1<3h; 1<5h; 1 6-22h)
2	1	4.96	2 (6-22 h)
	2	2.07	0
	3	1.18	0
3	1	5.55	3 (4-22 h)
	2	5.40	2 (4-22 h)
	3	4.64	1 (4-22 h)
4	1	0.88	0
	2	0.25	0
	3	1.33	0
5	1	0.66	0
	2	0.60	0
	3	1.29	0
6	1	3.02	3 (2 1-5 h; 1 5-24 h)
	2	3.36	3 (2 1-5 h; 1 5-24 h)
	3	3.27	3 (1-5 h)

Table S2. Feeding of *Paracentrotus lividus* on the red macroalga *Asparagopsis taxiformis* with *O. cf. ovata* as epiphyte: sea-urchin and macroalga wet weight (WW) and total epiphytic cells in each experiment (avg±SDV). Macroalgae were completely eaten in all cases. Asterisks indicate samples in which sea urchin tissues were weakly positive to the mouse bioassay (1-2 mice dead in less than 24 hours in 1-2 replicate samples). Gonads analysed separately were not toxic.

Experiment Number	<i>P. lividus</i> soft tissues WW (g)	<i>A. taxiformis</i> WW (g)	Total <i>O. cf. ovata</i> (cells · 10 ⁵)
1	40.6 ± 7.2 (*)	55.8 ± 0.6	3.3 ± 0.9
2	34.4 ± 7.4	131.5 ± 0.5	7.4 ± 1.3
3	28.5 ± 13.4 (*)	85.3 ± 2.5	8.7 ± 0.5
4a	42.2 ± 4.5	87.5 ± 0.7	8.3 ± 0.9
4b	42.2 ± 4.5	81.0 ± 1.9	8.6 ± 1.7
5	43.1 ± 6.2	81.2 ± 0.7	2.7 ± 0.3
6	49.5 ± 5.9	84.0 ± 0.1	2.9 ± 0.4
7	69.9 ± 8.8	101.3 ± 0.2	1.7 ± 0.3
8	55.5 ± 2.7	70.9 ± 0.2	1.2 ± 0.2