

## Mediterranean Marine Science

Vol 22, No 1 (2021)

Vol 22, No 1 (2021)



### The dual impact of *Ostreopsis cf. ovata* on *Mytilus galloprovincialis* and *Paracentrotus lividus*: Toxin accumulation and pathological aspects

ANGELA SARDO, RACHELE ROSSI, VITTORIO SOPRANO, PATRIZIA CIMINIELLO, ERNESTO FATTORUSSO, PAOLA CIRINO, ADRIANA ZINGONE

doi: [10.12681/mms.24160](https://doi.org/10.12681/mms.24160)

#### To cite this article:

SARDO, A., ROSSI, R., SOPRANO, V., CIMINIELLO, P., FATTORUSSO, E., CIRINO, P., & ZINGONE, A. (2021). The dual impact of *Ostreopsis cf. ovata* on *Mytilus galloprovincialis* and *Paracentrotus lividus*: Toxin accumulation and pathological aspects. *Mediterranean Marine Science*, 22(1), 59–72. <https://doi.org/10.12681/mms.24160>

## The dual impact of *Ostreopsis cf. ovata* on *Mytilus galloprovincialis* and *Paracentrotus lividus*: Toxin accumulation and pathological aspects

Angela SARDO, Rachele ROSSI, Vittorio SOPRANO, Patrizia Ciminiello, Ernesto FATTORUSSO,  
 Paola CIRINO and Adriana ZINGONE

*Mediterranean Marine Science*, 2021, 22 (1)

**Table S1.** Cell ingestion and mussel toxicity in the 24 h experiments in which *Mytilus galloprovincialis* were fed with *Ostreopsis cf. ovata* cultures. Toxins were always below the detection limit at the chemical analyses.

Experiment number	Replicate	Ingested <i>O. cf. ovata</i> (cells · 10 <sup>4</sup> g <sup>-1</sup> mussel WW)	N° dead mice (death times)
1	1	3.24	3 (<2h)
	2	3.15	3 (2 <5 h; 1 6-22h)
	3	3.43	3 (1<3h; 1<5h; 1 6-22h)
2	1	4.96	2 (6-22 h)
	2	2.07	0
	3	1.18	0
3	1	5.55	3 (4-22 h)
	2	5.40	2 (4-22 h)
	3	4.64	1 (4-22 h)
4	1	0.88	0
	2	0.25	0
	3	1.33	0
5	1	0.66	0
	2	0.60	0
	3	1.29	0
6	1	3.02	3 (2 1-5 h; 1 5-24 h)
	2	3.36	3 (2 1-5 h; 1 5-24 h)
	3	3.27	3 (1-5 h)

**Table S2.** Feeding of *Paracentrotus lividus* on the red macroalga *Asparagopsis taxiformis* with *O. cf. ovata* as epiphyte: sea-urchin and macroalga wet weight (WW) and total epiphytic cells in each experiment (avg±SDV). Macroalgae were completely eaten in all cases. Asterisks indicate samples in which sea urchin tissues were weakly positive to the mouse bioassay (1-2 mice dead in less than 24 hours in 1-2 replicate samples). Gonads analysed separately were not toxic.

Experiment Number	<i>P. lividus</i> soft tissues WW (g)	<i>A. taxiformis</i> WW (g)	Total <i>O. cf. ovata</i> (cells · 10 <sup>5</sup> )
1	40.6 ± 7.2 (*)	55.8 ± 0.6	3.3 ± 0.9
2	34.4 ± 7.4	131.5 ± 0.5	7.4 ± 1.3
3	28.5 ± 13.4 (*)	85.3 ± 2.5	8.7 ± 0.5
4a	42.2 ± 4.5	87.5 ± 0.7	8.3 ± 0.9
4b	42.2 ± 4.5	81.0 ± 1.9	8.6 ± 1.7
5	43.1 ± 6.2	81.2 ± 0.7	2.7 ± 0.3
6	49.5 ± 5.9	84.0 ± 0.1	2.9 ± 0.4
7	69.9 ± 8.8	101.3 ± 0.2	1.7 ± 0.3
8	55.5 ± 2.7	70.9 ± 0.2	1.2 ± 0.2