ePublishing

ΕΘΝΙΚΟ ΚΕΝΤΡΟ ΤΕΚΜΗΡΙΩΣΗΣ
EKT NATIONAL DOCUMENTATION CENTRE

Mediterranean Marine Science

Vol 22, No 2 (2021)

VOL 22, No 2 (2021)



New discoveries in Eastern Mediterranean mesophotic sponge grounds: updated checklist and description of three novel sponge species

TAL IDAN, SIGAL SHEFER, TAMAR FELDSTEIN, MICHA ILAN

doi: 10.12681/mms.25837

To cite this article:

IDAN, T., SHEFER, S., FELDSTEIN, T., & ILAN, M. (2021). New discoveries in Eastern Mediterranean mesophotic sponge grounds: updated checklist and description of three novel sponge species. *Mediterranean Marine Science*, *22*(2), 270–284. https://doi.org/10.12681/mms.25837

https://epublishing.ekt.gr | **e-Publisher:** EKT | **Downloaded at:** 29/03/2024 07:14:58

Mediterranean Marine Science
Indexed in WoS (Web of Science, ISI Thomson) and SCOPUS
The journal is available on line at http://www.medit-mar-sc.net
DOI: http://dx.doi.org/10.12681/mms.25837

New discoveries in Eastern Mediterranean mesophotic sponge grounds: updated checklist and description of three novel sponge species

Tal IDAN, Sigal SHEFER, Tamar FELDSTEIN and Micha ILAN

Mediterranean Marine Science, 2021, 22 (2)




Fig. S1: Phylogenetic tree used to support the morphological identification *Hemiasterella verae* sp. nov. (marked by '*'). The maximum likelihood tree was reconstructed with PhyML 3.0 (Guindon *et al*., 2010) based on 28S rDNA sequences, using the GTR model of sequence evolution. Bootstrap supports are given near the corresponding nodes.




Fig. S2: Phylogenetic tree based on 28S rDNA sequences, used to support the morphological identification of *Axinella venusta* sp. nov. (marked by '*'). The maximum likelihood tree was reconstructed with PhyML 3.0 (Guindon *et al*., 2010), using the GTR model of sequence evolution. Bootstrap supports (100 replicates) given near the corresponding nodes.