



# **Mediterranean Marine Science**

Vol 23, No 1 (2022)

VOL 23, No 1 (2022)



Sabellaria alveolata (Annelida, Polychaeta) bioconstructions and associated macroalgal community from Portopalo di Capo Passero (SE Sicily)

ROSSANA SANFILIPPO, DONATELLA SERIO, CLAUDIA DEIAS, ANTONIETTA ROSSO

doi: 10.12681/mms.27171

### To cite this article:

SANFILIPPO, R., SERIO, D., DEIAS, C., & ROSSO, A. (2021). Sabellaria alveolata (Annelida, Polychaeta) bioconstructions and associated macroalgal community from Portopalo di Capo Passero (SE Sicily). *Mediterranean Marine Science*, *23*(1), 150–156. https://doi.org/10.12681/mms.27171

## Short Communication

# Sabellaria alveolata (Annelida, Polychaeta) bioconstructions and associated macroalgal community from Portopalo di Capo Passero (SE Sicily)

#### Rossana SANFILIPPO<sup>1,2</sup>, Donatella SERIO<sup>2,3</sup>, Claudia DEIAS<sup>1</sup> and Antonietta ROSSO<sup>1,2</sup>

<sup>1</sup>Department of Biological, Geological and Environmental Sciences, University of Catania, Section of Earth Sciences, Corso Italia, 57, 95129, Catania, Italy

<sup>2</sup>CoNISMa, Piazzale Flaminio, 9, 00195, Roma

<sup>3</sup> Department of Biological, Geological and Environmental Sciences, University of Catania, Section of Plant Biology,

Via Empedocle, 58, 95128, Catania, Italy

Corresponding author: Rossana SANFILIPPO; sanfiros@unict.it

Contributing Editor: Konstantinos TSIAMIS

Received: 31 May 2021; Accepted: 6 December 2021; Published online: 14 March 2022

#### Abstract

The shallow-water reef building polychaete *Sabellaria alveolata* is recorded from a new locality in the southeastern corner of Sicily, off Portopalo di Capo Passero. Here, a particular habitat is formed on upper infralittoral rocky bottoms beaten by waves, with *S. alveolata* developing as veneer-like bioconstructions interspersed/covered with a dense macroalgal community including alien *Caulerpa* taxa and dominated by *Ericaria giacconei*. First insights about the development of this particular habitat, as well as observations on possible relationships between *Sabellaria* and the associated algae, are presented.

Keywords: Polychaetes; Sabellariidae; Fucales; *Ericaria giacconei;* infralittoral; bioconstruction; benthic ecology; alien species; Sicily Straits.

#### Introduction

The habitat-forming tubicolous polychaete Sabellaria alveolata (Linnaeus, 1767) is widely distributed in north-eastern Atlantic and the western Mediterranean Sea (e.g., La Porta & Nicoletti, 2009; Bertocci et al., 2017; Sanfilippo et al., 2020). The species occurs from midlittoral to upper infralittoral, where it lives solitary or more frequently aggregated, forming bioconstructions of a huge number of individuals adjacent to each other inside protective tubes built by agglutinating sand grains (Gruet, 1986; Sanfilippo et al., 2019). Depending on environmental conditions, the bioconstructions range from a few cm in thickness ("veneers" sensu Gruet, 1982) to ca. 50 cm (hummocks or pillows), up to more than 1 meter (reefs), with extensions of few square decimeters to several tens of square meters (Delbono et al., 2003; Noernberg et al., 2010).

Bioconstructions develop where the worm can catch sand particles suspended by turbulent waters (Gruet, 1972, 1982). For this reason, *Sabellaria* reefs seem restricted in Sicily to the sandy beaches that extensively occur along the southern coast (see Sanfilippo *et al.*, 2020).

Owing to its building activity, S. alveolata deeply transforms the colonised biotopes, increasing bottom

heterogeneity and providing additional substrates, space and shelter, thus promoting species richness (Dubois et al., 2002; Jones et al., 2018; Bonifazi et al., 2019, and references therein). However, little is known about associated animal taxa and their possible interactions with S. alveolata, especially for Mediterranean reefs. Papers often deal with single taxonomic groups (mostly mollusks, polychaetes and amphipod crustaceans), or even species (e.g., La Porta & Nicoletti, 2009; Iaciofano et al., 2015; Schimmenti et al., 2016), rather than the whole associated communities (Bertocci et al., 2017; Bonifazi et al., 2019). Even less is known about the associated flora, with only a few algae mentioned, including Ulva Linnaeus, Cladophora Kützing (Dubois et al., 2006), and Caulerpa J.V. Lamouroux (Musco et al., 2014; Vega Fernández et al., 2019). Analogously, the role of Sabellaria reefs as available substrate for colonization by algal communities is just mentioned (see Delbono et al., 2003).

In the southeastern corner of Sicily, near the village of Portopalo di Capo Passero, *S. alveolata* coexists with a rich algal community forming crustose bioconstructions on shallowly submerged rocky bottoms interspersed with a dense algal vegetation dominated by the rare species *Ericaria giacconei* Serio & G. Furnari. This species has been described from Porto Palo di Capo Passero (Serio & Furnari, 2021) to replace *Cystoseira hyblaea* Giaccone, in order to solve taxonomic and nomenclatural problems produced by the necessary genus displacement of an invalidly published species.

The present note aims at reporting and characterising these new *Sabellaria* bioconstructions from Portopalo di Capo Passero describing the assemblages they contribute to form, also focusing on macroalgal community and intergrowing algal species.

#### **Materials and Methods**

Veneer-like bioconstructions of *S. alveolata* inter growing with dense *Ericaria* community were discovered in September 2018 at the southeastern coast of Sicily, *ca*. 6 km SW of the Portopalo di Capo Passero village (Contrada Guardiana, 36.651056°N, 15.077067°E), in a small rocky shallow-water embayment constantly exposed to strong waves and winds. Cenozoic limestone constitutes both the colonized seafloor and emerged outcrops delimiting the embayment. The rocky bottom is gently sloping and locally covered by biogenic sandy sediments.

Since the first finding of these bioconstructions in 2018, several surveys have been carried out seasonally in the area. On the 2<sup>nd</sup> half of July 2020, scuba divers took pictures of the seabed through underwater GoPro 8Hero black camera to document the nature of the bottom and the distribution/extension of the *Sabellaria* bioconstructions and the algal cover (Fig. 1). In order to investigate the floristic biodiversity, sampling of macroalgal species growing on the rocky bottom and on *Sabellaria* crusts were carried out at -0.50 m depth in July and October

2020, detaching algae at their base using a small knife. Furthermore, the macroalgal species growing on two large-sized thalli of *E. giacconei* were studied (Table 1). The low number of replicates was forced by the need of minimizing the impact on a rare species.

Using a hammer, two small portions of the *Sabellaria* crust were sampled at the same depth (-0.50 m) in July 2020 to obtain living specimens for species identification and to describe the bioconstruction architecture. The *Sabellaria* samples PCP1 and PCP2 had sizes of *ca*. 8x11x4 cm and *ca*. 7x7x5 cm, respectively. The collected macroalgae and *Sabellaria* samples were preserved in 70% ethanol in seawater solution for subsequent analyses.

In the Palaeoecology laboratory of the Department of Biological, Geological and Environmental Science (Bi-GeA), Sabellaria individuals were extracted from their tubes and identified at species level under a Zeiss Axioplan II stereomicroscope using the morphological characters indicated by Fauvel (1927) and Kirtley (1994). The general architecture of the sampled Sabellaria bioconstruction was examined with the naked eye and photographed (Fig. 2). In the Phycology laboratory of the BiGeA, samples were studied for the identification of all macroalgal species under stereomicroscope. Morpho-anatomical observations were performed by optical microscope Zeiss Axioplan. Some specimens were stained with 1% aqueous aniline blue acidified with diluted HCl, which enhances visualization of pit connections. Sections were made by razor blade.

Sabellaria specimens and framework samples are housed in the Palaeontological Museum of the BiGeA (University of Catania). Algal material is housed in the



*Fig. 1: Sabellaria alveolata-Ericaria giacconei* habitat from Portopalo di Capo Passero (Sicily): A) Overview of the bottom with *Sabellaria* bioconstructions densely covered by *Ericaria* community. B) *Sabellaria* veneer-like crusts colonized by several thalli of macroalgae (*Laurencia* spp. and *Caulerpa* spp.). C) *Ericaria* thalli encircled by *Sabellaria* crusts. D) Portions of *Sabellaria* crusts covered by the invasive alien species *Caulerpa cylindracea* and *C. taxifolia* v. *distichophylla*.

Locality	Portopalo di Capo Passero (Syracuse)		
Station	Contrada Guardiana	a Contrada Guardiana	
Date	July 2020	October 2020	
Depth (m)	0.50	0.50	
RHODOPHYTA			
Ceramium codii (H. Richards) Mazoyer		+	Е
Ceramium comptum Børgesen	+	+	Е
Ceramium tenerrimum (G. Martens) Okamura	s		Е
Ceramium cfr brevizonatum H.E. Petersen	s	+	Е
Ceramium circinatum (Kützing) J. Agardh	+	+	Е
Ceramium nodosum (Kützing) A.W. Griffiths & Harvey		+	Е
Chondria dasyphylla (Woodward) C. Agardh		m	Е
Choreonema thuretii (Bornet) F. Schmitz	s		Е
Dasya rigidula (Kützing) Ardissone	+		Е
Dipterosiphonia rigens (C. Agardh) Falkenberg	+	+	Е
Erythrocystis montagnei (Derbès & Solier) P.C. Silva	+	+	Е
Gayliella mazoyerae T.O. Cho, Fredericq & Hommersand	+	+	Е
Herposiphonia secunda (C. Agardh) Ambronn	s	+	Е
Herposiphonia tenella (C. Agardh) Ambronn		+	Е
Heterosiphonia crispella (C. Agardh) M.J. Wynne	+	+	Е
Hydrolithon boreale (Foslie) Y.M. Chamberlain	+	+	Е
Jania pedunculata J.V. Lamouroux var. adhaerens (J.V. Lamouroux)		+	Е
A.S. Harvey, Woelkerling & Reviers			
Jania virgata (Zanardini) Montagne	s	+	ES
Laurencia microcladia Kützing	f	+	ES
Laurencia cfr obtusa (Hudson) J.V. Lamouroux		+	S
Laurencia intricata J.V. Lamouroux		+	S
Laurenciella marilzae (Gil-Rodríguez et al.) Gil-Rodríguez et al.	s	fs	S
Lithophyllum pustulatum (J.V. Lamouroux) Foslie	s	+	Е
Pneophyllum zonale (P. Crouan & H. Crouan) Y.M. Chamberlain	+	+	Е
Polysiphonia sertularioides (Grateloup) J. Agardh		+	Е
Rytiphlaea tinctoria (Clemente) C. Agardh	+	+	S
Taenioma nanum (Kützing) Papenfuss		+	Е
Womersleyella setacea (Hollenberg) R.E. Norris		+	Е
OCHROPHYTA			
Cystoseira compressa (Esper) Gerloff & Nizamuddin		j	S
Dictyota dichotoma (Hudson) J.V. Lamouroux	+	+	ES
Dictyota fasciola (Roth) J.V. Lamouroux	+	+	S
Dictyota mediterranea (Schiffner) G. Furnari	+	+	ES
Padina distromatica Hauck		+	S
Padina pavonica (Linnaeus) Thivy	+		S
Sphacelaria cirrosa (Roth) C. Agardh	+		Е
CHLOROPHYTA			
Caulerpa taxifolia (M. Vahl) C. Agardh var. distichophylla (Sonder)	+	+	S
Verlaque, Huisman & Procaccini			
Caulerpa cylindracea Sonder	+	+	S
Cladophora laetevirens (Dillwyn) Kützing	+	+	S
Phaeophila dendroides (P. Crouan & H. Crouan) Batters	+	+	Е
Pseudochlorodesmis furcellata (Zanardini) Børgesen	+		E
	nur	nher of species on Sabellaria	11

Table 1. Epiphytic macroalgal species on *Ericaria giacconei* (E) and *Sabellaria alveolata* (S). + = presence; f = female gametophyte; m = male gametophyte; s = sporophyte; j = juvenile stage.

number of species on Sabellaria 11

number of species on Ericaria 25

number of species on both Ericaria and Sabellaria 4

Total number of species 40

Rhodophyta 28

Ochrophyta 7

Chlorophyta 5



*Fig. 2: Sabellaria alveolata* from Portopalo di Capo Passero (Sicily): A) Outer surface of sample PCP1 showing tubes with intact apertural edges and well-developed rims. Scale bar: 5 mm. B) Outer surface of sample PCP2 with incomplete tubes disrupted in their distalmost part. Scale bar: 5 mm. C) Upper view of the veneer-like bioconstruction with prostrate and dorsally flattened tubes. Sample PCP2. Scale bar: 1 cm. D) Dorsal view of the anterior end of an individual with the opercular crown. Scale bar: 2 mm. E) Outer palea of the chitinous operculum figured in D bearing a smooth central tooth indicative of the species. Scale bar: 0.1 mm. F) Detail of the outer surface of sample PCP1 with a tube distal end hosting a stolon of *Caulerpa cylindracea*. Note the different nature, shape and size of the agglutinated sandy particles. Scale bar: 5 mm.

phycology laboratory of the BiGeA - Plant Biology Section of the University of Catania.

#### Results

In the study area, the *Sabellaria* bioconstructions cover a narrow rocky area subparallel to the coastline of about 100 square meters, from 0.3 m depth down to 1 m, growing on restricted bottom surfaces interspersed within a dense algal canopy dominated by *E. giacconei* (Fig. 1A). The *Sabellaria* bioconstructions are found below and around the *Ericaria* thalli and are locally colonized by some other algal species on their upper surface (Fig. 1 C-D). *E. giacconei* shows high specific coverage values, with largest thalli being up to 30 cm high.

The floristic study showed a total of 40 species (28 Rhodophyta, 7 Ochrophyta, 5 Chlorophyta) 25 of which found on *E. giacconei*, 11 on *Sabellaria* and 4 on both species (Table 1).

Both floristic study and field observations showed that the geniculate coralline red alga *Jania virgata* is very abundant on *E. giacconei*. Furthermore, numerous thalli of *Laurencia microcladia* and *Laurenciella marilzae*, as well as the invasive alien species *Caulerpa cylindracea* and *C. taxifolia* var. *distichophylla*, grow on the *Sabellaria* bioconstruction (Fig. 1B, D). *Sabellaria* patches no longer occur below one meter depth, where *E. giacconei* has lower coverage values. There, on rocky outcrops it is largely replaced by an association of species of the *Laurencia* complex associated with *Dictyota* spp., *Padina*  pavonica and Gongolaria elegans (Sauvageau) Molinari & Guiry, which become dominant. At this depth a sandy cover begins to be found on the bottom, hosting Caulerpa prolifera (Forsskål) J.V. Lamouroux, Dasycladus vermicularis (Scopoli) Krasser and Posidonia oceanica (Linnaeus) Delile.

The Sabellaria bioconstructions associated with the algal canopy consist of aggregates of adjacent sand-made tubes that form irregularly shaped and discontinuous veneer-like patches, each 5-20 cm thick and up to 60 cm wide (Fig. 1). Owing to their structure, consisting of tubes made by agglutinated sandy particles, these patches are barely noticeable, if not for their distinctive honeycomb-like surface, which derives from the regular pattern of adjacent tube apertures (Fig. 1B, C). Locally, large thalli of Ericaria are encircled and become immersed in between the Sabellaria crusts for much (up to the upper third) of the cauloid (Fig. 1C). In these sheltered portions below the algal fronds, the Sabellaria tubes show intact apertural edges with well-developed rims (Fig. 2A). Conversely, in portions of the bioconstruction exposed to hydrodynamism, unprotected by the *Ericaria* thalli, tubes are incomplete and more or less disrupted in their distalmost part (Fig. 2 B).

Observations performed in the field and on the two *Sabellaria* crust samples show that the bioconstruction corresponds to the veneer-like type, consisting of tubes irregularly bent in their initial part and then growing straight, subparallel to each other and prostrate in their distal part, with isoriented apertures facing landwards (Figs 1C, 2C). Tubes are circular to sub-quadrangular in

cross section and up to 7 cm long with a 3-6 mm wide lumen and an outer diameter up to 9 mm at the aperture (Fig. 2B, C). Each tube ends with an infundibular asymmetrical rim, usually much protruding on the flattened dorsal side, forming a sort of roof above the aperture (Fig. 2A, C). The tube wall is *ca*. 2 mm thick and typically three-layered. It is formed by agglutinate sandy particles mainly consisting of carbonate clasts and subordinate lithic grains of different shape and size (Fig. 2F). Living animals found inside tubes demonstrated they unequivocally belong to *S. alveolata* owing to the diagnostic morphology of the chitinous outer paleae of the operculum that show a smooth central tooth (Fig. 2D, E) compared to those of *S. spinulosa* that bear a central denticulate plume (Lezzi *et al.*, 2015: fig. 2).

#### **Discussion and Conclusions**

The new finding of S. alveolata from the Portopalo di Capo Passero coast integrates the known geographical distribution of the species along the coasts of Sicily. In the study area bioconstructions develop as veneers (see Gruet, 1982; Gruet & Bodeur, 1995, Curd et al., 2019) in contrast to what reported from other Sicilian areas, where hummocks, pillows and reefs have been identified (Schimmenti et al., 2016; Bertocci et al., 2017; Sanfilippo et al., 2019). This crustose bioconstruction, formed by tubes prostrate in their distal ends, and apertures directed towards the coastline, would indicate conditions of high hydrodynamic energy, as evidenced by Gruet & Bodeur (1995). This fits well with the environmental conditions of the investigated coastal sector that is beaten by strong waves and currents that are forced by dominant landwards directed winds (e.g., Ciani et al., 2019).

The association of macroalgal species with Sabellaria reefs has been very rarely reported (e.g., Dubois et al., 2006) since underwater imaging of these biostructures usually miss algal cover (e.g., Bonifazi et al., 2019). Although Sabellaria bioconstructions have been considered as biodiversity hotspots, very high values in overall abundances and species richness negatively correlate with the number of "sand crowns" considered as an index of health status for the worm (Bonifazi et al., 2019). Pending for results of ongoing monitoring, it seems that in the Portopalo di Capo Passero bioconstruction worm growth is not negatively affected by the presence of the numerous co-occurring algal species that grow on top and sideway. Instead, a beneficial influence could be suggested for this association because first observations after winter showed that on the upper surfaces of the Sabellaria crusts tube apertures are intact close to Ericaria fronds but disrupted far from the algal fronds.

The coexistence of *S. alveolata* and *E. giacconei* in the same biotope and their growth relationships deserve to be investigated, including the possible advantage for the perennial portion of *Ericaria* thalli to be encircled by the sandy crust formed by the worm, or likewise the positive effects of the presence of *E. giacconei* and other macroalgae on *Sabellaria* bioconstructions in mitigating too strong hydrodynamic action, thus favoring sediment capture by the worm. Although formally unreported, a similar interaction is visible on underwater photographs in a report of ARPA Sicilia (2014) for some *Sabellaria* bioconstructions located on bottoms *ca*. 100 km west of our study site, where algae seem to be absent but tufts of short *Posidonia oceanica* leaves are surrounded by the tube worm's aggregates. Seemingly, both *Ericaria* fronds and *Posidonia* leaves may perform the same positive function for *Sabellaria*. Further species found in epibiosis on both *E. giacconei* thalli and directly on the exposed surface of the *Sabellaria* bioconstruction are under examination in order to assess the total algal biodiversity, as well as the occurrence of associated sessile animal species.

Among macrophytes easily registered in the field, the invasive alien species *Caulerpa cylindracea* and *C. taxifolia* var. *distichophylla* are common and highly developed especially in summer months. The presence of stolons of *C. cylindracea*, even within some empty tubes of *Sabellaria* (Fig. 2F), highlights the invasive colonization strategy of this macroalga. Their possible impact on the growth of the *Sabellaria* veneers and/or the protection from dismantlement by environmental energy remains to be investigated, as well as the impact of these algae on the assemblage's biodiversity.

Although the occurrence of invasive *Caulerpa* species may be highly negative for particular species, such as the frame-builder coral *Cladocora caespitosa* (Linnaeus, 1767) (see Kružić *et al.*, 2008), other data point to the beneficial effect of populations of *C. cylindracea* intermixed with other algae on macrozoobenthos dispersion and in attracting some vagile taxa (Sinopoli *et al.*, 2020). The presence of *C. cilindracea* and *C. taxifolia* var. *distichophylla* growing on *Sabellaria* tubes does not seem to hinder the crust development. This is also obvious (even not described) for other *Sabellaria* bioconstructions in neighboring areas (Vega Fernández *et al.*, 2019: fig 1) and sabellarids have been considered tolerant worms persisting in *Posidonia* meadows impacted by *C. taxifolia* var. *distichophylla* (Musco *et al.*, 2014).

Finally, the rough Sabellaria surface seems to favour Ericaria settlement and the development of the new germlings. Pons-Fita et al. (2020) hypothesized a similar positive interaction between the habitat-forming species Cladocora caespitosa and Treptacantha ballesterosii (Orellana & Sansón) [corresponding to Gongolaria montagnei (J.Agardh) Kuntze] proposing a new habitat to be included in the list of the Spanish Mediterranean habitats. Pons-Fita et al. (2021) also assert that the close and long-lasting relationships they found between Cystoseira s.l. species and C. caespitosa are species- and site-specific, even though they are mainly neutral. The mechanisms that promote this coexistence are yet to be understood, but results by Pons-Fita et al. (2021) challenge the theory of competitive exclusion between C. caespitosa and fleshy macroalgae in the Mediterranean.

Likewise, *Cladocora-Gongolaria*, the *Sabellaria-Ericaria* co-occurrence and close intergrowth here first described represents an example of taxa that thrive together in a healty algal community falling in the "Biocoenosis of the Infralittoral Algae" (IA) (Habitat III.6.1 according to SPA/BIO Protocol), or the "Biocoenosis of the Photophilic Algae" (AP) *sensu* Pérès & Picard (1964).

Both the "*Cystoseira* complex" and the *Sabellaria* bioconstructions are protected owing to their vulnerability and the key functional role in coastal ecosystems. They are listed in Annex II ('endangered and threatened species') of the Barcelona Convention, Mediterranean Action Plan of the United Nations, and Annex I of the EC - Council Directive EEC/92/43 on the Conservation of Natural Habitats and of Wild Fauna and Flora, respectively. Consequently, the status and extension of the *Sabellaria-Ericaria* community deserve to be further investigated and monitored in the area, especially in relation to the anthropogenic impact and intense trampling in summer months, as suggested by Plicanti *et al.* (2016) for other bioconstructions.

The preliminary data here reported may constitute a baseline for future studies addressing the interactions between *Sabellaria* bioconstructions and *Ericaria* community through time, and for comparing data from an underway monitoring to investigate seasonal changes produced by human and/or natural disturbance.

#### Author contribution

RS, AR, DS designed the study. CD contributed to field and laboratory investigation; RS wrote the first draft of the manuscript. All authors interpreted data and wrote the agreed version of the manuscript.

#### Acknowledgements

Maria Flavia Gravina, University of Roma Tor Vergata is thanked for useful discussion on the taxonomy of *S. alveolata*. This study was financially supported by University of Catania (R. Sanfilippo, A. Rosso and D. Serio) through funds provided in the frame of the "Piano per la Ricerca 2016-18", grant n. 22722132118, and "Pia-CeRi-Piano Incentivi per la Ricerca di Ateneo 2020-22, linea di intervento 2". CD benefited from fellowship for doctorate studies from the University of Catania. This is the Catania Palaeoecological Group contribution n. 480.

#### References

- ARPA Sicilia U.O.C. Struttura territoriale di Ragusa, 2014. Progetto *Caulerpa*: Indagini ambientali e rilievi sulla diffusione della *Caulerpa* nel Canale di Sicilia, 80 pp.
- Bertocci, I., Badalamenti, F., Lo Brutto, S., Mikac, B., Pipitone, C. et al., 2017. Reducing the data-deficiency of threatened European habitats: Spatial variation of sabellariid worm reefs and associated fauna in the Sicily Channel, Mediterranean Sea. Marine Environmental Research, 130, 325-337.
- Bonifazi, A., Lezzi, M., Ventura, D., Lisco, S., Cardone, F. et al., 2019. Macrofaunal biodiversity associated with different developmental phases of a threatened Mediterranean

Sabellaria alveolata (Linnaeus, 1767) reef, Marine Environmental Research, 145, 97-111.

- Ciani, S., Rio, M-H., Menna, M., Santoleri R., 2019. A Synergetic Approach for the Space-Based Sea Surface Currents Retrieval in the Mediterranean Sea. *Remote Sensing*, 11, 1285.
- Curd, A., Pernet, F., Corporeau, C., Delisle, L., Firth, L.B. *et al.*, 2019. Connecting organic to mineral: How the physiological state of an ecosystem-engineer is linked to its habitat structure. *Ecological Indicators*, 98, 49-60.
- Delbono, I., Bianchi, C.N., Morri, C., 2003. Le biocostruzioni di Sabellaria alveolata come indicatori ambientali: area costiera fra Chiavari e Sestri Levante. In O. Ferretti (Ed.), Studi per la creazione di strumenti di gestione costiera. Golfo del Tigullio. (pp. 130-140) edizione ENEA S. Teresa Centro Ricerche Ambiente Marino, La Spezia.
- Dubois, S., Retière, C., Olivier, F., 2002. Biodiversity associated with Sabellaria alveolata (Polychaeta: Sabellaridae) reefs: effects of human disturbances. Journal of the Marine Biological Association of the U.K., 82, 817-826.
- Dubois, S., Commito, J.A., Olivier, F., Retière, C., 2006. Effects of epibionts on *Sabellaria alveolata* (L.) biogenic reefs and their associated fauna in the Bay of Mont Saint-Michel. *Estuarine Coastal and Shelf Science*, 68, 635-646.
- Fauvel, P., 1927. Polychètes sédentaires. Faune de France. 16. P. Lechevalier, Paris. 473 pp.
- Gruet, Y., 1972. Aspects morphologiques et dynamiques de constructions de l'Annélide Polychète Sabellaria alveolata (Linné). Travaux de l'Institut Scientifique et Technique des Pêches Maritimes, 36(2), 131-161.
- Gruet, Y., 1982. Recherches sur l'écologie des «récifs» d'hermelles édifiés par l'Annélide polychète Sabellaria alveolata (Linné). PhD Thesis, Universite de Nantes, 1-234.
- Gruet, Y., 1986. Spatio-temporal changes of sabellarian reefs built by the sedentary polychaete *Sabellaria alveolata* (Linné). *Marine Ecology*, 7, 303-319.
- Gruet, Y., Bodeur, Y., 1995. Ecological conditions of modern sabellarian reefs development: geological implications. *Publication du Service Géologique du Luxembourg*, 29, 73-80.
- Iaciofano, D., Musco, L., Vega Fernández, T., Lo Brutto, S., 2015. Crostacei anfipodi associati a reef di Sabellaria alveolata (L.) (Annelida) dello Stretto di Sicilia (Mar Mediterraneo). Biologia Marina Mediterranea, 22(1), 95-96.
- Jones, A.G., Dubois, S.F., Desroy, N., Fournier, J., 2018. Interplay between abiotic factors and species assemblages mediated by the ecosystem engineer *Sabellaria alveolata* (Annelida: Polychaeta). *Estuarine, Coastal and Shelf Science*, 200, 1-18.
- Kirtley, D.W., 1994. A review and taxonomic revision of the family Sabellariidae Johnston, 1865 (Annelida; Polychaeta). Sabecon Press, Florida. 223 pp.
- Kružić, P., Zŭljević, A., Nikolic, V., 2008. The highly invasive alga *Caulerpa racemosa* var. *cylindracea* poses a new threat to the banks of the coral *Cladocora caespitosa* in the Adriatic Sea. *Coral Reefs*, 27, 441.
- La Porta, B., Nicoletti, L., 2009. Sabellaria alveolata (Linnaeus) reefs in the central Tyrrhenian Sea (Italy) and associated polychaete fauna. Zoosymposia, 2, 527-536.
- Lezzi, M., Cardone, F., Mikac, B., Giangrande, A., 2015. Variation and ontogenetic changes of opercular paleae in a pop-

ulation of *Sabellaria spinulosa* (Polychaeta: Sabellaridae) from the South Adriatic Sea, with remarks on larval development. *Scientia Marina*, 79 (1), 137-150.

- Musco, L., Andaloro, F., Mikac, B., Mirto, S., Vega Fernandez, T. et al., 2014. Concern about the spread of the invader seaweed Caulerpa taxifolia var. distichophylla (Chlorophyta: Caulerpales) to the Western Mediterranean. Mediterranean Marine Science, 15 (3), 532-538.
- Noernberg, M.A., Fournier, J., Dubois, S., Populus, J., 2010. Using airborne laser altimetry to estimate *Sabellaria alveolata* (Polychaeta: Sabellariidae) reefs volume in tidal flat environments. *Estuarine, Coastal and Shelf Science*, 90 (2), 93-102.
- Pérès, J.M., Picard, J., 1964. Nouveau manuel de bionomie benthique de la mer Méditerranée. *Recueil des Travaux de la Station Marine d'Endoume*, 31 (47), 1-137.
- Plicanti, A., Domínguez, R., Dubois, S., Bertocci, I., 2016. Human impacts on biogenic habitats: Effects of experimental trampling on *Sabellaria alveolata* (Linnaeus, 1767) reefs. *Journal of Experimental Marine Biology and Ecology*, 478, 34-44.
- Pons-Fita, A., Verdura, J., Santamaría, J., Kersting, D.K., Ballesteros, E., 2020. Coexistence of the reef-building coral *Cladocora caespitosa* and the canopy-forming alga *Treptacantha ballesterosii*: Description of a new Mediterranean habitat. *Scientia Marina*, 84 (3), 263-271.
- Pons-Fita, A., Kersting, D.K., Ballesteros, E., 2021. Co-occurrence of a reef-building coral and canopy-forming macroalgae in the Mediterranean Sea. *Mediterranean Marine*

Science, 22 (3), 697-705.

- Sanfilippo, R., Rosso, A., Mastandrea, A., Viola, A., Deias, C., 2019. Sabellaria alveolata sandcastle worm from the Mediterranean Sea: new insights on tube architecture and biocement. Journal of Morphology, 80, 1839-1849.
- Sanfilippo, R., Guido, A., Insacco, G., Deias, C., Catania, G. et al., 2020. Distribution of Sabellaria alveolata (Polychaeta Sabellariidae) in the Mediterranean Sea: update and new findings. Zoosymposia, 19, 198-208. Proceedings of the 13° IPC Conference, Long Beach, 2-8 August 2019.
- Schimmenti, E., Musco, L., Lo Brutto, S., Mikac, B., Nygren, A., 2016. Mediterranean record of *Eulalia ornata* (Annelida: Phyllodocida) corroboratingits fidelity link with the *Sabellaria alveolata* reef habitat. *Mediterranean Marine Science*, 17 (2), 359-370.
- Serio, D., Furnari, G., 2021. Ericaria giacconei sp. nov. (Sargassaceae, Fucophyceae), the species to which the invalidly published Cystoseira hyblaea Giaccone should be referred. Cryptogamie, Algologie, 42 (10), 141-151.
- Sinopoli, M., Allegra, A., Andaloro, F., Consoli, P., Esposito, V. et al., 2020. Assessing the effect of the alien seaweed Caulerpa cylindracea on infralittoral rocky benthic invertebrate community: Evidence from a Mediterranean Marine Protected Area. Regional Studies in Marine Science, 38, 101372.
- Vega Fernández, T., Badalamenti, F., Bonaviri, C., Di Trapani, F., Gianguzza, P. *et al.*, 2019. Synergistic reduction of a native key herbivore performance by two non-indigenous invasive algae. *Marine Pollution Bulletin*, 141, 649-654.