

Mediterranean Marine Science

Vol 23, No 4 (2022)

VOL 23, No 4 (2022)



Species characterization of soft bottom habitats by depth and sediment particle size on the Catalan coast (NW Mediterranean): unexpected species composition of the assemblages

SUSANA PINEDO, ESTHER JORDANA, ENRIC BALLESTEROS

doi: [10.12681/mms.28127](https://doi.org/10.12681/mms.28127)

To cite this article:

PINEDO, S., JORDANA, E., & BALLESTEROS, E. (2022). Species characterization of soft bottom habitats by depth and sediment particle size on the Catalan coast (NW Mediterranean): unexpected species composition of the assemblages. *Mediterranean Marine Science*, 23(4), 789–804. <https://doi.org/10.12681/mms.28127>

Species characterization of soft bottom habitats by depth and sediment particle size on the Catalan coast (NW Mediterranean): unexpected species composition of the assemblages

Susana PINEDO, Esther JORDANA and Enric BALLESTEROS

Mediterranean Marine Science, 23 (3) 2022

Table S1. Results of dissimilarity analysis (SIMPER) between shallow and deep samples. The taxa contributing to 50 % of dissimilarity are included.

Taxa	Deep	Shallow	Contribution (%)	Cumulative contribution (%)
	Abundance (avg)	Abundance (avg)		
<i>Lucinella divaricata</i>	0.64	5.51	3.37	3.37
<i>Spisula subtruncata</i>	0.63	4.74	3.18	6.54
MACTRIDAE	0.10	3.24	2.10	8.64
<i>Paradoneis armata</i>	0.31	3.26	2.05	10.70
<i>Apseudopsis bacescui</i>	0.76	2.46	1.55	12.25
<i>Fabulina fabula</i>	0.06	2.23	1.46	13.71
<i>Spio decorata</i>	0.64	1.91	1.34	15.05
NEMATODA	0.55	1.86	1.33	16.38
<i>Centraloecetes dellavallei</i>	0.81	1.70	1.28	17.65
<i>Prionospio caspersi</i>	0.16	1.80	1.26	18.91
<i>Loripes orbiculatus</i>	0.36	1.82	1.23	20.14
<i>Thracia phaseolina</i>	0.45	1.92	1.22	21.36
<i>Mediomastus fragilis</i>	1.27	1.39	1.18	22.54
<i>Magelona johnstoni</i>	0.01	1.54	1.05	23.59
<i>Dosinia lupinus</i>	0.59	1.51	1.03	24.62
<i>Parvicardium cf pinnulatum</i>	0.05	1.52	1.02	25.64
<i>Ampelisca brevicornis</i>	0.05	1.33	0.95	26.59
<i>Peresiella clymenoides</i>	0.25	1.45	0.95	27.54
<i>Kirkegaardia heterochaeta</i>	1.32	0.08	0.93	28.47
<i>Urothoe intermedia</i>	0.18	1.30	0.91	29.39
<i>Lembos</i> sp1	0.00	1.28	0.90	30.28
<i>Bathyporeia guilliamsoniana</i>	0.07	1.28	0.88	31.16
<i>Chone durneri</i>	0.77	1.05	0.88	32.04
<i>Perioculodes longimanus</i>	0.21	1.23	0.85	32.89
NEMERTEA	0.60	1.28	0.84	33.73
<i>Magelona minuta</i>	1.20	0.14	0.84	34.58
<i>Levinsenia gracilis</i>	1.22	0.08	0.84	35.42
<i>Lumbrineris</i> sp1	1.09	0.12	0.83	36.24
VENERIDAE	0.11	1.05	0.81	37.05
<i>Scoletoma laurentiana</i>	0.53	0.98	0.79	37.85
<i>Donax trunculus</i>	0.00	1.08	0.77	38.62
<i>Pseudocuma longicorne</i>	0.13	1.04	0.74	39.36
<i>Ditrupa arietina</i>	0.90	0.29	0.68	40.05

Continued

Table S1 continued

Taxa	Deep	Shallow	Contribution (%)	Cumulative contribution (%)
	Abundance (avg)	Abundance (avg)		
<i>Nucula hanleyi</i>	0.95	0.13	0.68	40.72
<i>Megaluropus massiliensis</i>	0.01	0.94	0.67	41.39
<i>Moerella distorta</i>	0.64	0.69	0.66	42.05
<i>Micronephthys longicornis</i>	0.51	0.85	0.65	42.70
<i>Aponuphis bilineata</i>	0.94	0.32	0.65	43.35
<i>Leucothoe incisa</i>	0.32	0.92	0.63	43.98
<i>Goniadella galaica</i>	0.74	0.41	0.63	44.61
<i>Perioculodes aequimanus</i>	0.14	0.81	0.63	45.24
<i>Chamelea gallina</i>	0.00	0.89	0.61	45.86
<i>Nephtys kersivalensis</i>	0.68	0.47	0.61	46.47
TUBULANIDAE	0.27	0.87	0.60	47.06
<i>Owenia fusiformis</i>	0.40	0.68	0.58	47.64
<i>Scolaricia typica</i>	0.23	0.75	0.56	48.20
<i>Aspidosiphon muelleri</i>	0.77	0.04	0.55	48.75
<i>Sigambra parva</i>	0.59	0.54	0.54	49.30
<i>Aricidea claudiae</i>	0.72	0.05	0.54	49.83
<i>Medicorophium runcicorne</i>	0.21	0.70	0.52	50.35

Table S2. Contribution and average abundance of taxa responsible for most of similarity (SIMPER analysis) for shallow samples. The taxa contributing to 50 % of the similarity are included.

Taxa	Abundance (avg)	Similarity (avg)	Contribution (%)	Cumulative contribution (%)
<i>Spisula subtruncata</i>	4.74	2.35	8.75	8.75
<i>Lucinella divaricata</i>	5.51	2.27	8.45	17.20
<i>Paradoneis armata</i>	3.26	1.75	6.52	23.72
MACTRIDAE	3.24	1.24	4.60	28.32
<i>Apseudopsis bacescui</i>	2.46	0.92	3.42	31.74
<i>Fabulina fabula</i>	2.23	0.89	3.31	35.04
<i>Spio decorata</i>	1.91	0.85	3.15	38.19
<i>Thracia phaseolina</i>	1.92	0.68	2.52	40.71
<i>Prionospio caspersi</i>	1.80	0.64	2.38	43.09
NEMERTEA	1.28	0.56	2.07	45.17
NEMATODA	1.86	0.54	2.03	47.20
<i>Centraloecetes dellavallei</i>	1.70	0.54	2.02	49.21
<i>Magelona johnstoni</i>	1.54	0.54	1.99	51.21

Table S3. Contribution and average abundance of taxa responsible for most of similarity (SIMPER analysis) for deep samples. The taxa contributing to 50 % of the similarity are included.

Taxa	Abundance (avg)	Similarity (avg)	Contribution (%)	Cumulative contribution (%)
<i>Kirkegaardia heterochaeta</i>	1.32	1.02	6.35	6.35
<i>Levinsenia gracilis</i>	1.22	0.93	5.84	12.19
<i>Aponuphis bilineata</i>	0.94	0.77	4.79	16.98
<i>Lumbrineris</i> sp1	1.09	0.77	4.79	21.77
<i>Nucula hanleyi</i>	0.95	0.76	4.73	26.50
<i>Magelona minuta</i>	1.20	0.67	4.20	30.71
<i>Mediomastus fragilis</i>	1.27	0.59	3.66	34.37
<i>Goniadella galaica</i>	0.74	0.54	3.35	37.72
<i>Apseudopsis bacescui</i>	0.76	0.50	3.15	40.87
<i>Nephtys kersivalensis</i>	0.68	0.42	2.61	43.49
<i>Cossura soyeri</i>	0.59	0.36	2.23	45.72
<i>Aspidosiphon muelleri</i>	0.77	0.33	2.06	47.78
<i>Aricidea claudiae</i>	0.72	0.32	1.98	49.77
<i>Sigambra parva</i>	0.59	0.28	1.75	51.52

Table S4. Results of PERMANOVA main test for environmental variables differences among the six soft bottom habitats identified.

Environmental variable	Source	df	SS	MS	Pseudo-F	p
MPD	Habitat	5	66.17	13.83	86.12	0.001
	Residual	415	66.66	0.16		
	Total	420	135.82			
SC	Habitat	5	194.08	38.82	42.20	0.001
	Residual	415	381.70	0.92		
	Total	420	575.78			
OM	Habitat	5	309.12	61.82	113.93	0.001
	Residual	415	225.19	0.54		
	Total	420	534.31			