



Mediterranean Marine Science

Vol 23, No 4 (2022)

VOL 23, No 4 (2022)



Variation in the stable isotope trophic position of the bluefish Pomatomus saltatrix (Linnaeus, 1766) from two Mediterranean sites: insights from a global meta-analysis

FRANCESCO MANCINI, ANTONIO OSCAR LILLO, ROBERTA BARDELLI , SALVATRICE VIZZINI, GIORGIO MANCINELLI

doi: <u>10.12681/mms.29325</u>

To cite this article:

MANCINI, F., LILLO, A. O., BARDELLI, R., VIZZINI, S., & MANCINELLI, G. (2022). Variation in the stable isotope trophic position of the bluefish Pomatomus saltatrix (Linnaeus, 1766) from two Mediterranean sites: insights from a global meta-analysis. *Mediterranean Marine Science*, *23*(4), 850–863. https://doi.org/10.12681/mms.29325

Mediterranean Marine Science Indexed in WoS (Web of Science, ISI Thomson) and SCOPUS The journal is available on line at http://www.medit-mar-sc.net www.hcmr.gr DOI: http://doi.org/10.12681/mms.29325

Variation in the stable isotope trophic position of the bluefish *Pomatomus saltatrix* (Linnaeus, 1766) from two Mediterranean sites: insights from a global meta-analysis

Francesco MANCINI, Antonio Oscar LILLO, Roberta BARDELLI, Salvatrice VIZZINI and Giorgio MANCINELLI

Mediterranean Marine Science, 23 (4) 2022



Fig. S1: Global distribution of the bluefish. Records collated from the Global Information Biodiversity Facility – GBIF (https://www.gbif.org/). Data downloaded on September 12^{th,} 2021 - GBIF Occurrence Download https://doi.org/10.15468/dl.26nchn. For the Mediterranean Sea, occurrences recorded after 2001 are highlighted in red.



Fig. S2: Effect of the attack of a bluefish on a *Lithognathus mormyrus* specimen caught by a beach angler on the Ionian Sea at the mouth of Sinni River (40.152830°N, 16.691038°E), close to the town of Policoro in the Basilicata Region, Italy (Photo courtesy of Antonio Oscar Lillo).



Fig. S3: Location of the 30 investigations included in the database. Identification number (Ref#) of references reported in Table 2 are included.



Fig. S4: Allometric relationships of the type WW = a × TSL^b (Froese, 2006), linking the total standard length TSL (in cm) and total wet weight WW (in g) of *Pomatomus saltatrix* specimens from the Ionian and Adriatic Sea. The power curves are included, together with their Pearson *r* coefficient and statistical significance, and growth constant b. Noticeably, the relationship for Ionian bluefish was characterized by a growth constant significantly higher than that characterizing specimens from the Adriatic Sea (ANCOVA after log-transformation of data: $F_{2,}18 = 3.9$, P = 0.03).

References

Froese, R., 2006. Cube law, condition factor and weight-length relationships: history, meta-analysis and recommendations. *Journal of Applied Ichthyology 22*, 241-2.