

## Mediterranean Marine Science

Vol 23, No 3 (2022)

VOL 23, No 3 (2022)



### A molecular framework for the systematics of the Mediterranean spindle-shells (Gastropoda, Neogastropoda, Fascioliariidae, Fusiniinae)

GIULIA FASSIO, PAOLO RUSSO, GIUSEPPE BONOMOLO, ALEXANDER E. FEDOSOV, MARIA VITTORIA MODICA, ELISA NOCELLA, MARCO OLIVERIO

doi: [10.12681/mms.29935](https://doi.org/10.12681/mms.29935)


#### To cite this article:

Fassio, G., Russo, P., Bonomolo, G., Fedosov, A. E., Modica, M. V., Nocella, E., & OLIVERIO, M. (2022). A molecular framework for the systematics of the Mediterranean spindle-shells (Gastropoda, Neogastropoda, Fascioliariidae, Fusiniinae). *Mediterranean Marine Science*, 23(3), 623–636. <https://doi.org/10.12681/mms.29935>


## A molecular framework for the systematics of the Mediterranean spindle-shells (Gastropoda, Neogastropoda, Fasciolaridae, Fusininae)

Giulia FASSIO, Paolo RUSSO, Giuseppe BONOMOLO, Alexander E. FEDOSOV, Maria Vittoria MODICA, Elisa NOCELLA and Marco OLIVERIO

*Mediterranean Marine Science*, 23 (3) 2022




**Fig. S1:** Phylogenetic relationships among fasciolariaids as illustrated by the **Bayesian** majority consensus tree of the **16S** alignment. The tree is rooted on two buccinoideans (the Eosiphonidae *Manaria* sp., voucher MNHN-IM-013-19891, and an undetermined Buccinidae, voucher MNHN-IM-013-60365). Support values are given as posterior probabilities for the Bayesian analysis based on 107 generations, 25% burnin (only values  $\geq 0.95$  are shown).




**Fig. S2:** Phylogenetic relationships among fascioliids as illustrated by the **Bayesian** majority consensus tree of the **28S** alignment. The tree is rooted on two buccinoideans (the Eosiphonidae *Manaria* sp., voucher MNHN-IM-2013-19891, and an undetermined Buccinidae, voucher MNHN-IM-2013-60365). Support values are given as posterior probabilities for the Bayesian analysis based on 107 generations, 25% burnin (only values  $\geq 0.95$  are shown).




**Fig. S3:** Phylogenetic relationships among fascioliariids as illustrated by the **Bayesian** majority consensus tree of the **COI** alignment. The tree is rooted on two buccinoideans (the Eosiphonidae *Manaria* sp., voucher MNHN-IM-2013-19891, and an undetermined Buccinidae, voucher MNHN-IM-2013-60365). Support values are given as posterior probabilities for the Bayesian analysis based on 107 generations, 25% burnin (only values  $\geq 0.95$  are shown).




**Fig. S4:** Phylogenetic relationships among fascioliariids as illustrated by the **Bayesian** majority consensus tree of the **H3** alignment. The tree is rooted on two buccinoideans (the Eosiphonidae *Manaria* sp., voucher MNHN-IM-2013-19891, and an undetermined Buccinidae, voucher MNHN-IM-2013-60365). Support values are given as posterior probabilities for the Bayesian analysis based on 107 generations, 25% burnin (only values  $\geq 0.95$  are shown).



**Fig. S5:** Phylogenetic relationships among conoideans as illustrated by the **ML** majority consensus tree of the **16S** alignment. The tree is rooted on two buccinoideans (the Eosiphonidae *Manaria* sp., voucher MNHN-IM-2013-19891, and an undetermined Buccinidae, voucher MNHN-IM-2013-60365). Support values are given as ultrafast bootstrap support after ML analysis of 1000 pseudoreplicates (only values  $\geq 95\%$  are shown).




**Fig. S6:** Phylogenetic relationships among conoideans as illustrated by the **ML** majority consensus tree of the **28S** alignment. The tree is rooted on two buccinoideans (the Eosiphonidae *Manaria* sp., voucher MNHN-IM-2013-19891, and an undetermined Buccinidae, voucher MNHN-IM-2013-60365). Support values are given as ultrafast bootstrap support after ML analysis of 1000 pseudoreplicates (only values  $\geq 95\%$  are shown).



**Fig. S7:** Phylogenetic relationships among conoideans as illustrated by the ML majority consensus tree of the COI alignment. The tree is rooted on two buccinoideans (the Eosiphonidae *Manaria* sp., voucher MNHN-IM-2013-19891, and an undetermined Buccinidae, voucher MNHN-IM-2013-60365). Support values are given as ultrafast bootstrap support after ML analysis of 1000 pseudoreplicates (only values  $\geq 95\%$  are shown).





**Fig. S8:** Phylogenetic relationships among conoideans as illustrated by the ML majority consensus tree of the H3 alignment. The tree is rooted on two buccinoideans (the Eosiphonidae *Manaria* sp., voucher MNHN-IM-2013-19891, and an undetermined Buccinidae, voucher MNHN-IM-2013-60365). Support values are given as ultrafast bootstrap support after ML analysis of 1000 pseudoreplicates (only values  $\geq 95\%$  are shown).