

Mediterranean Marine Science

Vol 24, No 1 (2023)

VOL 24, No 1 (2023)



Spatio-temporal variation of the invasive copepod *Oithona davisae* in the zooplankton community of Kavala harbour

POLYXENI KOURKOUTMANI, KATERINA LOUFI,
GEORGIA KALANTARIDOU, AIKATERINI KARAGIANNI
, EVANGELIA MICHALOUDI

doi: [10.12681/mms.32127](https://doi.org/10.12681/mms.32127)

To cite this article:

KOURKOUTMANI, P., LOUFI, K., KALANTARIDOU, G., KARAGIANNI, A., & MICHALOUDI, E. (2023). Spatio-temporal variation of the invasive copepod *Oithona davisae* in the zooplankton community of Kavala harbour. *Mediterranean Marine Science*, 24(1), 174–181. <https://doi.org/10.12681/mms.32127>

Spatio-temporal variation of the invasive copepod *Oithona davisae* in the zooplankton community of Kavala harbour

Polyxeni KOURKOUTMANI, Katerina LOUFI, Georgia KALANTARIDOU, Aikaterini KARAGIANNI and Evangelia MICHALOUDI

School of Biology, Aristotle University of Thessaloniki, 54124 Thessaloniki, Greece

Corresponding author: Evangelia MICHALOUDI; tholi@bio.auth.gr

Contributing Editor: Stelios SOMARAKIS

Received: 29 November 2022; Accepted: 13 March 2023; Published online: 4 April 2023

Abstract

Non-indigenous species (NIS) are one of the anthropogenic effects greatly threatening biodiversity and ecosystem services. Ports and harbours are among the most susceptible environments for the arrival and establishment of these species since ballast waters represent a common pathway for NIS arrivals. However, relevant studies are limited. One of the few studies dealing with the zooplankton community of a harbour and reveals the presence of the widely distributed NIS *Oithona davisae* is presented in this paper, with special mention to the spatiotemporal variation of the species. The invader, *O. davisae*, a significant contributor to the zooplankton community, confirmed its thermophilic character and showed better development in the inner part of the harbour as well as patterns of coexistence with competing native species.

Keywords: Kavala Bay; port; zooplankton; *Oithona davisae*; non-indigenous species.

Introduction

Over the past few decades, as the rate of coastal urbanization is increasing rapidly and, combined with climate change, urban marine ecosystems are experiencing highly variable conditions influenced by several types of stressors, including land-based effluents, marine transport, fishing, biodiversity losses and extreme fluctuations in water properties (Todd *et al.*, 2019). These stressors include non-indigenous species (NIS) that are currently recognised as one of the important anthropogenic effects threatening ecosystem functioning (Molnar *et al.*, 2008). In several cases, the disruption of food web functioning has been shown to affect the maritime industry (Raaymakers, 2007). Planktonic copepods expand their distribution and are frequently recognised outside their indigenous environments. The copepod *Oithona davisae* is one of the most well-known NIS; it is a near-shore species, indigenous to the temperate coastal waters of the Indo West-Pacific, inhabiting eutrophic bays and transitional waters (Ferrari & Orsi, 1984; Uye & Sano, 1998; Uriarte *et al.*, 2016). Due to its high tolerance to temperature and salinity variations, *O. davisae* has successfully invaded a wide range of environments. These include the Black Sea, the Sea of Marmara and several sites across the Mediterranean Sea (e.g., Saiz *et al.*, 2003; Altukhov *et al.*, 2014; Razouls *et al.*, 2015-2022; Zagami

et al., 2018; Velasquez *et al.*, 2021;) while in the North Aegean Sea it has been recorded in the Thermaikos Gulf as well as the Kavala harbour (Dragicevic *et al.*, 2019; Kourkoutmani *et al.*, 2021). In some of these areas, it has established successful populations completely replacing *Oithona nana* (Gubanova *et al.*, 2014), while it has even been reported to successfully compete with other larger copepods (Gubanova *et al.*, 2019).

Ship ballast waters is the main vector recognised as the culprit for NIS introductions in the marine environment (Bailey, 2015). This makes ports and harbours among the most susceptible environments for new records of these species. However, the number of surveys relating to the structure and function of zooplankton communities in harbours is quite limited. This hampers the effective tracking of such bioinvasions, which can be further delayed when considering confusing taxonomy and small size. During the last three decades, several studies (e.g., Turner, 1994; Paffenhöfer, 1998; Zervoudaki *et al.*, 2006) have pointed to the under-sampling of the smaller-sized fractions when using a WP-2 net, which is traditionally used for mesozooplankton sampling. This results in an underestimation of the small-sized copepods, as well as their developmental stages, which need to be collected simultaneously, using a finer meshed net.

Here, we report on the presence of *Oithona davisae* in the small-sized fraction of the metazoan zooplankton

community of Kavala harbour. The investigation of its spatio-temporal distribution, along with differences in terms of biodiversity and abundance when using nets of different size selectivity, are reported for the first time in the study area and represents one of the few studies dealing with the zooplankton community in a harbour.

Materials and Methods

Study site

Kavala harbour is an important terminal for passenger ships in Greece, and since 2016 a number of floating platforms have been installed that provide numerous docking sites. It is located northwest of Kavala Bay (40.88056° N, 24.41667° E), a semi-enclosed area on the continental shelf of the northern Aegean Sea in the eastern Mediterranean Sea (Fig. 1). At the eastern part of the Bay, a commercial port, “Philip II” in New Karvali (40.95649° N, 24.50639° E), has been operating since 2002, serving mainly general cargo and bulk carrier vessels (www.marinetraffic.com). Kavala Bay is one of the most well-known fishing and nursery grounds in the northern Aegean Sea (Stergiou *et al.*, 1997), which is particularly important for small pelagic fish populations (Stergiou *et al.*, 2007; Sylaios *et al.*, 2010; Karagianni *et al.*, 2019). Furthermore, due to its location, Kavala Bay is influenced by the Black Sea Waters (BSW) since the surface current flowing through the Dardanelles along the Thracian Sea enters Kavala Bay (Sylaios *et al.*, 2005).

Sampling and microscopic analysis

Sampling was conducted monthly, from February 2017 to January 2018 (except October), from the floating platforms of the Kavala harbour basin (Fig. 1). Zooplankton samples were collected by vertical hauls throughout the water column, using both a $50\ \mu\text{m}$ and a $200\ \mu\text{m}$ mesh size plankton net for better representation of the different zooplankton size classes. All samples were preserved in 4% formalin-seawater solution for further microscopic analysis.

Laboratory analysis included species identification and abundance estimation under a microscope (Leica, Leitz Laborlux S). Taxonomic identification of individuals was based on appropriate taxonomic keys (Tregouboff & Rose, 1957; Conway, 2012a,b), as well as the classification system proposed by Razouls *et al.* (2005-2022). Adults, when possible, were identified all the way down to species level, whereas copepodites were grouped together and were not further identified. In the $50\ \mu\text{m}$ net samples, female adult individuals of the genus *Oithona* were identified based on morphological features (Fig. 2) of the rostrum, mandible, maxillule and urosome as well as the number of exopod spines of the swimming legs, as presented in Dragicevic *et al.* (2019). In the $200\ \mu\text{m}$ net samples, the presence of *O. davisae* was confirmed but *O. davisae* and *O. nana* adults were counted together; the number of males also includes all *Oithona* species. Quantitative samples were analysed according to Harris *et al.* (2000).




Fig. 1: A) Map of Greece, B) Map of the sampling stations in Kavala's harbour.




Fig. 2: *Oithona davisae* collected from Kavala's harbour A) dorsal view, B) side view, C) maxillule (scale 50 μm).

Data analysis

In order to identify the similarity of the zooplankton community during the sampling period, Hierarchical Cluster Analysis (CLUSTER) based on the Bray-Curtis similarity index was performed on the copepod's (nauplii, copepodites, adult copepods) abundance matrix, using group-average linking; abundance data were $\log(x+1)$ transformed in order to reduce bias due to highly abundant groups. Non-Metric Multidimensional Scaling (NMDS) was performed on the same Bray-Curtis similarity matrix in order to further visualize the clustering of the copepod's community. The similarity analysis routines, Analysis of Similarity (ANOSIM) and Similarity Percentage Analysis (SIMPER), were used to test the significance levels and sources of variance between the zooplankton assemblages of the different groups derived by hierarchical cluster analysis. The above analyses were performed using the Plymouth Routine in Multivariate Ecological Research (PRIMER) v.6 software package (Clarke & Gorley, 2006).

Results

In total, 60 zooplankton taxa were identified in samples obtained with both nets (50 and 200 μm) during the sampling period in Kavala harbour; 32 Copepoda, 6 Cladocera, 6 gelatinous taxa, and 16 meroplanktonic taxa (Table S1).

Total 50 μm net abundance ranged from 387.26 to 49037.09 ind m^{-3} while the 200 μm net abundance ranged from 1.27 to 10908.62 ind m^{-3} (Fig. 3). The mean abundance of the 200 μm net (1876.13 ind m^{-3}) during the sampling period was at least five times lower than the mean 50 μm net abundance (10763.52 ind m^{-3}) (t-test; $p = 0.01$). However, at Station 4 in March, the 200 μm net abundance was higher than that of the 50 μm net due to the higher numbers of Bryozoa and *Pleopis polyphemoides* in the 200 μm net (Fig. 3, Table S1). The maximum differences between the numbers of the 200 and 50 μm net were observed for copepodites of cyclopoida and nauplii, which showed 98 times higher values when collected with the 50 μm net, while almost identical abundances for adult copepods were recorded for both nets. Also, other smaller-sized fractions, e.g., rotifer taxa, were captured only when using the 50 μm plankton net (Table S1). During the sampling period, zooplankton communities collected with both nets presented similar distributions, exhibiting substantial temporal variability with abundance values being higher in late summer-early autumn (Fig. 3).

The developmental stages of copepods, nauplii and copepodites, dominated the 50 μm net samples throughout the sampling period (range 53-98%) contrary to the 200 μm net samples where their contribution was considerably lower, averaging 18% of the total community (Fig. S1). Within the copepod's community of the 50 μm net, *Oithona nana* and *Oithona davisae* dominated, contribut-




Fig. 3: Abundance (individuals m^{-3}) of the zooplankton community, grouped by sampling stations, during the sampling period 2017-2018 in Kavala harbour.

ing cumulatively up to 90% to adult copepod abundance, in contrast to the community collected with the 200 μm net where their contribution among adult copepods was rather low (<10%) (Fig. 4). Regarding their temporal distribution, *Oithona nana* showed a persistent occurrence throughout the sampling period while *O. davisae* was more abundant from late spring to early winter, although they both exhibited their peak abundance during the summer-autumn period. The native *O. nana* exhibited extended spatial presence in the sampling area, contrary to *O. davisae*, which reached maximum density at the stations situated further away from the open sea and closer to the area where ships dock (Stations 4 and 5) (Fig. 5).

Cluster analysis based on copepod (nauplii, copepodites, adult copepods) abundance data revealed three major clusters. Group 'a' consisted of only one sampling (September 2017; St.4). The samplings of September, November and August formed the cluster of Group 'c', whereas the rest of the samplings made up the cluster of Group 'b'. The dissimilarity between Groups 'b' and 'c'

(ANOSIM, $R = 0.819$, $p = 0.001$; Table S2) was mainly due to the contribution of the invasive copepod *Oithona davisae* (SIMPER average dissimilarity 9.78 %) (Fig. 6, Table S2).

Discussion

In our study we recorded the well-established fact that the small-sized fraction of crustacean zooplankton, is vastly underestimated due to the traditional sampling methods (Calbet, 2001; Gallienne & Robins, 2001; Turner, 2004; Zervoudaki *et al.*, 2006), with copepod developmental stages (nauplii and copepodites) (cumulative contribution reached up to 98%) mainly contributing to this difference.

In total, copepods dominated the zooplankton community of Kavala's harbour collected using a 50 μm net and contributed up to 98%, with the small-sized oithonoids (*O. nana* and *O. davisae*) dominating in terms of




Fig. 4: Contribution (%) of *Oithona davisae* and *Oithona nana* to the rest of the adult copepods of the zooplankton community collected with the 50 and 200 μm net.




Fig. 5: Abundances (ind m^{-3}) (mean \pm SD) of *Oithona davisae* and *Oithona nana* during the sampling period 2017-2018 in Kavala harbour.




Fig. 6: (A) Cluster Analysis and (B) Non-metric Multi-dimensional Scaling configuration (NMDS) on copepod's abundance data (50 μ m net). The groups were delineated at a 60% similarity level (dashed line).

adult copepod abundance. This is of no surprise considering that oithonoids have been described as the most widespread and abundant copepods worldwide (Gallienne & Robins, 2001). Among them, *Oithona davisae* and *Oithona nana* have been overlooked in zooplankton studies because, due to their small size, they escape from the classical sampling equipment used for zooplankton collection (Gallienne & Robins, 2001; Tseng *et al.*, 2007). Therefore, in many cases there is no clear picture of their distribution patterns or the time of *O. davisae* invasion in the newly established areas. In the Mediterranean Sea, it has so far been reported in the coastal lakes Faro & Ganzirri in Italy (Zagami *et al.*, 2018), Venice Lagoon (Vidjak *et al.*, 2019; Pansera *et al.*, 2021), Barcelona port in Spain (Saiz *et al.*, 2003), Thermaikos Bay in Greece (Dragicevic *et al.*, 2019), the Turkish coasts of the Aegean Sea (Terbiyik Kurt & Beşiktepe, 2019) and the Levant

Sea (Terbiyik Kurt *et al.*, 2022). Besides recording its presence, few of these studies investigated its seasonal and spatial variations. In every case, including the current study, *O. davisae* has confirmed its thermophilic character by exhibiting higher abundance during the period from summer to autumn (Ueda, 1991), which is the warm period in the Mediterranean. Its dominance in autumn could probably provide a niche partitioning in time, allowing for coexistence with the native competing *Oithona nana*. This could be a successful strategy at the beginning of its invasion phase that could later lead to a full overtake of *O. davisae* against *O. nana* as is the case recorded in the Black Sea where it occupied the niche of *O. nana* and resulted in its complete disappearance (Gubanov *et al.*, 2014). This cannot be confirmed herein since there is no clear time frame as regards the beginning of its invasion due to lack of proper monitoring. Of course, differences

in spatial distribution can also provide niche partitioning in space. In the studies mentioned above, including the current study, reporting spatial data, there was a clear spatial differentiation between *O. davisae* and *O. nana*. *Oithona davisae* was clearly in favour of the inner part of the harbour while *O. nana* was more abundant at the stations of the outer part (Fig. 5). *Oithona davisae* seems to find local conditions that promote its settlement and distribution, especially in the inner and more trophic sites (Pansera *et al.*, 2021). Future monitoring of the evolution of the non-indigenous *O. davisae* population will allow investigation of potential consequences of its introduction, such as exclusion of congeneric species and competition with other copepods for limited resources and microhabitats.

It is evident that underestimation of the small-sized fraction of zooplankton can result in erroneous estimates of biomass and secondary production since taxa of the metazoan microzooplankton community have a higher impact on the trophic coupling's effectiveness between primary producers, protozooplankton and higher trophic levels than larger taxa (Turner, 2004; Calbet & Saiz, 2005). This is of great importance for ecosystem functioning of ports and harbours that, apart from being susceptible to the arrival and establishment of NIS, are highly affected by nutrient effluents and subject to eutrophication. In eutrophic coastal areas, the link between the traditional food web and the microbial loop in eutrophic coastal areas can determine the energy flow and final trophic level reached (Sommer *et al.*, 2018). The developmental stages of copepods and small-sized oithonoids can contribute to this; and the same applies to rotifers, a group of taxa erroneously considered absent from nearshore/coastal environments (Fontaneto *et al.*, 2006) mainly because the majority of studies focus on the mesozooplankton community. Therefore, within the framework of the current study, we recorded the presence of *Synchaeta* spp., which had a notable contribution in the summer-autumn period community collected with a 50 µm net.

The current study is the first study on the zooplankton community in the Kavala's harbour region and one of the few on harbour areas in Greece and worldwide. We followed the approach of combined mesh size nets in order to investigate the under-represented size fraction of the smaller metazoan zooplankton community, reporting on the presence and distribution patterns of the invasive copepod *Oithona davisae* for the first time in the study area. The results of our study confirm the thermophilic character of *O. davisae* as well as the fact that it finds local conditions in harbours that promote its settlement and distribution, especially in the inner part of harbours. At the current point *O. davisae* exhibited coexistence patterns with the competing native species. However, since the time of its arrival is unknown, future monitoring of the harbour area will clearly frame the effects of its arrival on the structure and functioning of the zooplankton community in Kavala harbour.

References

- Altukhov, D.A., Gubanova, A.D., Mukhanov, V.S., 2014. New invasive copepod *Oithona davisae* Ferrari and Orsi, 1984: seasonal dynamics in Sevastopol Bay and expansion along the Black Sea coasts. *Marine Ecology*, 35, 28-34.
- Bailey, S.A., 2015. An overview of thirty years of research on ballast water as a vector for aquatic invasive species to freshwater and marine environments. *Aquatic Ecosystem Health & Management*, 18 (3), 261-268.
- Calbet, A., 2001. Annual Zooplankton Succession in Coastal NW Mediterranean Waters: The Importance of the Smaller Size Fractions. *Journal of Plankton Research*, 23 (3), 319-331.
- Calbet, A., Saiz, E., 2005. The ciliate-copepod link in marine ecosystems. *Aquatic Microbial Ecology*, 38 (2), 157-167.
- Clarke, K.R., Gorley, R.N., 2006. PRIMER v6: user manual/tutorial. PRIMER-e, Plymouth.
- Conway, D.V., 2012a. *Marine Zooplankton of Southern Britain-Part 1: Radiolaria, Heliozoa, Foraminifera, Ciliophora, Cnidaria, Ctenophora, Platyhelminthes, Nemertea, Rotifera and Mollusca*. Occasional Publication of the Marine Biological Association of the United Kingdom, United Kingdom, 138 pp.
- Conway, D.V., 2012b. *Marine Zooplankton of Southern Britain-Part 2: Arachnida, Pycnogonida, Cladocera, Facetotecta, Cirripedia and Copepoda*. Occasional Publication of the Marine Biological Association of the United Kingdom, United Kingdom, 163 pp.
- Dragicevic, B., Anadolli, O., Angel, D., Benabdi, M., Bitar, G. *et al.*, 2019. New Mediterranean Biodiversity Records (December 2019). *Mediterranean Marine Science*, 20 (3), 645-656.
- Ferrari, F.D., Orsi, J., 1984. *Oithona davisae*, new species, and *Limnoithona sinensis* (Burckhardt, 1912) (Copepoda: Oithonidae) from the Sacramento-San Joaquin Estuary, California. *Journal of Crustacean Biology*, 4 (1), 106-126.
- Fontaneto, D., De Smet, W.H., Ricci, C., 2006. Rotifers in saltwater environments, re-evaluation of an inconspicuous taxon. *Journal of the Marine Biological Association of the United Kingdom*, 86 (4), 623-656.
- Gallienne, C.P., Robins, D.B., 2001. Is *Oithona* the most important copepod in the world's oceans? *Journal of Plankton Research*, 23 (12), 1421-1432.
- Gubanova, A., Altukhov, D., Stefanova, K., Arashkevich, E., Kamburska, L. *et al.*, 2014. Species composition of Black Sea marine planktonic copepods. *Journal of Marine Systems*, 135, 44-52.
- Gubanova, A.D., Garbazey, O.A., Popova, E.V., Altukhov, D.A., Mukhanov, V.S., 2019. *Oithona davisae*: Naturalization in the Black Sea, interannual and seasonal dynamics, and effect on the structure of the planktonic copepod community. *Oceanology*, 59 (6), 912-919.
- Harris, R., Wiebe, P., Lenz, J., Skjoldal, H.R., Huntley, M., 2000. *ICES zooplankton methodology manual*. Academic Press, San Diego, 684 pp.
- Karagianni, A., Artemiou, A., Tsikliras, A.C., Michaloudi, E., 2019. Summer mesozooplankton assemblages in relation to environmental parameters in Kavala Gulf, northern Aegean Sea. *Scientia Marina*, 83 (1), 41.
- Kourkoutmani P., Kalantaridou G., Loufi A., Karagianni A.,

- Michaloudi E., 2021. Presence of the invasive cyclopoid *Oithona davisae* Ferrari & Orsi, 1984 in Kavala harbor, North Aegean Sea, In: *Joint ESENIAS and DIAS Scientific Conference*, 7-9 December 2021, Virtual Conference.
- Molnar, J.L., Gamboa, R.L., Revenga, C., Spalding, M.D., 2008. Assessing the global threat of invasive species to marine biodiversity. *Frontiers in Ecology and the Environment*, 6 (9), 485-492.
- Paffenhöfer, G.A., 1998. On the relation of structure, perception and activity in marine planktonic copepods. *Journal of Marine Systems*, 15 (1-4), 457-473.
- Pansera, M., Camatti, E., Schroeder, A., Zagami, G., Bergamasco, A., 2021. The non-indigenous *Oithona davisae* in a Mediterranean transitional environment: Coexistence patterns with competing species. *Scientific Reports*, 11 (1), 1-14.
- Raaymakers, S., 2007. Possible effects of climate change on the spread of invasive marine species and implications for maritime industries. *WMU Journal of Maritime Affairs*, 6 (2), 235-240.
- Razouls, C., Desreumaux, N., Kouwenberg, J., De Bovée, F. 2005-2022. *Biodiversity of Marine Planktonic Copepods* (morphology, geographical distribution and biological data) <http://copepodes.obs-banyuls.fr/en> (Accessed 28 January 2022)
- Saiz, E., Calbet, A., Broglio, E., 2003. Effects of small-scale turbulence on copepods: The case of *Oithona davisae*. *Limnology and Oceanography*, 48 (3), 1304-1311.
- Sommer, U., Charalampous, E., Scotti, M., Moustaka-Gouni, M., 2018. Big fish eat small fish: implications for food chain length? *Community Ecology*, 19 (2), 107-115.
- Stergiou, K.I., Christou, E.D., Georgopoulos, D., Zenetos, A., Souvermezoglou, C., 1997. The Hellenic Seas: Physics, chemistry, biology and fisheries. *Oceanography and Marine Biology*, 35, 415-538.
- Stergiou, K.I., Moutopoulos, D.K., Tsikliras, A.C., 2007. Spatial and temporal variability in Hellenic marine fisheries landings. p. 141-150. In: *State of Hellenic Fisheries*. Papaconstantinou, C., Zenetos, A., Vassilopoulou, V., Tserpes, G. (Eds). Hellenic Centre for Marine Research, Athens, Greece.
- Sylaios, G.K., Koutroumanidis, T., Tsikliras, A.C., 2010. Ranking and classification of fishing areas using fuzzy models and techniques. *Fisheries Management and Ecology*, 17 (3), 240-253.
- Sylaios, G., Stamatis, N., Kallianiotis, A., Vidoris, P., 2005. Monitoring water quality and assessment of land-based nutrient loadings and cycling in Kavala Gulf. *Water Resources Management*, 19 (6), 713-735.
- Terbiyik Kurt, T., Beşiktepe, Ş., 2019. First distribution record of the invasive copepod *Oithona davisae* Ferrari and Orsi, 1984, in the coastal waters of the Aegean Sea. *Marine Ecology*, 40 (3), e12548.
- Terbiyik-Kurt, T., Polat, S., Guy-Haim, T., 2022. New invasive copepod species from the Levantine Sea, north-eastern Mediterranean: *Oithona davisae* (Ferrari and Orsi, 1984) (Cyclopoida: Oithonidae). *BioInvasions Record*, 11 (1), 215-226.
- Todd, P.A., Heery, E.C., Loke, L.H.L., Thurstan, R.H., Kotze, D.J. *et al.*, 2019. Towards an urban marine ecology: Characterizing the drivers, patterns and processes of marine ecosystems in coastal cities. *Oikos*, 128 (9), 1215-1242.
- Tregouboff, G., Rose, M., 1957. *Manuel de Planctologie Méditerranéenne*. Centre National de la Recherche Scientifique, Paris, 587 pp.
- Tseng, L.C., Chen, Q.C., Hwang, J.S., Kumar, R., Dahms, H.U., 2007. Mesh size affects abundance estimates of *Oithona* spp. (Copepoda, Cyclopoida). *Crustaceana*, 80 (7), 827-837.
- Turner, J.T., 1994. Planktonic copepods of Boston Harbor, Massachusetts Bay and Cape Cod Bay, 1992. p. 405-413. In: *Ecology and Morphology of Copepods*. Ferrari, F.D., Bradley, B.P. (Eds). Springer, Dordrecht.
- Turner, J.T., 2004. The importance of small planktonic copepods and their roles in pelagic marine food webs. *Zoological Studies*, 43 (2), 255-266.
- Ueda, H., 1991. Horizontal distribution of planktonic copepods in inlet waters. *Bulletin of Plankton Society of Japan*, p. 143-160. In: *Proceedings of the Fourth International Conference on Copepoda, Karuizawa, Japan, 16-20 September 1990*. Plankton Society of Japan, Hiroshima.
- Uriarte, I., Villate, F., Iriarte, A., 2016. Zooplankton recolonization of the inner estuary of Bilbao: Influence of pollution abatement, climate and non-indigenous species. *Journal of Plankton Research*, 38 (3), 718-731.
- Uye, S., Sano, K., 1998. Seasonal variations in biomass, growth rate and production rate of the small cyclopoid copepod *Oithona davisae* in a temperate eutrophic inlet. *Marine Ecology Progress Series*, 163, 37-44.
- Velasquez, X., Morov, A.R., Kurt, T.T., Meron, D., Guy-Haim, T., 2021. Two-way bioinvasion: tracking the neritic non-native cyclopoid copepods *Dioithona oculata* and *Oithona davisae* (Oithonidae) in the Eastern Mediterranean Sea. *Mediterranean Marine Science*, 22 (3), 586-602.
- Vidjak, O., Bojanić, N., de Olazabal, A., Benzi, M., Brautović, I. *et al.*, 2019. Zooplankton in Adriatic port environments: Indigenous communities and non-indigenous species. *Marine pollution bulletin*, 147, 133-149.
- Zagami, G., Brugnano, C., Granata, A., Guglielmo, L., Minuto-li, R. *et al.*, 2018. Biogeographical distribution and ecology of the planktonic copepod *Oithona davisae*: Rapid invasion in Lakes Faro and Ganzirri (Central Mediterranean Sea). p. 59-82. In: *Trends in Copepod Studies - Distribution, Biology and Ecology*. Uttieri, M. (Ed). Nova Science Publishers, New York.
- Zervoudaki, S., Nielsen, T.G., Christou, E.D., Siokou-Frangou, I., 2006. Zooplankton distribution and diversity in a frontal area of the Aegean Sea. *Marine Biology Research*, 2 (3), 149-168.

Supplementary Data

The following supplementary information is available online for the article:

Table S1. List of recorded taxa in Kavala harbour. Mean abundance values (individuals m^{-3}) of all samplings per station. Adults include males and females.

Table S2. Results of ANOSIM and SIMPER analysis applied on Bray-Curtis similarity matrix of $\log(x+1)$ transformed copepods abundance data, collected with 50 μm plankton net.

Fig. S1: Contribution (%) of the main groups to the 200 μm and 50 μm net abundance in the Kavala harbour during 2017-2018. a) Contribution (%) of nauplii and copepodites on the rest zooplankton community. b) Contribution (%) of dominant mesozooplankton groups to total abundance.